This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

LANE MEDICAL LIBRARY STANFORD
2 45 0179 8392

LANE

MEDICAL LIBRARY

LEVI COOPER LANE FUND

LIBRARY
OF
COOPER MEDICAL COLLEGE
DATE Sept 1904
NO. 6199 CLASS 6 a
~~GIFT OF~~
Purchase




LIBRARY OF
Dr. C. S. Sargent.
Ho Jemite Building.
And please return it. You may think this a strange request, but I find that although many of my friends are poor arithmeticians, they are nearly all good bookkeepers.—SCOTT.

THE DISSECTION

OF THE

HUMAN BODY.

HOLDEN.

YEO'S

Manual of Physiology.

A Manual of Physiology for Students and Physicians.

By GERALD F. YEO, M.D., F.R.C.S.,
Professor of Physiology in King's College, London.

WITH OVER 300 CAREFULLY PRINTED ILLUSTRATIONS AND A GLOSSARY.

SMALL OCTAVO. 750 PAGES. CLOTH, $4.00; LEATHER, $5.00.

"The writer has endeavored to 'avoid theories which have not borne the test of time, and such details of methods as are unnecessary for junior students.' His experience as a teacher and as an examiner has led him to lay stress on the points which are hard to grasp and are commonly misunderstood, and he has directed attention more to those subjects which have a practical medical or surgical application, than to those relating to abstract physiological subjects. * * * We have pleasure in recommending this book, as a most excellent manual, being what it pretends to be—elementary, and yet containing all that is really of importance to the student of medicine. It is difficult to devise an original method of treating such a well worn subject as physiology, but the present volume undoubtedly has originality in its method. We think the author treats his subject in the best manner possible when he combines microscopical anatomy intimately with the discussion of strict physiology." *The Med. Times and Gazette, London*

P. BLAKISTON, SON & CO., 1012 Walnut Street, Philadelphia,
Medical Publishers, Booksellers and Importers.

HOLDEN'S ANATOMY.

A MANUAL OF DISSECTION OF THE HUMAN BODY.

BY

LUTHER HOLDEN,

LATE PRESIDENT OF THE ROYAL COLLEGE OF SURGEONS OF ENGLAND; CONSULTING SURGEON TO ST. BARTHOLOMEW'S AND THE FOUNDLING HOSPITALS.

FIFTH EDITION.

EDITED BY

JOHN LANGTON,

SURGEON TO, AND LECTURER ON ANATOMY AT, ST. BARTHOLOMEW'S HOSPITAL; MEMBER OF THE BOARD OF EXAMINERS, ROYAL COLLEGE OF SURGEONS OF ENGLAND.

WITH OVER TWO HUNDRED ILLUSTRATIONS.

PHILADELPHIA:
P. BLAKISTON, SON & CO.,
No. 1012 WALNUT STREET.
1885.

LANE LIBRARY

PRESS OF WM. F. FELL & CO., 1220-1224 SANSOM ST.

TO

THE STUDENTS

OF

ST. BARTHOLOMEW'S HOSPITAL,

IN THE HOPE THAT IT MAY ASSIST THEM IN THEIR

ANATOMICAL STUDIES,

This Manual is Dedicated,

BY THEIR FAITHFUL FRIEND AND WELL-WISHER,

THE AUTHOR.

PREFACE

TO

THE FIFTH EDITION.

In this edition the Editor has most carefully revised the entire work. The order of dissection has been here and there altered, and further illustrations and additional matter introduced, especially concerning the Anatomy of the Nervous System and the Organs of Special Sense.

The object throughout has been to be as concise as possible, and to put the subject in as clear and practical a light as is compatible with the faithful handling of its natural difficulties.

It is hoped that the work, in its present form, is adapted, not only for students, but for members of the profession who wish to refresh their anatomical knowledge.

The best thanks of the Editor are due to the Demonstrators and Assistant Demonstrators of Anatomy at St. Bartholomew's Hospital, for valuable suggestions.

2 HARLEY STREET, CAVENDISH SQUARE,
LONDON.

PREFACE

TO

THE FIRST EDITION.

If any apology be needed for the appearance of the present Manual, it may be stated, without any wish to disparage the labors of others, that the works of this kind hitherto published seem to the Author open to one or the other of two objections: either as being too systematic, and therefore not adapted for the dissecting room, or as obscuring the more important features of Anatomy by a multiplicity of minute and variable details.

In endeavoring to supply a presumed deficiency, the Author has made it his special aim to direct the attention of the student to the prominent facts of Anatomy, and to teach him the groundwork of the science; to trace the connection, and to point out the relative situation of parts, without perplexing him with minute descriptions.

A concise and accurate account is given of all the parts of the human body—the bones excepted, of which a competent knowledge is presupposed—and directions are laid down for the best method of dissecting it.

The several regions of the body are treated of in the order considered most suitable for their examination; and the muscles, vessels, nerves, etc., are described, as they are successively exposed to view in the process of dissection.

The Author has written the work entirely from actual observation: at the same time no available sources of information have been neglected, the highest authorities both English and Foreign having been carefully consulted. His acknowledgments are especially due to F. C. SKEY, Esq., F. R. S., Lecturer on Anatomy at St. Bartholomew's Hospital, for many valuable suggestions. He is also much indebted to his young friend, Mr. W. CLUBBE, for able assistance in dissections.

CONTENTS.

PAGE

DISSECTION OF THE SCALP, ... 17

DISSECTION OF THE FACE, ... 40

DISSECTION OF THE ORBIT, ... 64

DISSECTION OF THE NECK, ... 80

COURSE AND RELATIONS OF THE SUBCLAVIAN ARTERIES, . . 131

THE MUSCLES OF MASTICATION. TEMPORAL AND PTERYGO-MAXILLARY REGIONS, ... 145

BRANCHES OF THE INTERNAL MAXILLARY ARTERY IN THE THREE STAGES OF ITS COURSE, ... 153

BRANCHES OF THE INFERIOR MAXILLARY NERVE 157

DISSECTION OF THE THORAX, ... 174

DISSECTION OF THE HEART, ... 217

FŒTAL CIRCULATION, ... 232

STRUCTURE OF THE LUNGS, ... 235

DISSECTION OF THE PHARYNX, ... 244

DISSECTION OF THE LARYNX, ... 257

DISSECTION OF THE TONGUE, ... 272

DISSECTION OF THE SUPERIOR MAXILLARY NERVE, ... 277

DISSECTION OF THE NINTH, TENTH AND ELEVENTH CRANIAL NERVES AT THE BASE OF THE SKULL, ... 284

DISSECTION OF THE NOSE, ... 290

DISSECTION OF THE MUSCLES OF THE BACK, ... 297

LIGAMENTS OF THE SPINE, ... 314

DISSECTION OF THE UPPER EXTREMITY, ... 324

DISSECTION OF THE AXILLA, ... 331

PAGE

DISSECTION OF THE UPPER ARM, ... 343
DISSECTION OF THE FRONT OF THE FOREARM, ... 359
DISSECTION OF THE PALM OF THE HAND, ... 373
MUSCLES OF THE BACK CONNECTED WITH THE ARM, . . . 387
DISSECTION OF THE MUSCLES OF THE SHOULDER, . . . 398
DISSECTION OF THE BACK OF THE FOREARM, ... 408
DISSECTION OF THE LIGAMENTS, ... 421
DISSECTION OF THE ABDOMEN, ... 439
DISSECTION OF THE PARTS CONCERNED IN INGUINAL HERNIA, . 460
DISSECTION OF THE PELVIC VISCERA, ... 519
DISSECTION OF THE MALE PERINEUM, ... 525
DISSECTION OF THE FEMALE PERINEUM, ... 540
ANATOMY OF THE SIDE VIEW OF THE PELVIC VISCERA, . . 545
STRUCTURE OF THE BLADDER, PROSTATE, URETHRA AND PENIS, . 566
DISSECTION OF THE FEMALE PELVIC VISCERA, ... 581
DISSECTION OF THE ABDOMINAL VISCERA, ... 593
DISSECTION OF THE LOWER EXTREMITY, ... 626
ANATOMY OF THE PARTS CONCERNED IN FEMORAL HERNIA, . . 635
DISSECTION OF THE FRONT OF THE LEG, ... 659
DISSECTION OF THE GLUTEAL REGION, ... 669
DISSECTION OF THE BACK OF THE THIGH, ... 684
DISSECTION OF THE BACK OF THE LEG. 688
DISSECTION OF THE SOLE OF THE FOOT, ... 698
DISSECTION OF THE LIGAMENTS, ... 708
DISSECTION OF THE BRAIN, ... 729
DISSECTION OF THE SPINAL CORD, ... 792
DISSECTION OF THE EYE, ... 804
DISSECTION OF THE ORGAN OF HEARING, ... 826
DISSECTION OF THE MAMMARY GLAND, ... 845
DISSECTION OF THE SCROTUM AND TESTIS, ... 846

INDEX, ... 857

LIST OF ILLUSTRATIONS.

FIG.		PAGE
1.	Muscular and Aponeurotic Stratum of the Scalp,	18
2.	Sensory Nerves of the Scalp and Face,	21
3.	Branches of the Facial Nerve,	23
4.	Diagram to show the Formation of a Sinus,	27
5.	The Cranial Sinuses,	28
6.	The Venous Sinuses at the Base of the Skull,	31
7.	The Exit of the Cranial Nerves,	35
8.	The Nerves in the Foramen Jugulare,	36
9.	Relation of Structures in the Cavernous Sinus,	37
10.	Relations of the Nerves in the Sphenoidal Fissure,	38
11.	Relations of the Nerves and Muscles in the Orbit,	39
12.	The Geniculate Ganglion of the Facial Nerve,	40
13.	The Muscles of the Face,	42
14.	The Lachrymal Apparatus,	50
15.	The Muscles of the Pharynx,	54
16.	The Branches of the External Carotid Artery,	56
17.	The Branches of the Facial Nerve,	63
18.	The Sensory Nerves of the Scalp and Face,	64
19.	The Nerves of the Orbit,	67
20.	View of the Orbit from above,	69
21.	Lachrymal Duct,	71
22.	View of the Optic and Lower Nerves of the Orbit,	74
23.	Insertion of the Recti Muscles,	79
24.	The Superficial Nerves and Veins of the Neck,	83
25.	The Triangles of the Neck,	90
26.	Central Line of the Neck,	96
27.	Digastric Triangle and Contents,	109
28.	The Branches of the External Carotid Artery and their Branches.	114
29.	Muscles, Vessels, and Nerves of the Tongue,	120

FIG. PAGE
30. The Heart and Large Vessels, ... 133
31. The Inosculations of the Subclavian Artery, ... 141
32. The Formation of the Brachial Plexus and its Branches, . . 144
33. Pterygoid Muscles and the Internal Maxillary Artery, . . 151
34. Plan of the Internal Maxillary Artery, ... 154
35. Plan of the Branches of the Inferior Maxillary Nerve, . . 158
36. The Communications of the Facial, Glosso-pharyngeal, Pneumogastric, Spinal Accessory, Hypoglossal, Sympathetic, and the two Upper Cervical Nerves, ... 171
37. Form and Position of the Lungs, ... 175
38. The Reflections of the Pleural Sacs, ... 180
39. The Heart, showing Interpleural Space, ... 182
40. Form and Position of the Lungs, ... 183
41. Relative Position of the Heart and its Valves with regard to the Walls of the Chest, ... 187
42. Superior Vena Cava and its Tributaries, ... 192
43. Course and Relations of the Arch of the Aorta, ... 194
44. The Course of the Vena Azygos and the Thoracic Duct, . . 202
45. The Thoracic Portion of the Sympathetic Nerve, ... 208
46. Diagram of a Spinal Nerve, ... 211
47. The Constituents of the Root of each Lung and their relative position. 216
48. The Interior of the Right Auricle, ... 220
49. The relative Position of the Valves of the Heart, seen from above. 228
50. Scheme of the Fœtal Circulation, ... 233
51. Ultimate Air-cells of the Lung, ... 241
52. Side view of the Muscles of the Pharynx, ... 246
53. View of the Constrictor Muscles from behind, ... 248
54. View of the Pharynx laid open from behind, ... 250
55. Shape of the Glottis when at rest, ... 265
56. Diagram showing the Action of the Crico-thyroid Muscle, . . 266
57. Glottis dilated: Muscles dilating it, ... 267
58. Side view of the Muscles of the Larynx, ... 268
59. Glottis closed: Muscles closing it, ... 269
60. Upper Surface of the Tongue, with the Fauces and Tonsils, . 273
61. Diagram of the Superior Maxillary Nerve, ... 278
62. Deep view of the Spheno-palatine Ganglion, ... 280
63. Communications of the Facial, Glosso-pharyngeal, Pneumogastric,

FIG. PAGE

Spinal Accessory, Hypoglossal, Sympathetic, and the two Upper Cervical Nerves, ... 283

64. The Geniculate Ganglion of the Facial Nerve and its Communications, ... 287

65. Cartilages of the Nose, ... 291

66. Transverse Section through the Abdomen to show the attachment of the Lumbar Fascia, ... 298

67. The Superficial Muscles of the Back, ... 299

68. The Suboccipital Triangle, ... 307

69. The Cutaneous Nerves of the Back, ... 311

70. The Prevertebral Muscles, ... 313

71. The Odontoid and Transverse Ligaments, ... 318

72. Costo-vertebral Ligaments, ... 320

73. The Ligaments connecting the Rib with the Vertebra, . . . 321

74. Transverse Section to show the Ligaments and the Fibro-Cartilage of the Lower Jaw, ... 323

75. The Axilla, ... 335

76. Plan of the Branches of the Axillary Artery, ... 336

77. The Origins of the Triceps, ... 337

78. The Brachial Plexus of Nerves, ... 339

79. Distribution of the Cutaneous Nerves to the front of the Shoulder and Arm, ... 345

80. Superficial Veins and Nerves at the bend of the Left Elbow, . 346

81. Plan of the Chief Branches of the Brachial Artery, . . . 353

82. The Muscles of the Front of the Forearm, ... 362

83. The Superficial and Deep Palmar Arches, ... 376

84. Cutaneous Nerves of the Back, ... 389

85. The Superficial Muscles of the Back, ... 391

86. Cutaneous Nerves of the Left Shoulder and Arm (posterior view) . 399

87. Analysis of the Deltoid, ... 400

88. The Arteries of the Scapula, ... 404

89. The Anastomoses of the Arteries at the back of the Elbow and Wrist Joints, ... 416

90. The Dorsal Interossei, ... 420

91. The Palmar Interossei, and the Adductor Pollicis, ... 420

92. The Sterno-clavicular Ligaments, ... 423

FIG. PAGE

93. Anterior view of the Scapulo-clavicular Ligaments, and of the Shoulder-joint, ... 424
94. Ligaments of the Elbow-joint, ... 428
95. The Ligaments and Synovial Membranes of the Wrist-joint, . 432
96. The Abdominal Regions, ... 440
97. Superficial Vessels and Glands of the Groin, ... 442
98. Poupart's Ligament, and the External Abdominal Ring, . . 446
99. The Lower Fibres of the Internal Oblique and Transversalis, with the Cremaster Muscle, ... 448
100. Transverse Section to show the Formation of the Sheaths of the Rectus, the Quadratus Lumborum, and the Erector Spinæ, . 451
101. The Fascia Transversalis, seen from the front, ... 455
102. Varieties of Congenital Inguinal Herniæ, ... 464
103. Section to show the Ileo-cæcal Valve and Appendix Vermiformis. 471
104. Relative Position of the Kidneys and the Large Intestine, seen from behind, ... 472
105. The Peritoneum, ... 479
106. A Transverse Section through the Upper Part of the Abdominal Cavity, ... 481
107. A Transverse Section through the Lower Part of the Abdominal Cavity, ... 482
108. Branches of the Abdominal Aorta, ... 485
109. The Branches of the Cœliac Axis, ... 487
110. The Vena Portæ, ... 490
111. Plan of the Mesenteric Arteries, and their Communications, . 492
112. The Diaphragm, ... 499
113. The Diaphragm from its Upper Surface, ... 500
114. The Course and Relations of the Abdominal Aorta and Vena Cava Inferior 504
115. Plan of the Lumbar Plexus and Branches 517
116. The Relative Position of the Pelvic Viscera 522
117. Vertical Section through the Female Pelvic Viscera . . . 524
118. Framework of the Perineum 526
119. Muscles, with Superficial Vessels and Nerves, of the Perineum . 530
120. The Accelerator Urinæ in profile 532
121. The Triangular Ligament of the Urethra 533

FIG. PAGE
122. The Parts behind the Anterior Layer of the Triangular Ligament 534
123. The Relations of the Compressor Urethræ 538
124. Bulb of the Vagina 544
125. Vertical Section through the Perineum and Pelvic Viscera . 545
126. Transverse Section to show the Reflections of the Pelvic Fascia 546
127. Side View of the Pelvic Viscera 549
128. Posterior View of the Bladder 552
129. Plan of the Branches of the Internal Iliac Artery . . . 558
130. View of the Abnormalities of the Obturator Artery . . . 560
131. Plan of the Sacral Plexus and Branches 564
132. Bladder and Urethra exposed from the Upper Surface . . 568
133. Transverse Sections of the Urethra 575
134. Transverse Section through the Penis 578
135. The Uterus, the Ovaries and Fallopian Tubes 589
136. The Under Surface of the Liver 595
137. Transverse Sections of Lobules of the Liver 597
138. Longitudinal Sections of Lobules of the Liver 598
139. Section of the Kidney 608
140. The Course and Arrangement of the Uriniferous Tubes . . 610
141. A Tubulus Uriniferus 612
142. Section to show the Ileo-cæcal Valve 623
143. Saphenous Opening with the Cribriform Fascia 629
144. Fascia on the Outside of the Thigh 633
145. The Femoral Ring and the Saphenous Opening 634
146. Position of Parts under the Crural Arch 637
147. The Sheath of the Femoral Vessels 638
148. View of the Abnormalities of the Obturator Artery . . . 642
149. Scarpa's Triangle 644
150. Section through Hunter's Canal 653
151. Plan of the Inosculations of the Circumflex Arteries . . 656
152. Plan of the Sacral Plexus and Branches 676
153. Deep Muscles of the Gluteal Region 677
154. The Arteries of the Gluteal Region 679
155. Left Popliteal Space 681
156. Diagram showing Action of the Hamstring Muscles . . . 687
157. Diagram showing Action of the Gastrocnemius 691
158. Muscles, Vessels and Nerves of the Sole of the Right Foot . 701

FIG. PAGE
159. Plantar Arteries 703
160. View of the Third Layer of the Muscles of the Foot . . . 706
161. The Sacro-sciatic Ligaments 710
162. Vertical Section through the Hip 713
163. The Semilunar Cartilages and Lateral Ligaments of the Knee 717
164. Crucial Ligaments of the Knee 721
165. Ligaments of Ankle-joint 722
166. The External Lateral Ligament 723
167. Calcaneo-cuboid Articulation 726
168. Interosseous Ligaments of the Cuneiform Bones . . . 727
169. The Articulations of the Tarsus and the Tarso-metatarsus . 728
170. The Base of the Brain 732
171. The Front Surface of the Medulla Oblongata . . . 737
172. The Fourth Ventricle and the Restiform Bodies . . . 739
173. The Course of the Fibres through the Medulla Oblongata . 742
174. The General Division of the Brain 746
175. Convolutions and Fissures of the External Surface of the Brain 748
176. Convolutions of the Upper Surface of the Brain . . . 750
177. Convolutions of the Base of the Cerebrum 752
178. Convolutions and Fissures of the Median Surface of Right Hemisphere 754
179. Diagram of the Course of the Fibres through the Medulla and Pons 759
180. The Origins of the Olfactory and Optic Nerves 760
181. The Floor of the Fourth Ventricle 764
182. Upper Surface of the Corpus Callosum 767
183. Diagram of the Lamina Cinerea 767
184. Vertical Section through the Corpus Callosum, and parts below . 768
185. The Lateral Ventricles 770
186. Transverse Vertical Section through the Brain 771
187. The Fornix 773
188. The Lateral Ventricles and the Velum Interpositum . . . 776
189. The Cerebellum 781
190. Floor of the Fourth Ventricle 784
191. Superior Surface of the Cerebellum 787
192. Inferior Surface of the Cerebellum 789
193. The Spinal Veins (vertical section) 793

FIG. PAGE
194. The Spinal Veins (transverse section) 794
195. The Ligamentum Denticulatum 797
196. A Transverse Section through the Spinal Cord and its Membranes 799
197. Insertion of the Recti Muscles 807
198. A Vertical Section of the Eye 808
199. The Choroid, the Ciliary Muscle, and Nerves 812
200. The Various Layers of the Retina 818
201. Arteries of the Retina 824
202. The Ossicles of the Right Tympanum 832
203. Osseous Labyrinth of the Right Side 836
204. The Osseous Cochlea 838
205. Section of a Coil of the Cochlea 839
206. Vertical Section of the First Turn of the Cochlea . . . 841
207. A Vertical Section through the Testicle 848
208. Transverse Section through the Left Testicle 850

A MANUAL

OF THE

DISSECTION OF THE HUMAN BODY.

DISSECTION OF THE SCALP.

Dissection. An Incision should be made from the root of the nose along the mesial line of the vertex to the external protuberance of the occipital bone; another, horizontally round each half of the head, to join at right angles the two ends of the first incision. These incisions must not divide more than the skin, so that the subcutaneous vessels and nerves be not injured. It is well to dissect on one side of the head the muscles only, reserving the other side for the dissection of the vessels and nerves.

Strata composing the Scalp. The several strata of tissues covering the skull-cap are—1, the *skin;* 2, a thin layer of *connective tissue* and *fat* which contains the cutaneous vessels and nerves and the bulbs of the hair; and by which the skin is very closely connected to, 3; the *broad, thin aponeurosis* of the occipito-frontalis muscle (aponeurosis of the scalp); 4, an abundance of loose *connective tissue*, which permits the free motion of the scalp upon, 5, the *pericranium*, or periosteum of the skull-cap.

Immediately beneath the skin, then, we expose the thin stratum of connective and adipose tissue which firmly connects it with the aponeurosis of the scalp. This layer is continuous behind with the superficial fascia covering the muscles at the back of the neck, and laterally it passes over the temporal fascia. It forms a bed for the bulbs of the hair and for the ramifications of

the cutaneous arteries. The toughness of this tissue, in which the arteries ramify, does not permit them to retract when divided; hence the hemorrhage which follows incised wounds of the scalp; hence, also, the difficulty of drawing them out with the forceps.

Occipito-frontalis Muscle and Epicranial Aponeurosis.

This cutaneous muscle is closely connected to the scalp. It consists of two fleshy portions, one on the occiput, the other on the forehead, connected by a broad aponeurosis. The *occipital*

FIG. 1.

DIAGRAM SHOWING THE MUSCULAR AND APONEUROTIC STRATUM OF THE SCALP.

A. Attollens aurem.
B. Attrahens aurem.
C. Retrahens aurem.
D. Orbicularis palpebrarum.

portion of the muscle is thin, and takes origin from the outer two-thirds of the upper curved line of the occipital bone, and the adjoining part of the mastoid process of the temporal bone. The fibres ascend over the back of the head for about two inches, and then terminate in the epicranial aponeurosis. The *frontal portion*, commencing in an arched form from the epicranial aponeurosis below the coronal suture, descends over the forehead, and termi-

nates partly in the skin of the brow, partly in the orbicularis oculi and corrugator supercilii, while some of the inner fibres are continuous in front of the nose with the pyramidalis nasi muscle. The aponeurosis of the scalp covers the vertex of the skull, the two being continuous across the middle line. It is continued over the temples and side of the head, gradually changing from tendinous into connective tissue. This muscle enables us to move the scalp backward and forward. But its chief action is as a muscle of expression. It elevates the brows, and occasions the transverse wrinkles in the expression of surprise. The occipital portion is supplied by the posterior auricular branch of the facial; the frontal portion by the temporal branch of the same nerve.

Muscles of the Ear. There are several small muscles to move the cartilage of the ear. In man they are thin and pale, and require care to dissect them out satisfactorily. In animals who possess a more delicate sense of hearing, they are much more developed, for the purpose of quickly directing the cartilage of the ear toward the direction of the sound.

Attollens Aurem. To indicate the position of this muscle the student should draw down the upper part of the pinna of the ear, when it will be found immediately under the ridge of skin so produced. It is a thin, fan-shaped muscle, and *arises* from the epicranial aponeurosis, and is *inserted* into the cranial aspect of the upper part of the concha.

Attrahens Aurem. This muscle is the smallest of these muscles, and its situation is indicated by the prominence of skin produced by drawing backward the front part of the helix. It *arises* from the aponeurosis of the occipito-frontalis, and is *inserted* into the front of the helix.

Retrahens Aurem. This muscle is exposed by reflecting the skin from the ridge produced by drawing the pinna forward. Consisting of two or three fasciculi, it *arises* from the base of the mastoid process and is *inserted* into the lower part of the concha.

The retrahens and the attollens aurem are supplied by the posterior auricular branch of the facial nerve; the attrahens, by an offset from the temporal branch of the same nerve.

Arteries of the Scalp. The *arteries* of the scalp are derived, in front, from the *supra-orbital* and *frontal* arteries, branches of the ophthalmic artery, which is a branch of the internal carotid; on the sides, from the *temporal*; behind, from the *occipital* and *posterior auricular*, all branches of the external carotid.

The *frontal* emerges from the orbit at its inner angle; it runs upward for a short distance on the forehead and inosculates with the following artery.

The *supra-orbital* passes through the supra-orbital notch and then divides into a superficial and a deep branch. It distributes branches, some of which ascend toward the top of the head and communicate with the temporal and frontal arteries.

The *temporal*, about two inches above the zygoma, divides into two branches—an anterior and a posterior. The *anterior* runs forward in a tortuous course and anastomoses with the supra-orbital and frontal arteries; the *posterior* (usually the larger) arches backward over the temporal fascia, and its branches communicate with the corresponding branch of the opposite side and with the occipital and posterior auricular arteries.

The *posterior auricular* is a small vessel seen in the cleft between the ear and the mastoid process. It ascends, and divides into two branches: one, the mastoid or occipital, which passes backward and inosculates with the occipital; the other, the auricular, which runs forward above the ear and communicates with the posterior branch of the temporal artery.

The *occipital* may be noticed piercing the trapezius near to the external occipital protuberance; ascending over the back of the head, it divides into numerous branches which inosculate with the preceding arteries.

The frontal vein passes downward with its corresponding artery, and joins the supra-orbital vein, to form the angular vein. The other veins of the scalp accompany their respective arteries.

Nerves of the Scalp. The sensory nerves of the scalp are derived from each of the three divisions of the fifth cranial nerve, namely, the ophthalmic, the superior and inferior maxillary; also from the second cervical nerve. The nerves to

the muscles of the scalp and ear come from the facial, which is one of the divisions of the seventh cranial nerve.

In front will be found the supra-trochlear and supra-orbital nerves; in the temporal region, there are the temporal filament from the orbital branch of the superior maxillary, the auriculo-temporal, and the temporal branches of the facial nerve; and behind will be seen the posterior auricular branch of the facial,

FIG. 2.

DIAGRAM OF THE SENSORY NERVES OF THE SCALP AND FACE.

1. Great occipital.
2. Small occipital.
3. Auricular br. of the pneumogastric.
4. Great auricular.
5. Auriculo-temporal.
6. Temporal br of superior maxillary nerve.
7. Supra-orbital.
8. Supra-trochlear.
9. Malar br. of superior maxillary nerve.
10. Infra-trochlear.
11. Naso-lobular.
12. Infra-orbital.
13. Buccal br. of inferior maxillary nerve.
14. Mental.

the small and great occipital nerves, and, occasionally, a small filament from the posterior division of the sub-occipital nerve.

The *supra-trochlear nerve* is derived from the frontal branch of the ophthalmic division of the fifth. It appears at the inner angle of the orbit, and ascending beneath the orbicularis palpebrarum and occipito-frontalis, it finally supplies the skin of the forehead, and the upper eyelid.

The *supra-orbital nerve* is a continuation of the frontal branch of the fifth. It emerges from the orbit through the notch in the frontal bone, and subdivides into branches, which are covered at first by the fibres of the orbicularis and occipito-frontalis; but they presently become subcutaneous, and terminate in two branches—an *inner*, which ascends to supply the structures as high as the parietal bone; and an *outer* and larger, which may be traced over the vertex as far as the occipital bone.

The *temporal branch* of the *orbital branch* of the superior maxillary nerve pierces the temporal fascia about an inch above the zygoma, and is distributed to the skin of the temple, communicating with the facial nerve and occasionally with the following.

The *auriculo-temporal nerve*, a branch of the inferior maxillary nerve, after sending a small filament to the upper part of the pinna, divides into two branches, which accompany the divisions of the superficial temporal artery; of these, the posterior is the smaller. The anterior communicates with the facial nerve, and with the orbital branch of the superior maxillary.

The *temporal branches* of the facial nerve lie superficial to the temporal fascia, and supply the attrahens and attollens aurem, the orbicularis palpebrarum, the corrugator supercilii and the occipito-frontalis. These branches communicate with the temporal branch of the superior maxillary, the auriculo-temporal nerve, and with the lachrymal and supra-orbital branches of the ophthalmic.

The *posterior auricular nerve* is a branch of the facial, and divides, like its accompanying artery behind the pinna of the ear, into a *posterior* or *occipital* branch which supplies the posterior belly of the occipito-frontalis, and into an *anterior* or *auricular* branch which ends in the auricle, the retrahens and attollens aurem. It communicates with the great auricular and small occipital nerves, and with the auricular branch (Arnold's) of the pneumogastric.

The *auricular branch of the pneumogastric* (Arnold's) emerges from the auricular fissure immediately behind the pinna, and supplies the skin of the pinna and the neighborhood.

The *great occipital nerve* is the internal branch of the posterior division of the second cervical nerve. After piercing the com-

plexus it appears on the occiput with the occipital artery, and divides into wide-spreading branches which supply the skin. It communicates with the posterior auricular, the small occipital, and the third cervical nerves.

The *small occipital nerve*, a branch of the anterior division of the second cervical nerve, runs along the posterior border of the sterno-mastoid, and supplies the scalp behind the ear. It communicates with the great auricular, and with the two preceding nerves.

FIG. 3.



DIAGRAM OF THE BRANCHES OF THE FACIAL NERVE.

1. Branch to occipito-frontalis.
2. Posterior auricular.
3. Temporal brs.
4. Malar brs.
5. Infra-orbital.
6. Buccal.
7. Supra-maxillary.
8. Infra-maxillary.

Occasionally, though rarely, a cutaneous branch of the *sub-occipital nerve* is distributed to the back of the head.

Points of Surgical Interest. Raise the aponeurosis of the scalp, and observe the quantity of loose connective tissue which intervenes between it and the pericranium. This tissue never contains fat. There are some points of surgical interest concerning it:—1. Its looseness accounts for the extensive effusions

of blood which one often sees after injuries of the head. 2. It admits of large flaps of the scalp being detached from the skull-cap; but these flaps rarely slough, unless severely damaged, because they carry their blood vessels with them. 3. In phlegmonous erysipelas of the scalp, the connective tissue becomes infiltrated with pus and sloughs; hence the necessity of making incisions; for the scalp will not lose its vitality, and liberate the sloughs like the skin of other parts under similar conditions, because its vessels run above the diseased tissue, and therefore its supply of blood is not cut off.

Lymphatics of the Scalp. The lymphatics of the scalp run for the most part backward toward the occiput, to join the occipital and posterior auricular glands; a few run toward the root of the zygoma, where they enter the parotid lymphatic glands. It is in these situations, therefore, that one finds glandular enlargements when the scalp is diseased.

Dissection. To examine the brain and its membranes, the skull-cap must be removed about half an inch above the supra-orbital ridges in front, and on a level with the occipital protuberance behind. The student should remember that the bone in the temporal region is very thin, and that here especial care is needed that the brain be not injured by the saw. It is better to saw only through the outer table of the skull, and to break through the inner with a chisel. In this way the dura mater and the brain are less likely to be injured. On removing the skull-cap, which is more or less intimately attached to the subjacent membrane, we expose a tough fibrous layer, *the dura mater*, which forms the most external of the membranes of the brain.

The meningeal arteries ramify between the skull and the dura mater. We cannot, however, with the brain *in situ*, trace their course, at present, throughout; so their consideration must be deferred until the brain has been removed.

Dura Mater. This membrane is so called because it was thought to give rise to all the other fibrous membranes in the body. It is a dense white fibrous membrane, rough on its outer aspect, where it is more or less adherent to the inner

surface of the skull, forming its internal periosteum. On its inner surface it is smooth and shining, being lined by a layer of endothelial cells, which anatomists now describe as constituting a part of the dura mater. In consequence, the term "*subdural space*" is now substituted for the old one, "*the cavity of the arachnoid.*" The dura mater differs in its adhesion to the subjacent bones; its adhesion is firmest at the sutures, the petrous portion of the temporal bone, the basilar process, the body of the sphenoid, the cribriform plate of the ethmoid bone, the depressions for the Pacchionian bodies, and at the margin of the foramen magnum. In front it sends downward a prolongation into the foramen cæcum; also numerous small tubular sheaths through the foramina in the cribriform plate. It further sends a prolongation through the optic foramen, and another through the sphenoidal fissure into the orbit.

The dura mater is supplied with *nerves* by the recurrent branch of the fourth nerve, and by the fifth cranial nerve. Filaments have likewise been traced into it from the sympathetic and from the Gasserian ganglion.

Its remarkably tough and fibrous structure adapts it exceedingly well to the four purposes which it serves:—1. It forms the internal periosteum of the skull. 2. It forms, for the support of the lobes of the brain, three partitions—namely, the falx cerebri, the falx cerebelli, and the tentorium cerebelli. 3. It forms the sinuses or venous canals which return the blood from the brain. 4. It forms sheaths for the nerves as they leave the skull.

Of the partitions formed by the dura mater for the support of the lobes of the brain, two are vertical, and separate, respectively, the two hemispheres of the cerebrum and those of the cerebellum; the third arches backward, and supports the posterior lobes of the cerebrum.

Falx Cerebri. This partition is named, from its resemblance to the blade of a sickle, *falx cerebri*. It is received into the longitudinal fissure, and separates the two cerebral hemispheres. It begins in a point attached to the *crista galli*, and gradually becomes broader as it extends backward. Its upper

edge is convex, and attached to the median groove on the inner aspect of the vertex of the skull; its lower margin is concave and free, and runs along the upper aspect of the corpus callosum. From its base or broadest part proceeds the sloping arched partition named *tentorium cerebelli*. This forms an arch for the support of the posterior lobes of the cerebrum, so that they may not press upon the cerebellum beneath. The tentorium is attached to the transverse ridge of the occipital bone, to the superior border of the petrous portion of the temporal bone, and to the posterior and anterior clinoid processes of the sphenoid. In front there is a large oval opening, to allow of the passage of the crura cerebri. The small median partition which separates the lobes of the cerebellum is called the *falx cerebelli*. It is placed vertically in the same plane with the falx cerebri, and its point is downward toward the foramen magnum. As it approaches the foramen it usually divides into two small folds.

Tentorium Cerebelli.

Falx Cerebelli.

Glandulæ Pacchioni.

In the neighborhood of the superior longitudinal sinus, we meet with small, white, elevated granulations, sometimes arranged singly, sometimes in clusters, which are received into the depressions on the inner aspect of the skull-cap. They are termed *glandulæ Pacchioni*,[1] and are found in four situations:—1. On the outside of the dura mater, close to the superior longitudinal sinus, and so large as to occasion depressions in the bones. 2. Along the margin of the fissure of Sylvius. 3. On the surface of the pia mater. 4. In the interior of the superior longitudinal sinus, covered by its lining membrane. 5. On the posterior and antero-inferior parts of the posterior lobe of the cerebrum.

They are due to an increased growth of the villi, which are normally found in the arachnoid membrane, and make their way, through the dura mater or the pia mater, to the different situations in which they are found. The greatest growth takes place from the visceral layer, as may be seen in the dissection of the brain. These bodies are not found at birth, but usually commence their growth about the third year, and are always found at

[1] After the Italian anatomist who first described them, in 1705.

the seventh year, after which they gradually increase as life advances.[1]

Sinuses of the Dura Mater. It is one of the peculiarities of the cerebral circulation, that the blood is returned through canals or *sinuses* formed by the dura mater. These canals are produced by a splitting of the dura mater into two layers, as shown in Fig. 4, where 1 represents a vertical section through the superior longitudinal sinus. They are lined by the same smooth membrane continuous with that of the venous system. Since their walls consist of unyielding structure, and are always on the stretch, it is obvious that they are admirably adapted to resist the pressure of the brain. There are fifteen of these sinuses, and they are classified into two groups—a *supero-posterior* and an *infero-anterior*. The supero-posterior group comprises the superior longitudinal, the inferior longitudinal, the straight, the lateral and the occipital sinuses; while the infero-anterior group includes the cavernous, the circular, the superior and inferior petrosal, and the straight sinuses. Of these fifteen sinuses, five are pairs and five are single, as follows :—

FIG. 4.



DIAGRAM TO SHOW FORMATION OF A SINUS.

The five pairs of sinuses are—	The five single sinuses are—
The lateral.	The superior longitudinal.
The superior petrosal.	The inferior longitudinal.
The inferior petrosal.	The circular.
The cavernous.	The transverse.
The occipital.	The straight.

The blood from all these sinuses is eventually discharged into the internal jugular veins.

Superior Longitudinal Sinus. This runs along the upper attached border of the falx cerebri (Fig. 5). It begins very small at the foramen cæcum, gradually increases in size in its course backward, and opposite the internal protuberance of the occipital bone opens into a triangular dilatation, the *torcular Herophili*, or the *confluence of the sinuses*. It then divides into the right

[1] It is stated that when fluid is injected into the subarachnoid space, it passes into the Pacchionian bodies.

and left lateral sinuses, the right being generally the larger. Besides numerous veins from the cancellous texture of the skull-cap, the superior longitudinal sinus receives large veins from the upper part of each hemisphere of the cerebrum, and an emissary vein through the parietal foramen. It is interesting to observe that these veins run (as a rule) from behind forward, contrary to the current of blood in the sinus, and that they pass through the wall of the sinus very obliquely, like the ureter into the bladder. The probable object of this oblique entrance is to prevent regurgitation of blood from the sinus into the veins of the brain.

FIG. 5.



1. Superior longitudinal sinus.
2. Inferior longitudinal sinus.
3. Straight sinus.
4, 4. Venæ Galeni.

Cut open the superior longitudinal sinus; observe that it is triangular, with its base upward, and that its cavity is intersected in many places by slender fibrous cords, termed *chordæ Willisii*.[1] Their precise use is not understood.

The brain should now be removed, and preserved in spirit for future examination. Its anatomy, with that of its remaining membranes, will be described in a subsequent part of this work.

Dissection. The brain is to be removed in the following manner:—The dura mater should be cut through with a pair of scissors on a level corresponding with the sawn calvarium, care being taken to cut completely through the falx cerebri in the front part of the longitudinal fissure. When this has been done the dura mater can be easily turned back over the brain, leaving its smooth, convex surface exposed. Now lift up gently, with the fingers of the left hand, the frontal lobes from the anterior fossæ, taking care to raise with the brain the soft olfactory lobes from the cribriform plate of the ethmoid. Two white, flat nerves, the optic, come into view prior to their leaving the skull through the optic foramina; these must be divided with a sharp knife,

[1] So called after Willis, who first described them in his work, *De Cerebri Anatome*, 1664.

together with the ophthalmic arteries which lie beneath the corresponding nerves.[1] In the middle line, fixed firmly in the sella turcica, lies the pituitary body, attached to the brain by a process—the infundibulum. It is not easy to remove this body from the fossa in which it rests, owing to its being retained in its position by dura mater. When this is removed, two round white nerves—the third—are observed, one on each side, lying on the inner free border of the tentorium cerebelli, immediately behind the anterior clinoid process of the sphenoid. Divide these and then proceed to cut through the tentorium cerebelli close to its attachment to the posterior clinoid process and the upper border of the petrous portion of the temporal bone, as far back as the lateral sinus. If this be done with care, the nerves lying beneath the tentorium will not have been injured. Immediately external to the third nerves are the slender fourth nerves; and still further outside are the soft, flattened fifth nerves. Cut these through, still gently raising the brain from the skull base, when the seventh pair come into view as they pass backward and outward toward the internal auditory foramina. When these have been cut, we notice the two sixth nerves running directly forward to pierce the dura mater covering the basilar process of the occiput. Divide these as they pierce the dura mater, when the three divisions of the eighth are brought well into view, lying behind and internal to the seventh; the anterior one is the glosso-pharyngeal, the middle one is the pneumogastric, and the hindermost one is the spinal accessory, whose spinal portion can be traced coming up from the foramen magnum. These all emerge through the jugular foramina. Below and internal to these are the hypoglossal nerves, which usually pass through the dura mater in two fasciculi. Cut these and then pass down the knife as far into the spinal canal as possible, and cut through the spinal cord, the two vertebral arteries, and the spinal portions of the spinal accessory nerves. Now lay the knife aside, when by gentle traction the brain can be easily removed from the skull.

The other sinuses should now be examined.

[1] It is well that each pair of nerves should be cut through first on one side and then on the other, before passing on to the nerve next in numerical order.

Lateral Sinuses. These are the two great sinuses through which all the blood from the brain is returned to the jugular veins. Their course is well marked in the dry skull. The right is usually the larger. Each commences at the internal occipital protuberance, and proceeds at first horizontally outward, enclosed between the layers of the tentorium, along a groove in the occipital bone and the posterior inferior angle of the parietal; it then descends along the mastoid portion of the temporal bone, and again indenting the occipital bone, turns forward to the foramen lacerum posterius, and terminates in the bulb of the internal jugular vein,[1] where it is joined by the inferior petrosal sinus. It receives blood also from the inferior cerebral and cerebellar veins, from the diploë, and the superior petrosal sinus. It communicates with the veins of the scalp through emissary veins, which pass through the mastoid and posterior condylar foramina.

Inferior Longitudinal Sinus. This is of small size. It runs in the inferior free border of the falx cerebri, and terminates in the straight sinus at the anterior margin of the tentorium (Fig. 5).

Straight Sinus. This may be considered as the continuation of the preceding. It runs along the line of junction of the falx cerebri with the tentorium cerebelli, and terminates in the torcular Herophili at the divergence of the two lateral sinuses. It receives the inferior cerebral and the superior cerebellar veins, and also the two *venæ Galeni* (Fig. 5), which return the blood from the lateral and third ventricles of the brain.

Cavernous Sinus. This is so called because its interior is intersected by numerous cords. It extends along the side of the body of the sphenoid bone, outside the internal carotid artery. It receives the ophthalmic vein which leaves the orbit through the sphenoidal fissure and the anterior inferior cerebral veins; it communicates with the circular sinus which surrounds the pituitary body (Fig. 6). At the apex of the petrous portion of the temporal bone it divides into the superior and inferior petrosal sinuses.

[1] It has, in some subjects, another outlet, through the foramen mastoideum, or else through the posterior condylar foramen.

Circular Sinus. This surrounds the pituitary body (P in the diagram, Fig. 6.), and communicates on each side with the cavernous sinus. The posterior branch is sometimes absent.

Petrosal Sinuses. These lead from the cavernous to the lateral sinuses. There are two on each side. The *superior* runs along the upper portion of the pars petrosa, in the

FIG. 6.



DIAGRAM OF THE VENOUS SINUSES AT THE BASE OF THE SKULL.

attached border of the tentorium cerebelli; the *inferior*, the larger of the two, runs along the suture between the pars petrosa and the occipital bone, and ends in the lateral sinus just before this terminates in the internal jugular vein. The superior sinus receives the inferior cerebral, the superior cerebellar veins, and a small branch from the tympanum; the inferior sinus is joined by the inferior cerebellar and auditory veins.

Transverse Sinus. This extends from one inferior petrosal to the other, across the basilar process of the occipital bone. It communicates below with the anterior spinal veins.

Occipital Sinuses. These are very small. They commence around the margin of the foramen magnum, run in the falx cerebelli, and uniting to form a single sinus, open into the torcular Herophili.[1] They join inferiorly with the posterior spinal veins.

Meningeal Arteries. These arteries ramify between the skull and the dura mater. Their course may be traced by the grooves which they make in the bones. They are termed *anterior*, *middle*, and *posterior*, from the fossæ in which they ramify.

The *anterior meningeal* are derived from the ethmoidal branches of the ophthalmic artery and the cavernous portions of the internal carotid. They supply the dura mater in the neighborhood of the ethmoid bone.

The *middle meningeal* are three in number: the most important is the *arteria meningea media*, a branch of the internal maxillary artery. It enters the skull through the foramen spinosum, and divides into two principal branches: one, the anterior, runs in a groove near the anterior border of the parietal bone; the other, the posterior, curves backward over the temporal bone, and subsequently ramifies on the parietal bone. The artery gives off a small branch—the *petrosal*, which enters the hiatus Fallopii and anastomoses with the stylo-mastoid artery in the aquæductus Fallopii; one or more anastomosing branches which enter the orbit through the sphenoidal fissure to communicate with the ophthalmic artery; and some temporal branches which pierce the sphenoid bone to enter the temporal fossa. It is accompanied by two veins which empty themselves into the internal maxillary

[1] The junction of the several sinuses opposite the spine of the occipital bone is termed the *torcular Herophili*, after the celebrated anatomist who first described it. It is a kind of triangular reservoir, with the base below, and presents six openings—namely, that of the superior longitudinal sinus, those of the two lateral and of the two occipital, and that of the straight sinus. The term *torcular* is an incorrect version of the original word σωλήν (a canal or gutter) employed by Herophilus.

vein. The *arteria meningia parva*, which enters the skull through the foramen ovale, and a *meningeal branch* from the ascending pharyngeal artery, which comes up through the foramen lacerum medium, also supply the dura mater and bones of the middle fossa.

The *posterior meningeal* come from the occipital, the ascending pharyngeal and the vertebral arteries; the two former enter the skull through the foramen jugulare, and the latter through the foramen magnum. The meningeal veins, with the exception of the middle meningeal, open into the various sinuses.

The position of the meningeal arteries renders them liable to injury in fractures of the skull; hence, extravasation of blood between the skull and dura mater is one of the common causes of compression of the brain.

Dissection. The student should now examine the cranial nerves as they pass out through the foramina in the base of the skull, and then dissect the cavernous sinus.

Exit of the Cranial Nerves. The cranial nerves proceed in pairs through the foramina at the base of the skull; they are named first, second, third, fourth, etc., pairs, according to the order of succession from before backward. As they pass through the foramina, each receives a process from the three membranes of the brain, the dura mater, the pia mater, and the arachnoid; the two first are gradually lost upon the nerve, while the arachnoid is reflected back.

The *first* is the *olfactory nerve*. This cannot be seen, because the olfactory bulb has been removed with the brain. From the under aspect of the bulb proceed about twenty branches, which pass through the foramina in the cribriform plate of the ethmoid bone, and are arranged in three groups—inner, middle, and outer. The *inner* (larger) pass to the septum nasi; the *middle* to the roof of the nose; and the *outer* to the outer wall of the nose as low as the middle turbinated bone.

The *second* (*optic nerve*) passes through the foramen opticum into the orbit, accompanied by the ophthalmic artery.

In order to see the next three pairs of nerves, the dura mater must be carefully removed from the side of the body of the

sphenoid, and the nerves traced as they pass through the tentorium cerebelli.

The *third* (*motor oculi*) passes through the dura mater, close behind the anterior clinoid process, traverses the outer wall of the cavernous sinus, and enters the orbit through the sphenoidal fissure, where it receives some filaments from the cavernous plexus of the sympathetic. Before passing through the fissure, it divides into two branches, an upper and a lower, which enter the orbit between the two heads of the external rectus.

The *fourth* (*trochlearis*), a small nerve, passes through the dura mater a little behind the posterior clinoid process. It passes through the outer wall of the cavernous sinus, lying below the preceding nerve and above the first division of the fifth, and then runs forward through the sphenoidal fissure. Here it lies above the third nerve, and is finally distributed to the superior oblique muscle, on its orbital surface. In passing through the cavernous sinus it receives some branches from the sympathetic plexus. It also communicates here with the ophthalmic nerve, and sends back a recurrent branch to supply the tentorium cerebelli as far back as the internal occipital protuberance.

The *fifth* (*trifacial*) nerve passes through an aperture in the dura mater beneath the tentorium cerebelli, just above the apex of the petrous portion of the temporal bone. It consists of two parts —a larger or sensory root, and a smaller or motor. Upon its larger or sensory root is developed a large ganglion, the *Gasserian ganglion;* while the motor root lies below and unconnected with it. From this ganglion proceed the three primary divisions of the nerve—the *ophthalmic*, which passes through the outer wall of the cavernous sinus below the fourth nerve, and subsequently enters the orbit through the sphenoidal fissure; while in the cavernous sinus this nerve receives filaments of communication from the cavernous plexus, and also sends back a recurrent branch to supply the tentorium cerebelli (Arnold); the ophthalmic nerve is frequently intimately connected with a branch from the fourth nerve; it is also connected by a small branch with the sixth nerve, the *superior maxillary*, which gives off a small recurrent branch to the dura mater and middle meningeal artery, and then leaves

the skull through the foramen rotundum, and the *inferior maxillary*, which passes through the foramen ovale. The smaller or motor root of the fifth lies beneath the ganglion, with which it

FIG. 7.



DIAGRAM OF THE EXIT OF THE CRANIAL NERVES.

has no communication, and then joins the inferior maxillary division to supply the muscles of mastication with motor power.

The *sixth* (*abducens*) nerve pierces the dura mater behind the body of the sphenoid bone, which it grooves. It then passes along the inner wall of the cavernous sinus, external to the in-

ternal carotid artery, and enters the orbit through the sphenoidal fissure, to supply the external rectus, between the two heads of which it passes. It is connected, as it passes along the inner wall of the cavernous sinus, with the cavernous plexus, the ophthalmic nerve, and in the orbit with Meckel's ganglion.

The *seventh* or *facial nerve* passes through the meatus auditorius internus, together with the auditory nerve and artery. As it passes along the meatus it is separated from the auditory nerve, upon which it lies, by the portio intermedia. At the bottom of the auditory meatus, the facial nerve leaves the auditory to traverse a tortuous bony canal, the "aquæductus Fallopii." In

FIG. 8.



DIAGRAM SHOWING THE RELATIONS OF THE VESSELS AND NERVES PASSING THROUGH THE FORAMEN JUGULARE.

the meatus auditorius, the facial and the auditory nerves are connected by small filaments.

The *eighth* or *auditory nerve* passes outward through the internal auditory meatus in company with the preceding nerve. It is the larger of the two nerves, and lies below the facial, which lies in a groove on this nerve. In the meatus the auditory divides into two branches, cochlear and vestibular.

The *ninth* or *glosso-pharyngeal nerve* passes through the jugular foramen in front of the pneumogastric and spinal accessory nerves. This nerve has a separate tube of dura mater and arachnoid, and

lies in a groove in the lower border of the pars petrosa of the temporal bone, together with the two succeeding nerves (Fig. 8).

The *tenth* or *pneumogastric nerve* emerges through the jugular foramen behind and rather internal to the glosso-pharyngeal. It is enclosed in a common sheath of dura mater with the spinal accessory, but is separated from it by a thin septum of arachnoid membrane.

The *eleventh* or *spinal accessory* also passes through the foramen jugulare, lying behind the preceding nerve.

The *twelfth* or *hypoglossal nerve* passes through the anterior condylar foramen, piercing the dura mater by two fasciculi which unite external to the skull.[1]

Dissection. We must now examine the cavernous sinus, and the nerves which course along its walls to the orbit—namely, the third, the fourth, the ophthalmic division of the fifth and the sixth nerves.

Cavernous Sinus. This sinus (Fig. 6) lies by the side of the body of the sphenoid bone. In front it receives the ophthalmic vein, which passes backward through the sphenoidal fissure; while posteriorly it divides into the superior and inferior

FIG. 9.



RELATION OF THE VARIOUS STRUCTURES PASSING THROUGH THE CAVERNOUS SINUS.

petrosal sinuses, which have been already described; on the inner side it communicates with the circular sinus, which surrounds the pituitary body (P in the diagram, Fig. 6). The interior of the sinus is remarkable for the numerous fine bands of reticular tissue which interlace in all directions.

In the outer wall of the cavernous sinus we trace, from above

[1] According to the classification adopted by Willis, the cranial nerves are arranged in nine pairs; the seventh including the facial and the auditory; the eighth comprising the glosso-pharyngeal, the pneumogastric, and the spinal accessory. The arrangement adopted in this handbook is that by Sömmering.

downward, the third nerve, the fourth, and the ophthalmic division of the fifth, in their course to the orbit. On its inner wall are situated the internal carotid artery with the sixth nerve below and to its outer side. These structures are not actually within the sinus so as to be bathed by the blood, for they are separated from it by the lining membrane of the sinus (Fig. 9).

Relative Positions of Nerves in Sphenoidal Fissure.

These nerves should be traced from the cavernous sinus, forward, so as to see how they alter their relative positions before entering the sphenoidal fissure, and again, in their passage through it.

Just *before entering the sphenoidal fissure*, the fourth nerve (on

FIG. 10.



DIAGRAM OF THE RELATIONS OF THE NERVES AS THEY PASS THROUGH THE SPHENOIDAL FISSURE.

its way to the orbital surface of the superior oblique) gets above the third, which here divides into an upper and a lower branch (both proceeding to the ocular surface of the muscles they supply); lower still, we have the frontal, lachrymal and nasal divisions of the ophthalmic; lowest of all is the sixth nerve on its way to the external rectus.

In their passage through the sphenoidal fissure, we find that the fourth nerve, the frontal and lachrymal branches of the ophthalmic, lie at the top, on nearly the same level, and they enter the orbit above the muscles in this order, from within outward. Lower,

and in the following order from above downward, come the upper division of the third, the nasal branch of the ophthalmic, the lower division of the third, and the sixth; all of which (with the ophthalmic vein) enter the orbit between the two origins of the rectus externus (Fig. 10).

At the *back of the orbit* the relation of these nerves is further altered. The fourth, frontal and lachrymal, are still on the same level; the upper division of the third is below the superior rectus, and above the optic nerve is the nasal n.; the sixth is on the inner side of the external rectus, while the lower division of the third is below and to the outer side of the optic nerve, close to which is the lenticular ganglion (Fig. 11).

FIG. 11.



RELATIONS OF THE NERVES AND MUSCLES AT THE BACK OF THE ORBIT.

The dissector will better remember the varying relations of these nerves, when he has learned their respective destinations.

Curves of the Carotid Artery. After the removal of the cavernous sinus, a good view is obtained of the curves, like the letter S, made by the internal carotid artery on the side of the pituitary fossa. The vessel enters the cranium at the apex of the petrous portion of the temporal bone, makes its sigmoid curves within the cavernous sinus, and then passes through the dura mater, between the anterior clinoid process and the optic nerve, where it gives off the ophthalmic artery. Within the cavernous sinus, small branches, *arteriæ receptaculi*, arise from the carotid and supply the pituitary body and the walls of the sinus.

A careful dissection would show a plexus of sympathetic nerves on the outer side of the internal carotid artery, as it lies by the side of the body of the sphenoid. This is the CAROTID PLEXUS. It is connected by numerous filaments with the sixth nerve and the Gasserian ganglion. It further furnishes the large deep petrosal nerve which unites with the large superficial petrosal nerve of the facial to form the Vidian; and also the small deep petrosal nerve which joins probably the tympanic plexus. Those filaments of the sympathetic seen on the inner side of the artery in the upper part of the cavernous sinus constitute the CAVERNOUS PLEXUS, which is in communication with the third, the fourth, and the ophthalmic division of the fifth nerves, and gives a branch to the lenticular ganglion in the orbit.

FIG. 12.

THE GENICULATE GANGLION OF THE FACIAL NERVE.

1. The chorda tympani.
2. The geniculate ganglion of the facial nerve.
3. The great petrosal nerve.
4. The lesser petrosal nerve lying over the tensor tympani.
5. The external petrosal nerve communicating with the sympathetic plexus on the arteria meningea media (6).
7. The Gasserian ganglion.

On removing the Gasserian ganglion, three small nerves are seen lying on the anterior surface of the petrous portion of the temporal bone. One, the *large superficial petrosal nerve*, enters the hiatus Fallopii to join the facial; the second, immediately external to the preceding, is the *small superficial petrosal*, which passes from the facial to join the otic ganglion; the third, the *external superficial petrosal nerve* (not always present), passes from the facial to communicate with the sympathetic on the middle meningeal artery.

DISSECTION OF THE FACE.

Much practice is required to make a good dissection of the face, and it is well, therefore, to dissect this part before the skin and subjacent structures get dry and discolored.

The muscles of expression are numerous and complicated; they are interwoven with the subcutaneous tissue and closely united to the skin; their fibres are often pale and indistinct. The face is largely supplied with motor and sensory nerves, of which the ramifications extend far and wide. Therefore you must not be discouraged if, in a first attempt, you fail to make a satisfactory display of the parts.

The cheeks and nostrils should be distended with horse-hair, and the lips sewn together.

Make an incision down the mesial line of the face; another from the chin along the base of the lower jaw to the angle; then prolong it, in front of the ear, to the zygoma. Reflect the skin from below upward. Each muscle, to be properly cleaned, should be put on the stretch by hooks.

The student is recommended to make out the muscles and arteries on the one side, leaving the other side for the display of the nerves.

The motor nerve which supplies all the muscles of expression in the face is the "*portio dura*," or facial nerve. It emerges from the stylo-mastoid foramen, and divides into branches, which pass through the parotid gland, forming a plexus, termed the "*pes anserinus*."

The sensory nerves of the face are chiefly derived from the three divisions of the fifth cranial nerve; namely, the supra-orbital, the supra-trochlear, the lachrymal, the infra-trochlear, and naso-lobular, which latter supply the ala and the tip of the nose; the three sets of branches from the infra-orbital; and the mental. The other nerves which confer sensation upon the face are, the great auricular branch of the cervical plexus, which supplies the skin covering the parotid gland and part of the cheek.

It is convenient to arrange the muscles of the face under three groups; appertaining, respectively, to the mouth, the nose, the eyebrows and lids. Begin with those of the mouth.

The muscles of the mouth are arranged thus: there is an orbicular or sphincter muscle surrounding the lips; from this, as from a common centre, muscles diverge and are fixed into the surrounding bones. They are named elevators, depressors, sphincters, etc., according to their respective action.

Musculus Risorius (Santorini). This muscle is usually considered as a part of the platysma myoides, the large subcutaneous muscle of the neck. It arises by thin fasciculi from the fascia

FIG. 13.

1. Risorius.
2. Orbicularis oris.
3. Depressor anguli oris.
4. Depressor labii inferioris.
5. Levator menti.
6. Orbicularis palpebrarum.
7. Zygomaticus major.
8. Zygomaticus minor.
9. Pyramidalis nasi.
10. Compressor naris.
11. Levator labii superioris et alæ nasi.
12. Levator labii superioris.
13. Levator anguli oris.
14. Buccinator.
15. Occipito-frontalis.
16. Masseter.
17. Sterno-mastoid.
18. Trapezius.
19. Splenius capitis.
20. Digastric.
21. Levator proprius alæ nasi anterior, or dilatator naris anterior.
22. Levator proprius alæ nasi posterior, or dilatator naris posterior.
23. Platysma myoides.
24. Depressor alæ nasi.

over the masseter muscle, and passes horizontally forward to be inserted into the angle of the mouth, where it intermingles with

the orbicularis oris and depressor anguli oris. It produces the smile, not of good humor, but of derision.

Orbicularis Oris. This muscle, nearly an inch in breadth, surrounds the mouth, forming a kind of sphincter. Its size and thickness in different individuals produce the variety in the prominence of the lips. Observe that its fibres, except the most internal, do not surround the mouth in one unbroken series, but that those of the upper and lower lip decussate at the angles of the mouth, and intermingle with the fibres of the buccinator and other muscles which converge from different parts of the face.

The orbicularis consists of two parts, an inner or *labial* part, and an outer or *facial;* the difference in appearance of the fibres being very marked. The *labial* part consists of pale, thin fibres, forming more or less of the inner part of the orbicularis, and has no attachment to bone; the *facial* part is thinner but broader, and besides being connected with other muscles, is attached to bone thus: in the upper lip by two fasciculi on each side, one to the septum nasi, the other to the alveolar border opposite the incisor teeth; in the lower lip by a single fasciculus to the lower jaw on each side opposite the canine tooth. The cutaneous surface of the muscle is intimately connected with the lips and the surrounding skin; the deep surface is separated from the mucous membrane by the labial glands and the coronary vessels.

The orbicularis is the antagonist of all the muscles which move the lips. Upon a nice balance of their opposite actions depends the play and infinitely varied expression of the mouth.[1]

Depressor Anguli Oris. This muscle *arises* broadly from the oblique line of the lower jaw behind the foramen mentale, and is *inserted* narrowly into the angle of the mouth, intermingling with the zygomatici, the risorius, and orbicularis oris. It is an important muscle in the expression of sorrowful emotions. We see its action when children cry.

[1] In strong, muscular lips the upper part of the orbicularis sends a small subcutaneous slip of muscle from each side along the septum nasi nearly to the apex. The interval between the two slips corresponds to the furrow which leads from the nose to the lip. This is the *naso-labialis* or *depressor septi narium* of Haller and Albinus.

Depressor Labii Inferioris, or Quadratus Menti. This muscle *arises* from the oblique line of the lower jaw below the foramen mentale, and is *inserted* into the lower lip, its fibres intermingling with those of its fellow of the opposite side and the orbicularis. It covers the vessels and nerves which emerge from the foramen.

Levator Menti, or Levator Labii Inferioris. This muscle *arises* from the lower jaw, from the fossa below the incisor teeth, and passing down, is *inserted* into the skin of the chin. To see it, evert the lower lip and remove the mucous membrane on either side of the frænum. There are two of them, one for each side. Their action is well seen when we shave the chin, or protrude the lower lip.

Zygomaticus Major and Minor. The zygomaticus major *arises* from the outer surface of the malar bone close to its suture with the zygoma, passes obliquely downward and inward, and is *inserted* into the angle of the mouth, joining the depressor anguli and orbicularis oris.

The zygomaticus minor *arises* from the outer surface of the malar bone, in front of the preceding, and is *inserted* into the outer border of the levator labii superioris, near the corner of the mouth. The zygomaticus minor is often absent. The zygomaticus major is the muscle of laughing; the minor expresses sadness.

Before examining the orbicularis palpebrarum, notice the tendo oculi. To make the tendon more apparent, the tarsal cartilages should be drawn outward.

Tendo Oculi, or Palpebrarum. This tendon is a thin cord about two lines[1] in length and is readily felt at the inner angle of the eye by drawing the eyelids outward. It is fixed to the nasal process of the superior maxillary bone, in front of the lachrymal groove, passes horizontally outward, and divides into two diverging portions, one of which is attached to the upper, the other to the lower tarsal cartilage. The tendon crosses the lachrymal sac a little above the centre, and furnishes a tendinous expansion which covers the sac and is attached to the margin of the bony

[1] A line is the twelfth part of an inch.

groove which contains it. To see this expansion we must reflect that portion of the orbicularis palpebrarum which covers the sac.

In puncturing the lachrymal sac the knife is introduced below the tendon, in a direction downward, outward, and a little backward. We have to divide the skin, a few fibres of the orbicularis, and the fibrous expansion from the tendo palpebrarum. The angular artery and vein are situated on the inner side of the incision.

Orbicularis Palpebrarum. This thin, broad muscle surrounds the margin of the orbit and the eyelids, forming a sphincter. It is attached on the inner side to the tendo palpebrarum, to the nasal process of the superior maxillary bone, to the internal angular process of the frontal bone, and to the lower margin of the orbit. From this attachment the fibres form a series of oval curves, taking a wide sweep, and pass uninterruptedly round the eyelids and orbit.

The fibres which belong to the eyelids (orbicularis palpebrarum) are thin and pale, and form, over each eyelid, a series of elliptical curves which meet at the external canthus of the lids, and are loosely attached to the external tarsal ligament. The degree of their curvature becomes less as they approach the margin of the lids, so that some fibres proceed close to the lashes. This was first pointed out by Riolanus,[1] and described as the *musculus ciliaris*.[2] The fibres which spread over the orbital margins (*orbital portion*) are thicker and redder, and mingle, on the forehead, with the occipito-frontalis and corrugator supercilii, on the cheek, with the elevators of the upper lip and nose and the zygomaticus minor.

No fat is found on the eyelids; nothing intervenes between the skin and the muscles but loose connective tissue, that there may be no impediment to the free play of the lids.

The orbicular muscle not only closes the eyelids, but protects

[1] *Anthropologia*, lib. v, cap. 10.

[2] Strictly speaking, the musculus ciliaris arises from the two little divisions of the tendo oculi, and is inserted at the external canthus, into the fibrous tissue which unites the two tarsal cartilages.

the eye. When the eye is threatened, the muscle suddenly contracts, presses the eye back into the orbit, and contracts the skin of the brow and cheek so as to form a soft cushion in front of it. The cushion itself may be severely bruised, as is seen in a "black eye;" but the globe itself is rarely injured. When the eye is closed, as in winking, the palpebral portion of the muscle contracts. Observe this movement, and notice that the lids are drawn slightly inward as well as closed. The object of this inward motion is to direct the tears toward the inner angle of the eyelids, where they are absorbed by the puncta lachrymalia.

The tensor tarsi muscle is described in the dissection of the orbit.

Since the orbicular muscle is supplied by the facial nerve, it is affected in facial palsy, and the patient cannot close the lids.

Corrugator Supercilii. This *arises* from the inner end of the superciliary ridge of the frontal bone, and is *inserted* into the under surface of the orbicularis palpebrarum and occipito-frontalis. It lies concealed beneath these two muscles, and is the proper muscle of frowning. Its nerve is derived from the facial.

The present being a good opportunity to examine the appendages (*tutamina oculi*) of the eyes, postpone for the present the dissection of the remaining muscles of the face.

The Eyelids. The *eyelids* are two movable elliptical folds, consisting of strata of different tissues. The *upper lid* is large and more movable than the *lower*, so that when the eye is closed, it is mainly by this fold. The interval between the two lids is called the *fissura palpebrarum*, which terminates on the inner and outer sides in two angles, the *canthi*. The lids are thickest at their borders, are somewhat curved, and near the inner canthus each presents a slight elevation, the *papilla lachrymalis*, at the top of which is a small opening, the *punctum lachrymale;* this is the commencement of a small canal, *canaliculus*, which receives the tears and conveys them to the lachrymal sac, and thence through the nasal duct to the nose. At the inner canthus the two lids are separated by an oval space, the *lacus lachrymalis*, where the mucous membrane is raised into a rounded eminence, the *caruncula*.

Caruncula Lachrymalis. The *caruncula lachrymalis* is the red, rounded eminence situated at the inner canthus and formed by the conjunctiva. It is composed of an aggregation of sebaceous and sweat glands covered by mucous membrane; on the surface of it are minute hairs. Resting upon the eyeball, external to the caruncle, is a slight vertical, triangular fold of conjunctiva, *plica semilunaris*, which is the rudimentary membrana nictitans (the third eyelid found in birds). Both in the caruncle and plica semilunaris unstriped muscular tissue has been demonstrated.

The *conjunctiva* is the mucous membrane which covers the inner surface of the lids and the front of the eyeball. The portion lining the lids is termed the *palpebral;* that portion covering the front of the eye the *ocular*. The angle of its reflection from the lids to eyeball is called the *fornix conjunctivæ*, where are situated a number of racemose glands; there is also some lymphoid tissue found in other parts of the conjunctiva. The palpebral conjunctiva is more vascular than the ocular, and it presents a number of minute papillæ, which, when enlarged and aggregated by inflammation, give rise to the disease called "granular lids." The conjunctiva will be more fully described with the anatomy of the eye.

The *eyelashes* (cilia) are placed in two or more rows along the edges of the tarsal cartilages. The eyelashes of the upper lid are longer and more numerous than in the lower; and their convexity is directed downward, while those of the lower lid present an opposite curve. The bulbs of the lashes are situated between the tarsal cartilage and the fibres of the orbicularis palpebrarum. They are supplied with blood by the palpebral branches of the ophthalmic artery, which run parallel and close to the free borders of the lids beneath the orbicular muscle.

Structure of the Eyelids. The *eyelids* are composed of different tissues, arranged in successive strata, one beneath the other. They are—1. The *skin*; 2. The *orbicularis palpebrarum*; 3. The *palpebral ligament*, which extends from the margin of the orbit to the cartilage; 4. The expanded *tendon of the levator palpebræ* (in the upper lid only); 5. The *tarsal cartilage*; 6. A thin *layer of fascia*, in which are seen the blood-vessels; 7. The *Mei-*

bomian glands, which lie imbedded in the tarsal cartilage; 8. *Conjunctiva.* These structures are severally connected by areolar tissue, which never contains fat.

Such, in outline, is the structure of the eyelids. Their use is best described by Socrates, who, in answer to the question whether animals were made by chance or design, replies: "Think you not that it looks like the work of *forethought*, because the sight is delicate, to guard it with eyelids as with shutters, which open when we want to see, and shut during sleep; and, that even winds may not hurt them, to make eyelashes in the lids for a sieve; and to furnish the parts over the eyes with eyebrows, as with eaves, so that even the sweat from off the head may do them no mischief?" [1]

The *skin* of the eyelids is remarkably smooth and delicate, and destitute of fat. It is abundantly supplied with sensory nerves by branches of the fifth pair—namely, by the supra-orbital, supra-trochlear, infra-trochlear, lachrymal, and infra-orbital nerves.

The *orbicularis palpebrarum* has been already described (p. 45). It is supplied by the facial nerve.

The *levator palpebræ* arises from the lesser wing of the sphenoid above the optic foramen, gradually becomes broader, and terminates in a thin aponeurosis, which unites with the broad tarsal ligament, and is lost on the upper surface of the superior tarsal cartilage.

Tarsal Cartilages and Ligaments. These are plates of dense connective tissue, which support and give shape to the eyelids. There is one for each lid, and they are connected at the angles (*commissures* or *canthi*) of the lids through the medium of fibrous tissue. They can best be examined by everting the lids. Each cartilage resembles its lid in form. The upper is the larger, is broad in the middle, and gradually becomes narrower at either

[1] Xenophon's *Memorabilia*, b. I, c. vi, § 6: *οὐ δοκεῖ σοι καὶ τόδε προνοίας ἔργον ἐοικέναι, τό, ἐπεὶ ἀσθενὴς μέν ἐστιν ἡ ὄψις, βλεφάροις αὐτὴν θυρῶσαι, ἃ, ὅταν μὲν αὐτῇ χρῆσθαί τι δέῃ, ἀναπετάννυται, ἐν δὲ τῷ ὕπνῳ συγκλείεται; ὡς δ' ἂν μηδὲ ἄνεμοι βλάπτωσιν, ἡθμὸν βλεφαρίδας ἐμφῦσαι· ὀφρύσι τε ἀπωγεισῶσαι τὰ ὑπὲρ των ὀμμάτων, ὡς μηδ' ὁ ἐκ τῆς κεφαλῆς ἱδρὼς κακουργῇ.*

end. The lower is nearly of uniform breadth throughout. Both are thicker on the nasal than on the temporal side. They are connected to the margin of the orbit, and maintained in position by the *broad tarsal* or *palpebral ligament;* this is a continuation from the periosteum of the orbit to the tarsal cartilage, and is denser at the outer part of the orbit. There are two of them—upper and lower—and they pass to each cartilage respectively. When an abscess forms in the connective tissue of the lids, these ligaments prevent the matter from making its way into the orbit.

Each tarsal cartilage is attached on its outer side to the malar bone by the external tarsal ligament, and on its side to the nasal process of the superior maxillary by the tendo palpebrarum or the internal tarsal ligament.

The free or ciliary margin is straight, and is the thickest part of the tarsal cartilages. It is generally stated that the inner edge of each is sloped or beveled off; and that, when the lids are closed, there is formed, with the globe of the eye, a triangular channel. This channel is said to conduct the tears to the puncta lachrymalia. According to our observation, this channel does not exist; for when the lids are closed, their margins are in such accurate apposition, that not the slightest interspace can be discovered between them.

Puncta Lachrymalia.

The *puncta lachrymalia* are two pin-hole apertures, easily discovered on the margin of the lids, close to the inner angle. They are the orifices of the canals, called *canaliculi*, which pass inward, and convey the tears into the lachrymal sac. Observe that their orifices are directed backward. The upper canaliculus, the longer and narrower of the two, ascends for a short distance nearly vertically, and then dilating into a small pouch makes a sharp bend inward for about a quarter of an inch, to join the lachrymal sac; the lower canal descends perpendicularly, and, like the upper, makes a sharp bend, after which it pursues a direction upward and inward to the sac. The two canals open separately into the sac (sometimes by a common orifice). In facial palsy, the tensor tarsi being affected, the puncta lose their proper direction, and the tears flow over the cheek.

In the introduction of probes for the purpose of opening the contracted puncta, or of slitting up the lachrymal ducts, it is necessary to know the exact direction of these canals. (*See* diagram.) When from any cause the tears are secreted in greater quantity than usual, they overflow and trickle down the cheek.

Meibomian Glands. These long, compound, sebaceous glands, so called after the anatomist[1] who first described them, are situated on the under surface of each of the tarsal cartilages. In the upper lid there are between twenty and thirty; not quite so many in the lower. On everting the lid, they are seen running in longitudinal parallel rows, in grooves, in the cartilage. Under the microscope, each is seen to consist of a straight central tube, round the sides of which are a number of openings leading to short cæcal dilatations. The orifices of these glands are situated on the free margin of the lids behind the lashes. They are lined with flattened epithelial cells which, in the cæcal dilatations and ducts, become cubical and filled with fat. Their function is to secrete a sebaceous material, which prevents the lids from sticking together.

FIG. 14.



Tensor Tarsi. This muscle is only a deeper part of the orbicularis palpebrarum, and lies just behind the tendo palpebrarum. To expose it, cut perpendicularly through the middle of the upper and lower lids, and turn the inner halves toward the nose. After removing the mucous membrane, the muscle will be seen *arising* from the ridge of the lachrymal bone. It passes nearly horizontally outward, for about three lines, and then divides into two portions, which are *inserted* into the upper and lower tarsal cartilages, close to the orifices of the lachrymal ducts. It is probable that the tensor tarsi draws backward the open mouths of the ducts, so that they may receive the tears at

[1] H. Meibom, *De Vasis Palpebrarum novis.* Helmstedt, 1666.

the inner angle of the eye. It is supplied by a small branch from the facial nerve.

Let us now examine the muscles in connection with the nose, namely, the pyramidalis nasi, the compressor naris, the depressor alæ nasi, and the smaller intrinsic muscles of the nose. All are supplied by the facial nerve.

Pyramidalis Nasi. This is situated on the bridge of the nose, one on each side of the mesial line, and is usually regarded as a continuation of the inner part of the occipito-frontalis (p. 42). The two muscles diverge as they descend, and their fibres blend with those of the compressor naris. Their action produces transverse wrinkles of the skin at the root of the nose, as in the expression of an aggressive feeling.

Compressor Naris. This muscle is triangular, and *arises* by its apex from the inner side of the canine fossa of the superior maxilla, and is *attached* to a broad, thin aponeurosis, which spreads over the dorsum of the nose and joins its fellow. The origin of this muscle is concealed by the levator labii superioris alæque nasi.

When this muscle is reflected from its junction with its fellow, a small nerve is seen running down toward the tip of the nose. This nerve is the *superficial branch* of the nasal nerve (called also *naso-lobular*). It becomes subcutaneous between the nasal bone and the cartilage, and supplies the tip and lobule of the nose. It is joined by a branch of the facial nerve at its termination.

Depressor Alæ Nasi. This *arises* from the superior maxilla, above the second incisor tooth, and is *inserted* into the septum and ala of the nose. It is situated between the mucous membrane and the muscular structure of the upper lip; so that, to expose it, the upper lip must be everted, and the mucous membrane removed.

Besides the muscles above described, we find in connection with the cartilages of the alæ of the nose, pale, muscular fibres which have no very definite arrangement and require a lens for their detection. The *dilatator naris posterior* arises from the nasal process of the superior maxilla and the sesamoid cartilages, and is inserted into the skin of the margin of the nostril; the *dilatator*

naris anterior descends vertically from the cartilage of the aperture to its free margin. The action of these small muscles is to raise and evert the ala of the nose, and to counteract its tendency to be closed by atmospheric pressure. In dyspnœa, and in certain mental emotions, they contract with great energy.

Levator Labii Superioris Alæque Nasi. This *arises* from the nasal process of the superior maxillary bone near its orbital margin, and passing downward divides into two portions: an inner *inserted* into the side of the ala of the nose; an outer, into the upper lip, where its fibres blend with the orbicularis oris and levator labii superioris. It acts chiefly in expressing the smile of derision. Its habitual use occasions the deep furrow which, in most faces, runs from the ala of the nose to the corner of the mouth.

Levator Labii Superioris Proprius. This *arises* from the lower margin of the orbit, *i. e.*, from the superior maxilla and malar bone, above the infra-orbital foramen, and is *inserted* into the upper lip, where its fibres blend with the orbicularis oris. It is nearly an inch in breadth at its origin, which covers the infra-orbital vessels and nerves, and is itself overlapped by the orbicularis palpebrarum.

Levator Anguli Oris. This muscle, which is covered by the levator labii superioris, *arises* from the canine fossa of the superior maxilla, below the infra-orbital foramen, and is *inserted* into the angle of the mouth, superficial to the buccinator, its fibres blending with those of the orbicularis oris, the zygomatici, and the depressor anguli oris.

Buccinator. The buccinator *arises* from the outer surface of the alveolar borders of the upper and lower jaws corresponding to the molar teeth, and behind from the pterygo-maxillary ligament. The fibres pass forward and converge, to be inserted into the angle of the mouth and the muscular structure of the lips; the central fibres decussate, while the upper fibres pass to the upper lip, and the lower fibres pass to the lower lip. The muscle is covered on its inner aspect by the mucous membrane of the cheek, and on its outer by a thin fascia which passes backward, and is continuous with that covering the pharynx.

The buccinator is the principal muscle of the cheek. It forms,

with the superior constrictor of the pharynx, a continuous muscular wall for the side of the mouth and pharynx. The bond of connection between the buccinator and the superior constrictor is a tendinous band, the *pterygo-maxillary ligament.* This ligament (*see* diagram) extends from the hamular process vertically to the posterior extremity of the mylo-hyoid ridge of the lower jaw near the last molar tooth. It is simply a fibrous intersection between the two muscles.

The duct of the parotid gland pierces the buccinator obliquely, and opens into the mouth opposite the second molar tooth of the upper jaw.

The chief use of the buccinator is to keep the food between the teeth during mastication. It can also widen the mouth. Its power of expelling air from the mouth, as in whistling or playing on a wind instrument, has given rise to its peculiar name. It is supplied by the facial nerve, and is, therefore, affected in facial paralysis.

The buccinator is in relation, externally and behind, with a large amount of buccal fat, with the masseter and temporal muscles; in front with the risorius, the levator anguli oris, depressor anguli oris, the zygomatici, the duct of the parotid gland, the facial artery and vein, and the facial and buccal nerves; internally with the mucous membrane of the mouth and buccal glands; and posteriorly with the pterygo-maxillary ligament.

Buccal Fascia. The buccinator muscle is covered by a thin layer of fascia, which adheres closely to its surface, and is attached to the alveolar border of the upper and lower jaws. This structure is thin over the anterior part of the muscle, but more dense behind, where it is continuous with the aponeurosis of the pharynx. It is called the *bucco-pharyngeal fascia*, since it supports and strengthens the muscular walls of these cavities. In consequence of the density of this fascia, abscesses do not readily burst into the mouth or the pharynx.

Buccal and Molar Glands. The *buccal* glands, in structure compound racemose, like the salivary, are situated between the buccinator and the mucous membrane. They resemble the labial glands found beneath the mucous membrane of the lips, though

somewhat smaller. Three or four other glands, about the size of a little split pea, should be made out, as they lie between the masseter and buccinator: these are the *molar* glands. Their secretion, said to be mucous, is conveyed to the mouth by separate ducts near the last molar teeth.

FIG. 15.



MUSCLES OF THE PHARYNX.

Between the buccinator and the masseter, there is, in almost all subjects, an accumulation of fat. It is found, beneath the zygoma especially, in large round masses, and may be turned out with the handle of the scalpel. It helps to fill up the zygomatic fossa, and being soft and elastic, presents no obstacle to the free movements

of the jaw. Its absorption in emaciated individuals occasions the sinking of the cheek.

Facial Artery.

The *facial* (external maxillary) artery is the third branch of the external carotid. It ascends tortuously beneath the posterior belly of the digastricus and the stylo-hyoideus, next *through* or under the substance of the submaxillary gland; it then rests upon the mylo-hyoideus, and subsequently mounts over the base of the jaw at the *anterior edge of the masseter* muscle. This part of the course of the facial will be fully examined further on in the dissection of the neck. It now ascends tortuously near the corner of the mouth and the ala of the nose, toward the inner angle of the eye, where, much diminished in size, it inosculates with the terminal branch of the ophthalmic, a branch of the internal carotid. In the first part of its course on the face, the artery is covered by the platysma and the deep fascia; above the corner of the mouth it is crossed by a few fibres of the risorious and the zygomatici; still higher it is covered by some of the fibres of the elevator of the upper lip.[1] It lies successively upon the buccinator, levator anguli oris, and levator labii superioris alæque nasi muscles. In its course along the face it gives off the following branches:—

a. The *inferior labial* artery passes inward under the depressor anguli oris and inosculates with the mental branch of the inferior dental, the inferior coronary, and the submental arteries.

b. The *inferior coronary* artery comes off near the angle of the mouth, either directly from the facial, or in common with the superior coronary. It runs tortuously along the lower lip, beneath the depressor anguli oris; it then pierces the orbicularis, running between this muscle and the mucous membrane of the lip. It inosculates largely with its fellow, and the inferior labial and the mental arteries.

c. The *superior coronary*, larger than the preceding, is given off beneath the zygomatici. It proceeds along the upper lip close to the mucous membrane, and inosculates with its fellow; thus is formed round the mouth a complete **arterial circle**, which can be felt pulsating on the inner side of the lip, near the free border.

[1] Not infrequently the artery lies superficial to this muscle.

From this circle numerous branches pass off to the papillæ of the lips, and the labial glands. The superior coronary gives off a

FIG. 16.



BRANCHES OF THE EXTERNAL CAROTID ARTERY.

1. External Carotid.
2. Lingual.
3. Facial.
4. Inferior labial.
5. Inferior coronary.
6. Superior coronary.
7. Lateral nasal.
8. Angular.
9. Superior thyroid.
10. and 16. Occipital.
11. Posterior auricular.
12. Anterior auricular.
13. Internal maxillary.
14. Transverse facial.
15. Middle temporal.
17. Anterior temporal.
18. Posterior temporal.
19. Supra-orbital.
20. Frontal.

branch, *the artery of the septum*, which ascends along the septum to the apex of the nose; also a small one to the ala nasi.

d. The *lateral artery of the nose*, a branch of considerable size, arises opposite the ala nasi, ramifies upon the external surface of

the nose, and inosculates with the nasal branch of the ophthalmic artery, the infra-orbital, and the artery of the septum.

e. The *angular artery*, which may be regarded as the termination of the facial, inosculates on the inner side of the tendo palpebrarum with the nasal branch of the ophthalmic artery.

The facial artery supplies numerous branches to the muscles of the face, and inosculates with the transversalis faciei, infra-orbital, the mental, the sublingual branch of the lingual, the nasal branches of the internal maxillary and the ophthalmic, the ascending pharyngeal and descending palatine arteries.

The facial artery and its branches are surrounded by a minute plexus of nerves (nervi molles), invisible to the naked eye. They are derived from the superior cervical ganglion of the sympathetic, and exert a powerful influence over the contraction and dilatation of the capillary vessels, and thus occasion those sudden changes in the countenance indicative of certain mental emotions, *e. g.*, blushing or sudden paleness.[1]

The *facial vein* does not run with the artery, but takes a straight course from the inner angle of the eye to the anterior border of the masseter. In this course it descends upon the levator labii superioris, then passes beneath the zygomatic muscles, over the termination of the parotid duct, and at the anterior border of the masseter passes over the jaw, behind the facial artery, and joins the internal jugular.

The facial vein is a continuation of the frontal, which descends over the forehead, and after receiving the supra-orbital, takes the name of *angular* at the corner of the eye. It communicates with the ophthalmic vein, receives the veins of the eyelids, the external part of the nose, the coronary veins, and others from the muscles of the face. Near the angle of the mouth it is increased in size by a communicating branch from the infra-orbital vein, and by a large vein which comes from the temporo-maxillary vein. The other veins which empty

[1] MM. Bernard and Brown-Séquard have proved by experiment, that if the branches of the sympathetic which accompany the facial artery be divided, the capillary vessels of the face, being deprived of their contractile power, become immediately distended with blood, and the temperature of the face is raised.

themselves into the facial correspond with the branches given off from the facial artery.

Arteria Transversalis Faciei. This artery arises from the temporal, or occasionally from the external carotid in the substance of the parotid gland. It runs forward across the masseter between the parotid duct and the zygoma, and is distributed to the glandula socia parotidis and the masseter. It anastomoses with the infra-orbital, buccal and facial. It is seldom of large size, except when it supplies those parts which usually receive blood from the facial. We have seen it as large as a goose-quill, furnishing the coronary and the nasal arteries; the facial itself not being larger than a sewing thread.

The parotid gland is now to be examined. Its boundaries, its deep relations, the course of its duct, and the objects contained within the gland, must be carefully observed.

Parotid Gland. The parotid,[1] the largest of the salivary glands, occupies the space between the ramus of the jaw and the mastoid process, and weighs between five and eight drachms. It is bounded above by the zygoma; below, by the sterno-mastoid and digastric muscles; behind, by the meatus auditorius externus and the mastoid process; in front, it lies over the ascending ramus of the jaw, and is prolonged for some distance over the masseter. It is separated from the submaxillary gland by the stylo-maxillary ligament; sometimes the two glands are directly continuous.

The superficial surface of the gland is flat, and covered by a strong layer of fascia, a continuation of the cervical, and has one or two lymphatic glands lying on it. It not only surrounds the gland, but sends down numerous partitions which form a framework for its lobes. The density of this sheath explains the pain caused by inflammation of the gland, the tardiness with which abscesses within it make their way to the surface, and the propriety of an early opening.

The deep surface of the gland is irregular, and moulded upon the subjacent parts. Thus it sends a prolongation which passes inward between the neck of the jaw and the internal lateral liga-

[1] From παρά, near; οὖς, the ear.

ment; another process which passes in front of the styloid process, and extends upward and occupies the posterior part of the glenoid cavity; a third process passes behind the styloid process, below the mastoid process and behind the sterno-mastoid muscle, and sometimes penetrates deep enough to be in contact with the internal jugular vein.

The internal carotid artery and internal jugular vein are in contact with the gland behind.

On carefully removing the substance of the parotid gland, the following structures are seen in its interior, proceeding in the order of their depth from the surface:—

1. Two or more small lymphatic glands.

2. The pes anserinus, or primary branches of the facial nerve, which emerge at its anterior border.

3. Branches from the great auricular and auriculo-temporal nerves, which communicate in its substance with the facial nerve.

4. The external jugular vein, formed by the junction of the internal maxillary and temporal veins.

5. The external carotid artery, which, after distributing many branches to the gland, divides, opposite the neck of the jaw, into the internal maxillary and temporal; the latter giving off in the gland the posterior auricular and transverse facial arteries.[1]

That portion of the gland which lies on the masseter muscle is called *glandula socia parotidis*. It varies in size in different subjects; and is situated chiefly above the parotid duct, into which it pours its secretion by one or two smaller ducts.

The *duct* of the parotid gland (ductus Stenonis[2]), about two inches and a half long, is very thick and strong. In this respect it differs from the duct of the submaxillary gland, which is less

[1] Reviewing the intimate and deep connections of the parotid gland, one cannot but conclude that it is almost impracticable to remove it entirely during life. If this conclusion be correct, even in the normal condition of the gland, what must it be when the gland is enlarged by disease? John Bell, however, relates a case in which he was induced to attempt the extirpation of a diseased parotid (*Principles of Surgery*, vol. iii, p. 262). Other surgeons, too, of more modern date, have attempted the same thing. It is not unlikely that they have mistaken a tumor in the substance of the parotid for disease of the parotid itself.

[2] Nic. Steno, *De Glandulis Oris*, etc. Bat. 1661.

exposed to injury. It runs transversely forward over the masseter, about an inch below the zygoma, through the fat of the cheek, then perforates the buccinator obliquely, and opens into the mouth, opposite the second molar tooth of the upper jaw. Near its termination it is crossed by the zygomaticus major and the facial vein. After perforating the buccinator, the duct passes for a short distance between the muscle and the mucous membrane. Its orifice is small and contracted, compared with the diameter of the rest of the duct, which will admit a crow-quill; it is not easily found in the mouth, being concealed by a fold of mucous membrane.

The direction of the parotid duct corresponds with a line drawn from the middle of the lobule of the ear to a point midway between the nose and the mouth.

The blood supply of the parotid is derived from the external carotid and its branches, which are accompanied by their respective veins. Its nerves are supplied from the sympathetic plexus around the external carotid, the auriculo-temporal, the great auricular and the facial nerves.

The *lymphatic* glands about the parotid deserve notice, since they are liable to become enlarged, and simulate disease of the parotid itself. A lymphatic gland lies close to the root of the zygoma, in front of the cartilage of the ear; this gland is sometimes affected in disease of the external tunics of the eye; *e.g.*, in purulent ophthalmia; also in affections of the scalp.

The parotid belongs to the compound racemose form of glands. Tracing its main duct into the substance of the gland, we find that it divides into smaller ones, which again divide into the smaller ramuscules which open into cæcal dilatations called *alveoli*. Each alveolus, about $\frac{1}{150}$ of an inch in diameter, has a more or less defined basement membrane, upon which the cells rest. The cells are flattened and spheroidal, enclosing nuclei, some of them having outstanding processes from the bases of the cells. An aggregation of these alveoli forms a small lobule, from which a small excretory duct proceeds; these lobules are united by intervening connective tissue, which is a continuation inward of the dense fascia covering the gland. The connective tissue varies

much in thickness in different situations, and where it is most abundant it is distinctly lamellar (Klein),[1] and contains numerous granular plasma cells and lymphoid tissue. The small ramuscules have only a basement membrane with flattened cells, which change in the smaller ducts to a columnar form, while in the larger ducts the epithelium is squamous.

The parotid gland secretes an alkaline watery fluid, containing solids which amount to 6 parts in 1000. The solids consist of ptyalin, a special ferment, mucus, and certain salts, chiefly sulpho-cyanide of potassium, chlorides and phosphates of potassium, sodium, lime and magnesia. Apart from its mechanical properties, the parotid secretion has the power, by means of its ptyalin, of converting starch into dextrine and grape sugar.

To display the plexus of nerves (pes anserinus), formed by the branches of the facial, cut into the parotid gland by a vertical incision until the main trunk of the nerve is reached.

Portio Dura, or Facial Nerve. This is the seventh cranial nerve, and is the motor nerve of the face. It supplies all the muscles of expression, the platysma, and the buccinator. Through some of its branches it supplies other muscles, the description of which will be deferred till the facial nerve is dissected in the temporal bone. It arises immediately below the pons Varolii, from the lateral tract of the medulla oblongata, between the olivary and restiform bodies. The nerve enters the meatus auditorius internus, lying upon the auditory nerve, traverses a tortuous bony canal (*aqueductus Fallopii*) in the petrous portion of the temporal bone, and leaves the skull at the stylo-mastoid foramen. Its course and connections in the temporal bone will be studied hereafter; at present we must trace the facial part of the nerve.

Having emerged from the stylo-mastoid foramen, the nerve enters the parotid gland, and divides behind the ramus of the jaw into two primary branches, named, from their distribution, *temporo-facial* and *cervico-facial*. These primary branches cross over the external carotid artery and the external jugular vein, and form, by their communications within the substance of the parotid, the

[1] Klein, *Quarterly Journal of Microscopical Science*, 1882.

plexus called *pes anserinus*, from its fancied resemblance to the skeleton of a goose's foot. (Diagram, p. 63.)

Close to the stylo-mastoid foramen, the facial nerve gives off its *posterior auricular* branch (p. 22), which ascends behind the ear and divides into two, an *auricular* and an *occipital;* the former supplies the retrahens and attollens aurem, the latter the posterior belly of the occipito-frontalis. This branch communicates with the deep branch of the great auricular n., with the small occipital, and with the auricular branch of the pneumogastric. Its two next *branches* supply the stylo-hyoideus and the posterior belly of the digastricus. The digastric nerve enters the muscle by many filaments; the nerve to the stylo-hyoid is long, and enters the muscle about the middle. The stylo-hyoid branch communicates with the sympathetic on the external carotid a.; the digastric branch with the glosso-pharyngeal near the base of the skull. These two muscular nerves are frequently given off from a common branch.

The *temporo-facial* division, the larger of the two, in passing through the parotid gland, crosses the external carotid and the neck of the jaw, receives two or more communications from the auriculo-temporal (branch of the fifth) and subdivides into temporal, malar, and infra-orbital branches.

The *temporal* branches ascend over the zygoma, supply the frontalis, the attrahens aurem, the orbicularis palpebrarum, the corrugator supercilii, and tensor tarsi, and communicate with filaments of the supra-orbital nerve, with the temporal branch of the superior maxillary n., with the auriculo-temporal n., and with the lachrymal n.

The *malar* branches cross the malar bone, supply the orbicular muscle, and communicate with filaments of the lachrymal, the supra-orbital, the superior maxillary, and the malar branch of the superior maxillary.

The *infra-orbital* branches are the largest, and proceed transversely forward over the masseter, beneath the zygomatici, to supply the orbicularis oris, the elevators of the upper lip, and the muscles of the nose. The superficial branches join with the nasal and infra-trochlear branches of the ophthalmic along the side of

the nose; the deep branches communicate beneath the levator labii superioris with the infra-orbital branches of the superior maxillary nerve forming the *infra-orbital plexus*, and also with the buccal branches of the facial.

The *cervico-facial* division, joined in the parotid gland by filaments from the great auricular (branch of the cervical plexus), descends toward the angle of the jaw, and subdivides into buccal, supra- and infra-maxillary branches.

FIG. 17.

DIAGRAM OF THE BRANCHES OF THE FACIAL NERVE.

1. Branch to occipito-frontalis.
2. Posterior auricular.
3. Temporal branches.
4. Malar branches.
5. Infra-orbital.
6. Buccal.
7. Supra-maxillary.
8. Infra-maxillary.

The *buccal* branches pass forward over the masseter, parallel with the parotid duct, and supply the buccinator; they communicate with the buccal branch of the inferior maxillary nerve (third division of the fifth), and with the infra-orbital nerve.

The *supra-maxillary* branches advance over the masseter and facial artery, and run under the platysma and the depressor muscles of the lower lip, all of which they supply. Some of the filaments communicate with the mental branch of the inferior dental nerve.

The *infra-maxillary* or cervical branches, one or more in number, arch forward below the jaw, covered by the platysma, as low as the hyoid bone, and communicate with the superficial cervical (branch of the cervical plexus).

Sensory Nerves of the Face. These are the supra-orbital, the supra- and infra-trochlear, the naso-lobular, the temporo-malar, the infra-orbital, and the mental, all branches of the fifth pair.

FIG. 18.

DIAGRAM OF THE SENSORY NERVES OF THE SCALP AND FACE.

1. Great occipital.
2. Small occipital.
3. Auricular br. of the pneumogastric.
4. Great auricular.
5. Auriculo-temporal.
6. Temporal br. of superior maxillary nerve.
7. Supra-orbital.
8. Supra-trochlear.
9. Malar br. of superior maxillary nerve.
10. Infra-trochlear.
11. Naso-lobular.
12. Infra-orbital.
13. Buccal br. of inferior maxillary nerve.
14. Mental.

The *supra-orbital nerve* is the continuation of the frontal, which is a branch of the first division of the fifth pair. It leaves the orbit through the supra-orbital notch and ascends upon the forehead, at first covered by the orbicularis and occipito-frontalis. It presently divides into two sets of branches—an *outer*, the larger, which passes backward as far as the occipital bone, and an *inner*, which ascends as far as the parietal bone. It distributes sensory

muscular branches also to the orbicularis palpebrarum, corrugator supercilii, the occipito-frontalis, to the pericranium and branches which supply the skin of the forehead, upper eyelid and scalp. It communicates with the facial nerve on the forehead. The supra-orbital artery is a branch of the ophthalmic.

The *supra-trochlear n.*, or internal frontal, appears at the inner angle of the orbit between the supra-orbital foramen and the pulley of the superior oblique, and sends down in front of the pulley a loop to communicate with the infra-trochlear branch of the nasal. The main trunk of the nerve ascends to the forehead. Its further course has been described (p. 21).

The *infra-trochlear n.* issues from the orbit below the pulley, and supplies branches to the eyelids, the conjunctiva, lachrymal sac, and the side of the nose.

The *infra-orbital nerve* is the terminal branch of the superior maxillary or second division of the fifth nerve. It emerges with its artery from the infra-orbital foramen, covered by the levator labii superioris. The nerve immediately divides into several branches, *palpebral*, *nasal*, and *labial*; the palpebral, ascending beneath the orbicularis, supply the lower eyelid, and communicate with the facial and the malar branch of the orbital nerve; the nasal pass inward to supply the nose, and join the nasal branch (naso-lobular) of the ophthalmic; the labial, by far the most numerous, descend into the upper lip, beneath the levator labii superioris, and eventually terminate in lashes of filaments, which endow the papillæ of the lip and the mucous membrane of the mouth with exquisite sensibility. Close to the infra-orbital foramen is the infra-orbital plexus, before alluded to (p. 63).

The *infra-orbital artery* is the terminal branch of the internal maxillary; it supplies the muscles, the skin, and the front teeth of the upper jaw, and inosculates with the transverse facial, buccal, facial and coronary arteries.

The *naso-lobular nerve* is the external branch of the nasal nerve, and is distributed to the tip and lobule of the nose, and is joined by filaments from the facial nerve.

The *temporal branch* of the *orbital nerve* (branch of the superior maxillary nerve, running along the outer wall of the orbit, and

which divides into a temporal and a malar branch) issues through the temporal fascia about a finger's breadth above the zygoma, and supplies the skin of the temple. It communicates with the facial and the auriculo-temporal nerves.

The *malar nerve*, a branch also of the orbital nerve, issues through a foramen in the malar bone, and, after piercing the orbicularis palpebrarum, supplies the skin of the cheek over the malar bone. It communicates with the facial and the palpebral branches of the infra-orbital nerve.

The *mental nerve* is a branch of the inferior maxillary or third division of the fifth. It emerges from the mental foramen in the lower jaw, in a direction upward and backward, beneath the depressor anguli oris. It soon divides into a number of branches beneath the depressor labii inferioris, some of which supply the skin of the chin, but the greater number terminate in the papillæ of the lower lip. It communicates with the facial nerve.

The *mental artery* is a branch of the inferior dental. It supplies the gums and the chin, and inosculates with the sub-mental, the inferior labial and inferior coronary arteries.

Dissection. To expose the contents of the orbit, remove that portion of the orbital plate which forms the roof of the orbit, as far back as the optic foramen, making one section, with a saw, on the outer side, the other on the inner side of the roof, so that the two sections converge at the optic foramen. In doing this, be careful not to injure the little pulley on the inner side for the superior oblique. If the bone be sufficiently sawn through, a gentle tap with the saw on the front of the orbital plate will fracture its thin wall transversely. The anterior fourth of the roof should be turned forward and downward, and kept in this position by hooks; the remainder is to be removed by bone forceps nearly as far as the optic foramen, so as to leave a ring of bone from which most of the ocular muscles have their origin. The eyeball should be made tense by blowing air through a blowpipe passed well into the globe through the cranial end of the optic nerve.

Periosteum of the Orbit. The roof being removed we expose the fibrous membrane which lines the walls of the orbit. It

is a continuation of the dura mater through the sphenoidal fissure. Traced forward, we find that it is loosely connected to the walls of the orbit, and that at the margin of the orbit it divides into two layers, one of which is continuous with the periosteum of the forehead, the other forms the broad tarsal ligament which fixes the tarsal cartilage.

The periosteum is now to be removed, and the fascia of the orbit made out. The following objects should then be carefully traced: in the middle are seen the *frontal artery and nerve*, lying on the levator palpebræ; on the outer side, the *lachrymal nerve*

FIG. 19.



DIAGRAM OF THE NERVES OF THE ORBIT.

and artery pass forward on the external rectus to the lachrymal gland, which lies under cover of the external angular process; on the inner side is the *fourth nerve*, lying on and supplying the superior oblique.

Fascia of the Orbit and Capsule of Tenon.

The fascia of the orbit provides the lachrymal gland and each of the muscles with a loose sheath, thin and delicate at the back of the orbit, but stronger near the eyeball. It is pierced behind by the optic nerve and by the arteries and nerves of the orbit, while in front it is connected with the ocular conjunctiva close to the cornea. The sheaths are firmly adherent to the muscles, and their tendinous

insertions into the globe are connected by the fascia. From the insertions of the muscles it is reflected as a double layer backward over the globe, so that it resembles a serous membranous sac—a tunica vaginalis—one layer being loosely connected with the globe, the other lining the fat in which the globe is set. These layers are lined with epithelium, and are separated by an external lymph space. This reflection of the orbital fascia is called the *capsule of Tenon*, its use being to allow free movement of the globe.

The orbit contains a large quantity of granular fat, which forms a soft bed for the eye, and prevents its being retracted too far by its muscles. Upon the amount of this fat depends, in some measure, the prominence of the eyes. Its absorption in disease or old age occasions the sinking of the eyeballs.

Contents of the Orbit. In the middle of the orbit is the eyeball, surrounded by more or less fat, and attached to it are six muscles which move it; four, running forward in a straight direction, are called the *recti*, and are arranged one above, one below, and one on each side of the globe; the remaining two are called, from their direction, *obliqui*, one superior, the other inferior. There is also a muscle to raise the upper eyelid, termed *levator palpebræ*. The *nerves* are: the optic, which passes through the optic foramen; the third, the fourth, the first division of the fifth, the sixth, and some filaments of the sympathetic, all of which pass through the sphenoidal fissure. The third supplies all the muscles with motor power, except the superior oblique, which is supplied by the fourth, and the external rectus, which is supplied by the sixth. The first or ophthalmic division of the fifth divides into a frontal, lachrymal and nasal branch. The ophthalmic artery, a branch of the internal carotid, passes into the orbit through the optic foramen; its vein passes backward through the sphenoidal fissure to join the cavernous sinus.

Frontal Nerve. The *ophthalmic*, or first division of the fifth—a sensory nerve—after giving off from its inner and lower side, whilst within the cavernous sinus, the nasal nerve, divides into the frontal and lachrymal nerves, of which the former is the larger. It is the smallest division of the fifth, and runs for-

ward for the distance of about an inch; in its course it is connected with the cavernous plexus of the sympathetic, with the third, fourth and sixth nerves, and close to its origin from the Gasserian ganglion it sends off a small recurrent branch to the tentorium cerebelli. One of its divisions, the frontal nerve, runs forward upon the upper surface of the levator palpebræ, on which, about midway in the orbit, it divides into two branches—the supra-trochlear and the supra-orbital.

a. The *supra-trochlear*, the smaller of the two (Fig. 20) runs obliquely inward above the pulley of the superior oblique to the

FIG. 20.



VIEW OF ORBIT FROM ABOVE.

inner angle of the orbit. Here it gives off a small communication downward to the infra-trochlear branch of the nasal, and then divides, after passing between the bone and the orbicularis palpebrarum, into filaments which supply the skin of the upper eyelid, forehead and nose. One or two small filaments may be traced through the bone to the mucous membrane of the frontal sinuses.[1]

b. The *supra-orbital* is the continuation of the frontal nerve, and runs forward on the levator palpebræ to the supra-orbital notch, through which it ascends to supply the skin of the upper eyelid,

[1] These filaments have been described by Blumenbach, *De Sinibus Frontalibus.*

forehead, pericranium and scalp. Its cutaneous branches, an inner and an outer, which run upward beneath the occipito-frontalis, have been described in the dissection of the scalp (p. 22). It supplies with common sensation the orbicularis palpebrarum, the occipito-frontalis, and the corrugator supercilii, where it joins the facial nerve.

Lachrymal Nerve. This is the smallest of the three branches of the ophthalmic nerve. It runs along the upper border of the external rectus, on the outer side of the orbit, with the lachrymal artery, through the lachrymal gland, which it supplies, as well as the upper eyelid. Its branches within the orbit are: 1, a branch which passes down behind the lachrymal gland to communicate with the orbital branch of the superior maxillary nerve; 2, filaments to the lachrymal gland. It then pierces the palpebral ligament, to supply the skin of the upper eyelid.

Fourth Cranial Nerve. This nerve enters the orbit through the sphenoidal fissure above the other nerves. It runs along the inner side of the frontal nerve, and enters the upper or orbital surface of the superior oblique, to which it is solely distributed. This nerve is joined in the outer wall of the cavernous sinus by filaments from the sympathetic. It communicates occasionally with the lachrymal, and the ophthalmic division of the fifth. Here also it sends backward two or more filaments to supply the tentorium cerebelli.

Lachrymal Gland. This gland is situated below the external angular process of the frontal bone. It is about the size and shape of an almond. Its upper surface is convex, in adaptation to the roof the orbit; its lower is concave, in adaptation to the eyeball, and rests upon the external and superior rectus. The anterior part of the gland lies sometimes separated from the rest, close to the back part of the upper eyelid, and is covered by the conjunctiva. The gland is invested by a capsule[1] formed by the fascia of the orbit.

[1] This capsule, being a little stronger on the under surface of the gland, is described and figured by Sömmering as a distinct ligament, *Icones Oculi Humani*, tab. vii.

The lachrymal gland consists of an aggregation of small lobes composed of smaller lobules, connected by fibro-cellular tissue, and resembles the structure of the salivary glands. The excretory ducts, seven to ten in number, run parallel, and perforate the conjunctiva in a row, about a quarter of an inch above the edge of the tarsal cartilage (Fig. 21). They are not easily discovered in the human eye; in that of the horse or bullock they are large enough to admit a small probe. The secretion of the gland keeps the surface of the cornea constantly moist and polished; but if dust or any foreign substance irritate the eye, the tears flow in abundance and wash it off.

FIG. 21.



All the muscles of the orbit, with the exception of the inferior oblique, arise from the margin of the foramen opticum, and pass forward, like ribands, to their insertions.

Levator Palpebræ. This muscle *arises* from the roof of the orbit, above and in front of the optic foramen. It gradually increases in breadth, and terminates in a broad, thin aponeurosis, which is *inserted* into the upper surface of the tarsal cartilage beneath the palpebral ligament. It is constantly in action when the eyes are open, in order to counteract the tendency of the lids to fall. As sleep approaches, the muscle relaxes, the eyes feel heavy, and the lids close. Its nerve comes from the superior division of the third nerve, and enters it on its under or ocular aspect.

Obliquus Superior. This muscle *arises* from the inner side of the optic foramen. It runs forward along the inner and upper side of the orbit, and terminates in a round tendon, which passes through a fibro-cartilaginous pulley—*trochlea*—attached to the trochlear fossa in the frontal bone. From the pulley the tendon is reflected outward and backward, beneath the superior rectus, and is *inserted* by an expanded tendon into the outer part of the sclerotic coat, midway between the cornea

and the entrance of the optic nerve. The pulley is lined by a synovial membrane, which is continued over the tendon. The *action* of this muscle will be considered with that of the inferior oblique. It is supplied by the fourth nerve, which enters the back part of its upper surface.

The frontal nerve and levator palpebræ are now to be cut through the middle and reflected, the front part forward and the hind part backward. On its under aspect is seen the twig from the upper division of the third nerve entering it. On reflecting this muscle the superior rectus is exposed.

The *superior rectus arises* by a tendinous origin from the upper margin of the optic foramen and from the sheath of the optic nerve, and is *inserted* by a broad, thin tendon into the sclerotic coat, about a quarter of an inch from the margin of the cornea (p. 78).

Dissection. Reflect this muscle by cutting through the middle, and, in doing so, observe a filament from the third nerve entering its under aspect. After the removal of a quantity of granular fat, the following objects are exposed: beneath the muscle are the optic nerve, the ophthalmic artery and vein, the nasal nerve and its ciliary branches crossing over the optic nerve, and further forward is the reflected tendon of the superior oblique; on the outer side of the optic nerve, and close to the ophthalmic artery, is the lenticular ganglion, with numerous ciliary filaments passing forward from it to enter the sclerotic. The student should now trace backward the two roots which enter the upper and lower angle respectively of the ganglion, the upper being its sensory branch from the nasal, the lower its motor root from the lower division of the third nerve. Further back should be traced the third, the nasal branch of the ophthalmic, and the sixth nerves, passing between the two heads of the external rectus to their respective destinations. The ophthalmic artery and its branches may also at this stage be exposed and cleaned.

Nasal Nerve. This is one of the three divisions of the ophthalmic branch of the fifth, and is usually the first branch given off (Fig. 19, p. 67). It enters the orbit through the sphenoidal fissure, between the two origins of the external

rectus, and between the two divisions of the third n. It then crosses obliquely over the optic nerve, beneath the levator palpebræ and the superior rectus, toward the inner wall of the orbit. After giving off the *infra-trochlear* branch, the nerve passes out of the orbit between the superior oblique and internal rectus, through the anterior ethmoidal foramen, into the cranium, where it lies beneath the dura mater, upon the cribriform plate of the ethmoid bone. It soon leaves the cranium through the nasal slit near the crista galli, and enters the nose. Here it divides into two branches—an *inner* or *septal*, which supplies the mucous membrane of the front of the septum; and an *outer*, the main continuation of the nerve—which runs in a groove on the under surface of the nasal bone, and distributes branches to the pituitary membrane of the outer part of the nose and the two lower turbinated bones; it also gives off a superficial branch, which emerges between the nasal bone and the cartilage, under the name of the *naso-lobular*, and is distributed to the skin of the tip and ala of the nose (p. 65).

The nasal nerve gives off the following branches in the orbit:—

a. One slender filament to the *lenticular ganglion* (forming its upper or long root) is given off from the nasal nerve as it passes between the heads of the external rectus close to the optic nerve. It is about half an inch long, and enters the posterior-superior angle of the ganglion.

b. Two or three *long ciliary nerves*. They run along the inner side of the optic nerve to the back of the globe of the eye. They are joined by filaments from the lenticular ganglion, and pass through the sclerotic coat to supply the iris (Fig. 22).

c. Infra-trochlear nerve.—This runs forward, along the inner side of the orbit, below the pulley of the superior oblique, where it communicates with the supra-trochlear branch of the frontal nerve. It passes to the inner angle of the orbit, and divides into filaments, which supply the skin of the eyelids, the caruncle, the lachrymal sac, and the side of the nose.

Optic Nerve.

This nerve, having passed through the optic foramen, proceeds forward and a little outward, for about an inch, to the globe of the eye, which it enters on the

nasal side of its axis. It pierces the sclerotic and choroid coats, and then expands to form the retina. The nerve is invested by a dense fibrous coat derived from the dura mater, and by a thin one from the arachnoid, both of which pass forward as far as the sclerotic. At the optic foramen it is surrounded by the tendinous origins of the recti; in the rest of its course, by loose fat and by the ciliary nerves and arteries. It is pierced in its course through the orbit by the arteria centralis retinæ, which runs along with its vein in the middle of the nerve, to the eyeball.[1]

FIG. 22.



VIEW OF OPTIC AND LOWER NERVES OF ORBIT.

Ophthalmic Artery. This artery arises from the internal carotid, close by the anterior clinoid process. It enters the orbit through the optic foramen, outside and below the optic nerve; occasionally through the sphenoidal fissure. Its course in the orbit is remarkably tortuous. Situated at first on the outer side of the optic nerve, it soon crosses over it, and runs along the inner side of the orbit between the superior and internal recti, to inosculate with the internal angular artery (the terminal branch of the facial). Its branches arise in the following order:—

a. Lachrymal artery.—This branch proceeds along the outer

[1] A small branch from Meckel's ganglion, ascending through the spheno-maxillary fissure, is described by Arnold as joining the optic nerve.

wall of the orbit, above the external rectus, in company with the nerve of the same name, to the lachrymal gland. After supplying the gland, it terminates in the conjunctiva and eyelids. In the orbit it gives off some *malar* branches which pierce the malar bone to get to the temporal fossa, and anastomose with the deep temporal arteries. It also sends a branch backward through the sphenoidal fissure, to anastomose with the arteria meningea media.

b. Supra-orbital artery.—This branch runs forward with the frontal nerve, under the roof of the orbit and upon the levator palpebræ. It emerges on the forehead through the supra-orbital foramen, where it communicates with the superficial temporal, frontal and angular arteries.

c. Arteria centralis retinæ.—This small branch enters the optic nerve obliquely on the outer aspect, close to the optic foramen. It runs in the centre of this nerve to the interior of the eye.

d. Ciliary arteries.—These branches may be arranged in three groups. The *short ciliary*, twelve to fifteen in number, proceed tortuously forward with the optic nerve, and pierce the sclerotic coat at the back of the eye, to supply the choroid coat and the iris. The *long ciliary*, two in number, run on each side of the optic nerve, enter the sclerotic, and pass horizontally forward, one on each side of the globe, between the sclerotic and the choroid, nearly as far as the iris, where each divides into an upper and a lower branch. These branches of the two long ciliary arteries anastomose with the anterior ciliary and form two vascular circles, an outer at the circumference of the iris, the *circulus major*, and an inner at the free margin of the iris, the *circulus minor*. The *anterior ciliary* are branches of the muscular and lachrymal arteries, and proceed with the tendons of the recti, and enter the front part of the sclerotic coat. In inflammation of the iris the vascular zone round the cornea arises from enlargement and congestion of the anterior ciliary arteries.

e. Ethmoidal arteries.—Of these arteries, two in number, the *anterior* and larger passes through the anterior ethmoidal foramen with the nasal nerve; the *posterior* enters the posterior ethmoidal foramen with the spheno-ethmoidal nerve. The anterior gives

off branches to the frontal and anterior ethmoidal cells, and a nasal branch to the nose; it likewise gives off an anterior meningeal branch to the dura mater in the anterior fossa. The posterior is distributed to the posterior ethmoidal cells and upper part of the nose.

f. Muscular branches.—There is an upper and a lower branch, supplying respectively the upper and lower muscles; besides these, there are irregular branches, from the lachrymal and supra-orbital arteries.

g. Palpebral arteries.—These branches, a *superior* and an *inferior*, proceed from the ophthalmic artery near the front of the orbit. They are distributed to their respective eyelids, forming arches near the margins of the lids between the tarsal cartilages and the orbicularis palpebrarum, with branches from the lachrymal and the infra-orbital arteries.

h. Nasal artery.—This branch may be considered one of the terminal divisions of the ophthalmic. It leaves the orbit on the nasal side of the eye, above the tendon of the orbicularis, and inosculates with the angular and nasal arteries of the facial. It supplies the side of the nose and the lachrymal sac.

i. Frontal artery.—This is the other terminal branch of the ophthalmic. It emerges at the inner angle of the eye, ascends, and inosculates with the supra-orbital artery.

Ophthalmic Veins. There are two ophthalmic veins. The *superior* commences at the inner angle of the eye by a communication with the frontal and angular veins. It runs backward above the optic nerve, in a straighter course than the artery, receives the veins corresponding to the arteries of the upper and inner part of the orbit, and finally passes between the two heads of the external rectus, through the inner part of the sphenoidal fissure, to terminate in the cavernous sinus. The *inferior ophthalmic* vein is formed by the union of branches from the lower and outer part of the orbit, and proceeding backward along the floor of the orbit, opens into the superior vein, or directly into the cavernous sinus. In front it sends a communicating vein through the spheno-maxillary fissure to join the pterygoid plexus.

Ophthalmic or Lenticular Ganglion. This small ganglion[1] (Fig. 19, p. 67), of reddish color, and about the size of a pin's head, is situated at the back of the orbit, between the optic nerve and the external rectus, on the outer side of, and usually closely adherent to, the ophthalmic artery. It is somewhat quadrilateral in shape, and receives its *sensory* or *long root* from the nasal nerve, which joins its posterior superior angle; its *motor* or *short root*, from the branch of the third nerve, going to the inferior oblique, which enters its posterior inferior angle; and its *sympathetic root* from the cavernous plexus, which joins it at its posterior border, or in conjunction with its sensory root. The ganglion, thus furnished with motor, sensory and sympathetic roots, gives off the *short ciliary nerves*. These, from eight to twelve in number, issue from the anterior upper and lower angles of the ganglion, usually four or five from the upper, the remainder from the lower. They run very tortuously with the optic nerve, pass through the back of the sclerotic coat, where they are joined by the long ciliary (from the nasal), and are distributed to the iris and the ciliary muscle. Since the ciliary nerves derive their motor influence from the third nerve, the iris must lose its power of contraction when this nerve is paralyzed.

Third Nerve, Motor Oculi. The third nerve passes forward in the outer wall of the cavernous sinus, and here receives one or two filaments from the cavernous plexus of the sympathetic. Just before it enters the inner end of the sphenoidal fissure it divides into two branches, both of which pass between the two heads of origin of the external rectus, separated from each other by the nasal nerve. The *upper* and smaller division has been already traced into the superior rectus and levator palpebræ. The *lower* division, after a short course, divides into three branches, one passing inward under the optic nerve to supply the internal rectus, another passes to the inferior rectus, and a third runs along the floor of the orbit to the inferior oblique (Fig. 19). This last-named branch sends a small twig upward to the len-

[1] W. Marshall regards this ganglion, from its mode of development and from its relations in some of the lower vertebrates, to be connected more with the third nerve than the ophthalmic.

ticular ganglion, mentioned in the description of this ganglion, and another to the inferior rectus.

What is the result of paralysis of the third nerve? Falling of the upper eyelid (ptosis), external squint, dilatation and immobility of the pupil.

Sixth Nerve, Abducens. This nerve lies in the inner wall of the cavernous sinus external to the internal carotid artery, passes through the sphenoidal fissure, and enters the orbit between the two heads of the external rectus. Here it lies below the lower division of the third and above the ophthalmic vein. The nerve terminates in fine filaments, which are distributed to the ocular surface of the external rectus. In the cavernous sinus it is joined by filaments from the carotid plexus, and in the orbit by a branch from Meckel's ganglion and from the ophthalmic nerve.

Respecting the motor nerves in the orbit, observe that they all enter the ocular surface of the muscles, with the exception of the fourth, which enters the orbital surface of the superior oblique.

Recti Muscles. The *internal* and *inferior recti* arise from a fibrous band—the *ligament of Zinn*—attached to the inner and lower borders of the optic foramen. The *external rectus* arises by two heads, the lower from the ligament of Zinn and the lower border of the sphenoidal fissure, the upper from the margin of the foramen opticum; between these heads pass, in the following order, from above downward—the upper division of the third, the nasal, the lower division of the third, the sixth nerves, and the ophthalmic vein.

The four recti diverge from each other, one above, one below, and one on each side of the optic nerve. Their broad, thin tendons are inserted into the sclerotic coat of the eye, about a quarter of an inch from the margin of the cornea (Fig. 23).

The recti muscles enable us to direct the eye toward different points; hence the names given to them by Albinus—attollens, depressor, adductor, and abductor oculi. It is obvious that by the single action of one, or the combined action of two, the eye can be turned toward any direction.

The rectus superior is supplied by the upper division of the

third nerve; the rectus internus, the rectus inferior and obliquus inferior, by the lower division. The rectus externus is supplied by the sixth.

Follow the recti to the eye, in order to see the tendons by which they are inserted. Notice also the anterior ciliary arteries, which run to the eye along the tendons. The congestion of these little vessels occasions the red zone round the cornea in iritis. It has been already mentioned that the tendons are invested by a fascia, which passes from one to the other, forming a loose tunic—*capsule of Tenon*—over the back of the eye. This tunic consists of two layers with an intermediate space, lined with flat cells, thus allowing free mobility of the globe. It is this fascia which resists the passage of the hook in the operation for the cure of squinting. Even after the complete division of the tendon, the eye may still be held in its faulty position, if this tissue, instead of possessing its proper softness and pliancy, happen to have become contracted and unyielding. Under such circumstances it is necessary to divide it freely with the scissors.

FIG. 23.



INSERTION OF THE RECTI MUSCLES WITH THE ANTERIOR CILIARY ARTERIES.

By removing the conjunctival coat of the eye, the tendons of the recti are soon exposed. The breadth and the precise situation of their insertion deserve attention in reference to the operation for strabismus. The breadth of their insertion is about three-eighths of an inch, but the line of this insertion is not, at all points, equidistant from the cornea. The centre of the insertion is nearer to the cornea by about one line than either end. Taking the internal rectus, which has most frequently to be divided in strabismus, we find that the centre of its tendon is, upon an average, three lines only from the cornea, the lower part nearly five lines, and the upper four. It is, therefore, very possible that the lower part may be left undivided in the operation, being

more in the background than the rest. The tendon of the internal rectus is nearer to the cornea than either of the others.

Inferior Oblique. This muscle *arises*, by a flat tendon, from the orbital plate of the superior maxilla, on the outer side of the lachrymal groove. It runs outward and backward between the orbit and the inferior rectus, then curves upward between the globe and the external rectus, and is *inserted* by a broad thin tendon into the outer and back part of the sclerotic, close to the tendon of the superior oblique.[1] It is supplied by the lower division of the third nerve.

The tensor tarsi muscle has been described in the dissection of the face (p. 50).

Orbital Branch of the Superior Maxillary Nerve. This is always very small, and is sometimes absent. It comes from the trunk of the superior maxillary and the spheno-maxillary fossa, enters the orbit through the spheno-maxillary fissure, and divides into two branches. Of these, one, the *temporal*, lies in a groove in the outer wall of the orbit, and after sending a small branch to the lachrymal nerve in the orbit, passes through a foramen in the malar bone to the temporal fossa. It then pierces the temporal aponeurosis an inch above the zygoma, and supplies the skin of the temple communicating with the facial, and joining frequently with the auriculo-temporal branch of the inferior-maxillary. The other branch, the *malar*, passes along the outer part of the floor of the orbit, imbedded in fat, and makes its exit through a foramen in the malar bone, to supply the skin of the cheek over the malar bone (p. 66).

DISSECTION OF THE NECK.

Surface Marking. Before the student reflects the skin of the neck he should examine the skin surface, which in some places is raised, in others depressed, indicating thereby unevenness of the subjacent structures. The neck is bounded above by a well-marked transverse ridge, indicating the lower border of the inferior maxilla, and at its lower part the neck is bounded by

[1] On the action of the recti and oblique muscles, consult M. Foster, *Text-Book of Physiology*.

another ridge which corresponds with the clavicle. Crossing obliquely from the centre of the neck below to the mastoid process above, is the rounded prominence caused by the sterno-mastoid, and crossing this muscle diagonally from its anterior to its posterior border is the external jugular vein, which varies in size in different subjects. In front and behind the sterno-mastoid are two triangular depressions; the posterior one, it will be seen, has its base at the clavicle, the anterior one at the lower jaw. The posterior triangle has the trapezius as its outer boundary, but this border is only well defined inferiorly, where the hollow becomes most marked, and takes the name of the supra-clavicular or Mohrenheim's fossa. In this is placed deeply the subclavian artery, the posterior belly of the omo-hyoid, and the brachial plexus. In front of the sterno-mastoid is another triangular hollow space with its base upward; this is called the carotid triangle, for in it lies the carotid artery immediately beneath the anterior border of the sterno-mastoid. The body of the hyoid bone can always be felt in the middle line below the symphysis of the lower jaw. About a finger's breadth below the hyoid is the prominent pomum Adami of the thyroid cartilage, and a short distance below this cartilage is the cricoid, separated from the cartilage above by the crico-thyroid membrane. The cricoid cartilage corresponds with the fifth cervical vertebra, and from it the trachea passes down, gradually receding from the surface, so that there is, especially in emaciated subjects, a deep hollow—fonticulus gutturis—immediately above the sternum. In front of the second, third, and fourth rings of the trachea is the isthmus of the thyroid gland, and there are usually another four rings below these, above the sternum, covered more or less by the depressor muscles of the os hyoides.

Dissection. The head must be slightly raised, and the face turned from the side on which the dissection is to be made. Then make a vertical incision through the skin, down the middle of the neck from the symphysis of the lower jaw to the sternum; a second along the clavicle to the acromion; a third along the base of the jaw as far as the mastoid process. Reflect the skin and subcutaneous fat, and expose the cutaneous

muscle, called the *platysma myoides*. Between the platysma and the skin is a layer of adipose tissue, called the *superficial fascia*. It varies in thickness in different subjects, but is generally more abundant at the upper part of the neck, especially in corpulent individuals, in whom it occasions a double chin.

Platysma Myoides.

The *platysma myoides* is the thin cutaneous muscle covering the front and side of the neck. It arises from the subcutaneous tissue over the pectoralis major, trapezius and deltoid muscles;[1] thence proceeding obliquely over the clavicle and the side of the neck, its fibres become more closely aggregated, and terminate thus:—The anterior cross those of the opposite platysma, immediately below the symphysis of the jaw, and are lost in the skin of the chin; the middle are attached along the base of the jaw; the posterior cross the masseter muscle, and terminate, partly in the subcutaneous tissue of the cheek, partly in the muscles at the corner of the mouth blending with the depressor anguli oris and orbicularis.[2]

The platysma forms a strong muscular defence for the neck. It is also a muscle of expression.[3] It is supplied with nerves by the cervical plexus, and by the cervical branch of the facial nerve.

[1] Some anatomists describe it as having a slender origin from the clavicle and the acromion.

[2] Some of the uppermost fibres of this part of the platysma take the name of *musculus risorius*: this has been described among the muscles of the face.

[3] If the entire muscle be permanently contracted it may occasion wry-neck, though distortion from such a cause is an exceedingly rare occurrence. A case in point is related by Mr. Gooch (*Chirurg. Works*), in which a complete cure was effected, after the failure of all ordinary means of relief, by the division of the platysma a little below the jaw.

The platysma myoides belongs to a class of muscles called *cutaneous*, from their office of moving the skin. There are not many in man, except upon the neck and face, and there is a little one (*palmaris brevis*) in the palm of the hand. To understand their use thoroughly we must refer to the lower orders of animals, in whom they fulfill very important functions, by moving not only the skin, but also its appendages. For instance, by muscles of this kind, the hedgehog, porcupine, and animals of that family can roll themselves up and erect their quills; we are all familiar with the broad "*panniculus carnosus*" on the sides of herbivorous quadrupeds, which enables them to twitch their skins, and thus rid themselves of insects. In birds, too, these cutaneous muscles are extremely numerous, each feather having appropriate muscles to move it.

Dissection. Cut through the platysma near the clavicle and turn it upward. Beneath it lies the general investment of the neck, called the *deep cervical fascia.* Upon this fascia we trace the superficial branches of the cervical plexus of nerves, the external jugular vein, and a smaller vein in front, called the anterior jugular. These superficial veins are so variable in size and course, that a general description only is applicable.

External Jugular Vein. The *external jugular vein* is formed within the substance of the parotid gland by the junction of the temporal and internal maxillary veins. After receiving the

FIG. 24.



DIAGRAM OF THE SUPERFICIAL NERVES AND VEINS OF THE NECK.

transverse facial and posterior auricular veins, it appears at the lower border of the gland, crosses obliquely over the sterno-mastoid muscle (Fig. 24), running along its posterior border, nearly as low down as the clavicle, where it pierces the deep cervical fascia and terminates in the subclavian vein. Accompanying the vein in its upper part is the auricularis magnus nerve, and crossing it, about the middle, is the superficial cervical nerve, both being branches of the superficial cervical plexus. It is usually provided with two pairs of valves—the lower, imperfect,

close to its termination in the subclavian vein; the upper, placed about an inch and a half above the clavicle. A line drawn from the angle of the jaw to the middle of the clavicle would indicate its course. To trace the vein, during life, press upon it just above the clavicle; but do not be surprised if you fail to find it; it is sometimes wanting, and frequently very small.

Near the angle of the jaw the external jugular vein communicates by a large branch with the internal jugular, and about its middle it is joined by a large vein—*posterior external jugular*—from the occipital region.

Before its termination the external jugular vein generally receives the supra-scapular, posterior scapular, and other unnamed veins; a disposition very embarrassing to the surgeon, because there is a *confluence of veins immediately over the subclavian artery* in the place where it is usually tied.

Anterior Jugular Vein. The *anterior jugular vein* is situated more in the middle of the neck, and is much smaller than the external jugular. It commences by small branches below the chin, and runs down the front of the neck, nearly to the sternum; it then curves outward, beneath the sterno-mastoid muscle, and opens either into the external jugular or the subclavian vein. We commonly meet with two anterior jugular veins, one on either side; immediately above the sternum they communicate by a transverse branch.

The size of the anterior jugular vein is inversely proportionate to that of the external jugular. When the external jugular is small, or terminates in the internal jugular, then the anterior jugular becomes an important supplemental vein, and attains considerable size. It is not uncommon to find it a quarter of an inch in diameter, and we have seen it nearly half an inch. These varieties should be remembered in tracheotomy.

Superficial lymphatic glands are sometimes found near the cutaneous veins of the neck. From four to six in number, they are small and escape observation unless enlarged by disease. One or two are situated over the sterno-mastoid muscle; others, near the mesial line.

Cutaneous Nerves of the Neck. The cutaneous nerves of the neck are the superficial branches of the cervical plexus; the plexus itself cannot at present be seen. It is formed by the communications of the anterior divisions of the four upper cervical nerves, and lies under the sterno-mastoid muscle, close to the transverse processes of the four upper cervical vertebræ, resting on the levator anguli scapulæ and the scalenus medius. The superficial branches of the plexus emerge from beneath the posterior border of the sterno-mastoid, and take different directions. They are named thus (Fig. 24):—

Superficial branches of the cervical plexus.	Ascending branches	Great auricular. Small occipital.
	Transverse branch	Superficial cervical.
	Descending branches	Sternal. Clavicular. Acromial.

The *great auricular n.* comes from the second and third cervical nerves, winds round the posterior border of the sterno-mastoid, and ascends obliquely over that muscle, near the external jugular vein, toward the parotid gland. Near the gland it divides into two principal branches, of which the *anterior* or *facial* branches are distributed to the skin over the parotid gland, where they join branches from the facial nerve, and to the side of the cheek; the *posterior* or *auricular*, after ascending a short distance, give off a branch, which ramifies mainly upon the cranial aspect of the cartilage of the ear; and a smaller branch, the *mastoid*, which supplies the skin over the mastoid process. Other filaments of this nerve communicate, in the substance of the parotid gland, with branches of the facial nerve.

The *small occipital n.* comes from the second cervical nerve, and is occasionally double. It ascends along the posterior border of the sterno-mastoid muscle to the occiput, where it supplies the back of the scalp, and communicates with the great occipital, the great auricular, and the posterior auricular nerves. It also sends off one branch, which is distributed to the skin of the temporal region, and another *auricular* to the pinna of the ear. Beneath the sterno-mastoid this nerve commonly forms a loop, which embraces the nervus accessorius, and sends a branch to it.

The *superficial cervical n.* comes from the second and third cervical nerves. It passes transversely forward over the sterno-mastoid muscle, and supplies the front of the neck. Some of its filaments ascend toward the jaw, and join the cervical branch of the facial nerve; other filaments descend and supply the skin in front of the neck as low as the sternum.

The *descending* or *supra-clavicular* branches are derived from the third and fourth cervical nerves, and divide into three branches, which cross over the clavicle, and supply the skin of the front of the chest and shoulder. Of these, one, called the *internal* or *sternal*, supplies the skin over the inner end of the clavicle and the upper part of the sternum; another, the *middle* or *clavicular*, passes over the middle of the clavicle, and is distributed to the skin over the pectoral muscle, the mammary gland, and the nipple; the third, named *external* or *acromial*, crosses over the trapezius and acromion to supply the skin of the shoulder.

Reviewing these cutaneous branches of the cervical plexus, we find that they have a very wide distribution, for they supply the skin covering the following parts—viz., the ear, the back of the scalp, the side of the cheek, the parotid gland, the front and side of the neck, the upper and front part of the chest and shoulder.

Cervical Branch of the Facial Nerve. Look for this branch beneath the fascia near the angle of the jaw (p. 83). It leaves the parotid gland, and, piercing the deep cervical fascia, divides into filaments which curve forward below the jaw; some of these, forming arches, join the superficial cervical branch of the cervical plexus; others supply the platysma and skin.

Deep Cervical Fascia. Now turn your attention to the membranous investment called the *deep cervical fascia*, which encloses the several structures of the neck. In some subjects the fascia is very thin; in others, with strong muscles, it is proportionally dense and resisting. It is always stronger in particular situations, for the more effective protection of the parts beneath; for instance, in front of the trachea, in the fossa above the clavicle, and below the angle of the jaw. It not only covers the soft parts of the neck collectively, but, by its inflections, forms separate

sheaths for the muscles, vessels, and glands. It isolates them, and keeps them in their proper relative position. A lengthened description of its numerous layers would be not only extremely tedious, but unintelligible, without considerable knowledge of the anatomy of the neck. We propose, therefore, to give only a general outline of the fascia, and of its principal layers, commencing from behind.

Tracing it from behind, we find that the cervical fascia (sometimes called *deep cervical* or *muscular fascia* of the neck) is attached to the ligamentum nuchæ and to the spinous and transverse processes of the cervical vertebræ. From these attachments it passes forward over the posterior triangle of the neck to the posterior border of the sterno-mastoid, where it splits into two layers, superficial and deep, which invest that muscle and reunite at its anterior border. The superficial layer passes toward the mesial line, where it becomes continuous with the corresponding fascia of the opposite side. The layer which lies in front of the sterno-mastoid is attached above to the base of the inferior maxilla, and passes over the parotid gland to the zygoma, to the mastoid process, and the superior curved line of the occipital bone. Traced downward, we find it attached to the clavicle and to the upper border of the sternum. In the middle line it is closely connected to the hyoid bone, and below the thyroid body divides into two layers, one being attached to the front of the upper border of the sternum, the other to the back of the upper border of the same bone. Between these layers there is a well-marked interval, containing more or less fat, and one or two small lymphatic glands. This layer forms investing sheaths for the depressor muscles of the os hyoides and larynx.

The deep layer—viz., that which passes beneath the sterno-mastoid—forms the common sheath for the carotid artery, internal jugular vein, and the pneumogastric nerve, which lie behind this muscle; the structures contained in the carotid sheath are separated from each other by delicate septa. The fascia is continued behind the pharynx (constituting the *prævertebral fascia*) to join the fascia of the opposite side, while another prolongation passes in front of the trachea beneath the sterno-thyroid muscle. Below,

it is attached to the first rib, to which it binds down the intermediate tendon of the omo-hyoid; and still further down it is continuous in the chest with the pericardium. It may also be traced under the clavicle, along the axillary vessels and nerves into the axilla. Above, it is attached to the angle of the lower jaw, from which it extends backward to the styloid process, and forms the *stylo-maxillary* ligament. Thence it is attached to the base of the skull, the petrous portion of the temporal bone, and the basilar process of the occipital bone.

A correct knowledge of the attachments of the principal layers of the cervical fascia is essential to a right understanding of the course which pus takes when it forms in the neck. For instance, suppose the pus to be formed at the lower part of the neck. If it be seated under the superficial layer (which is attached to the clavicle), it may burrow beneath the clavicle into the axilla. But if it be seated beneath the deep layer (which is attached to the first rib), then it becomes more serious, since the pus may travel through the loose tissue by the side of the pharynx, and make its way into the chest, where it may burrow down the anterior or the posterior mediastinum, and burst into the trachea or the œsophagus.

Besides forming sheaths for the several structures of the neck, there are other purposes to which the cervical fascia is subservient. The firm attachment of its layers to the sternum, the first rib, and the clavicle, forms a fibrous barrier at the upper opening of the chest, which supports the soft parts, and prevents their yielding to the pressure of the atmosphere during inspiration. Dr. Allan Burns[1] first pointed out this important function of the cervical fascia, and has recorded a case exemplifying the results of its destruction by disease.

Moreover, the great veins at the root of the neck, namely, the internal jugular, subclavian, and innominate, are so closely united by the cervical fascia to the adjacent bones and muscles, that when divided they gape. They are, as the French express it, "*canalisés*," and are, therefore, better able to resist the pressure of the atmosphere, which tends to render them flaccid and impervious

[1] *Surgical Anatomy of the Head and Neck.*

during inspiration. But this anatomical disposition of the great veins makes them more liable to the entrance of air when wounded. Instances of death have been recorded, resulting from the sudden entrance of air into the veins during operations about the neck, or even the axilla.

Sterno-cleido-mastoideus. The sterno-cleido-mastoideus is the large muscle which passes obliquely across the neck. It *arises* by a rounded tendon from the upper part of the sternum, and by fleshy fibres from the sternal third of the clavicle. It is *inserted* by a thick tendon into the external surface of the mastoid process, and by a thin aponeurosis into about the outer half of the superior curved ridge of the occipital bone.

The sternal origin of the muscle is at first separated from the clavicular by a slight interval; subsequently the sternal fibres gradually overlap the clavicular. The muscle is confined by its strong sheath of fascia, in such a manner that it forms a slight curve, with the convexity forward. Observe especially that its front border overlaps the common carotid artery; along this border we make the incision in the operation of tying the vessel.

Action of Sterno-mastoid. When both sterno-mastoids act simultaneously they draw the head and neck forward and downward, and are, therefore, especially concerned in raising the head from the recumbent position. When one sterno-mastoid acts singly, it turns the head obliquely toward the opposite shoulder; in this action it co-operates with the splenius of the other side.[1] On emergency, the sterno-mastoid acts as a muscle of inspiration, by raising the sternum; its fixed point being, in this case, at the head.

The sterno-mastoid is supplied by three nutrient arteries—an upper, a middle, and a lower. The upper sterno-mastoid artery, a branch of the occipital, enters the muscle with the n. accessorius, close to the mastoid process of the temporal bone; the middle

[1] The single action of the muscle is well seen when it becomes rigid and causes a wry-neck. Other means of relief failing, the division of the muscle near its origin is sometimes beneficial in curing the distortion. In deciding as to the propriety of this operation, we should be careful to examine the condition of the other muscles, lest, after having divided the sterno-mastoid, we should be disappointed in removing the deformity.

mastoid is a branch of the superior thyroid, and enters the under surface of the muscle, crossing over the common carotid on a level with the thyroid cartilage; the lower mastoid is a branch of the supra-scapular, and supplies the clavicular portion of the muscle, close to its origin.

The sterno-mastoid is supplied with nerves by the n. accessorius, and by branches from the deep cervical plexus: these branches come from the second and sometimes the third cervical nerves.

FIG. 25.



DIAGRAM OF TRIANGLES OF THE NECK.

1. Superior carotid triangle.
2. Inferior " "
3. Occipital triangle.
4. Supra-clavicular triangle.
5. Submaxillary triangle.

Triangles of the Neck. Anatomists avail themselves of the oblique direction of the sterno-mastoid muscle to divide the neck on each side into two great triangles, an anterior and a posterior (Fig. 25). The base of the anterior triangle is formed by the jaw, its sides by the mesial line and the front border of the sterno-mastoid. The posterior has the clavicle for the base, while the sides are defined by the hind border of the sterno-mastoid, and the front border of the trapezius.

The omo-hyoid muscle, which crosses the neck under the

sterno-mastoid, subdivides these primary triangles into four smaller ones (Fig. 25) of unequal size; an anterior superior, an anterior inferior, a posterior superior, and a posterior inferior. The direction of the omo-hyoid muscle renders their boundaries at once obvious.

Contents of Posterior Triangle. The fat and connective tissue must now be carefully removed from the posterior triangle. The following muscles will be seen forming its floor, viz.: beginning from above, the splenius capitis, the levator anguli scapulæ, the scalenus medius and posticus, and a small portion of the serratus magnus. The posterior belly of the omo-hyoid crosses this triangle about an inch above the clavicle, and subdivides it into two unequal parts—an upper or *occipital*, and a lower or *supra-clavicular*. In the occipital triangle, the larger of the two, besides the muscles just mentioned (with the exception of the serratus magnus), are found the descending branches of the cervical plexus; and, passing obliquely downward from beneath the sterno-mastoid is the spinal accessory nerve, which enters the under part of the trapezius. Curving round the posterior border of the sterno-mastoid, and becoming superficial, are the ascending and transverse branches of the superficial cervical plexus. The transversalis colli (posterior scapular) artery and vein, and its branch, the superficialis colli (which chiefly supplies the trapezius), cross transversely outward the lower part of the space. A chain of lymphatic glands is also found along the posterior border of the sterno-mastoid.

Nervus Accessorius. The upper part of the sterno-mastoid is traversed obliquely by a large nerve called the spinal accessory or n. accessorius. This nerve, the eleventh cranial, consists of two parts: one, the *accessory*, arises from the side of the medulla oblongata below the pneumogastric nerve; the other, the *spinal* part, arises from the cervical portion of the spinal cord by a series of filaments from the lateral tract as low down as the sixth cervical vertebra. The spinal portion ascends between the ligamentum denticulatum and the posterior roots of the spinal nerves, through the foramen magnum into the skull. Within the cranium the two parts unite and form a single nerve, which leaves

the skull through the foramen jugulare. Here the *accessory* portion is connected with the ganglion of the root of the pneumogastric by several filaments; and lower down it again joins the pneumogastric at the ganglion of the trunk, below which the two nerves become blended. The accessory and spinal portions communicate in the foramen jugulare. Below the foramen the *spinal* part runs behind the internal jugular vein, the digastric and stylo-hyoid muscles, and then pierces obliquely the upper third of the sterno-mastoid. Emerging beneath its outer border, it crosses the posterior triangle of the neck to the under surface of the trapezius, to which it is distributed. The nervus accessorius supplies also the sterno-mastoid in its passage through it, and here it joins some branches from the third cervical. After leaving the muscle it is joined by branches from the second and third cervical nerves. Beneath the trapezius it forms a plexus with the third and fourth cervical nerves. The upper mastoid artery, a branch of the occipital, enters the sterno-mastoid with the nerve.

Supra-Clavicular Triangle. The *supra-clavicular* or *subclavian triangle* is bounded below by the clavicle, in front by the outer border of the sterno-mastoid, and above by the posterior belly of the omo-hyoid muscle. The area of the triangle thus formed will vary in proportion to the obliquity of the omo-hyoid muscle, and the extent to which the sterno-mastoid and trapezius are attached to the clavicle. The depth of the vessels and nerves contained in this space depends, not only upon the degree to which the clavicle arches forward, but varies with the elevation and depression of the shoulder.

Dissection. The descending branches of the cervical plexus, together with some fat, should now be cut through and turned aside, when a layer of a fascia which binds down the omo-hyoid muscle to the clavicle will be exposed. Beneath this is a deeper layer of fascia, which covers the subclavian vessels and brachial plexus of nerves, and descends with them under the clavicle into the axilla. Between these two layers we meet with more or less fat and connective tissue and lymphatic glands continuous with those in the axilla. It will be easily understood how a collection of pus in the axilla may ascend in front of the vessels

and point above the clavicle, or, *vice versâ*, how matter formed in the neck may travel under the clavicle and point in the axilla.

Near the posterior border of the sterno-mastoid muscle the external jugular vein passes through both layers of the deep fascia, and terminates in the subclavian; but before its termination it is commonly joined by the supra-scapular, the posterior scapular, and other unnamed veins proceeding from the surrounding muscles; so that there is in this situation a *confluence of veins*, which, when large or distended, is exceedingly embarrassing.

The fascia and the glands should be removed, and the following objects carefully dissected. Behind and nearly parallel with the clavicle is the supra-scapular (transversalis humeri) artery, a branch of the thyroid axis. A little higher is the transversalis colli, or posterior scapular (commonly a branch of the thyroid axis), which crosses the lower part of the neck toward the posterior superior angle of the scapula. Both these arteries are very irregular in respect to their origin, *the last particularly* being often given off from the subclavian in the third part of its course Search for the outer border of the scalenus anticus, which descends from the transverse processes of the cervical vertebræ to the first rib: running down longitudinally upon it may be seen the phrenic nerve. The subclavian vein lies upon the first rib in front of the insertion of the anterior scalene muscle behind the clavicle, so that it is not usually seen in this triangle. The subclavian artery rises up into the neck as high as an inch above the clavicle, and sometimes on the right side as high as an inch and a half. It appears higher than the vein, emerging beneath the outer border of the scalenus anticus, and care must be taken to preserve the small branch from the brachial plexus, which crosses the artery and proceeds to the subclavius muscle. The large nerves constituting the brachial plexus come out between the scalenus anticus and medius, higher than the subclavian artery, and on a plane posterior to that vessel. These different objects will be described in detail hereafter.

Dissection of the Anterior Triangle. The *anterior triangle* must now be dissected. In doing so, notice, before the deep cervical fascia is removed, the arching forward of the anterior border of the

sterno-mastoid muscle, which is connected to the lower jaw by the fascia, so that the common carotid artery is concealed from view before the parts are disturbed. The anterior triangle is bounded behind by the anterior border of the sterno-mastoid, in front by the middle line of the neck, and above by the lower border of the inferior maxilla. Covering the triangle are the superficial and deep cervical fasciæ, and the platysma; passing across it are the superficial cervical n., the infra-maxillary branch of the facial nerve; and descending in front is the anterior jugular vein. This space is subdivided by the anterior belly of the omo-hyoid into a *superior* and an *inferior carotid* triangle, and above them is a third triangle mapped out by the converging bellies of the digastric muscle and the lower jaw, and is called the *sub-maxillary* or *digastric* triangle (Fig. 25, p. 90).

The *inferior carotid triangle* is bounded above and below by the omo-hyoid and sterno-mastoid muscles, and in front by the middle line. The muscles forming its floor are the sterno-hyoid and sterno-thyroid muscles, and lying on them is the anterior jugular vein; in the middle line is the thyroid body covering the trachea.[1]

The *superior carotid triangle* has for its boundaries the sterno-mastoid, the omo-hyoid, and the posterior belly of the digastricus. Its muscular floor is formed by the hyo-glossus, the middle and inferior pharyngeal constrictors, and the thyro-hyoid. In this space are found the bifurcation of the common carotid into its external and internal divisions, and the following branches of the external carotid—the superior thyroid, lingual, facial, the occipital, and the ascending pharyngeal arteries—their accompanying veins and the internal jugular vein. The nerves seen are the hypo-glossal, crossing over the external carotid, the infra-maxillary branch of the facial, the spinal accessory, the superior and external laryngeal nerves, and in front of the carotid sheath is the descendens noni.

The *digastric* triangle will be described subsequently (p. 109).

Now examine the flat muscles in front of the neck, which pull

[1] The vessels and nerves lying within and upon the carotid sheath are not seen, as they are situated beneath the anterior border of the sterno-mastoid.

down the larynx and os hyoides—namely, the sterno-hyoid, sterno-thyroid, omo-hyoid, and thyro-hyoid.[1] Remove the fascia which covers them, disturbing them as little as possible, and take care of the nerves (branches of the descendens noni), which enter their outer borders.

Sterno-hyoid. The sterno-hyoid *arises* from the back part of the sternum and posterior sterno-clavicular ligament, from the clavicle and occasionally from the cartilage of the first rib, and is *inserted* into the lower border of the body of the os hyoides. This is the most superficial of the muscles in front of the neck. We cut in the mesial line between these muscles in laryngotomy.

Sterno-thyroid. The sterno-thyroid *arises* from the back part of the sternum, below and internal to the origin of the sterno-hyoid, and the cartilage of the first rib, and is *inserted* into the oblique ridge on the ala of the thyroid cartilage. This muscle is situated immediately under, and is much broader than, the sterno-hyoid.

The two sterno-hyoid muscles converge as they ascend to their insertions, and opposite the cricoid cartilage and the two or three upper rings of the trachea they are in contact with one another. The sterno-thyroid, however, diverge to their insertions, but are in contact below, the result of which is that the trachea is completely covered in front by muscular fibres.

Omo-hyoid. The omo-hyoid consists of two fleshy portions connected by a tendon. It *arises* from the upper border of the scapula, and sometimes from the ligament over the notch, and is *inserted* into the lower border of the body of the os hyoides just external to the sterno-hyoid. From the scapula it comes nearly horizontally forward across the lower part of the neck, and passes beneath the sterno-mastoid, *over* the sheath of the great vessels of the neck on a level with the cricoid cartilage; then, changing its direction, it ascends nearly vertically close to

[1] The sterno-hyoid and sterno-thyroid muscles often present slight transverse tendinous lines. These tendinous intersections are quite rudimentary in man; but in some animals with long necks, *e.g.*, the giraffe, they are so developed that each depressor muscle is composed of alternations of muscle and tendon.

the outer border of the sterno-hyoid. Thus the muscle does not proceed straight from origin to insertion, but forms an obtuse angle beneath the sterno-mastoid muscle. The intermediate

FIG. 26.



CENTRAL LINE OF NECK.—COURSE AND RELATIONS OF COMMON CAROTID ARTERY.

tendon is situated at the angle and is bound down to the first rib and the sternum by a process of the deep cervical fascia. The object of this peculiar direction of the omo-hyoid appears to be to keep tense that part of the cervical fascia which covers the apex of the pleura, and thus to resist atmospheric pressure.

Relations of the Omo-hyoid. At its origin the omo-hyoid is covered by the trapezius, then by the clavicle and subclavius, and lastly, by the sterno-mastoid and platysma myoides. It lies on the scalenus medius and anticus, the brachial plexus, the phrenic nerve, then on the internal jugular vein, pneumogastric nerve and common carotid artery enclosed within their common sheath, on the descendens noni, the sterno-thyroid, and thyro-hyoid muscles.

These depressor muscles are all supplied with nerves (Fig. 26, p. 96) by the *descendens noni* (a branch of the twelfth or hypoglossal), and by the *communicantes noni* (branches of the second and third cervical nerves). The descendens noni sends a separate branch to each belly of the omo-hyoid. They are supplied with blood by the superior and inferior thyroid arteries.

Thyro-hyoid. The thyro-hyoid *arises* from the oblique line on the ala of the thyroid cartilage, and runs up to be *inserted* into the lower border of the body and the inner half of the great cornu of the hyoid bone. This muscle is a continuation of the sterno-thyroid. It is supplied by a special branch of the hypoglossal nerve which enters the muscle close by its posterior border, in company with the hyoid branch of the lingual artery. In front of the muscle, are the omo-hyoid and sterno-hyoid muscles, and it covers the thyro-hyoid membrane, the thyroid cartilage, and the superior laryngeal vessels and nerve as they enter the larynx.

Action of the Depressor Muscles. The sterno-hyoid, sterno-thyroid, omo-hyoid, and thyro-hyoid muscles co-operate in fixing the larynx and os hyoides, *e. g.*, in sucking, or they depress the larynx after it has been raised in deglutition. Again, they depress it in the utterance of low notes. That the larynx is raised or depressed, according to the height of the note, may be ascertained by placing the finger on it while singing through an octave. The omo-hyoid, in addition, is a tensor of the cervical fascia, and draws down the hyoid bone to its own side. The thyro-hyoid depresses the hyoid bone, or elevates the thyroid cartilage, according as the one or the other is the fixed point.

Dissection. The sterno-mastoid muscle must now be cut transversely through the middle, and the two ends turned upward and downward, so that they may be replaced if necessary. This done, notice the strong layer of fascia which lies under the muscle and forms part of its sheath. It is attached to the angle of the jaw, thence descends over the large vessels of the neck, and is firmly connected to the clavicle and first rib. This fascia prevents matter coming to the surface, when suppuration takes place by the side of the pharynx.

Remove the fascia, and clean the various structures beneath the sterno-mastoid, taking care not to cut away the descendens noni and communicantes noni nerves, which lie in front of the sheath of the common carotid. Dissect out the lymphatic glands which lie along the sheath of the large vessels.

Parts exposed beneath the Sterno-Mastoid. The objects exposed to view, when the muscle is reflected, are very numerous. Among these the more important are: the sterno-clavicular articulation, the splenius capitis and colli, the posterior belly of the digastricus, the levator anguli scapulæ, scalenus medius and anticus, omo-hyoid, sterno-hyoid and sterno-thyroid muscles; the occipital artery, the common carotid artery and its division, the internal jugular vein, the subclavian artery and the branches of the first part of its course, the cervical plexus, and the lower cervical nerves which form the brachial plexus; the phrenic, pneumogastric, hypoglossal and spinal accessory nerves, the descendens and communicantes noni nerves; the subclavian vein and its tributaries; and lastly, a small part of the parotid gland, and the three sterno-mastoid arteries. On the left side, in addition, we find the thoracic duct; on the right side, the right lymphatic duct.

Course and Relations of the Common Carotid. The *common carotid artery* is now exposed in the whole extent of its course in the neck. It arises, on the right side, from the arteria innominata, behind the upper part of the right sterno-clavicular articulation; on the left, from the arch of the aorta. It ascends in front of the bodies of the cervical vertebræ, by the side of the trachea, thyroid gland and larynx, as high as the upper border of the

thyroid cartilage, and then divides into the external and internal carotids. Thus, a line drawn from the sternal end of the clavicle to a point midway between the mastoid process and the angle of the jaw, will nearly indicate its course. It is contained in a sheath of the deep cervical fascia, together with the internal jugular vein and the pneumogastric nerve. The vein lies on the outer side of, and parallel with, the artery; the nerve lies behind and between the artery and the vein. The structures contained within this sheath are separated from each other by a thin septum of fascia, so that each has a separate investment. Owing to the increasing breadth of the larynx, the two common carotid arteries, which at their origin lie near together, are separated by a wide interval at their point of division.

At the lower part of the neck the carotid artery is deeply placed, but as it ascends it becomes more superficial, although it has the appearance of being deeply situated, owing to the prominence of the thyroid cartilage. *In front* the artery is covered by the skin, superficial fascia, platysma myoides, deep fascia, the sternal portion of the sterno-mastoid, the sterno-hyoid and thyroid muscles, and on a level with the cricoid cartilage, it is crossed by the omo-hyoid. Above this point the artery becomes more superficial, and is covered by the integument, platysma, the cervical fasciæ, the middle sterno-mastoid artery, and only slightly overlapped by the sterno-mastoid. Lying upon the sheath of the artery, we find the descendens noni joined by the communicantes noni nerves. The sheath is crossed by the facial, the superior and middle thyroid veins, and lower down by the anterior jugular vein, all of which empty themselves into the internal jugular. This is the general rule, and especial attention should be directed to it, because the veins are liable to be overlooked and injured in the operation of tying the carotid. To the *inner side* of the artery we find the trachea, the thyroid body, the recurrent laryngeal nerve, the inferior thyroid artery, the external laryngeal nerve, the inferior constrictor of the pharynx and the larynx. On the *outer side* are the pneumogastric nerve and the internal jugular vein. *Behind* the artery are the sympathetic nerve, the inferior thyroid artery, the recurrent laryngeal nerve;

and lastly, the carotid sheath lies successively upon the longus colli and the rectus capitis anticus major muscles.[1]

The common carotid may be ligatured, either above or below the omo-hyoid. It is most accessible above the point where this muscle crosses; and therefore, if the surgeon has his choice, he would prefer to tie the vessel in this situation. In the higher operation we make an incision, three inches in length, along the inner border of the sterno-mastoid, the centre of the incision being opposite the cricoid cartilage; we cut through the skin, superficial fascia, platysma, deep cervical fascia, when we come to the anterior border of the sterno-mastoid. The overlapping edge of this muscle must be drawn outward, and the muscle at the same time relaxed by turning the head to the same side. The sheath of the vessel is then exposed, and a small opening is to be made on its inner side large enough to admit the aneurism needle, which should be passed round the artery on its outer side, so as to avoid wounding the internal jugular vein. The vessel is then to be ligatured, care being taken not to separate more of the sheath than is necessary from the artery, and not to include in the ligature the pneumogastric or descendens noni nerves.

After ligation of the artery, the collateral circulation is maintained by the following vessels: between the branches of the external and internal carotid arteries of the opposite side with the corresponding branches of the ligatured side; between the vertebral and the posterior communicating of the same side; between the inferior and superior thyroids of the same side; between the profunda cervicis and the princeps cervicis of the occipital of the same side.

[1] It is important that we should be aware that the common carotids vary occasionally in their origin. Thus the right may arise in common with the left carotid, or the right may arise separately from the arch of the aorta, in which case the right subclavian is usually transposed. The left may be given off from the innominate artery of the right side, or it may arise in common with the left subclavian, and thus form a left innominate. In transposition of the aorta there is a left innominate, which is given off first, the right carotid and the right subclavian arising as separate branches from the arch. The place of division of the common carotid is subject to considerable variation; it may divide higher or lower than usual, the former being the more frequent. Rarely there has been no common carotid artery, the external and internal arising as separate branches from the arch of the aorta.

In what respects the Left Carotid differs from the Right.

In the first part of its course the left carotid differs from the right in the following particulars:—

1. It arises from the arch of the aorta, is therefore longer and deeper seated than the right, and is covered by the first bone of the sternum.

2. It is crossed by the left brachio-cephalic vein.

3. It is in close relation with the œsophagus and the trachea.

4. It is in close relation with the left recurrent laryngeal nerve.

5. It is in close relation posteriorly with the thoracic duct.

6. It is covered by the thymus gland in early life.

The artery has *in front* the sternum, the sterno-hyoid and sterno-thyroid muscles, the left innominate vein, and the remains of the thymus gland; to the *left side* it has the left subclavian artery and the left pneumogastric nerve; to the *right side* the arteria innominata; and *behind*, the trachea, œsophagus, and thoracic duct.

The common carotid, as a rule, gives off no branch in its course; but, occasionally, the middle sterno-mastoid, the superior thyroid, or, more rarely, the vertebral, arise from it prior to its division. At its bifurcation it usually presents a slight bulbous enlargement, which is sometimes so marked that it might be mistaken for an incipient aneurism. It is necessary to know that the carotid sometimes divides as low as the level of the cricoid cartilage, and that not infrequently the division takes place as high as the hyoid bone.

Internal Jugular Vein.

The internal jugular vein is the continuation of the lateral sinus, and returns the blood from the brain. Leaving the skull through the foramen jugulare, it receives the inferior petrosal sinus, and at the junction it presents a slight enlargement, the *sinus*. The vein descends on the outer side of the internal carotid, and subsequently the common carotid arteries, in the same sheath, and joins the subclavian vein at a right angle to form the brachio-cephalic or innominate vein. In its course down the neck it receives the pharyngeal, occipital, facial, lingual, superior and middle thyroid veins.

Previous to their terminations the internal jugular veins incline

somewhat to the right side to meet the corresponding subclavian veins; thus, on the right side, there is a triangular interval between the artery and vein, in which is seen the pneumogastric nerve and vertebral artery; on the left side the vein slightly overlaps the artery, thus rendering ligature of the left carotid more difficult than of the right. The internal jugular veins, moreover, advance slightly to meet the subclavian veins, so that they lie on a plane a little anterior to their accompanying arteries. A little before their termination the internal jugulars have a double valve.

Descendens Noni and Communicantes Noni Nerves.

The *descendens noni* (p. 109), a branch of the hypoglossal, runs down obliquely over the sheath of the carotid to supply the depressor muscles of the os hyoides. Trace the nerve upward to see that it leaves the hypoglossal where this nerve curves round the occipital artery. For a short distance the descendens noni lies within the carotid sheath; but, about the level of the os hyoides, it comes through the sheath, and crosses obliquely over the carotid, from the outer to the inner side. The descendens noni is reinforced by one or more nerves termed *communicantes noni*, derived from the second and third cervical nerves. These communicating branches descend on the outer side of the internal jugular vein, and form generally two loops in front of the carotid sheath, constituting a triangular plexus called the "*ansa hypoglossi*." From these loops the nerves proceed to the anterior and posterior bellies of the omo-hyoid, to the sterno-hyoid, and sterno-thyroid muscles. A small branch may sometimes be traced proceeding from the descendens noni into the chest, to join the cardiac and phrenic nerves.

In some subjects the descendens noni seems to be wanting, in which case it will probably be concealed *within* the carotid sheath; when this happens the reinforcing loops from the cervical nerves will be found behind the internal jugular vein.[1]

[1] By many anatomists the descendens noni is regarded as the combination of filaments from the hypoglossal and pneumogastric nerves; by some, it is looked upon as a branch of the pneumogastric; and lastly, which is most probable, it is considered by others to be mainly derived from a branch which is sent to the hypoglossal from the first and second cervical nerves.

Dissection. The thyroid body should now be examined. To expose it, reflect the sterno-hyoid and thyroid muscles from their insertions, so that they can be replaced if necessary. Next observe the lymphatic glands of the neck, and lastly survey the objects in the central line of the neck, from the jaw to the sternum.

Thyroid Body. This very vascular, gland-like body lies over the front and sides of the upper part of the trachea, and extends upward on each side of the larynx. It consists of *two lateral lobes*, connected a little below the cricoid cartilage by a transverse portion called the *isthmus*, and weighs from one to two ounces. Each lobe is conical, about two inches in length, and an inch and a quarter in breadth. Its base is opposite the fifth or sixth ring of the trachea, and the apex by the side of the thyroid cartilage. Its anterior surface is convex, and is covered by the sterno-hyoid, sterno-thyroid, and omo-hyoid muscles; its deep surface—concave—embraces the sides of the trachea and larynx, and usually extends so far backward as to be in contact with the pharynx. Its external border overlaps, in most cases partially, but sometimes completely, the common carotid artery, particularly on the right side; and there are instances in which the lobe is deeply grooved by the vessel.

The isthmus lies over the second and third rings of the trachea. This portion of the organ varies much in its dimensions. In some instances there is no transverse portion. This corresponds with the normal disposition in most of the lower orders of mammalia; but in man, it is a failure in the union of the two halves by which the organ is originally developed.[1] Generally, the vertical measurement is about half an inch. Between its upper border and the cricoid cartilage is a space about one-third of an inch in extent, where the trachea is free; this space, therefore, is the more preferable situation for tracheotomy. But the vertical measurement of this isthmus is sometimes of very considerable

[1] Concerning the development of the lateral halves and central portion of the thyroid body, see a paper by Callender, in the *Proceedings of the Royal Society*, 1867.

length, so that it has been seen covering the trachea almost down to the sternum.[1]

The thyroid body is closely connected, by areolar tissue, to the sides of the trachea, to the cricoid and thyroid cartilages. Hence it rises and falls with the larynx in deglutition.

The thyroid varies in size in different individuals and at different periods of life. It is relatively larger in the child than the adult, in the female than the male. In old age it diminishes in size, becomes firmer, and occasionally contains earthy matter.[2]

By far the most notable considerations in respect to the thyroid body are the number, the large size, and the free inosculations of its *arteries*. The superior thyroid arteries come from the external carotid and enter the front surface of the apex of each lobe; the inferior thyroid come from the subclavian, and enter the under surface of the base. An artery, called the middle thyroid (thyroidea ima), is observed in some subjects; it is given off from the arteria innominata, or the arch of the aorta, and ascends directly in front of the trachea to the isthmus.

Its *veins* are equally large and form a plexus upon it. The superior and middle thyroid veins cross the common carotid, and open into the internal jugular. The inferior thyroid veins, two in number, descend over the front of the trachea, communicate freely with each other, and terminate in the left brachio-cephalic vein. When you perform tracheotomy, bear in mind the size of these

[1] From the upper part of the isthmus, or from the adjacent border of either lobe, most commonly the left, a conical prolongation of the thyroid body, called the *pyramid*, frequently ascends in front of the crico-thyroid membrane, as high as the pomum Adami, and is attached to the body of the os hyoides by fibrous tissue. In some subjects we may observe a few muscular fibres passing from the os hyoides to the pyramid. This constitutes the *levator glandulæ thyroideæ* (see preparation in Museum of St. Barth. Hosp., Patholog. Series, No. 14) of some anatomists. There are instances in which the pyramid is double; and lastly, we have seen a considerable portion of this thyroid substance lying over the crico-thyroid membrane, completely isolated from the rest of the organ. These varieties deserve notice, because any one portion of this structure may become enlarged independently of the rest, and occasion a bronchocele.

[2] The thyroid body is, primarily, developed as a pouch from the anterior wall of the pharynx; the lateral lobes are first formed, and are subsequently united by the isthmus. W. Müller, *Jenaisch. Zeitsch.*, 1871.

inferior thyroid veins, and the possible existence of a middle thyroid artery.

Its *nerves* are furnished by the middle and inferior cervical ganglia of the sympathetic. They accompany the arteries.

The *lymphatics* of the thyroid body are both numerous and large. They form a dense network on the surface, and pass into the connective tissue of the gland, and eventually "enclose the primary lobes in complete rings or more or less perfect arches." On the right side they open into the right lymphatic duct, on the left side into the thoracic duct.

Structure of the Thyroid Body. The thyroid body belongs to the class of duct-less glands, since no excretory duct has been discovered. It is invested by a thin covering of dense areolar tissue, which connects it with the surrounding structures and also penetrates it, imperfectly dividing it into lobes and supporting the vessels as they enter it. It consists of a multitude of closed vesicles, which are imbedded in a delicate reticulum. The cells vary in size, from $\frac{1}{150}$ of an inch to that of a pin's head, and do not communicate with each other. In hypertrophy of the gland we sometimes see them as large as a horse-bean, or even larger. The vesicles are oval and are lined by a single layer of endothelial columnar cells containing a glairy yellow fluid, in which are found a large number of nuclei, nucleated cells, and not infrequently red blood-cells in various stages of disintegration and decolorization.[1] The blood-vessels pass into the connective tissue surrounding the vesicles, and form a dense capillary plexus, which comes into close relation with the vesicular epithelium cells, and with the endothelium of the lymph-spaces; from these lymph-spaces, which are placed between the vesicles, the lymphatics have their commencement. The function of the gland is probably that of disintegration of the red blood-cells, and of the conveyance into the general lymphatic system of the products of these degenerative changes.

An enlargement of the thyroid body is termed a "bronchocele." If the relation of its lobes to the trachea and œsophagus

[1] Baber, "Researches on the Minute Structure of the Thyroid Gland," *Philos. Trans.*, 1881.

be properly understood, it is easy to predicate the consequences which may result from their enlargement. The nature and severity of the symptoms will, to a certain extent, be determined by the part of the organ affected. An enlargement of the left lobe is more likely to produce a difficulty in swallowing, on account of the inclination of the œsophagus toward the left side. If the isthmus be enlarged, difficulty in breathing will probably be the prominent symptom, and in order to remove this danger the isthmus has been divided, and in part removed.

An instance is related by Allan Burns, in which the isthmus was placed between the trachea and the œsophagus. It must be obvious that enlargement of a part so situated would occasion great difficulty in swallowing. I have seen two cases in which the lateral lobes projected so far inward that they completely embraced the back of the œsophagus.

Small lymphatic glands are observed about the thyroid body, especially in front of the trachea; one is often situated over the crico-thyroid membrane. These glands, if enlarged by disease, might be mistaken for a small bronchocele.

Deep Cervical Lymphatic Glands.

In the connective tissue which surrounds the great vessels of the neck, we meet with a series of lymphatic glands, called the deep cervical. They form an uninterrupted chain (whence their name *glandulæ concatenatæ*), from the base of the skull, along the side of the neck, to the clavicle, beneath which they are continuous with the thoracic and the axillary glands. Some of these glands lie anterior to the common carotid artery; others, between it and the spine. This disposition explains the well-known fact, that, when these glands are enlarged, the great vessels and nerves of the neck are liable to become imbedded in their substance.

The glands are particularly numerous near the division of the common carotid, by the side of the pharynx, and the posterior belly of the digastricus. The lymphatics connected with them come from all parts of the head and neck. These vessels unite, to form, on both sides of the neck, one or more absorbent trunks, called the jugular. On the left side this jugular trunk joins the thoracic duct, or opens by a separate orifice into the junction of

the left internal jugular and subclavian veins; on the right side it opens into the right lymphatic duct, a short trunk about half an inch in length, which terminates at the angle of the junction of the right internal jugular and subclavian veins. The terminations of the thoracic duct and the right lymphatic duct are guarded by two small semilunar valves, in order to prevent regurgitation of blood back from the veins.

The contiguity of the glands to the great vessels and nerves of the neck explains the symptoms produced by their enlargement. The tumor may be so situated as to be raised and depressed by the pulsation of the carotid, and thus simulate an aneurism. A careful examination, however, will distinguish between an inherent and a communicated pulsation. By grasping the tumor we become sensible that the pulsation does not depend upon any variation of its magnitude, but upon the impulse derived from the artery; consequently, if the tumor be lifted from the vessel, all feeling of pulsation ceases.

Survey of the Central Line of the Neck.

The parts in the central line of the neck should now be well studied (Fig. 26, p. 96). Beginning at the chin, we observe the insertions of the digastric muscles. Below these is the junction, or raphé, of the mylo-hyoid muscles. Then comes the os hyoides. Below the os hyoides is the thyro-hyoid membrane, attached above to the posterior and upper border of the hyoid bone, and below to the thyroid cartilage. Next is the pomum Adami, or projection of the thyroid cartilage, which is apparent between the contiguous borders of the sterno-hyoidei. Below the thyroid cartilage is the cricoid. These two cartilages are connected by the crico-thyroid membrane, across which runs the crico-thyroid artery to join its fellow. Below the cricoid cartilage is the trachea. This is crossed by the isthmus of the thyroid body, and lower down it recedes from the surface, covered by the inferior thyroid veins.

Now the chief surgical interest lies just above and just below the cricoid cartilage. This cartilage can be felt very plainly in the living subject at any age, no matter how fat. In laryngotomy, the crico-thyroid membrane is divided transversely. The membrane should be divided *close* to the edge of the cricoid c., for two

reasons: 1. In order to be further from the vocal cords. 2. To avoid the crico-thyroid artery which crosses the middle of the membrane. If more room be required, the cricoid cartilage should be divided longitudinally.

In tracheotomy, the trachea may be opened by a perpendicular incision, above the isthmus of the thyroid body, or below it. The operation above the isthmus, if there be space enough for the introduction of the tube, is the easier and safer of the two; for here the trachea is nearer to the surface, and no large blood vessels are, generally speaking, in the way. The space available measures from a quarter to half an inch; and the isthmus is not so firmly adherent to the trachea as to prevent its being drawn downward for a short distance. However, it is right to state that, in one case out of every eight or ten, there is *no* available space.

Tracheotomy below the isthmus is neither an easy nor a safe operation, for many reasons: 1. The trachea recedes from the surface as it descends, so that just above the sternum it is nearly an inch and a half from the skin. 2. The large inferior thyroid veins are in the way. 3. A middle thyroid artery may run up in front of the trachea, direct from the arteria innominata. 4. The arteria innominata itself lies sometimes upon the trachea higher than usual, and may, therefore, be in danger. 5. The left brachio-cephalic vein in some cases crosses the trachea above the edge of the sternum, instead of below it. The celebrated French surgeon, Béclard, used to relate in his lectures the following occurrence: A student had fallen into the Seine, and was nearly drowned. As he was recovering very gradually, some kind friends attempted to accelerate the process by making an opening into the trachea. In so doing they wounded the brachio-cephalic vein. Blood poured into the trachea, and the result was instantly fatal.

Whoever pays attention to this subject in the dissecting-room will soon be convinced of the fact that, not only large veins, but large arteries, occasionally cross the crico-thyroid membrane as well as the trachea, thus showing the necessity of cutting *cautiously* down to, and fairly exposing, the air tube before we venture to open it. It is preferable, after making the first incision through the skin, to lay aside the sharp knife and to use a blunt one, so

that the tissues may be torn rather than cut; by this proceeding the liability to hemorrhage is materially lessened.

Dissection of the Submaxillary Region, or the Digastric Triangle.

When the platysma and the cervical fascia have been removed from their attachment to the jaw, the most conspicuous object is the submaxillary gland. Observe that the fascia is here very strong, and forms for the gland a complete cap-

FIG. 27.



DIGASTRIC TRIANGLE AND CONTENTS.

sule. Beneath the jaw are several lymphatic glands, from six to ten in number, of which some lie superficial to the salivary gland, others beneath it. These glands receive the lymphatics of the face, the tonsils and the tongue.

A little dissection will expose a muscle called the digastricus, consisting of two distinct fleshy portions connected by a tendon.

They form, with the body of the jaw, a triangle called the *digastric*, of which we propose to examine the contents. The muscles forming its floor are the mylo-hyoideus and hyo-glossus. Under the submaxillary gland is the facial artery, which here runs a tortuous course, and finally turns up over the lower jaw in front of the masseter muscle. Lying on the mylo-hyoideus, under cover of the lower jaw, is the submental artery, accompanied by the mylo-hyoid nerve and artery. Behind the submaxillary gland, and separating it from the parotid, which also is contained within this triangle, is the stylo-maxillary ligament. Ascending and then entering the parotid is the external carotid artery, in front of which is the infra-maxillary branch of the facial nerve. Deep in this space are situated the internal jugular vein, the internal carotid artery, and the pneumogastric nerve; and running obliquely forward between the internal and external carotid arteries, are the stylo-glossus, stylo-pharyngeus, glosso-pharyngeal nerve, and the stylo-hyoid ligament.

Digastricus.

The *digastricus* consists of two muscular bellies united by an intermediate tendon. The posterior belly *arises* from the digastric fossa of the temporal bone, passes obliquely downward, forward and inward, and then ascends to be *inserted* by its anterior belly close to the symphysis of the lower jaw. Raise the submaxillary gland to see the intermediate tendon of the digastricus piercing the stylo-hyoid muscle, the angle which it forms, and how it is fastened by aponeurosis to the body and the greater cornu of the os hyoides. Observe also that this aponeurosis—*supra-hyoid aponeurosis*—is connected in the mesial line with its fellow of the opposite side, so that a fibrous expansion occupies the interval between the anterior portions of the digastrici.

The chief action of the digastricus is to depress the lower jaw. But if the lower jaw be fixed, then the muscle raises the os hyoides, as in deglutition.

The posterior belly of the digastricus is supplied by a nerve from the facial; the anterior belly by a branch from the mylo-hyoidean nerve (which comes from the third division of the fifth pair).

Stylo-hyoideus. The stylo-hyoideus *arises* from the middle of the styloid process of the temporal bone, and passing downward and forward is *inserted* into the body of the os hyoides. This muscle at first runs above the posterior belly of the digastricus, and near its insertion is pierced by the digastric tendon. Its nerve is derived from the facial, close to its exit from the stylo-mastoid foramen, in common with the branch to the posterior belly of the digastricus.[1] Its action is to raise and draw back the os hyoides.

The digastric triangle is bounded above by the horizontal ramus of the lower jaw, and mastoid process of the temporal bone; behind by the posterior belly of the digastricus; and in front by the anterior belly. The objects to be examined in this triangle are twelve in number, as follows:—

1. Submaxillary salivary gland.
2. Facial vein.
3. Facial artery.
4. Submental artery.
5. Mylo-hyoidean nerve.
6. Submaxillary lymphatic glands.
7. Stylo-maxillary ligament.
8. Part of the parotid gland.
9. Part of the external carotid artery.
10. Mylo-hyoideus muscle.
11. Hypoglossal nerve.
12. Part of the hyo-glossus muscle.

Submaxillary Salivary Gland. In the ordinary position of the head, the submaxillary gland is partially concealed by the jaw, but when the head falls back the gland is more exposed. It is about the size of a chestnut, weighs about two drachms, and is divided into several lobes. Its upper margin is covered by the body of the jaw; its lower margin overlaps the side of the os hyoides. Its cutaneous surface is flat, being covered only by the skin, platysma, and deep cervical fascia; but the lobes on its deep surface are irregular, and often continuous with those of the sublingual gland. By raising the gland we find that it lies upon the mylo-hyoideus, the hyo-glossus, the stylo-glossus, the tendon of the digastricus, and a portion of the hypoglossal nerve, seen above the tendon. Part of the gland passes beneath the posterior border of the mylo-hyoid, and not infrequently becomes continuous with the sublingual gland. The facial artery lies in a groove on its

[1] In many, if not in most subjects, a small filament from the hypoglossal nerve is distributed to this muscle.

deeper surface, and subsequently upon its upper border; and it is separated from the parotid gland, which is situated behind it, by the stylo-maxillary ligament. Mark these relations well, because they are of importance, as will be presently explained in tying the lingual artery.

The duct of the gland cannot, at this stage of the dissection, be traced further, for it runs forward, under cover of the mylo-hyoideus, to end in the floor of the mouth, by the side of the frænum linguæ. The description of its course and relations had better, therefore, be deferred till it can be dissected in its whole length with the gustatory nerve in the pterygoid region.

Facial Vein. The facial vein does not accompany the facial artery, but runs nearly a straight course. It leaves the face at the anterior edge of the masseter m., then runs *over* the submaxillary gland, the digastricus and stylo-hyoideus and the carotid artery, to join the internal jugular. This is the rule—but there are frequent exceptions. Before it empties itself into the internal jugular it is joined by a large branch from the external jugular vein. The principal point to remember is, that the vein runs superficial to the gland, and that we must be cautious in opening abscesses under the jaw.

Course and Relations of the External Carotid Artery. The course and relations of the external carotid artery, and its branches in the neck, should now be made out as far as the parotid gland. In preparing a view of them, observe that nearly all the veins lie *in front* of their corresponding arteries. In removing the connective tissue, fat and lymphatic glands, the student must take care of the nerves and other structures which are liable to be injured.

The *external carotid* arises from the common carotid about the level of the upper border of the thyroid cartilage. It ascends to the interval between the ear and the jaw in a slightly curved direction, at first forward and then backward. The external and the internal carotids are in the adult nearly of equal size; but the external rapidly diminishes in size, owing to the large branches it gives off within a short distance. At first it *lies beneath* the skin, superficial fascia, platysma myoides, deep cervical fascia, some of

the superficial cervical nerves, and the sterno-mastoid muscle. It is next crossed by the hypoglossal nerve, the facial and lingual veins, the posterior belly of the digastricus and stylo-hyoideus; it then enters the parotid gland, where it lies beneath the facial nerve and the external jugular vein, and terminates between the external auditory meatus and the neck of the jaw, by dividing into the temporal and internal maxillary arteries. *Internally* the artery is in relation with the hyoid bone, the pharynx, the parotid gland, and the posterior border of the ascending ramus of the lower jaw.

Behind the external carotid, and separating it from the internal, are the stylo-glossus, the stylo-pharyngeus, the glosso-pharyngeal nerve, and the stylo-hyoid ligament. The superior laryngeal nerve and part of the parotid gland are also placed behind the artery.

Notice the relative position which the external and internal carotids bear to each other. The external lies at first on the same plane with, but nearer to the side of, the pharynx than the internal. It soon, however, changes its position, and crosses obliquely in front of the internal, to reach the space between the angle of the jaw and the mastoid process. The internal carotid ascends perpendicularly by the *side of the pharynx* to the base of the skull.

The external carotid gives off the following branches—

1. The superior thyroid.
2. The lingual.
3. The facial.
4. The occipital.
5. The posterior auricular.
6. The internal maxillary.
7. The temporal.
8. The ascending pharyngeal.

Superior Thyroid Artery. The superior thyroid, the first branch of the external carotid, arises just below the great cornu of the os hyoides. It lies in the superior carotid triangle, and, curving downward and inward, runs beneath the omo-hyoid, sterno-hyoid, and sterno-thyroid muscles to the upper and front surface of the thyroid body, in which it terminates. Its branches are the four following:—

1. The *hyoid*, a small muscular branch, runs horizontally in-

ward below the greater cornu of the os hyoides, and anastomoses with its fellow.

2. The *superior laryngeal* branch, accompanied by the superior

FIG. 28.



DIAGRAM OF THE BRANCHES OF THE EXTERNAL CAROTID ARTERY AND THEIR BRANCHES.

laryngeal nerve, runs inward beneath the thyro-hyoid muscle, pierces the thyro-hyoid membrane (sometimes the thyroid cartilage), supplies the muscles and the mucous membrane of the larynx, and anastomoses with its fellow of the opposite side.

3. The *middle sterno-mastoid*, a small branch, variable as to origin, descends over the sheath of the common carotid artery, and enters the under aspect of the sterno-mastoid muscle.

4. The *crico-thyroid*, an artery of great interest in reference to the operation of laryngotomy, crosses the crico-thyroid membrane, and communicates with a corresponding branch on the opposite side (Fig. 26, p. 96). One or two small branches pass through the membrane to the interior of the larynx. It is important to know that the crico-thyroid artery often varies in direction and size. In most cases it is small, and runs across the centre of the membrane; we should, therefore, be least likely to wound it in laryngotomy, by dividing the membrane close to the cricoid cartilage. But it is by no means infrequent to find this artery of considerable size, taking an oblique or even a perpendicular direction in front of the membrane, and finally distributed to one of the lobes of the thyroid body. We have seen several instances in which the membrane was crossed by the main trunk of the superior thyroid. These facts should establish the practical rule in laryngotomy, not to make an opening into the larynx until it has been fairly exposed.

Among the many arterial inosculations about the thyroid body are two which deserve notice: the one is formed between the two superior thyroid arteries along the upper border of the isthmus; the other takes place along the back part of the lateral lobe between the superior and inferior thyroid arteries of the same side.

The *superior thyroid vein* leaves the upper part of the thyroid body, crosses over the common carotid artery, and joins the internal jugular or the facial vein.

Superior Laryngeal Nerve. The *superior laryngeal nerve*, mentioned as accompanying the superior laryngeal artery, is given off from the inferior ganglion of the pneumogastric nerve. It descends by the side of the pharynx, behind both carotid arteries, and divides into two branches—the internal and external laryngeal nerves. The *internal branch* enters the larynx through the thyro-hyoid membrane, accompanied by the superior laryngeal artery, and supplies the mucous membrane of the larynx

with its exquisite sensibility. Some of its branches may be traced upward in the ary-epiglottidean fold, to supply the epiglottis and the base of the tongue; others descend to the rima glottidis; a large branch passes down behind the ala of the thyroid cartilage to join the recurrent laryngeal nerve; and a small branch pierces the arytenoideus to supply the mucous membrane beneath it. The *external branch*, descending beneath the depressors of the larynx, accompanies the crico-thyroid artery, and after distributing filaments to the pharyngeal plexus, supplies the thyroid body, the inferior constrictor, and the crico-thyroid muscles. It receives a branch from the superior cervical ganglion of the sympathetic, and sends off a cardiac filament to join the superior cardiac branch of the sympathetic behind the common carotid artery.

Lingual Artery. The lingual artery and its branches will be described in the dissection of the submaxillary region.

Facial Artery. The *facial artery* is the third branch of the external carotid. It runs tortuously *under* the hypoglossal nerve, the posterior belly of the digastricus and stylo-hyoideus, and beneath or through the substance of the submaxillary gland to the face, where it appears at the anterior border of the masseter. Below the jaw the facial rests on the mylo-hyoideus, and gives off the four following branches:—

1. The *ascending* or *inferior palatine artery* runs up between the stylo-glossus and the stylo-pharyngeus m., and behind the internal pterygoid m. to the pharynx, to which and the neighboring parts it gives branches. Ascending as far as the levator palati, it divides into two branches: one courses along the tensor palati to supply the soft palate; the other enters the tonsil, and anastomoses with the descending palatine of the internal maxillary, and with the tonsillar branches of the ascending pharyngeal.

2. The *tonsillar* runs up between the internal pterygoid and the stylo-glossus m.; then, perforating the superior constrictor, it supplies the tonsil and root of the tongue.

3. *Glandular* branches to the submaxillary gland and side of tongue.

4. The *submental* arises from the facial behind the submaxillary gland, and runs forward upon the mylo-hyoideus, beneath the inferior maxilla, distributing branches in its course to the gland and the adjacent muscles. It then curves over the bone and divides into two branches: a superficial one, which supplies the skin and lip; and a deep one, which runs between the muscles and the bone, and inosculates with the mental and inferior labial arteries. Beneath the inferior maxilla it usually inosculates with the sublingual artery.

The remaining branches of the external carotid artery will be described later on.

Mylo-hyoidean Nerve. Look for the mylo-hyoidean nerve near the submental artery. The nerve comes from the inferior dental (before its entrance into the dental foramen), and running along a groove on the inner side of the inferior maxilla, advances between the bone and the internal pterygoid m., to supply the mylo-hyoideus and the anterior belly of the digastricus.

Submaxillary Lymphatic Glands. The submaxillary lymphatic glands receive the lymphatics of the face and the tongue. They are often enlarged in cancerous diseases of the tongue or the lower lip. It should be remembered also that there are lymphatic glands in the mesial line below the chin.

Mylo-hyoideus. The mylo-hyoideus, a triangular muscle, *arises* from the mylo-hyoid ridge of the lower jaw, from the symphysis, as far back as the last molar tooth. Its posterior fibres are *inserted* into the body of the os hyoides, the anterior being attached to a medium tendinous line, termed the *raphé*. Thus the muscles of opposite sides form a muscular floor for the mouth. Superficially, it is in relation with the anterior belly of the digastricus, the submaxillary gland, the submental artery, and the mylo-hyoidean n. By its deep surface, it is in relation with part of the hyo-glossus, the stylo-glossus, the genio-hyoideus, Wharton's duct, the gustatory and hypoglossal nerves with their communications, and the sublingual gland. It is supplied with nerves by the mylo-hyoid branch of the inferior dental; with blood by the submental artery. The muscles of opposite sides conjointly elevate the os hyoides and the floor of the mouth—as in deglutition.

Stylo-maxillary Ligament. This is a layer of the deep cervical fascia, extending from the angle of the jaw to the styloid process. It is a broad sheet of fascia, and separates the submaxillary gland from the parotid. It is continuous with the fascia covering the pharynx; this gives it a surgical interest, because it prevents accumulations of matter formed near the tonsils and upper part of the pharynx from coming to the surface.

The remaining objects seen in the submaxillary triangle—namely, the parotid gland, the hypoglossal nerve, the hyo-glossus muscle—will be described presently when they can be better seen. Your attention should now be directed to a piece of surgical anatomy, which will enable you readily to find and tie the lingual artery. It is this:—

A curved incision, about two inches in length, being made from the lesser cornu along the upper border of the great cornu of the os hyoides, through the skin, the platysma, and the cervical fascia, you will come upon the lower edge of the submaxillary gland. Lift up the gland, which is easily done, and underneath it you will observe that the tendon of the digastricus makes two sides of a triangle, of which the base is formed by the hypoglossal nerve crossing the hyo-glossus muscle. Within this little triangle, cut transversely through the fibres of the hyo-glossus: under them is the lingual artery, lying on the middle constrictor. The first time you perform this operation on the dead subject, you will not unlikely miss the artery and cut through the middle constrictor into the pharynx.

Dissection. The facial vessels must now be divided immediately below the jaw. Reflect the anterior belly of the digastricus from its insertion; detach the mylo-hyoideus from the middle line and the os hyoides, and turn it over the body of the jaw, taking care not to injure the muscles and structures beneath. The lower jaw must now be sawn through, a little to the dissector's side of the symphysis, and the bone drawn upward by hooks. The tongue should then be drawn out of the mouth, and fastened by hooks. The os hyoides should be drawn by means of hooks, so as to put the parts on the stretch. All

this done, we have to make out, by carefully cleaning away the fat and connective tissue, the following objects represented in Fig. 29, p. 120—

1. Genio-hyoideus.
2. Hyo-glossus.
3. Stylo-glossus.
4. Genio-hyo-glossus.
5. Submaxillary duct.
6. Sublingual gland.
7. Hypoglossal nerve.
8. Gustatory nerve.
9. Submaxillary ganglion.
10. Lingual artery.

Genio-hyoideus. The genio-hyoideus *arises* from the inferior tubercle behind the symphysis of the jaw, and passes downward and backward to be *inserted* into the front of the body of the os hyoides. This round muscle is situated in the mesial line, parallel to its fellow. Its nerve comes from the hypoglossal, and its blood from the lingual artery. Its action is to draw the os hyoides forward and upward; and if the hyoid bone be fixed, it depresses the lower jaw.

Hyo-glossus. The hyo-glossus *arises* from the body, the greater and lesser cornua of the os hyoides, and is *inserted* into the posterior two-thirds of the side of the tongue, its fibres blending with the stylo-glossus and palato-glossus. It is a square and flat muscle, and its fibres ascend nearly perpendicularly from origin to insertion. The fibres arising from the body of the hyoid bone, termed the *basio-glossus*, are directed backward and upward and overlap the fibres which have their origin from the greater cornu and are termed the *kerato-glossus*. Those that arise from the lesser cornu are termed the *chondro-glossus*. The nerve to the hyo-glossus comes from the hypoglossal, and its blood from the lingual. Its action (with that of its fellow) is to depress the tongue. Observe the objects which lie *upon* the hyo-glossus; namely, the hypoglossal and gustatory nerves (which at the anterior border form one or more loops of communication with one another), the chorda tympani nerve, the submaxillary ganglion, the submaxillary gland and its duct, the hyoid branch of lingual artery, the lingual vein, the sublingual gland, the digastricus, stylo-hyoid, stylo-glossus, and mylo-hyoid muscles. *Beneath* the hyo-glossus muscle lie the lingual artery and vein, part of the middle constrictor of the pharynx, part of

the genio-hyo-glossus, the lingualis and the glosso-pharyngeal nerve.

Genio-hyo-glossus. The genio-hyo-glossus *arises* by a tendon from the upper tubercle behind the symphysis of the lower jaw, and is *inserted* as follows: the lower fibres into the body of the os hyoides; the upper fibres into the tongue, from the base to the apex. It is the largest and most important of the muscles of the tongue. It is fan-shaped, with the apex attached

FIG. 29.



MUSCLES, VESSELS, AND NERVES OF THE TONGUE.

to the symphysis; thence its fibres radiate into the entire length of the tongue. *Externally*, the muscle is in relation with the stylo-glossus, lingualis, and hyo-glossus, the lingual artery, the sublingual gland, Wharton's duct, the hypoglossal and gustatory nerves; *inferiorly*, by its lower border it is in contact with the genio-hyoid; *above*, by its anterior border with the mucous membrane of the mouth; and *internally*, it is in contact with its fellow and the fibrous septum of the tongue. It derives its nerves from the hypoglossal, and its blood from the lingual artery. Its action

is various. The posterior fibres, by raising the os hyoides and drawing forward the base of the tongue, protrude the tongue out of the mouth; the anterior draw the tongue back again. When every part of the muscle acts, it draws down the whole tongue, and is therefore one of the chief muscles concerned in suction.

Stylo-glossus. The stylo-glossus, a long and slender muscle, *arises* from the outer side of the styloid process near its apex and from the stylo-maxillary ligament; its fibres pass downward and forward, and then nearly horizontal, and are *inserted* along the side of the tongue. It runs outside the hyo-glossus nearly to the tip of the tongue, and blends with the fibres of this muscle, as well as with the palato-glossus. Its nerve comes from the hypoglossal. Its action is to retract the tongue.

Hypoglossal Nerve. The hypoglossal, or twelfth cranial nerve, is the *motor* nerve of the muscles of the tongue. It arises by several filaments, twelve to fifteen, from the front of the medulla oblongata between the anterior pyramid and the olivary body. It pierces the dura mater in two fasciculi which leave the skull through the anterior condylar foramen; these subsequently blend to form a single nerve trunk. It lies deeply beneath the internal jugular vein and internal carotid artery, where it is intimately connected with the lower ganglion of the pneumo-gastric nerve; it then comes up between the artery and vein, and, immediately below the posterior belly of the digastricus, curves forward over the occipital, the internal and external carotid and facial, arteries. Next it crosses the hyo-glossus muscle, and passing beneath the mylo-hyoid, divides into branches which supply the following muscles; namely, the stylo-glossus, hyo-glossus, genio-hyo-glossus, lingualis, and the genio-hyoideus.

As it curves round the occipital artery, the hypoglossal nerve sends the *descendens noni* to the depressors of the os hyoides (p. 120). It also sends a nerve to the thyro-hyoideus, which proceeds from it where it crosses over the external carotid, accompanied by the hyoid branch of the lingual artery. Near the anterior border of the hyo-glossus, it communicates by several loops with the gustatory nerve. (Fig. 29.)

The hypoglossal at its origin is purely a *motor* nerve. But after

leaving the skull, it receives communications from the first two cervical nerves. These communications are important physiologically for two reasons: 1. They account for the hypoglossal nerve containing sensory fibres. 2. They contribute the greater part of the filaments of the descendens noni. It is also connected by small branches with the pneumogastric nerve and the superior cervical ganglion of the sympathetic at the base of the skull.

Sublingual Gland. The sublingual gland lies immediately beneath the mucous membrane of the floor of the mouth. Its shape is oblong, with the long axis (about an inch and a half) directed from before backward, and it weighs about a drachm. Its relations are as follows: *above*, it is covered with mucous membrane: *below*, it rests upon the upper surface of the mylo-hyoid muscle; *internally*, it is in contact with the hyo-glossus, genio-hyo-glossus, stylo-glossus, the gustatory nerve and Wharton's duct; *posteriorly*, with the submaxillary gland; and in *front*, it rests in a depression behind the symphysis of the lower jaw.

The ducts of the sublingual gland (ducts of Rivinus[1]) vary in number from eight to twenty. They terminate by minute openings behind the orifice of the submaxillary duct, along the ridge felt upon the floor of the mouth. One or more ducts terminate in the submaxillary duct: one of these takes the name of the *duct of Bartholin*.

The duct of the submaxillary gland may now be traced across the hyo-glossus, and under the gustatory nerve to the floor of the mouth.

Lingual or Gustatory Nerve. This nerve is a branch of the inferior maxillary or third division of the fifth pair of cranial nerves. Emerging beneath the external pterygoid muscle, in company with, but in front of, the inferior dental nerve, it rests upon the internal pterygoid muscle. It descends between this latter muscle and the ramus of the lower jaw, and curves forward toward the side of the tongue over the superior constrictor of the pharynx, along the upper part of the hyo-glossus, at the anterior border of which it crosses, superficially, the duct of the submaxillary gland (Fig. 29). Having reached the under part of the tongue, the nerve

[1] Aug. Quirin. Rivinus, *De Dyspepsia*, Lips., 1678.

divides into numerous branches which pierce the muscular structure of the tongue, and then break up into filaments which supply the mucous membrane and the fungiform and filiform papillæ on its anterior three-fourths. Beneath the external pterygoid it is joined at an acute angle by the chorda tympani, a branch of the facial nerve; in its course it gives off some communicating branches to the hypoglossal nerve near the anterior border of the hyo-glossus. It supplies also the mucous membrane of the mouth, gums, and the sublingual gland, one or more branches to the submaxillary ganglion, and at the apex of the tongue the terminal branches of this nerve and the hypoglossal are connected.

Submaxillary Ganglion. At the lower border of the gustatory nerve as it lies upon the hyo-glossus muscle, and before it crosses the submaxillary duct, you will find a small, convex, triangular ganglion, about the size of a pin's head. Like the other ganglia in connection with the branches of the fifth pair, it receives filaments of *communication* of three different kinds—viz., motor, sensory and sympathetic. Its motor root is the chorda tympani, derived from the facial nerve; its sensory branches proceed from the gustatory; and its connection with the sympathetic system is established by a branch which comes from the nervi molles round the facial artery. The ganglion supplies five or six branches of *distribution* to the submaxillary gland, its duct, and the mucous membrane of the floor of the mouth. Meckel describes a small branch of the ganglion which sometimes passes forward to join a branch of the hypoglossal, on the hyo-glossus m., and ends in the genio-hyo-glossus.

Lingual Artery. The lingual artery is generally the second branch of the external carotid. Curving slightly upward and inward from its origin, the artery soon runs forward round the great cornu of the hyoid bone, beneath the posterior belly of the digastricus and stylo-hyoideus, and then passes beneath the hyo-glossus m. parallel to the os hyoides. At the anterior edge of the hyo-glossus it ascends to the under surface of the tongue, and is continued forward to the apex of the tongue under the name of *ranine*. Before the artery passes beneath the

hyo-glossus, it is crossed by the hypoglossal nerve, but it immediately after becomes separated from the nerve by this muscle. Under the hyo-glossus the artery lies upon the middle constrictor of the pharynx, and the genio-hyo-glossus; in the substance of the tongue, it lies betwen the genio-hyo-glossus and the inferior lingualis. The curves made by the artery are for the purpose of allowing the elongation of the tongue. Its branches are:—

1. The *hyoid*, a small artery which runs along the upper border of the hyoid bone, supplying the muscles and anastomosing with its fellow, and with the hyoid branch of the superior thyroid artery. The nerve to the thyro-hyoid muscle, which is derived from the hypoglossal, accompanies this artery.

2. The *dorsales linguæ*, two or more, run under the hyo-glossus to the back of the tongue, the mucous membrane, tonsil, and soft palate.

3. The *sublingual*, arising near the anterior border of the hyo-glossus, supplies the sublingual gland, the mylo-hyoideus, and the mucous membrane of the mouth and gums. This artery generally gives off the little artery of the frænum linguæ, which is sometimes wounded in cutting the frænum in children who are tongue-tied; especially when we neglect the rule of pointing the scissors downward and backward.

4. The *ranine* is the termination of the lingual artery. As it runs forward to the tip of the tongue along the outer side of the genio-hyo-glossus, along with the gustatory nerve, it distributes branches to the tongue, and at the tip inosculates slightly with its fellow of the opposite side.

The *ranine vein*, commencing at the tip of the tongue, after joining with the venæ comites of the lingual artery and the dorsal veins of the tongue, runs along its under surface *over* the hyoglossus, and terminates in the internal jugular or facial vein.

The best place for finding and tying the lingual artery has been mentioned (p. 118). The rule laid down is trustworthy only when the artery runs its normal course. We have known an instance in which a good anatomist failed in an attempt to tie the lingual artery, because the vessel arose from the facial behind the

submaxillary gland, and then passed through the mylo-hyoideus to reach the tongue.

Occipital Artery. The occipital artery arises from the posterior part of the external carotid, usually opposite the facial artery, and runs upward and backward along the lower border of the digastricus toward the mastoid process. It passes then under the posterior belly of the digastricus, and further on in its course it lies in the interval between the transverse process of the atlas and the mastoid process, close to the rectus capitis lateralis; it now changes its direction, for it runs horizontally backward in the occipital groove of the temporal bone, under all the muscles attached to the mastoid process—namely, the sterno-mastoid, the splenius capitis, the trachelo-mastoid, and the digastricus, and it lies on the superior oblique and the complexus. Arrived at the back of the head, the artery pierces the cranial attachment of the trapezius, and ascending, divides into wide-spreading branches for the supply of the scalp.

In the first part of its course, the occipital artery crosses over the internal carotid artery, the internal jugular vein, the pneumogastric and the spinal accessory nerves, and is itself crossed by the hypoglossal nerve. It sends off the seven following branches:—

1. *Muscular* branches to the digastric, stylo-hyoid, splenius, and trachelo-mastoid muscles.

2. The *superior sterno-mastoid*, which enters the muscle with the nervus accessorius.

3. The *auricular* ramifies on the cranial aspect of the concha.

4. The *posterior meningeal* ascends with the internal jugular vein, and enters the cranium through the foramen jugulare to supply the dura mater of the posterior fossa.

5. The *princeps cervicis*, which we shall see better hereafter, is a short trunk which runs down the back of the neck, and divides into two branches—a superficial, lying beneath the splenius, and supplying also the trapezius, and a deep branch lying under the complexus, and anastomosing with branches of the vertebral and with the deep cervical branch of the superior intercostal artery between this muscle and the semi-spinalis colli.

6. The *mastoid* enters the foramen in the mastoid process, and supplies the dura mater.

7. The *cranial* branches supply the scalp on its posterior aspect, and anastomose freely with the corresponding artery of the opposite side, the posterior auricular and the superficial temporal arteries.

The *occipital vein* accompanies the artery, and is connected with the lateral sinus through a small vein running through the mastoid foramen. It subsequently terminates in the internal jugular, occasionally in the external jugular vein.

Posterior Auricular Artery. The posterior auricular artery, the fifth branch, is given off from the posterior part of the external carotid. It arises above the digastricus, lies on the styloid process, and under cover of the parotid gland reaches the furrow between the cartilage of the ear and the mastoid process. Before it reaches the furrow it is crossed by the facial nerve,[1] and just beneath it is the spinal accessory. Above the mastoid process it divides into two branches, a posterior inosculating with the occipital, and an anterior communicating with the temporal. It supplies the back of the scalp and the cartilage of the ear. It gives off—

1. Small branches to the digastricus, stylo-hyoid, and the parotid gland.

2. The *stylo-mastoid*, a very constant little artery, which runs through the stylo-mastoid foramen to supply the mastoid cells, the vestibule, and the membrana tympani. In young subjects, one of these latter branches forms a vascular circle around the circumference of the membrane with the tympanic branch of the internal maxillary.

3. The *auricular* branch runs along the cranial surface of the auricle, and anastomoses with the superficial temporal and occipital arteries. Some of the branches pierce the cartilage of the ear and ramify on its anterior surface.

4. The *mastoid* branch is distributed to the structures over the mastoid process.

The *posterior auricular* vein is rather large, and running over the mastoid process, terminates in the external jugular vein.

[1] The posterior auricular artery frequently runs superficial to the facial nerve.

Posterior Auricular Nerve. The posterior auricular nerve lies close to the artery of the same name. It is the first branch of the seventh or facial nerve after its exit from the stylo-mastoid foramen. It runs behind the ear and divides into an *auricular* branch to the retrahens and the attollens aurem, and an *occipital* branch to the posterior belly of the occipito-frontalis, which communicates with the small occipital nerve. The nerve is connected with the great auricular nerve of the cervical plexus, and with the auricular branch of the pneumogastric nerve.

Ascending Pharyngeal Artery. This long and straight branch arises about half an inch above the division of the common carotid. It ascends between the internal carotid and the side of the pharynx to the base of the skull, lying upon the rectus capitis anticus major. It gives off numerous branches; among them are:—

1. *Small external branches* which pass outward to supply the anterior recti muscles, the superior cervical ganglion, the pneumogastric and hypoglossal nerves, and the prevertebral lymphatic glands. They anastomose with the ascending cervical artery.

2. *Pharyngeal branches*, some of which pass to the two lower pharyngeal constrictors and the stylo-pharyngeus: one, the largest of all, enters the pharynx above the superior constrictor, and terminates in the soft palate, the Eustachian tube, and the tonsils.

3. *Meningeal branches.*—One passes through the foramen lacerum posticum, with the internal jugular vein, and is distributed to the dura mater of the occipital fossa; another through the foramen lacerum medium, and one through the anterior condylar foramen.

The *pharyngeal vein* receives some meningeal branches, also small veins from the soft palate, Eustachian tube, and uniting, form the pharyngeal plexus which opens into the internal jugular or common facial vein.

The examination of the two remaining branches of the external carotid, the internal maxillary and temporal, must for the present be postponed. Meanwhile the student should make out the deep cervical plexus and its branches.

Cervical Plexus of Nerves. This plexus is formed by the anterior branches of the four upper cervical nerves. It consists of a series of loop-like communications, between these nerves, close to the transverse processes of the four upper cervical vertebræ; each nerve dividing into an ascending and a descending branch, with the exception of the first. The plexus rests on the levator anguli scapulæ and scalenus medius, and is situated behind the sterno-mastoid m. and the internal jugular vein.

The plexus gives off *superficial* and *deep* branches; the superficial coming from the second, third and fourth nerves, the deep from the third and fourth n. The superficial branches have been already described (p. 85).

The deep branches may be divided into an *internal* and an *external* series.

INTERNAL SERIES.—1. The *phrenic* arises from the third, fourth, and fifth cervical nerves, descends obliquely inward over the scalenus anticus, and then crosses over the first part of the subclavian artery. Near the thorax it is joined by the sympathetic, and frequently by a looped branch from the nerve to the subclavius muscle. Its course through the thorax to its destination in the diaphragm will be described p. 130.

2. The *communicantes noni* come from the second and third cervical nerves, wind round the internal jugular vein, and join the descendens noni in front of the carotid sheath, forming the "ansa hypoglossi." They supply the depressor muscles of the os hyoides and larynx.

3. *Muscular* branches which proceed from the first cervical and the loop between it and the second cervical, to the recti antici, the rectus lateralis, and longus colli muscles.

4. *Branches* which communicate with the pneumogastric, hypoglossal and sympathetic nerves, and one to join the fifth cervical.

EXTERNAL SERIES—1. One or more *communicating branches* to the nervus accessorius: firstly in the sterno-mastoid, then in the occipital triangle, and lastly beneath the trapezius.

2. *Muscular branches* to supply the trapezius, levator anguli scapulæ, scalenus medius and sterno-mastoid. The branches to the trapezius, levator anguli scapulæ, and scalenus medius, come

from the third and fourth; the branch to the sterno-mastoid from the second cervical nerve.

Dissection. The clavicle should now be sawn through the middle, and the sternal half raised with the sterno-mastoid attached, so that the bone can be replaced, to study its relation to the subjacent parts. The scalene muscles and the subclavian artery throughout its whole course must next be carefully dissected. While this is being done, the student must be careful not to injure the branches of the subclavian artery, the lymphatic duct on the right, and the thoracic duct on the left side, the nerve to the subclavius m., the phrenic nerve, the cervical and the brachial plexuses of nerves, and their small branches.

Scalene Muscles. The scalene muscles, so called from their resemblance to a scalene triangle, extend from the transverse processes of the cervical vertebræ to the first and second ribs. They may be considered as intercostal muscles, since the transverse processes of the cervical vertebræ are but rudimentary ribs. Anatomists describe them as three separate muscles—an anterior, a middle and a posterior; the anterior and middle are attached to the first rib, the posterior to the second. In plan and purpose these three muscles are one.

Scalenus Anticus. The scalenus anticus is attached above to the anterior tubercles of the transverse processes of the third, fourth, fifth and sixth cervical vertebræ, and below by a flat tendon to the tubercle on the *inner* border and upper surface of the first rib in the front of the groove for the subclavian artery.

Scalenus Medius. The scalenus medius is attached above to the posterior tubercles of the transverse processes of all the cervical vertebræ except the first, and below to the first rib behind the scalenus anticus, extending from the tubercle forward, for an inch and a half.

Scalenus Posticus. The scalenus posticus is attached above to the posterior tubercles of the transverse processes of the two or three lowest cervical vertebræ, and below to the second rib between its tubercle and angle, anterior to the levator costæ, and behind the serratus magnus.

The scaleni are supplied by branches derived from the lower cervical nerves.

The scalene muscles are important agents in raising the thorax, in a deep inspiration. Take a deep breath, and you can easily feel them contracting. They can bend the cervical portion of the spine, if their lower attachment be the fixed point, as in rising from the recumbent position.

The scalenus anticus is one of those muscles about which we ought to know well all that lies in front of it, and all that lies behind it. In the front of it are, the clavicle, the subclavius, the clavicular origin of the sterno-mastoid, the omo-hyoid, the phrenic nerve, the subclavian vein, the supra-scapular, the posterior scapular, and the ascending cervical arteries. Behind it are the subclavian artery, the five nerves which form the brachial plexus, and the pleura; to its inner side is the internal jugular vein, and the vertebral artery separates it from the longus colli.

Make your finger familiar with the feel of the tubercle on the first rib, to which the scalenus anticus is attached. This tubercle is the guide to the subclavian artery, for it enables you to find the *outer edge* of the scalenus anticus, where you must look for the vessel. Is the scalenus anticus entirely concealed from view by the sterno-mastoid or not? This will depend upon the breadth of the clavicular attachment of the sterno-mastoid. As a general rule, it may be said that the scalene muscle *is* concealed by the sterno-mastoid, and that consequently, in tying the subclavian artery, it may be necessary to divide partially the clavicular origin of the muscle.

Phrenic Nerve. The phrenic nerve runs down in front of the scalenus anticus, from the outer to the inner border. It arises from the third, fourth, and fifth cervical nerves, but chiefly from the fourth. It enters the chest between the subclavian artery and vein, crosses in front of the internal mammary artery, and continues its course between the pericardium and pleura, in front of the root of the lung, to the diaphragm, which it supplies.

When the spinal cord is injured above the fourth cervical vertebra, the origin of the phrenic is implicated; therefore the dia-

phragm, as well as the other muscles of inspiration, are paralyzed. Death is the immediate result.[1]

COURSE AND RELATIONS OF THE SUBCLAVIAN ARTERIES.

The left subclavian artery differs from the right, not only in its origin, but in the relations of the first part of its course. The right should, therefore, be examined first, and then the differences between it and the left.

Right Subclavian Artery. The right subclavian artery is one of the two great branches into which the arteria innominata divides behind the sterno-clavicular joint. It runs outward behind the scalenus anticus, then inclines downward over the first rib, at the outer border of which it takes the name of axillary. The artery describes a curve, of which the greatest convexity is between the scalene muscles. The height to which the arch ascends varies. Generally, it rises higher in women than in men, on the right side than on the left.

To study its relations more precisely, the course of the subclavian is divided into three parts: 1. The part which intervenes between its origin and the inner border of the scalenus anticus. 2. That which lies behind the scalenus. 3. That which intervenes between the outer border of the scalenus and the outer border of the first rib.

The *first* portion of the artery lies deeply in the neck and passes upward and outward to the inner border of the scalenus anticus. It is *covered* by the skin, platysma, superficial and deep fasciæ, the sternal end of the clavicle, the sterno-mastoid, sterno-hyoid, and sterno-thyroid muscles, and a layer of deep fascia,

[1] The phrenic nerve is joined by a filament from the sympathetic, and frequently by a filament from that branch of the brachial plexus which supplies the subclavius muscle. This is sometimes a branch of considerable size, and forms the greater portion of the phrenic itself. We have met with many instances in which this accessory branch was larger than the regular trunk; in all of them it crossed over the subclavian artery in the third part of its course, and would probably have been injured in the operation of tying this vessel. That such an accident has actually happened is reported by Bransby Cooper in his surgical lectures. He speaks of having injured this accessory branch of the phrenic in tying the subclavian artery. The patient had incessant spasm of the diaphragm till he died.

continued from the inner border of the scalenus anticus. It is *crossed* by the internal jugular and vertebral veins, by the pneumogastric and phrenic nerves, and by some cardiac filaments of the sympathetic. *Inferiorly* it rests upon the pleura. *Behind* the artery are the recurrent branch of the pneumogastric, the sympathetic nerve, the longus colli, the transverse process of the seventh cervical vertebra and the apex of the lung covered with the pleura. The subclavian vein lies below the artery. Three branches arise from this portion of the subclavian—viz., the vertebral, internal mammary, and thyroid axis.

In the *second* (the highest) part of its course, the artery lies between the scalene muscles. It is *covered* by skin, platysma, and superficial fascia, by the clavicular origin of the sterno-mastoid, the deep cervical fascia, and by the scalenus anticus and phrenic nerve, which separate it from the subclavian vein. *Behind* the artery is the scalenus medius; *above* it, is the brachial plexus; *below* it, is the pleura. Only one branch, the superior intercostal, is given off from this part of the artery.

In the *third* part of its course, the artery passes downward and outward, and lies in the supra-clavicular triangle upon the surface of the first rib. Here it is most superficial, and is *covered* by the skin, platysma, the two layers of the cervical fascia, and the clavicular branches of the superficial cervical plexus; subsequently by the suprascapular artery and vein, the clavicle, the subclavius muscle, with its nerve; and, what is of much more consequence, it is here crossed by the external jugular and (often) the supra and posterior scapular veins; so that there is here a confluence of large veins in front of the artery. The subclavian vein is situated below the artery, but on a plane anterior to it. *Below* it, is the first rib, and behind it the scalenus medius. *Above* the artery, and to its outer side, are the trunk nerves of the brachial plexus and the omo-hyoid m. One of these nerves (the conjoined fifth and sixth cervical) runs so nearly parallel with the artery, and *on a plane anterior* to it, that it is quite possible to mistake the nerve for the artery, in the operation of tying it. We have heard a hospital surgeon of great experience say, that he had seen this mistake committed

on three separate occasions. In this part of its course, the artery, as a rule, gives off no branches; the most frequent exceptions are the posterior scapular and supra-scapular.

FIG. 30.



Left Subclavian Artery. The left subclavian is the last of the three great branches which arise from the arch of the aorta. It ascends nearly vertically out of the chest, and then arches in front of the apex of the lung and pleura to reach the inner border of the scalenus anticus, behind which it runs over the first rib.

In the first part of its course the left subclavian lies deeply in the chest, near the spine. On its *outer* or left side it is covered by the pleura; on its *inner* or right side are at first the trachea, then the œsophagus and thoracic duct; in *front* are the left lung, covered with its pleura, the pneumogastric and phrenic nerves, and the cardiac branches, all of which lie parallel with the artery, the left common carotid, and the left brachio-cephalic vein; at the level of the upper part of the chest it has in front the sterno-thyroid, sterno-hyoid, the sterno-mastoid muscles, the left internal jugular and vertebral veins, and the sternal end of the clavicle; *behind* it are the longus colli, the vertebral column, the inferior cervical ganglion of the sympathetic, the œsophagus, and the thoracic duct.

Behind the scalenus anticus, and on the surface of the first rib, the relations of the left subclavian are similar to those of the right (p. 132).

The left subclavian, then, differs from the right only in the first part of its course. Now, what are these differences?

1. The left subclavian comes direct from the arch of the aorta, and is therefore longer, deeper in the chest, and more vertical than the right, which comes from the arteria innominata.

2. The left subclavian is in close relation with the œsophagus and the thoracic duct; the right is not.

3. The left subclavian is crossed by the left brachio-cephalic vein.

4. The left subclavian has the phrenic, pneumogastric, and cardiac nerves nearly parallel with it; on the right side, these nerves cross the artery at a nearly right angle.

5. The left subclavian is not embraced by the recurrent laryngeal nerve, like the right subclavian.

The thoracic duct bears an important relation to the left subclavian. It ascends from the chest to the left of the œsophagus and *behind* the artery; then arching behind the internal jugular vein as high as the seventh cervical vertebra, it curves downward and forward in front of the scalenus anticus to terminate in the subclavian vein at its junction with the jugular. The duct is so thin and transparent that it easily escapes observation; it is most

readily found by raising the subclavian vein near its junction with the jugular, and searching with the handle of the scalpel on the inner side of the scalenus anticus, in front of the vertebral vein.

Before tracing the branches of the subclavian artery, consider some points relating to the operation of tying it.

To tie the artery in the first part of its course, namely, on the inner edge of the scalenus anticus, is an operation of great difficulty and danger, even with the parts in a normal position. The great depth at which the artery is placed, the size and close proximity of its numerous branches, the large veins by which it is covered, its connection with the pneumogastric, recurrent laryngeal, phrenic and sympathetic nerves, and, above all, its close contiguity with the pleura, form a combination of circumstances so formidable that one cannot be surprised the operation has never been performed with a favorable result. On the left side the operation is more difficult to perform than on the right, owing to the difference in the anatomical relation of the two sides.

In the second part of its course, between the scalene muscles, the artery is more accessible, although it is rarely ligatured in this situation. It would be necessary to divide the clavicular origin of the sterno-mastoid, the cervical fascia, and the scalenus anticus, to reach the vessel; the phrenic nerve and the subclavian vein would be the chief objects exposed to injury. This operation was performed first and with success by Dupuytren in the year 1819. More recently it has been performed by Dr. Warren, of Boston. The patient recovered, though the pleura was wounded.[1]

But in the last part of its course, that is, on the outer side of the scalenus, the artery may be tied with comparative facility. The incision should be made from three to four inches in length, parallel with the upper border of the clavicle. We divide the platysma, some of the supra-clavicular nerves, and the cervical fascia. The external jugular vein and its tributaries must be drawn to the outer side, or divided and tied at both ends.

The connective tissue should now be carefully cut through,

[1] *Med. Chirurg. Trans.*, vol. xxix, p. 25.

and the posterior belly of the omo-hyoid sought for, as it runs just above the clavicle. After clearing away some fat and cellular tissue, the outer border of the scalenus anticus must be felt for, behind which the artery will be found lying upon the first rib. The operator now passes his finger downward along the outer border of this muscle, as far as its insertion into the tubercle of the first rib, which can always be distinctly felt. The artery having been exposed by carefully dividing a layer of fascia immediately covering the vessel, the ligature is to be passed round the artery from above downward, care being taken not to include in the ligature one of the cords of the brachial plexus.

Mr. Ramsden, of St. Bartholomew's Hospital, was the first who tied the subclavian in the third part of its course, in the year 1809; since that time the operation has been repeatedly performed, with very favorable results.

In the hands of a surgeon possessed of a practical knowledge of anatomy the operation is easy, provided all circumstances be favorable; but circumstances are often very unfavorable. Anatomical deviations are by no means rare, and it often happens that the aneurismal or other tumor, on account of which the operation is performed, raises the clavicle beyond its natural level, and so disturbs the parts, that to expose the artery and place a ligature around it becomes exceedingly difficult. Under such circumstances one cannot be surprised that even distinguished anatomists have committed mistakes. Sir Astley Cooper[1] failed in one instance. Dupuytren perforated the artery with the point of the needle, and included one of the nerves in the ligature: fatal hemorrhage was the result.[2] We were present at an operation in which the large nerve (a branch of the brachial plexus) which runs parallel with and on a plane anterior to the artery, was mistaken for it and tied, the surgeon being deceived by the pulsation communicated to the nerve.

The description of the means whereby the collateral circulation is maintained is deferred until the branches of the subclavian have been made out and described.

[1] *London Medical Review*, vol. ii, p. 300.

[2] *Edinburgh Med. and Surg. Journal*, vol. xvi, 1820.

Branches of the Subclavian Artery.

The branches of the subclavian extend so widely, that in the present dissection we can trace them only for a short distance. They are four in number—

1. The vertebral.

2. The thyroid axis, a short, thick trunk, which gives off the inferior thyroid, supra-scapular and posterior scapular.

3. The internal mammary.

4. The superior intercostal, which gives off the deep cervical.

As a rule, the vertebral, the thyroid axis and the internal mammary are given off from the subclavian in the first part of its course, and the superior intercostal in the second part. The most frequent deviation is, that the posterior scapular (transversalis colli) arises from the subclavian in the third part of its course.[1] On the left side the superior intercostal is frequently given off in the first part of the course of the subclavian.

Vertebral Artery.

This, the first and largest branch, arises from the upper and back part of the subclavian. It ascends in the neck, and for a short distance lies in the interval between the scalenus anticus and the longus colli. Here it enters the foramen in the transverse process of the sixth cervical vertebra, and ascends through the foramina in the transverse processes of the succeeding vertebræ. In the interval between the axis and the atlas, the artery makes a sigmoid curve, that it may not be stretched in the rotation of the head. Having traversed the foramen of the atlas, the artery curves backward along the groove in its arch, perforates the posterior occipito-atlantal ligament and the dura mater, then enters the skull through the foramen magnum, and unites with its fellow near the lower border of the pons Varolii to form the basilar artery.

Directly after the artery is given off from the subclavian, it lies behind the internal jugular vein, the inferior thyroid artery, and

[1] With reference to the origin of the posterior scapular (transversalis colli) artery, we made special observations during the winter session of 1858–59. We found that this artery was given off most frequently, not by the thyroid axis, but by the subclavian in the *third* part of its course. Under these circumstances the superficialis colli a. generally came from the thyroid axis.

the vertebral vein, and, on the left side, behind the thoracic duct. As it lies upon the groove on the neural arch of the atlas, it is separated from it by the suboccipital nerve, and is situated within the suboccipital triangle. After it has passed through the foramen magnum, the artery turns round the medulla oblongata, and is placed between the hypoglossal nerve and the anterior root of the suboccipital nerve.

The vertebral artery is accompanied by slender nerves from the inferior cervical ganglion of the sympathetic. These nerves communicate with the spinal nerves forming the brachial plexus.

Destined for the brain, the vertebral gives off no branches in the neck, except a few small muscular ones to the deeply-seated muscles, and which anastomose with the deep cervical, ascending cervical, and occipital arteries; it furnishes, however, *lateral spinal* branches to the spinal cord and its membranes which pass through the intervertebral foramina.

Each spinal branch divides into two branches, one passing along the root of the spinal nerve, is distributed to the spinal cord and its membranes; the other ramifies over the posterior surface of the body of the vertebra.

The cranial branches of the vertebral artery are mentioned at length in the description of the arteries of the brain.

The *vertebral vein* is formed by small branches from the muscles near the foramen magnum. It descends in front of the artery through the foramina in the transverse processes, and emerging through the transverse process of the sixth, crosses the subclavian artery and joins the brachio-cephalic vein, its orifice being guarded by a single or a double valve. It receives the veins from the neighboring muscles—the dorsi-spinal veins, veins from the spinal canal, the deep and ascending cervical, and the first intercostal veins. In some subjects it communicates with the lateral sinus by a branch through the posterior condylar foramen.

The cervical nerves pass through the intervertebral foramina behind the vertebral artery, so that the artery runs *behind* its vein, and in *front* of the nerves.

Thyroid Axis. The *thyroid axis* arises from the subclavian near the inner edge of the scalenus anticus, and after a course of a quarter of an inch divides into three branches, which take different directions; namely, the inferior thyroid, the supra-scapular, and the posterior scapular.

1. The *inferior thyroid* artery ascends tortuously behind the sheath of the common carotid and the sympathetic nerve, to the deep surface of the thyroid body, in which it communicates freely with the superior thyroid and with its fellow. Besides small branches to the trachea, the œsophagus, and the larynx, it gives off—

The *ascending cervical* artery which runs up close to the spine, between the scalenus anticus and the rectus capitis anticus major, and terminates in small branches, some of which supply these muscles; others enter the intervertebral foramina, and supply the spinal cord and its membranes. It anastomoses with the vertebral and ascending pharyngeal arteries.

2. The *supra-scapular* artery (transversalis humeri) runs outward over the scalenus anticus, covered by the sterno-mastoid m., then directly *beneath* and parallel with the clavicle; crossing over the third part of the subclavian artery, it passes beneath the posterior belly of the omo-hyoid to the superior border of the scapula. Here it is covered by the trapezius, passes *above* the transverse ligament which bridges over the notch; it gives off some branches which ramify in the supra-spinous fossa, and a large communicating branch which passes behind the neck of the scapula to reach the infra-spinous fossa, and inosculates freely in the infra-spinous fossa with the dorsalis scapulæ, a branch of the sub-scapular, and with the posterior scapular artery. Near the notch, it is joined by the supra-scapular nerve, which runs *through* it. The branches of this artery are numerous but small, and are as follows: the *inferior sterno-mastoid* (p. 90); the *supra-acromial*, which anastomoses with the acromio-thoracic artery; *articular* branches to the shoulder-joint; the *infra-spinous*, which ramifies in the infra-spinous fossa; and the *sub-scapular*, which ramifies in the substance of the subscapularis muscle.

3. The *transversalis colli* artery, of which the normal origin is

said to be from the thyroid axis, very frequently arises from the subclavian in the last part of its course. It is larger than the preceding artery, and runs tortuously across the side of the neck (higher than the supra-scapular), over the scalene muscles and the great nerves of the brachial plexus (sometimes between them), and divides into two branches, the superficial cervical and the posterior scapular. The *posterior scapular* disappears beneath the trapezius and the levator anguli scapulæ to reach the superior angle of the scapula. It then runs beneath the rhomboid muscles, which it supplies, down to the inferior angle of the scapula, anastomosing freely with the terminations of the supra- and sub-scapular arteries, and with the posterior branches of some of the intercostal arteries. The *superficial cervical* is given off in the space between the sterno-mastoid and trapezius. This vessel proceeds tortuously across the posterior triangle of the neck to the under surface of the trapezius, to which, with the levator anguli scapulæ, it is principally distributed.

The *superficialis colli* often comes *direct* from the thyroid axis.

The *veins* corresponding to the supra-scapular and posterior scapular arteries terminate in the external jugular, sometimes in the subclavian. The middle thyroid vein crosses in front of the common carotid artery, and joins the internal jugular.

Internal Mammary. This artery arises from the subclavian opposite to the thyroid axis. It descends slightly inward behind the clavicle and the subclavian vein, and enters the chest between the cartilage of the first rib and the pleura. It then passes behind the costal cartilages about half an inch from the border of the sternum. Its further course will be examined in the dissection of the chest. The corresponding vein, which results from the union of the two venæ comites, most frequently terminates in the brachio-cephalic vein.

Superior Intercostal. This artery is given off by the subclavian behind the scalenus anticus on the right side, and to its inner side on the left, so that you must divide the muscle to see it. It enters the chest behind the pleura, to the outer side of the first dorsal ganglion of the sympathetic. It runs over the necks of the first and second ribs, and furnishes the arteries of

the two upper intercostal spaces, and a posterior branch which is distributed to the muscles of the back and the spinal cord. It usually inosculates with the first intercostal branch of the aorta. The corresponding *vein* terminates on the right side in the vena azygos major; on the left in the brachio-cephalic.

Deep Cervical Artery. This artery arises from the superior intercostal, seldom direct from the subclavian. It goes to the back of the neck between the first rib and the transverse process of the seventh cervical vertebra, and ascends between the complexus and the semi-spinalis colli, both of which it supplies. It sometimes inosculates with the princeps cervicis, a branch of the occipital (p. 125).

FIG. 31.



DIAGRAM TO SHOW THE INOSCULATIONS OF THE SUBCLAVIAN ARTERY.

To test your knowledge of the branches of the subclavian artery, reflect upon the answer to the following question: "If the artery were tied in the *first* part of its course before it gives off any branches, how would the arm be supplied with blood?" The answer is, by six collateral channels, as follows: 1. By the communications between the superior and inferior thyroid; 2. Between the two vertebral: 3. Between the internal mammary and the intercostals and the epigastric; 4. Between the thoracic branches of the axillary and the intercostal branches of the

aorta; 5. Between the superior intercostal and the aortic intercostals; 6. Between the princeps cervicis and the deep cervical. Most of these inosculations are shown in the diagram (p. 141).

Again, if the subclavian were tied in the *third* part of its course, the circulation would be carried on by the communications: 1. Between the supra-scapular and the dorsalis scapulæ, a branch of the subscapular; 2. Between the supra-acromial branch of the supra-scapular and the acromio-thoracic; 3. Between the posterior scapular and the subscapular and dorsalis scapulæ; 4. Between the internal mammary, the aortic intercostals and superior intercostal on the one hand, and the long and short thoracic branches of the axillary, on the other.

Subclavian Vein. The subclavian vein does not form an arch like the artery, but proceeds in a nearly straight line over the first rib to join the internal jugular. It extends from the outer margin of the first rib to midway between the inner border of the scalenus anticus and the sterno-clavicular articulation, where it joins the internal jugular to form the brachio-cephalic vein. Throughout its whole course the vein is situated on a plane anterior to and a little lower than the artery, from which it is separated by the scalenus anticus, the phrenic and pneumogastric nerves. It has a pair of valves just before its junction with the internal jugular. It receives the anterior jugular, the external jugular, and through it, the supra-scapular and posterior scapular veins.

Brachial Plexus of Nerves. The large nerves forming the plexus which supplies the upper extremity are the anterior divisions of the four lower cervical and the larger portion of the first dorsal, with a small fasciculus derived from the fourth cervical nerve. Emerging from the intervertebral foramina the nerves appear between the anterior and middle scalene muscles, and pass with the subclavian artery into the axilla. In the neck the nerves have no plexiform arrangement, and it is only in the axilla that they branch and communicate largely with each other, and form the *brachial plexus* of nerves. The nerves in the neck are wide and are situated higher than the subclavian artery, and nearly on the same plane; but as they descend beneath the

clavicle, they converge and form large communications with each other, thus constituting the brachial plexus which completely surrounds the artery; one cord lying to the outer side, a second lying to the inner side, and a third behind the vessel.

The plexus is crossed superficially by the omo-hyoid muscle, and by the supra-scapular and posterior scapular arteries, and their corresponding veins.

The arrangement of the nerves in the formation of the plexus is very variable, and often not alike on both sides. The most usual arrangement is, that at the outer border of the scalenus anticus the fifth and sixth cervical nerves unite to form an upper trunk; the eighth and the first dorsal n. form a lower trunk; the seventh cervical runs for some distance alone, and forms a middle trunk. Now each of these four upper primary nerves divides into an anterior and a posterior branch; the anterior branches given off from the fifth, sixth, and seventh form the *outer cord* of the plexus; the anterior branches given off from the eighth cervical and first dorsal form the *inner cord;* while the posterior branches of all the nerves (namely, the fifth, sixth, seventh, and eighth cervical) unite to form the *posterior cord.*[1]

The branches arising from the plexus are best arranged into those given off above the clavicle, and those given off below it. The following are those given off *above* the clavicle :—

a. The branch forming one of the roots of the phrenic arises from the fifth cervical. (Not in the diagram.)

b. Nerve to the subclavius m.—This proceeds from the fifth and sixth cervical, and crosses the subclavian artery in the third part of its course. It frequently sends a filament, which passes in front of the subclavian vein to join the phrenic nerve.

c. Nerves to the *scaleni* and the *longus colli* muscles are given off from the lower cervical nerves as they leave the intervertebral foramina.

[1] Very frequently the posterior branch of the eighth cervical nerve does not, strictly speaking, form part of the posterior cord, but is continued on as a separate fasciculus to form part of the musculo-spiral nerve. For a description of the arrangement of the nerves constituting the plexus, see a paper, by Lucas, *Guy's Hospital Reports*, 1875; also Turner in the *Journal of Anatomy*, 1872.

d. Nerve to the rhomboid muscles.—This arises from the fifth cervical nerve, passes through the scalenus medius, and accompanies the posterior scapular artery, beneath the levator anguli scapulæ, which, as well as the rhomboid muscles, it supplies.

e. The *supra-scapular* nerve arises from the cord formed by the fifth and sixth cervical n., runs to the upper border of the scapula, where it meets with the corresponding artery, and then passes

FIG. 32.

DIAGRAM OF THE FORMATION OF THE BRACHIAL PLEXUS AND ITS BRANCHES.

C 4–8. Anterior trunks of the cervical nerves.
D 1. Anterior trunk of the first dorsal n.
9. N. to the rhomboid m.
10. Supra-scapular.
11. N. to subclavius m.
12–13. Anterior thoracic.
14, 15, 16. Subscapular n.
17. Lesser int. cutaneous.
18. Musculo-cutaneous.
19. Circumflex.
20. Median.
21. Musculo-spiral.
22. Ulnar.
23. Int. cutaneous.
24. Ext. respiratory of Bell.

through the notch in the scapula. In the supra-spinous fossa it gives off two branches to the supra-spinatus m. and an upper articular branch to the shoulder; it then descends behind the acromion process to the infra-spinous fossa, distributing a branch to the infra-spinatus muscle, and a lower articular filament to the shoulder-joint.

f. The *posterior thoracic nerve* (called *external respiratory* by Sir C. Bell) to the *serratus magnus* arises from the fifth and sixth cervical (sometimes also from the seventh) in the substance of the scalenus medius. It passes through this muscle and subsequently emerges below the rhomboid nerve; it then descends behind the brachial plexus and the subclavian vessels to the outer surface of the serratus magnus, to the several digitations of which it is exclusively distributed.

g. An *articular branch* is distributed to the shoulder-joint; besides some filaments to the constituent bones.

It only remains to be observed that the upper cord of the brachial plexus receives a branch from the lower cord of the cervical, and that each of its component nerves communicates by slender filaments with the sympathetic.

Below the clavicle the plexus gives off branches for the supply of the arm; namely, from the outer cord, the external anterior thoracic (to the pectoralis major), the musculo-cutaneous, and the outer head of the median; from the inner cord, the internal anterior thoracic n. (to the pectoralis minor), the inner head of the median, the ulnar, the internal cutaneous, and the lesser internal cutaneous (nerve of Wrisberg) nerves; from the posterior cord, the three subscapular (to the subscapularis, the latissimus dorsi, and teres major), the circumflex (to the deltoid and teres minor) and the musculo-spiral nerves; all of which will be described more fully in the dissection of the upper extremity.

TEMPORAL AND PTERYGO-MAXILLARY REGIONS.

In this dissection, the parts should be examined in the following order:—

1. Superficial and deep fasciæ.
2. Superficial arteries and nerves of the temple.
3. Masseter muscle.
4. Temporal aponeurosis.
5. Temporal muscle.
6. Pterygoid muscles.
7. Internal maxillary artery and branches.
8. Inferior maxillary nerves and branches.

To expose the temporal region, the skin of the temple should

be reflected from below upward. Beneath the skin you come upon a layer of tough connective tissue, continuous, above, with the aponeurosis of the scalp; below, with the fascia covering the masseter and the parotid gland. In this tissue are contained the superficial temporal vessels and nerves.

Temporal Artery. This is the smaller of the two terminal branches of the external carotid. Arising in the substance of the parotid gland near the neck of the jaw, it passes *over* the root of the zygoma, close to the meatus auditorius externus, ascends for about 1½ inches on the temporal fascia, and there divides into an anterior and a posterior branch. Above the zygoma it is superficial, being covered only by the attrahens aurem and a strong layer of fascia; here it is accompanied by branches of the facial nerve, and by the auriculo-temporal branch of the inferior division of the fifth nerve. It gives off the following branches:—

a. Several small branches to the parotid gland, the temporo-maxillary articulation, and the masseter.

b. The *transversalis faciei* (p. 58).

c. The *anterior auricular* branches, two in number, *superior* and *inferior*, ramify on the front of the pinna of the ear, inosculating with branches of the posterior auricular.

d. The *middle temporal*, a small vessel given off while the artery is still in the parotid gland, pierces the temporal fascia above the zygoma, and running in the substance of the temporal muscle, anastomoses with the temporal branches of the internal maxillary.

Of the two branches into which the temporal divides, the *anterior* runs tortuously toward the external angle of the frontal bone, distant from it about an inch. Its ramifications extend over the forehead, supplying the orbicularis and occipito-frontalis m., and inosculate with the supra-orbital and frontal arteries. The *posterior* runs toward the back of the head, and inosculates freely with the occipital and posterior auricular. The anterior branch, although the smaller, is usually selected for arteriotomy, the posterior being covered by a strong and unyielding fascia.

The *temporal vein* is formed by the junction of the veins accom-

panying the terminal branches of the temporal artery, which are situated superficial to the arteries; just above the zygoma it is joined by the middle temporal vein which takes its origin from a plexus in the temporal fossa. The common temporal vein, formed by the union of these three veins, passes over the zygoma, enters the parotid gland, and joins the internal maxillary vein to form the temporo-maxillary vein.

Auriculo-temporal Nerve. This nerve supplies the temple and side of the head with common sensation. It arises, close to the foramen ovale, from the third division of the fifth pair by two roots (between which the middle meningeal artery runs). From its origin it proceeds outward beneath the external pterygoid, between the neck of the jaw and the internal lateral ligament. It then ascends beneath the parotid, over the root of the zygoma, where it accompanies the temporal artery, and divides, like it, into an *anterior* and a *posterior* branch.

The posterior branch is the smaller of the two; the anterior forms communications with the temporal branches of the facial, and the orbital branch of the superior maxillary. The ramifications of the nerve correspond with those of the artery.

Near their origin the roots of the nerve are connected by fine filaments with the otic ganglion, and close to the condyle of the jaw the nerve sends round the external carotid artery *two communicating branches* to the temporo-facial branch of the facial nerve. It here distributes *parotid branches* to the gland; *articular branches* to the temporo-maxillary articulation, to the meatus auditorius and the membrana tympani. Above the zygoma it gives off *two auricular filaments;* the *upper* ramifies in the skin of the outer aspect of the ear, mainly on the tragus and upper half of the auricle; the *lower* supplies the lobule and lower part of the pinna.

Lastly, in the subcutaneous tissue of the temple, we find the temporal branches of the facial nerve, which supply the frontalis, the attrahens aurem, the orbicularis palpebrarum, tensor tarsi, and corrugator supercilii.

Masseter Muscle. This muscle *arises* from the lower edge of the zygoma, and is *inserted* into the outer side of the

ramus and coronoid process of the jaw. The masseter is composed of superficial and deep fibres which cross like the letter X. The *superficial* fibres, constituting the principal part of the muscle, arise from the anterior two-thirds of the zygoma and the malar process of the superior maxilla, by tendinous fibres which occupy the front border of the muscle, and send aponeurotic partitions into its substance. These fibres pass downward and backward, this direction giving them greater advantage, and are inserted into the angle and part of the ramus of the jaw. The *deep* fibres, mainly muscular (which are concealed by the parotid gland), arise from the posterior third of the zygoma, incline forward, and are inserted into the upper half of the ramus and the coronoid process. Besides these, a few fibres, arising from the inner surface of the zygoma, are inserted into the coronoid process and the tendon of the temporal muscle. Its *action* is to raise the jaw and help to masticate the food. Its nerve comes from the inferior maxillary.

The following objects lie superficial to the masseter: 1. Zygomatici major and minor; 2. Orbicularis palpebrarum; 3. Glandula socia parotidis and parotid duct; 4. Transversalis faciei artery; 5. Facial artery and vein; 6. Branches of the facial nerve.

Temporal Fascia. This strong, shining, aponeurotic membrane covers the temporal muscle; its chief use being to give additional origin to its fibres. It is attached above to the temporal ridge, and increasing in thickness as it descends, divides near the zygoma into two layers, which are attached to the outer and inner borders of the zygomatic arch. These layers are separated by fat, in which is found a filament from the orbital branch of the superior maxillary nerve, and the orbital branch of the temporal artery. The density of this aponeurosis explains why abscesses in the temporal fossa rarely point outward; the pus generally makes its way beneath the zygoma, into the mouth.

Reflect the aponeurosis, and notice that it is separated from the temporal muscle, near the zygoma, by fat. The absorption of this fat and the wasting of the muscle occasion the sinking of the temple in emaciation and old age.

Dissection. Divide the zygomatic arch on each side of the masseter, and turn it downward, taking care of the masseteric nerve and artery which enter its under aspect. Observe the direction of the superficial and deep fibres, and the tendinous partitions which augment the power of the muscle by increasing its extent of origin. The masseteric nerve and artery enter the under surface of the muscle near to its posterior border, through the sigmoid notch of the jaw; the artery comes from the internal maxillary, the nerve from the motor division of the inferior maxillary.

Temporal Muscle. This broad, fan-shaped muscle *arises* from the whole of the temporal fossa (except the malar surface) and the deep surface of the temporal fascia. Its fibres converge to a strong tendon, which is inserted into the inner surface, the apex, and anterior border of the coronoid process, as far forward as the last molar tooth.

The fibres of the muscle, converging from their wide origin, pass under the zygomatic arch, and terminate upon their tendon, the outer surface of which is partially concealed by the insertion of those fibres which come from the temporal aponeurosis; remove them, and see how this tendon radiates into the muscle like the ribs of a fan. Its nerves (two deep temporal) are branches of the inferior maxillary (p. 158).

Between the posterior border of this muscle and the neck of the inferior maxilla, the masseteric nerve and artery pass to their destination; in front of the muscle, the buccal branch of the inferior maxillary nerve descends to the buccinator with its companion artery.

The temporal muscle is in relation on its deeper surface with the external pterygoid and buccinator muscles, the internal maxillary artery and vein, and the deep temporal arteries and nerves.

Pterygo-maxillary Region. The zygomatic arch having been already divided, the structures should be cleaned so as to expose the coronoid process of the jaw, the insertion of the temporal muscle, and the loose fat which surrounds it. Next, saw through the coronoid process in a direction downward and

forward, so as to include the insertion of the muscle, and reflect it upward without injuring the subjacent vessels and nerves.

Dissection. To gain a good view of the muscles, nerves, and vessels of the pterygo-maxillary region, a portion of the ascending ramus of the jaw must be removed with a Hey's saw, as shown in the diagram on the next page.

In this region we have to examine the two pterygoid muscles, the trunk and branches of the internal maxillary artery, the inferior maxillary nerve, and the internal lateral ligament of the lower jaw. All these structures are imbedded in loose, soft fat, which must be cautiously removed without injuring them.

External Pterygoid. This muscle *arises* by two heads, one, the upper, from the great wing of the sphenoid and from the ridge below it; the lower, from the outer surface of the external pterygoid plate, a few fibres taking origin from the outer side of the tuberosities of the palate and superior maxillary bones. The muscle passes horizontally backward and is *inserted* into the neck of the jaw, and slightly into the border of the inter-articular fibro-cartilage of the temporo-maxillary articulation.

The advantage of the insertion of some of its fibres into the inter-articular cartilage is, that the cartilage follows the condyle in all its movements. When the jaw is dislocated, it is chiefly by the action of this muscle, which draws the condyle forward into the zygomatic fossa; the inter-articular cartilage being dislocated with the condyle.

By its deep surface the muscle is in relation with the internal pterygoid m., the internal lateral ligament, the arteria meningea media, the auriculo-temporal, the gustatory, the inferior dental, and chorda tympani nerves, and occasionally with the internal maxillary artery. Between its two heads of origin the buccal and anterior deep temporal nerves emerge.

Internal Pterygoid. This muscle *arises* by musculo-tendinous fibres from the inner surface of the external pterygoid plate of the sphenoid bone and from that portion of the tuberosity of the palate bone which forms the lower part of the pterygoid

fossa, also by a smaller slip in front of the external pterygoid from the external surface of the tuberosities of the palate and superior maxillary bones. It is *inserted* into the rough surface on the inner side of the angle of the lower jaw, as high as the dental foramen.

The internal pterygoid is in relation superficially with the external pterygoid, the internal lateral ligament, the internal maxillary artery and vein, the inferior dental vessels and nerve,

FIG. 33.



PTERYGOID MUSCLES AND INTERNAL MAXILLARY ARTERY.

the mylo-hyoid artery and nerve, the chorda tympani and the buccal nerves; by its deep surface, with the tensor palati and superior constrictor muscles.

Notice particularly the direction of the fibres of the pterygoid muscles. The fibres of the external run horizontally outward and backward from their origin; the fibres of the internal run

downward, backward and outward from their origin. The internal pterygoid has tendinous septa like the masseter. Both the pterygoids get their nerves from the motor division of the inferior maxillary nerve.

The internal pterygoid raises the lower jaw, acting in concert with the temporal and masseter muscles; it moreover assists the external pterygoid and anterior part of the masseter to draw the jaw forward. The external pterygoid draws the jaw forward and somewhat to the opposite side, and also in conjunction with the internal pterygoid produces the lateral movements of the jaw essential to the mastication of the food. Consequently they are enormously developed in all ruminants and comparatively feebly in carnivorous animals. The antagonistic muscles of the forward action of the two pterygoids are the temporal m. and the deep fibres of the masseter.

Dissection. Saw through the neck of the jaw, disarticulate the condyle with its fibro-cartilage from the glenoid cavity, and turn it forward with the external pterygoid, so that the condyle can be replaced if desirable. A little dissection will bring into view the internal lateral ligament, the internal maxillary artery and vein, the inferior maxillary nerve and its branches, and the chorda tympani nerve.

Internal Maxillary Artery. This is the larger of the two terminal branches into which the external carotid divides, opposite the neck of the jaw in the parotid gland. It passes horizontally forward between the neck of the jaw and the internal lateral ligament, then runs tortuously, in some cases above, in others beneath, the external pterygoid, enters the spheno-maxillary fossa between the two heads of the external pterygoid, where it terminates by dividing into numerous branches.

The course of this artery is divided into three stages. In the first, the artery lies between the neck of the jaw and the internal lateral ligament; in the second, it lies either over or under the external pterygoid; in the third, it lies in the spheno-maxillary fossa.

BRANCHES OF THE INTERNAL MAXILLARY ARTERY IN THE THREE STAGES OF ITS COURSE.

BRANCHES IN THE FIRST STAGE.	BRANCHES IN THE SECOND STAGE.	BRANCHES IN THE THIRD STAGE.
a. Tympanic.	Six to the five muscles of mastication, namely:—	*i*. Superior dental
b. Meningea media.	*e*. Masseteric.	*j*. Infra-orbital.
c. Meningea parva.	*f*. Anterior and posterior deep temporal.	*k*. Descending palatine.
d. Inferior dental.	*g*. External and internal pterygoid.	*l*. Vidian.
	h. Buccal.	*m*. Pterygo-palatine.
		n. Nasal or spheno-palatine.

Branches in the First Part.

a. The *tympanic* ascends behind the articulation of the jaw, and passes through the Glaserian fissure to the tympanum. It supplies that cavity and the membrana tympani, and anastomoses with the stylo-mastoid and Vidian arteries. It occasionally gives off a *deep auricular* branch which pierces the anterior wall of the external auditory meatus, supplying the skin of this canal. This artery is not infrequently given off from a branch of the internal maxillary artery.

b. The *middle meningeal* artery ascends between the two roots of the auriculo-temporal nerve, behind the external pterygoid, and enters through the foramen spinosum into the cranium, where it ramifies between the dura mater and the bones. In the skull it gives off small branches to the Gasserian ganglion; a *petrosal* branch passing through the hiatus Fallopii; *orbital* branches entering the orbit through the sphenoidal fissure; and *temporal* branches which pierce the great wing of the sphenoid to enter the temporal fossa. Its further course is described at p. 32.

c. The *meningea parva* (not marked in the plan) ascends through the foramen ovale into the skull, and supplies chiefly the ganglion of the fifth cranial nerve. It often comes from the meningea media.

d. The *inferior dental* artery descends behind the neck of the jaw to the dental foramen, which it enters with the dental nerve. It then proceeds through a canal in the diploë to the symphysis, where it minutely inosculates with its fellow. In this canal, which runs beneath the roots of all the teeth, the artery gives branches

which ascend through the little foramina in the fangs, and supply the pulp in their interior. Opposite the foramen mentale arises the mental branch already described (p. 66). Before entering the dental foramen the artery furnishes a small branch—*mylo-hyoid*—which accompanies the nerve proceeding to the mylo-hyoid muscle.

Branches in the Second Part.

e. The *masseteric* branch passes through the sigmoid notch of the jaw behind the temporal muscle to the under surface of the masseter, with the masseteric nerve, and inosculates with the facial and transverse facial arteries.

FIG. 34.



PLAN OF INTERNAL MAXILLARY ARTERY.

f. The *anterior* and *posterior deep temporal* arteries ascend to supply the temporal muscle, ramifying between the muscle and the bone, one near the front, the other near the posterior border of the muscle. They communicate with the superficial and middle temporal arteries, with the terminal branches of the lachrymal a., and with the temporal branches of the arteria meningea media.

g. The *pterygoid* branches supply the internal and external pterygoid muscles.

h. The *buccal* branch runs forward with the nerve to the buccinator, where it anastomoses with the facial artery.

Branches in the Third Part.

i. The *superior dental* branch runs along the tuberosity of the superior maxillary bone, and sends small arteries through the foramina in the bone to the pulps of the molar and bicuspid teeth. It also supplies the gums, and the mucous membrane of the antrum.

j. The *infra-orbital* branch ascends through the spheno-maxillary fissure, then runs forward along the infra-orbital canal with the superior maxillary nerve, and emerges upon the face at the infra-orbital foramen, beneath the levator labii superioris. In the infra-orbital canal the artery sends branches, *anterior dental*, downward through little canals in the bone to the incisor and canine teeth, and upward into the orbit to the lachrymal gland, the inferior oblique and inferior rectus. After issuing from the foramen it sends upward branches to the lachrymal sac, and descending branches to the upper lip. The former anastomose with the nasal branches of the ophthalmic and facial arteries; the latter with the superior coronary, transverse facial, and buccal arteries.

k. The *descending palatine*, a branch of considerable size, runs down the posterior palatine canal with the palatine nerve (a branch from Meckel's ganglion), and then along the roof of the hard palate, toward the anterior palatine canal, in which, much diminished in size, it inosculates on the septum nasi with a branch of the spheno-palatine artery. It supplies the gums, the glands and mucous membrane of this part, and furnishes branches to the soft palate.

l. The *Vidian*, an insignificant branch, runs backward through the Vidian canal with the Vidian nerve, and is distributed to the Eustachian tube, the pharynx, and the tympanum.

m. The *pterygo-palatine* is a small but constant branch which runs backward through the pterygo-palatine canal with the pharyngeal nerve from Meckel's ganglion, and ramifies upon the upper part of the pharynx and the Eustachian tube.

n. The *nasal* or *spheno-palatine* branch enters the nose through the spheno-palatine foramen in company with the nasal nerve from Meckel's (spheno-palatine) ganglion, and ramifies upon the spongy bones, the ethmoidal cells and the antrum. One large

branch, the *artery of the septum*, runs along the septum nasi toward the anterior palatine canal, where it joins the descending palatine artery.

Observe that all the branches of the internal maxillary artery in the first and third parts of its course traverse bony canals; while the branches in the second part go directly to muscles.

The *internal maxillary vein* is formed by the veins corresponding to the branches of the artery. As the vein lies between the temporal and external pterygoid muscles it forms a plexus—*pterygoid plexus*—which communicates, above, with the cavernous sinus by branches which come through the foramina at the base of the skull; in front it communicates with the facial vein. It joins the temporal in the substance of the parotid gland, and thus communicates with the external jugular vein.

Pterygoid Plexus of Veins.

Inferior Maxillary Nerve and Branches.

This great nerve is the largest of the three divisions of the fifth cerebral nerve. It differs from the other two divisions, *i. e.* the ophthalmic and the superior maxillary, in that it contains motor as well as sensory filaments; the motor being furnished by the small non-ganglionic root of the fifth nerve. It is necessary to remember this point of its physiology, in order to understand its extensive distribution; for the sensory portion supplies the parts to which it is distributed with common sensation only, while the motor portion supplies all the muscles concerned in mastication.

The nerve, composed of sensory and motor filaments, emerges from the skull through the foramen ovale as a thick trunk, under the name of the inferior maxillary. It lies directly external to the Eustachian tube, and is covered by the external pterygoid muscle, which must be turned on one side to expose it. Immediately after its exit from the skull, the nerve divides into two parts, an anterior, or motor division, and a posterior or sensory division. From the *anterior* portion (chiefly motor) are derived branches distributed to the muscles of mastication and the buccal nerve. From the *posterior* (mainly sensory) come the following branches: the auriculo-temporal, gustatory, and inferior dental; there are also motor branches to the mylo-hyoid and anterior

belly of the digastricus. This apparent anomaly will be presently explained.

BRANCHES OF THE INFERIOR MAXILLARY NERVE.

ANTERIOR PORTION.	POSTERIOR PORTION.
Auriculo-temporal.	To temporal muscle.
Inferior dental.	To masseter.
Gustatory or lingual.	To external pterygoid.
Mylo-hyoideus.	To internal pterygoid.
Anterior belly of digastricus.	To buccal.

The *deep temporal branches*, two in number, *anterior* and *posterior*, pass outward close to the great wing of the sphenoid bone, and ascend with the temporal arteries to the temporal muscle. A *middle temporal* nerve is not infrequently present, and ascends beneath the temporal muscle to enter its deeper aspect. The posterior branch is occasionally joined with the masseteric nerve, the anterior with the buccal nerve.

The branch to the *masseter* runs outward above the external pterygoid, through the sigmoid notch of the jaw, to the under surface of the muscle.

The branch to the *external pterygoid* comes, apparently, from the buccal nerve in its passage through this muscle.

The branch to the *internal pterygoid* muscle proceeds from the inner side of the main trunk, close to the otic ganglion, and descending between the internal pterygoid and the tensor palati, enters the inner and deeper aspect of the muscle.

The *buccal* branch, a sensory nerve, united at its origin with the anterior deep temporal and external pterygoid nerves, passes either above or between the fibres of the external pterygoid to the buccinator, where it spreads out into filaments, which form a plexus with the buccal branches of the facial nerve, and then supply the skin, mucous membrane, and glands of the cheek with common sensation. The motor power of the buccinator, remember, is derived from the facial nerve. That this buccal branch is mainly sensory is proved by the action of the muscle still continuing when the motor division of the fifth nerve is paralyzed. The evidence is corroborated by a case in which this

buccal branch proceeded from the second division of the fifth nerve; no communication being discovered, after very careful dissection, between it and the motor root of the third division.[1]

The *auriculo-temporal* branch arises by two roots which embrace the middle meningeal artery before it enters the skull. The nerve runs backward behind the external pterygoid and the neck of the jaw, ascends at first beneath the parotid gland, then over

FIG. 35.



PLAN OF THE BRANCHES OF THE INFERIOR MAXILLARY NERVE.

the root of the zygoma with the temporal artery, and divides, like it, into an anterior and a posterior branch. The posterior branch supplies the pinna and surrounding tissues; the anterior is distributed to the skin covering the vertex and temporal region, communicating with the temporal branches of the facial nerve and the orbital branch of the superior maxillary.

[1] Turner, "On the Variation of the Buccal Nerve." *Journal of Anat. and Phys.*, No. 1, 1866.

The auriculo-temporal communicates at its origin with the otic ganglion, and then ascends behind the jaw with the temporal branches of the facial n.; it also gives off an *articular branch* to the temporo-maxillary joint; *two branches* to the meatus auditorius and the membrana tympani; *parotid branches* to the gland; *auricular branches*, two in number—an *inferior*, which is distributed to the ear below the auditory meatus, and a *superior* to the tragus and auricle. Its branches have been described (p. 22).

The *inferior dental* branch emerges beneath the external pterygoid, and descends between the ramus and the internal lateral ligament of the jaw to the dental foramen, which it enters with the dental artery. It then runs in the canal in the diploë of the jaw and furnishes filaments which ascend through the canals in the fangs of the teeth to the pulp in their interior. Opposite the foramen mentale it divides into two branches, the *mental* and *incisor*. Observe that the same nerve which supplies the teeth supplies the gums; hence the sympathy between them.

a. The *mylo-hyoid* branch, apparently arising from the dental, is derived from the motor root of the fifth, and may, with careful dissection, be traced to it. It leaves the sheath of the inferior dental nerve near the foramen in the jaw, and runs in a groove on the inner side of the ramus to the lower surface of the mylo-hyoid, which muscle it supplies together with the anterior portion of the digastricus.

b. The *dental* branches pass upward to the fangs of the molar and bicuspid teeth.

c. The *incisor* branch is the continuation of the nerve, and passes to the symphysis, supplying the canine and incisor teeth.

d. The *mental* branch (sometimes called *labial*) emerges through the foramen mentale, and soon divides into numerous branches; some ascend to the lower lip beneath the depressor labii inferioris, and communicate with the facial nerve; others pass inward to the skin of the chin.

The *gustatory* or *lingual nerve* lies at first behind the external pterygoid m., then descends obliquely forward between the ramus of the jaw and the internal pterygoid m., and subsequently for a short distance between the jaw and the superior constrictor of

the pharynx. Here it lies close under the mucous membrane of the mouth near the last molar tooth of the lower jaw. Division of it in this situation relieves pain in cancer of the tongue. The gustatory n. then rests upon the stylo-glossus and the hyo-glossus m., and after crossing Wharton's duct passes to the tip of the tongue.

The nerve at first lies in front of the inferior dental nerve (with which it is frequently connected), and beneath the internal maxillary a. Beneath the external pterygoid, the gustatory n. is joined at an acute angle by the *chorda tympani* (a branch of the facial). This branch emerges through a small canal, *canal of Huguier*, by the side of the Glaserian fissure, and passing behind the dental n., meets the gustatory, and runs along the lower border of this nerve to supply the submaxillary gland; part of it joins the submaxillary ganglion, and it is then eventually distributed to the lingualis muscle.

The gustatory nerve in its course gives off—

a. Communicating branches to the hypoglossal n., forming two or more loops at the anterior border of the hyo-glossus muscle.

b. Communicating branches to the submaxillary ganglion.

c. Branches to the mucous membrane of the mouth, gums, and sublingual gland.

d. Lingual branches which pass to the papillæ of the sides and tip of the tongue; here also we find communications between this nerve and the hypoglossal.

The duct of the submaxillary gland (p. 112), *Wharton's duct*, can now be traced to its termination. It passes from its under surface, runs forward under the mylo-hyoideus and upon the hyo-glossus muscle; it then passes beneath the gustatory nerve, and subsequently runs between the sublingual gland and the genio-hyo-glossus, to open into the floor of the mouth, by the side of the frænum linguæ. Its length is about two inches; its dimensions are not equal throughout; it is dilated about the middle, and contracted at the orifice. Saliva, collected in the dilated portion, is sometimes spirted to a considerable distance out of the narrow orifice, in consequence of the sudden contraction of the neighboring muscles.

The gland is supplied with nerves by branches from the submaxillary ganglion, from the sympathetic, and the mylo-hyoid nerves.

In the floor of the mouth there occasionally exists a cystic tumor, called a *ranula*, with semi-transparent walls, perceptible beneath the tongue. By some of the older writers it was looked upon as an abnormal dilatation of the submaxillary duct. There is, however, no reason for believing this swelling (except very rarely) to be connected with the duct. It is rather a cyst formed in the loose areolar tissue under the tongue, or is an enlargement of one of the small bursæ which normally exist in this situation. The character of the saliva presents no agreement with the fluid contained in these cysts, which is thickly glairy, like the white of an egg.

Internal Lateral Ligament of the Lower Jaw.

This so-called ligament (which is more like a layer of fascia) passes from the spinous process of the sphenoid bone to the inner side of the foramen dentale. Between this ligament and the neck of the jaw, we find the internal maxillary artery and vein, the auriculo-temporal nerve, the middle meningeal artery, the inferior dental nerve and artery, and a portion of the parotid gland.

At this stage of the dissection you will be able to trace the course and relations of the internal carotid artery. But before doing this, examine the several objects which intervene between the external and internal carotids. These are—1. The stylo-glossus; 2. The stylo-pharyngeus; 3. The glosso-pharyngeal nerve; 4. The stylo-hyoid ligament.

Stylo-glossus.

This *arises* from the front of the styloid process near the apex, and from the stylo-maxillary ligament. It passes at first downward and then horizontally forward, and is *inserted* along the side of the tongue as far as the tip, some of its lower fibres decussating with those of the hyo-glossus. Its action is to retract the tongue. Its nerve is a branch of the hypo-glossal.

Stylo-pharyngeus.

This *arises* from the inner side of the styloid process near the base, and is *inserted* into the

upper and posterior edges of the thyroid cartilage. It descends along the side of the pharynx between the superior and the middle constrictors; some of its fibres blend with the constrictor muscles; others join those of the palato-pharyngeus at its insertion. Curving round its lower border is seen the glosso-pharyngeal nerve, from which its nerve supply is derived. Its *action* is to raise the larynx with the pharynx in deglutition.[1]

Between the stylo-glossus and stylo-pharyngeus, and nearly parallel with both, is the *stylo-hyoid ligament.* It extends from the apex of the styloid process to the lesser cornu of the os hyoides. It is often more or less ossified.

The *ascending palatine artery*, a branch of the facial (p. 116), runs up between the stylo-glossus and the stylo-pharyngeus, and divides into branches which supply these muscles, the palate, the side of the pharynx, and the tonsils. It inosculates with the descending palatine, a branch of the internal maxillary.

Glosso-pharyngeal Nerve. The glosso-pharyngeal nerve is observed curving forward round the lower border of the stylo-pharyngeus (*see above*). It is the ninth cranial nerve; arises by five or six filaments from the groove between the olivary body and the restiform tract of the medulla oblongata, leaves the skull through the middle part of the foramen jugulare in a separate sheath of dura mater, in front of the pneumogastric and spinal accessory nerves, and descends between the internal jugular vein and the internal carotid artery. It then crosses in front of the artery below the styloid process, and proceeds along the lower border of the stylo-pharyngeus. At this point it curves forward over that muscle and the middle constrictor of the pharynx, and disappears beneath the hyo-glossus, where it divides into its terminal branches, which supply the mucous membrane of the pharynx, the back of the tongue, and the tonsils.

The glosso-pharyngeal is, at its origin, purely a sensory nerve. But soon after its exit from the skull it receives *communications*

[1] Varieties of this muscle are frequently met with, chiefly as supernumerary muscles. They arise variably from neighboring parts of the base of the skull close to the styloid process, and are inserted either into the pharyngeal constrictors or into the aponeurosis of the pharynx.

from the facial, the pneumogastric and the sympathetic, so that it soon becomes a compound nerve—*i. e.* composed of both sensory and motor filaments. At the base of the skull it presents two ganglia—the *jugular* and the *petrous* (ganglion of Andersch). The branches given off by the petrous ganglion will be dissected hereafter; at present the student can only make out the branches which this nerve gives off in the neck, namely:—

Carotid branches, which surround the internal carotid artery as far as its origin, and communicate with the pharyngeal branch of the pneumogastric and with the sympathetic.

Pharyngeal branches, three or four in number, which form by the side of the middle constrictor of the pharynx a plexus, the **pharyngeal plexus,** supplemented by filaments derived from the pneumogastric, the nervus accessorius, the external laryngeal, and the sympathetic. Its branches supply the constrictor muscles and the mucous membrane of the pharynx, the back of the tongue, and the tonsils.

Muscular branches which enter the stylo-pharyngeus m.

Tonsillar branches which are given to the soft palate and the fauces, and to the tonsils, forming a plexus (circulus tonsillaris).

Lingual branches, two in number, which are distributed to the base and lateral aspects of the tongue; one branch turns upward and is distributed to the papillæ circumvallatæ, and the mucous membrane of the posterior third of the tongue as far backward as the epiglottis; the other passes to the middle of the side of the tongue, communicating with the gustatory nerve.

The styloid process must now be cut through at its base, and turned forward, with the muscles arising from it. The internal carotid artery will thus be exposed in the cervical region, as far as the carotid canal. The part of the artery contained within the carotid canal will be described hereafter.

Internal Carotid Artery.

The *internal carotid artery* proceeds from the bifurcation of the common carotid at the upper border of the thyroid cartilage, and ascends vertically to the base of the skull *by the side of the pharynx*, in front of the transverse processes of the three upper cervical vertebræ. It enters the skull through the carotid canal in the temporal bone, runs tortuously

by the side of the body of the sphenoid, and terminates in branches which supply the orbit and the brain. It is divided into four portions—the cervical, petrous, cavernous, and cerebral. In the *cervical part* of its course, it is situated immediately to the outer side of the external carotid artery, behind the inner border of the sterno-mastoid. It soon gets beneath the external carotid, and lies deeply beneath the parotid gland, and ascends by the side of the pharynx and tonsil. It *lies upon* the rectus capitis anticus major, the superior laryngeal nerve, and the superior cervical ganglion of the sympathetic; to its *outer side* is the internal jugular vein and the pneumogastric nerve; to its *inner side* is the pharynx, the tonsil, and the ascending pharyngeal artery; it is *crossed*, successively, by the hypoglossal nerve, the occipital artery, the digastricus and stylo-hyoid muscles; higher up it is crossed by the styloid process, the stylo-glossus and stylo-pharyngeus muscles, by the glosso-pharyngeal nerve and the stylo-hyoid ligament, all of which last-named structures intervene between it and the external carotid.

The most important relation of the artery, in a surgical point of view, is, that it ascends close by the *side of the pharynx* and *tonsil*. In opening an abscess, therefore, near the tonsil, or at the back of the pharynx, be careful to introduce the knife with its point inward toward the mesial line; observe this caution the more, because in some subjects the internal carotid makes a curve, or even a complete curl upon itself, in its ascent near the pharynx. In such cases an undue deviation of the instrument in an outward direction might injure the vessel.

Ascending Pharyngeal Artery. This artery generally arises from the back part of the external carotid, about half an inch above the angle of the common carotid. It ascends in a straight course between the internal carotid artery and the side of the pharynx, toward the base of the skull, resting upon the rectus capitis anticus major. It gives off three sets of branches:—

a. Pharyngeal branches, three or four in number; the two lower supply the inferior and middle constrictors, and stylo-pharyngeus, anastomosing with the superior thyroid a.; the upper branch, the *palatine*, ascends upon the superior constrictor,

runs down with the levator palati, above the superior constrictor, and supplies the muscles of the palate, the Eustachian tube, and the tonsil.

b. Prevertebral branches, which supply the prevertebral muscles, the superior cervical ganglion of the sympathetic, the lymphatic glands, and the pneumogastric and hypoglossal nerves.

c. Meningeal branches, which supply the dura mater; passing through the foramen lacerum medium, the anterior condylar foramen, and the foramen jugulare with the internal jugular vein.

Pneumogastric Nerve.

The *pneumogastric nerve* is the tenth cranial nerve, and has the longest course of all the cerebral nerves. It arises from the medulla oblongata by a series of roots, from twelve to fifteen in number, from the front of the restiform body. It passes out of the skull in a common sheath of dura mater and arachnoid, with the nervus accessorius through the foramen jugulare.

Within the foramen jugulare a small ganglion—*ganglion of the root* (Arnold's ganglion)—about two lines in length, is situated upon the pneumogastric nerve, and is joined by a branch from the nervus accessorius. This ganglion will be described hereafter. About half an inch below the preceding the pneumogastric nerve swells out, and forms a second ganglion—*ganglion of the trunk*—(inferior ganglion), of a reddish-gray color. This ganglion occupies about an inch of the nerve, but does not involve the whole of its fibres, the branch from the spinal accessory joining the pneumogastric below the ganglion. It is united to the hypoglossal nerve, from which it receives filaments; it also receives filaments from the first and second spinal nerves, and from the superior cervical ganglion of the sympathetic.

Thus, the pneumogastric, at its origin probably a nerve of sensation only, becomes, in consequence of the connecting filaments from these various branches, a compound nerve, and in all respects analogous to a spinal nerve.

Leaving the skull at the foramen jugulare, the nerve descends in front of the cervical vertebræ, lying successively upon the rectus capitis anticus major and the longus colli. In the upper part of the neck it is situated, lying in the same sheath, between

the internal carotid artery and the internal jugular vein; lower down, it lies between and behind the common carotid and the internal jugular vein. It enters the chest, on the right side, crossing in front of the first part of the subclavian artery, nearly at a right angle; on the left, running nearly parallel with it.

In their course through the chest, the pneumogastric nerves have not similar relations. The *right* nerve lies beneath the subclavian vein, and then descending behind the right brachio-cephalic vein by the side of the trachea, is continued behind the right bronchus to the posterior part of the œsophagus. The *left* nerve passes behind the left brachio-cephalic vein, then crosses in front of the arch of the aorta, and behind the left bronchus to the anterior part of the œsophagus. Both nerves subdivide on the œsophagus into a plexus; the right nerve forming the *posterior œsophageal plexus*, the left the *anterior*. Each plexus again collects its fibres together to form a single trunk; thus two main nerves are formed which pass with the œsophagus through the diaphragm; of these the right is distributed over the posterior, the left over the anterior surface of the stomach.[1]

In their long course from the medulla oblongata to the abdomen, the pneumogastric nerves supply branches to most important organs: namely, to the pharynx, the larynx, the heart, the lungs, the œsophagus, the stomach, and the liver.

The branches of the pneumogastric are those of communication and those of distribution:—

1. The *branches of communication* are those in connection with the ganglion of the root and the ganglion of the trunk.

a. The *ganglion of the root* has connecting filaments with the accessory portion of the spinal accessory, the superior cervical ganglion of the sympathetic, and with the petrous ganglion of the glosso-pharyngeal.

b. The *ganglion of the trunk* has communicating filaments with the hypoglossal, the loop between the first two cervical nerves, and the superior cervical ganglion of the sympathetic.

[1] The differences in the course and destination of the right and the left pneumogastric nerves may be explained in the process of development. The student is therefore referred to works which treat of this subject.

2. The *branches of distribution* are—

a. The *auricular* (Arnold), which cannot at present be seen, will be made out in the dissection of the nerve at the base of the skull.

b. The *pharyngeal* arises from the upper part of the ganglion of the trunk, and, receiving a filament from the accessory part of the spinal accessory, descends either in front of or behind the internal carotid. The nerve, after passing to the inner side of the internal carotid, divides into branches, which with the other filaments (described p. 163) upon the middle constrictor muscle form the *pharyngeal plexus*. From this plexus branches are distributed to the muscles and the mucous membrane of the pharynx.[1]

c. The *superior laryngeal*, derived from the middle of the ganglion of the trunk, descends behind the internal carotid, and divides into two branches, the internal and the external laryngeal.

The *internal laryngeal* passes to the interval between the os hyoides and the thyroid cartilage, and enters the larynx (with the superior laryngeal a.), through the thyro-hyoid membrane, to be distributed, as a nerve of sensation, to the mucous membrane of the larynx and epiglottis. The *external laryngeal*, the smaller, gives off some branches to the pharyngeal plexus, the inferior constrictor, and the thyroid body, and then descends by the side of the larynx, beneath the depressors of the os hyoides, to supply the crico-thyroid muscle; it communicates with the superior cardiac nerve of the sympathetic.

d. The *cervical cardiac branches* descend behind the sheath of the carotid artery to the cardiac plexus. The *upper branches*, one or two in number, are small, and proceed from the ganglion of the trunk; they join the cardiac branches of the sympathetic and the deep cardiac plexus; the *lower* comes from the trunk of the pneumogastric before it enters the chest. Subsequently, the right lower cardiac nerve descends by the side of the innominate artery to join the deep cardiac plexus; the left passes over the arch of the aorta to join the superficial cardiac plexus.

[1] A branch from the plexus is described by Luschka as receiving filaments from the pharyngeal branches of the glosso-pharyngeal and pneumogastric nerves, and joining with the hypoglossal as it winds round the occipital artery.

e. The *inferior* or *recurrent laryngeal nerve* turns, on the right side, under the subclavian artery (p. 131), and ascends obliquely inward to the larynx behind the common carotid and the inferior thyroid arteries; it lies subsequently in the groove between the œsophagus and the trachea. On the left side, it turns under the arch of the aorta, just on the outer side of the remains of the ductus arteriosus; after which it runs up between the trachea and the œsophagus. On both sides the nerves enter the larynx beneath the lower border of the inferior constrictor, and supply all the intrinsic muscles of the larynx, except the crico-thyroid. These nerves as they turn under their respective vessels give off cardiac branches to the deep cardiac plexus. They supply also filaments to the trachea, œsophagus and inferior constrictor muscle.

The remaining branches of the pneumogastric nerve to the lungs, heart, œsophagus and stomach will be examined in the dissection of the chest.

Spinal Accessory Nerve.

The *spinal accessory nerve* issues through the middle part of the foramen jugulare, in a sheath of dura mater common to it and the pneumogastric nerve. It arises by numerous filaments from the side of the medulla oblongata below the pneumogastric, and from the lateral column of the spinal cord as low down as the sixth cervical vertebra. The filaments which arise from the medulla oblongata join to form the *accessory portion* of the nerve; the spinal filaments ascend between the ligamentum denticulatum and the posterior roots of the cervical spinal nerves, and form the *spinal portion* of the nerve. These portions converge to the jugular foramen, where they communicate with each other more or less, and are then continued onward below the jugular foramen as two portions—the internal or accessory, which joins the pneumogastric n.; the external or spinal, which is distributed to muscles.

The *accessory part*, within the foramen jugulare, sends one or more filaments to the ganglion of the root of the pneumogastric. It lies close to the pneumogastric nerve at the ganglion of the trunk, and is finally incorporated with the nerve below the ganglion. It sends filaments to the pharyngeal and superior laryngeal branches of the pneumogastric.

The *spinal part* separates from the accessory part below the foramen jugulare. It then takes a curved course backward and outward, lying in front of the internal jugular vein and the transverse process of the atlas, and behind the digastric and stylo-hyoid muscles. It pierces the upper part of the sterno-mastoid muscle accompanied by the superior sterno-mastoid artery, a branch of the occipital, and supplies the muscle, joining in its substance with branches from the third cervical n. The nerve then crosses obliquely the occipital triangle, where it communicates with the second and third cervical nerves. It is eventually distributed to the under aspect of the trapezius, where it is joined by branches from the third and fourth cervical nerves.

Hypoglossal Nerve. This nerve arises, by from ten to fifteen filaments, from the groove between the anterior pyramid and the olivary body. It passes through the dura mater in two fasciculi, which emerge from the skull through the anterior condylar foramen, and then unite to form a single nerve. It comes forward between the internal jugular vein and the internal carotid artery, where it is intimately connected with the pneumogastric nerve. Its further course has been described (p. 121).

In the anterior condylar foramen the hypoglossal gives off a small filament to the diploë and to the dura mater around the foramen magnum. At the base of the skull it gives off several branches, which connect it with the ganglion of the trunk of the pneumogastric nerve. These two nerves are sometimes almost inseparably united. It gives off also several delicate filaments to the superior cervical ganglion of the sympathetic, and communicates with the loop formed by the first two spinal nerves in front of the atlas.

Sympathetic Nerve. Now examine the cervical ganglia of the sympathetic system of nerves. This system consists of a series of ganglia arranged on each side of the spine, from the first cervical to the last sacral vertebra. The successive ganglia of the same side are connected by intermediate nerves, so as to form a continuous cord on each side of the spine; this constitutes what is called the trunk of the sympathetic system, and is connected with all the spinal nerves. Its upper or cephalic

extremity enters the cranium through the carotid canal, surrounds the internal carotid artery, communicates with the third, fourth, fifth and sixth cranial nerves, and joins its fellow of the opposite side upon the anterior communicating artery.[1] Its sacral extremity joins its fellow by means of the little *ganglion impar*, situated in the mesial line, upon the coccyx.

The ganglia are connected together by branches composed of gray and white nerve fibres; they are also connected with the spinal nerves by two filaments—one, of white nerve fibres which passes from the spinal nerve to the ganglion; the other, of gray, from the ganglion to the spinal nerve. Branches of distribution are also given off by the ganglia, some to the various blood vessels and viscera, forming intricate plexuses upon them; others to the various ganglia of the viscera—the cardiac and semilunar ganglia.

The different portions of the sympathetic gangliated cord receive, respectively, the distinguishing names of the cervical, dorsal, lumbar, sacral, and coccygeal. At present we have only to consider the cervical portion of it.

To expose the cervical ganglion of the sympathetic, the internal carotid artery, the pneumogastric, glosso-pharyngeal and hypoglossal nerves should be cut through, near the base of the skull; then by careful dissection the superior cervical ganglion can be traced out.

Cervical Ganglia of Sympathetic.

In the cervical portion of the sympathetic are three ganglia, named from their position, superior, middle and inferior.

The *superior cervical ganglion*, the largest of the three, is situated near the base of the skull, opposite the second and third cervical vertebræ, upon the rectus capitis anticus major, and lies behind and on the inner side of the internal carotid artery. It is of a reddish-gray color, like the other ganglia, of an elongated oval shape, varying in length from one to two inches.

To facilitate the description of its several branches, we divide them into an upper, a lower, an external, an internal, and an anterior set:—

[1] Here is situated the so called *Ganglion of Ribes*.

a. Its *upper* or *cranial* branch runs with the internal carotid a. into the carotid canal of the temporal bone, and there divides into two branches, an outer and an inner. The *outer* and larger branch accompanies the artery through its bony canal, ramifies

FIG. 36.



DIAGRAM OF THE COMMUNICATIONS OF THE FACIAL, GLOSSO-PHARYNGEAL, PNEUMOGASTRIC, SPINAL ACCESSORY, HYPOGLOSSAL, SYMPATHETIC, AND THE TWO UPPER CERVICAL NERVES.

1. Great petrosal nerve.
2. Lesser do.
3. External do.
4. Nerve to Stapedius muscle.
5. Spheno-palatine ganglion.
6. Otic ganglion.

upon it by the side of the body of the sphenoid, and so constitutes the "CAROTID PLEXUS."[1] From this outer branch a filament

[1] A small ganglion, the *carotid ganglion*, is sometimes met with in this plexus, on the under aspect of the artery.

proceeds to the Gasserian ganglion; another to the sixth cranial nerve; a third joins the great petrosal branch of the facial, and forms the Vidian nerve, and thus communicates with the spheno-palatine ganglion. It also communicates in the carotid canal with the tympanic branch of the glosso-pharyngeal. The *inner* branch, running on with the artery to the cavernous sinus, there forms another plexus, called from its position the "CAVERNOUS PLEXUS." Here the sympathetic is seen to communicate with the third, the fourth, and the ophthalmic branch of the fifth and sixth cranial nerves, and with the ophthalmic ganglion. Lastly, from both these plexuses secondary plexuses proceed, of which the minute filaments ramify on, and supply the coats of, the terminal branches of the internal carotid.

b. The *lower branch* descends and joins the middle cervical ganglion of the sympathetic.

c. The *external branches* are numerous, and connect the ganglion with the ganglion of the pneumogastric and hypoglossal nerves, and with the four upper cervical spinal nerves. A small twig also joins the petrosal ganglion of the glosso-pharyngeal and the upper ganglion of the pneumogastric in the foramen jugulare.

d. The *internal branches* are distributed to the pharynx, larynx, and the heart. The pharyngeal branches join the pharyngeal plexus on the middle constrictor of the pharynx; the laryngeal join the superior laryngeal nerve; the cardiac nerves, one or more in number—*superior cardiac*—descend behind the sheath of the carotid in front of the inferior thyroid artery and recurrent laryngeal nerve, and, entering the chest, join the superficial and deep cardiac plexuses.

e. The *anterior branches* lie in front of the external carotid artery and ramify around this vessel and its branches, forming the various plexuses, and named, on account of their delicacy, the *nervi molles.* In some of these plexuses are occasionally seen several ganglia, the intercarotic,[1] lingual, temporal, and pharyngeal ganglia. They are connected with the several gan-

[1] Situated in the angle at the bifurcation of the common carotid into the external and internal carotid arteries; it corresponds in structure with the coccygeal gland.

glia about the head and neck; namely, the ophthalmic, sphenopalatine, otic and submaxillary.

The *middle cervical ganglion*, the smallest of the three ganglia, is something less than a barleycorn in size. It is situated behind the carotid sheath, about the fifth or sixth cervical vertebra, on or near the inferior thyroid artery.

a. It is connected by branches with the superior ganglion above, and with the inferior cervical ganglion below.

b. Its *external branches* usually pass outward to join the fifth and sixth cervical spinal nerves.

c. Its *internal branches* are distributed to the thyroid body and the heart. The *branches to the thyroid body* accompany the inferior thyroid artery, and join the superior cardiac nerve, and in the gland they communicate with the external and recurrent laryngeal nerves. The *middle cardiac nerve*, the largest of the three cardiac nerves, descends, on the right side behind the common carotid a., usually in front of the first part of the subclavian artery, into the chest, when it lies on the trachea. It is joined by some cardiac filaments from the recurrent laryngeal nerve and superior cardiac nerve, and joins the deep cardiac plexus. On the left side this cardiac nerve enters the chest between the left carotid and subclavian arteries.

In cases where the middle cervical ganglion is absent, the preceding nerves are supplied by the sympathetic cord connecting the superior and inferior ganglia.

The *inferior cervical ganglion* is of considerable size, and is situated in the interval between the base of the transverse process of the seventh cervical vertebra and the neck of the first rib, immediately behind the vertebral artery, and to the inner side of the superior intercostal artery. Not infrequently it is coalesced with the first dorsal sympathetic ganglion.

Its branches are as follows:—

a. Superior branches, which pass upward and connect it with the middle cervical ganglion.

b. Inferior branches, which descend, some in front of and some behind the subclavian a., to join the first dorsal ganglion. One of these, the *inferior cardiac nerve*, passes behind the subclavian a.

in front of the trachea, to join the deep cardiac plexus, beneath the arch of the aorta, and communicates with the recurrent laryngeal and middle cardiac nerves.

c. External branches, which communicate with the seventh and eighth cervical nerves; others form a plexus around the vertebral artery, which join with the fourth, fifth and sixth cervical nerves.

DISSECTION OF THE THORAX.

Before the several organs contained in the thorax are examined, the student should have some knowledge of its framework. The ribs with their cartilages describe a series of arcs increasing in length from above downward, and form, with the dorsal vertebræ behind and the sternum in front, a barrel of a conical shape, broader in the lateral than in the antero-posterior diameter. The spaces between the ribs are occupied by intercostal muscles. In each intercostal space there are two layers of these muscles, arranged like the letter X. The fibres of the outer layer run obliquely from above downward and forward; those of the inner layer in the reverse direction. The base is closed in the recent state by a muscle—the diaphragm—which forms a muscular partition between the chest and abdomen. This partition is arched upward, so that it constitutes a vaulted floor for the chest, and by its capability of alternately falling and rising, it increases and diminishes the capacity of the thorax.

In front, the diaphragm is attached to the ensiform cartilage, but it slopes posteriorly, to become attached to the last rib. The circumference of the diaphragm is convex and muscular; in the centre it is flattened and aponeurotic. On the right side it corresponds, in front, with the upper border of the cartilage of the fifth rib; on the left side it corresponds with the upper border of the sixth rib.

The upper opening of the osseous thorax is bounded posteriorly by the body of the first dorsal vertebra, laterally by the first ribs, and in front by the upper border of the manubrium sterni.[1]

[1] That the student may have some knowledge of the diameters of the chest at different situations, the following measurements have been taken from a well articu-

Such, in outline, is the framework of the thorax, which contains the heart with its large vessels and the lungs. Its walls are composed of different structures—bone, cartilage, muscles and

FIG. 37.



FORM OF THE LUNGS, AND THE EXTENT TO WHICH THEY OVERLAP THE HEART AND ITS VALVES.

ligaments, which fulfill two important conditions: 1st, by their solidity and elasticity they protect the important organs contained

lated male skeleton of the average height:—The *antero-posterior* diameter at the upper opening of the thorax is 2¼ inches, at the articulation of the manubrium with the gladiolus it is 4½ inches, and at the junction of the gladiolus with the ensiform cartilage it has increased to 5⅝ inches. The *transverse diameter* of the upper opening was found to be 4⅜ inches; between the second ribs, 7 inches; between the third, 8⅛ inches; the diameter increased in regular proportion as far as the ninth rib, where it attained a measurement of 10⅝ inches; below this it gradually decreased. The upper border of the manubrium corresponds to the second dorsal vertebra. The articulation of the manubrium and the gladiolus is on a level with the fourth dorsal vertebra; and, lastly, the junction of the ensiform cartilage with the gladiolus is on a level with the border of the ninth or tenth dorsal vertebra.

in the thorax; 2d, by their alternate expansion and contraction they act as mechanical powers of respiration. For they can increase the capacity of the chest in three directions; in height, by the descent of the diaphragm; in width, by the rotation of the ribs; and in depth, by the elevation of the sternum.

The chest of the female differs from that of the male in the following points:—Its general capacity is less; the sternum is shorter; the upper opening is larger in proportion to the lower; the upper ribs are more movable, and therefore permit a greater enlargement of the chest at its upper part, in adaptation to the condition of the abdomen during pregnancy.

The *upper opening of the thorax* gives passage to the trachea, the œsophagus, the large vessels of the head and neck and upper extremities, viz., the innominate, the left carotid and subclavian arteries, with the left innominate and right subclavian and internal jugular veins, the superior intercostal and internal mammary arteries, the inferior thyroid veins, the sterno-hyoid, sterno-thyroid and longus colli muscles of each side, the pneumogastric, the left recurrent laryngeal, the phrenic and the sympathetic nerves; the cardiac branches of the sympathetic, and the cardiac branches of the pneumogastric; also to the anterior branch of the first dorsal nerve as it passes up to join the brachial plexus, the thoracic duct, the thymus gland (in early life), and, lastly, to the apices of the lungs, which, with their pleural coverings, rise up on each side into the neck for about one inch and a half above the first rib; the interspaces between these various structures being occupied by a dense fibro-cellular tissue, continuous with the deep cervical fascia.

The diaphragm, which forms the base of the thorax, is pierced by the following foramina:—the *aortic opening*, for the passage of the aorta, vena azygos major, thoracic duct; the *œsophageal opening* for the œsophagus, pneumogastric nerves, and œsophageal branch of the coronaria ventriculi artery; the *foramen quadratum*, for the vena cava inferior, a branch of the right phrenic nerve and lymphatics from the liver; the *right crus* transmits the greater and lesser splanchnic nerves; the *left crus*, in addition, transmits the vena

azygos minor. In front there are the narrow intervals for the passage of the internal mammary arteries.

Dissection. An opening must be made into the chest, by carefully removing the upper four-fifths of the sternum, and the cartilages of all the true ribs.[1] In doing this, care must be taken not to wound the pleura, which is closely connected with the cartilages. On one side the internal mammary artery should be dissected; on the other, removed.

In the dissection of the chest let us take the parts in the following order:—

1. Triangularis sterni, with the internal mammary artery.
2. Mediastina, anterior, middle, and posterior.
3. Pleura.
4. Position and form of the lungs.
5. Pericardium.
6. Position and relations of the heart.
7. Posterior mediastinum and its contents; namely, the aorta, the thoracic duct, the vena azygos, the œsophagus, and pneumogastric nerves.
8. Right and left brachio-cephalic veins and superior vena cava.
9. Course of the phrenic nerves.
10. Course and relations of the arch of the aorta,
11. The three great branches of the arch.
12. Sympathetic nerve.
13. Intercostal muscles, vessels and nerves.
14. Nerves of the heart; cardiac plexuses.

Triangularis Sterni. On the under surface of the sternum and cartilages of the ribs is a thin, flat muscle, named the *triangularis sterni*. It *arises* from the ensiform cartilage, the lower part of the side of the sternum, and the cartilages of two or three lower true ribs. Its fibres ascend obliquely outward, and are *inserted* by fleshy digitations into the lower borders of the cartilages of the true ribs—from the sixth to the second. Its lowest digitation runs transversely outward, each successive one, however, becomes more oblique, so that the highest one is nearly vertical in direction. The muscle is evidently a continuation upward

[1] Those who are more proficient in dissection should not remove the whole of the sternum, but leave a quarter of an inch of its upper part with the first rib attached to it. This portion serves as a valuable landmark, although it obstructs, to a certain extent, the view of the subjacent vessels.

of the anterior portion of the transversalis abdominis. Its action is to draw down the costal cartilages, and thus it acts in expiration. Its nerves come from the intercostal nerves, its arteries from the internal mammary.

Internal Mammary Artery. This artery is given off from the subclavian in the first part of its course opposite the thyroid axis. It passes down behind the clavicle, and on entering the chest it lies between the cartilage of the first rib and the pleura, and is crossed by the phrenic nerve. It then descends perpendicularly, about half an inch from the sternum, lying on the pleura and behind the costal cartilages; lower down it gets between the cartilages of the ribs and the triangularis sterni, as far as the seventh costal cartilage, where it divides into two branches, the *musculo-phrenic* and the *superior epigastric*. The latter branch then enters the wall of the abdomen behind the rectus abdominis, and finally inosculates with the deep epigastric (a branch of the external iliac). The branches of the internal mammary are as follows:—

a. Arteria comes nervi phrenici.—A very slender artery, which accompanies the phrenic nerve between the pleura and pericardium to the diaphragm, and anastomoses with the phrenic branches of the abdominal aorta, and internal mammary.

b. Mediastinal, pericardiac, sternal, and thymic.—These branches supply the cellular tissue of the anterior mediastium, the pericardium, and the triangularis sterni. The *thymic* are only visible in childhood, and disappear with the thymus gland.

c. Anterior intercostal.—Two for each intercostal space are distributed to the five or six upper intercostal spaces. They pass outward, and lie at first between the pleura and the internal intercostal muscle, and subsequently between the two intercostals. They inosculate with the intercostal arteries from the aorta.

d. The *perforating arteries* pass through the same number of intercostal spaces as the preceding branches, and supply the pectoral muscle and skin of the chest. In the female they are of large size (especially the third), to supply the mammary gland.

e. The *musculo-phrenic* branch runs outward behind the cartilages of the false ribs, pierces the attachment of the diaphragm,

and terminates near the last intercostal space. It supplies small branches to the diaphragm, to the sixth, seventh, and sometimes the eighth intercostal spaces.

Two venæ comites accompany the artery, and form a single trunk at the upper part of the chest, which terminates in the brachio-cephalic vein of its own side.

Lymphatic Glands. There are several *lymphatic glands* in the neighborhood of the internal mammary artery. They receive the lymphatics from the upper part of the abdominal wall, the diaphragm, the inner portion of the mammary gland, and the intercostal spaces. On the right side they terminate in the right lymphatic duct, on the left in the thoracic duct. In disease of the inner portion of the mamma, these glands may enlarge without any enlargement of those in the axilla.

Pleura. As the lungs are constantly gliding to and fro within the chest they are provided with a serous membrane to facilitate their motion. This membrane is termed the *pleura*. There is one for each lung. Each pleura forms a completely closed sac, and, like all other serous sacs, has a *parietal* and a *visceral layer*—that is, the first layer lines the containing walls, the latter is reflected over the contained organ or viscus. Its several parts are named after the surface to which they adhere: the parietal layer, which lines the ribs and intercostal muscles, is called *pleura costalis;* the visceral layer, which invests the lungs, *pleura pulmonis;* between these two layers is a space which is termed *the cavity of the pleura.*

Each pleura occupies its own half of the thorax; they do not communicate with one another, nor do they come into contact with each other, except for the short distance of about two inches in front, behind the sternum.

Unlike the peritoneum, the pleura forms no folds except a small one, called *ligamentum latum pulmonis*, which extends from the root of the lung to the diaphragm.

The *pleura costalis* (Fig. 38), in front, lines part of the back of the sternum and the inner surfaces of the costal cartilages; laterally, it is reflected over the ribs and the intercostal muscles; posteriorily, it is traced over the sides of the bodies of the dorsal

vertebræ; thence it passes to the back of the pericardium, over the posterior aspect of the root of the lung. It may now be traced, as the *pleura pulmonalis*, over the surface of the lung, to which it is intimately adherent, into the fissures between the lobes, as far as the anterior border of the lung; thence round its pericardial aspect to the front of the root of the lung, passing forward over the pericardium to the back of the sternum. Its only reflection, the ligamentum latum pulmonis, has been already alluded to. Below, the pleura covers the diaphragm.

The pleura rises as a conical dome into the base of the neck,

FIG. 38.



DIAGRAM OF THE REFLECTIONS OF THE PLEURAL SACS IN DOTTED LINES.

about an inch above the first rib, and is strengthened in this situation by expansions from the scaleni muscles.[1]

The thickness of the pleura differs; on the lung it is thin, semi-transparent and firmly adherent; on the ribs and diaphragm it is thick, and may be easily separated from its osseous and muscular connections.[2]

[1] A slip is described by Sibson as passing from the transverse process of the last cervical vertebra, and, spreading out, is inserted into the pleural dome and the inner margin of the first rib.

[2] From the prevertebral fascia, a ligamentous band passes downward along the inner border of the lung to be attached to the pericardium and the central tendon of

The spaces called *anterior* and *posterior mediastina*, formed by the separation of the pleuræ, will be described further on.

In health the internal surface of the pleura is smooth, polished, and lubricated by moisture sufficient to facilitate the sliding of the lung.[1] When this surface is thickened and roughened by inflammation, the moving lung produces a friction sound. When the pleural sac is distended by serum, it constitutes hydro-thorax; when by pus, empyema; when by air, pneumo-thorax; when by blood, hæmo-thorax.

Introduce your hand into the pleural sac, and ascertain that the reflection of the pleura on to the diaphragm corresponds with an imaginary line commencing at the lower part of the sternum, and sloping along the cartilages of the successive ribs down to the lower border of the last rib. Supposing a ball to lodge in the pleural sac, it might fall upon the dome of the diaphragm, and roll down to the lowest part of the pleural cavity. The place, therefore, to extract it, would be in the back, at the eleventh intercostal space. This operation has been done during life with success.

If a transverse section were made through the chest (*see* Fig. 38), you would observe that as the pleuræ nowhere come into actual contact, a space is left between them extending from the sternum to the spine, and which is larger in the middle than in front or behind. This interval is called by anatomists the *inter-pleural space* or the mediastinum, and for convenience sake is subdivided into three parts—an *anterior*, *middle* and *posterior mediastinum*.

Mediastina, Anterior, Middle and Posterior.

The *mediastina* are the spaces which the two pleural sacs leave between them in the antero-posterior plane of the chest, and which contain all the thoracic viscera except the lungs. There is an anterior, a middle and a posterior mediastinum. To put these spaces in

the diaphragm. As it passes downward it embraces the root of the lung, and supports it in its proper position. This band has been described as the "suspensory ligament of the diaphragm," by Teutleben.

[1] The pleura costalis is covered with flattened epithelial cells; the pleura pulmonalis with polyhedral granular cells. (Klein.)

the simplest light, let us imagine the heart and lungs to be removed from the chest, and the two pleural sacs to be left in it by themselves. The two sacs, if inflated, would then appear like two bladders, in contact only in the middle, as shown by the dotted outlines in the annexed scheme (Fig. 39). The interval marked *a*, behind the sternum, would represent the *anterior mediastinum;* the interval *b*, the *posterior mediastinum.* Now let us introduce the heart again, between the two pleural sacs: these must give way to make room for it, so that the two sacs are largely separated in the middle line of the chest; and the space thus occupied by the heart and large vessels takes the name of the *middle mediastinum.*

Fig. 39.

Looking at the chest in front, the *anterior mediastinum* appears as shown in the diagram (Fig. 40). It is not precisely vertical in its direction, for it inclines slightly toward the left, owing to the position of the heart. Its area varies; thus it is very shallow from before backward; it is extremely narrow in the middle where the edges of the lungs nearly meet; it is wider above, and widest of all below, where the lungs diverge. Posteriorly it is limited by the pericardium covering the heart, aorta and its branches, and the pulmonary artery.

What parts are contained in the anterior mediastinum? The remains of the thymus gland, the origins of the sterno-hyoid, sterno-thyroid, and triangularis sterni muscles, the left brachio-cephalic vein (which crosses behind the first bone of the sternum), a few lymphatic glands, and the left internal mammary artery and vein.

The *posterior mediastinum* (Fig. 38) is triangular in shape, placed in front of the dorsal vertebræ; it contains the œsophagus, the two pneumogastric nerves, the descending aorta, the thoracic duct, the greater and smaller azygos veins, the left superior inter-

costal vein, and some lymphatic glands. This space will be described in detail at a later stage.

The *middle mediastinum* is the largest of the mediastina, and contains the heart enclosed in the pericardium, the vena cava superior, the ascending aorta, the pulmonary arteries and veins, the phrenic nerves with their accompanying arteries, and the bifurcation of the trachea.

FIG. 40.



FORM OF THE LUNGS, AND THE EXTENT TO WHICH THEY OVERLAP THE HEART AND ITS VALVES.

A *superior mediastinum* has also been described comprising that part of the interpleural space which lies above a horizontal plane, extending behind from the lower part of the body of the fourth dorsal vertebra to the articulation between the manubrium and gladiolus in front. The contents of this mediastinum include all those structures found above this nearly horizontal plane, and are the transverse portion of the arch of the aorta and its three large branches, the trachea, œsophagus, and thoracic duct, the inno-

minate veins, superior vena cava, left recurrent laryngeal nerve, phrenic, pneumogastric, and cardiac nerves, lymphatic glands, and the thymus or its remains.

Before passing to the dissection of the contents of the thorax, the student should carefully trace the outline of the free borders of the pleuræ as seen in front of the chest. As the margins of the lungs for all practical purposes correspond with the borders of the pleuræ, we shall confine our description to the more important of the two structures, viz. the lungs. The value of this investigation is, that we are enabled to trace upon a living chest the outlines of the lungs, and know what parts are naturally resonant on percussion.

Commencing from above (Fig. 40, p. 183), we find that the apex of the lung extends into the neck, from an inch to an inch and a half above the clavicle. This part of the lung ascends behind the subclavian artery and the scalenus anticus muscle, and deserves especial attention, because it is, more than any other, the seat of tubercular disease. From the sternal end of the clavicles the lungs converge toward the middle line, where their borders nearly meet opposite the junction of the second rib. There is thus little or no lung behind the manubrium sterni.

From the level of the second costal cartilage to the level of the fourth, the inner margins of each lung run nearly parallel and almost in contact behind the middle of the sternum; consequently they overlap the great vessels at the root of the heart.

Below the level of the fourth costal cartilage the margins of the lungs diverge from each other, but not in an equal degree. The *left* presents the notch for the heart, and follows nearly the course of the fourth costal cartilage; at the lower part of its curve it projects more or less over the apex of the heart, like a little tongue. The *right* descends almost perpendicularly behind the sternum as low as the attachment of the ensiform cartilage, and then turning outward corresponds with the direction of the sixth costal cartilage. Hypertrophy of the heart, or effusion into the pericardium, will not only raise the point where the lungs diverge above the ordinary level, but also increase their divergence; hence the greater dullness on percussion.

Position and Form of the Lungs. The two lungs are situated in the chest; each in its own half of the thorax, with the heart, enclosed in its pericardium, between them. Each fits accurately into the cavity which contains it. Each, therefore, is conical in form; the *apex* projects into the root of the neck, a little more than an inch above the sternal end of the clavicle; the *base* is broad and rests on the diaphragm, the posterior part being thin and extending as far as the eleventh rib. Its *outer surface* is convex and adapted to the ribs; its *inner surface* is excavated, to make room for the heart in front; and behind presents a deep fissure—*hilum pulmonis*—for the attachment of the root of the lung. Its *posterior* surface is convex, and fits into the concavity of the thorax, on each side of the spinal column. The best way to see the shape of the lungs is to inject them through the trachea with wax, which is tantamount to taking a cast of each thoracic cavity. In such a preparation, besides the general convexities and concavities alluded to, you would find in the right lung a little indentation for the right brachio-cephalic vein; in the left an indentation for the arch of the aorta and the left subclavian artery.

Each lung is divided into an *upper* and a *lower lobe* by a deep fissure, which commences behind, about three inches from the apex, and proceeds obliquely downward and forward to the junction of the sixth rib with its cartilage (Fig. 40). Speaking broadly, nearly the whole of the anterior portion of the lung is formed by the upper lobe; nearly the whole of the posterior portion by the lower lobe. It should be noticed, however, that the upper lobe of the right lung is divided by a second fissure which marks off, from its lower part, a triangular portion called its *middle lobe*.

The dimensions of the right lung are greater than those of the left in all directions except the vertical; the reason of this exception is the greater elevation of the diaphragm on the right side by the liver. On an average the right lung weighs 24 ounces, the left 21 ounces.

The constituents of the root of the lung will be described hereafter, when they can be more satisfactorily displayed.

Præcordial Region. The *præcordial region* is the outline of the heart traced upon the front wall of the chest. It is important for auscultatory purposes that we should know how much of the heart is covered and separated from the wall of the chest by intervening lung (Fig. 40). The following will give a fair indication:—"Make a circle of two inches in diameter round a point midway between the nipple and the end of the sternum. This circle will define, sufficiently for all practical purposes, that part of the heart which lies immediately behind the wall of the chest, and is not covered by lung or pleura." [1]

This part of the præcordial region is naturally less resonant to percussion, for it is here uncovered, except by pericardium and loose connective tissue, and lies close behind the thoracic wall. In the rest of the præcordial region the heart is covered and separated from the chest wall by intervening lung.

Where should we put the stethoscope when we listen to the valves of the heart? For practical purposes it is enough to remember that the mouth of an ordinary-sized stethoscope will cover a portion of them all, if it be placed a little to the left of the mesial line of the sternum opposite the third intercostal space (Fig. 40, p. 183). They are all covered by a thin portion of lung; for this reason we ask a patient to stop breathing while we listen to his heart.

Position and Form of the Heart. The heart is situated obliquely in the chest, between the lungs. Its base, *i. e.*, the part by which it is attached, and from which its great vessels proceed, is directed upward toward the right shoulder; its apex points downward and to the left, between the fifth and sixth costal cartilages. It is supported, toward the abdomen, by the tendinous centre of the diaphragm. It is maintained in its position by a membranous bag termed the pericardium, which is lined by a serous membrane to facilitate its movements. The pericardium must first claim our attention.

Pericardium. The *pericardium* is the conical membranous bag which encloses the heart and the large vessels at its base. It is broadest *below*, where it is attached to the tendinous

[1] Latham's *Clinical Lectures*.

centre of the diaphragm, and to the muscular part in connection with the tendon, further to the left side than to the right; *above*, it is prolonged over the great vessels of the heart, about two inches from their origin, and is connected with the deep cervical fascia. *On each side*, it is in contact with the pleura; the phrenic nerve

FIG. 41.



RELATIVE POSITION OF THE HEART AND ITS VALVES WITH REGARD TO THE WALLS OF THE CHEST.

The valves are denoted by curved lines. The *aortic valves* are opposite the third intercostal space on the left side, close to the sternum. The *pulmonary valves* are just above the aortic, opposite the junction of the third rib with the sternum. The *mitral valves* are opposite the third intercostal space, about one inch to the left of the sternum. The *tricuspid valves* lie behind the middle of the sternum, about the level of the fourth rib. *Aortic murmurs*, as shown by the arrow, are propagated up the aorta; *mitral murmurs*, as shown by the arrow, are propagated toward the apex of the heart.

running down between them. *In front* of it is the anterior mediastinum; *behind* it is the posterior. Of the objects in the posterior mediastinum, that which is nearest to the pericardium is the œsophagus and the left pneumogastric nerve. It should be

remembered that the œsophagus is in close contact with the back of the pericardium and left auricle for nearly two inches; this fact accounts for what is sometimes observed in cases of pericarditis where there is much effusion: namely, pain and difficulty in swallowing.

The pericardium is a *fibro-serous* membrane, and consists of two layers—an external or fibrous, and an internal or serous. Its *fibrous* layer, a dense membrane, constitutes its chief strength, and is attached, below, to the central tendon and the adjoining muscular part of the diaphragm. Above, it forms eight tubular sheaths for the great vessels at the base of the heart; namely, one for the vena cava superior, four for the pulmonary veins, two for the pulmonary arteries, and one for the aorta. The *serous* layer forms a shut sac. Its parietal layer lines the fibrous layer to which it is intimately attached, and is reflected over the great vessels and the heart to form its visceral layer. To see where the serous layer is reflected over the vessels, distend the pericardium with air. Thus you will find that this layer is reflected over the aorta as high as the commencement of the transverse portion of the arch of the aorta. It is reflected over the front and sides of the vena cava superior.

The serous layer of the pericardium covers the large vessels to an extent greater than is generally imagined; though the extent is not precisely similar in all bodies. The aorta and pulmonary artery are enclosed in a complete sheath, two inches in length, so that these vessels are covered all round by the serous layer, except where they are in contact. Indeed, you can pass your finger behind them both, through a foramen bounded, in front, by the two great vessels themselves; behind, by the upper part of the auricles; and above, by the right pulmonary artery. Again, the back of the aorta, where it lies on the auricles, is covered by the serous pericardium. The superior cava is covered all round, except behind, where it crosses the right pulmonary artery. The inferior cava within the pericardium is partly covered in front. The left pulmonary veins are covered nearly all round; the right less so. Behind the auricles, chiefly the left, the serous layer extends upward in the form of a pouch, rising above their upper

border, so as to be loosely connected to the left bronchus. The object of these serous reflections is to facilitate the free action of the heart and the great vessels at its base.

In the healthy state, the capacity of the pericardium nearly corresponds to the size of the heart when distended to its utmost. The healthy pericardium, with the heart *in situ*, may be made to hold, in the adult, about ten ounces of fluid. The pericardium is not extensile. When an aneurism bursts into it, death is caused, not by loss of blood, but by compression of the heart in consequence of the inextensibility of the pericardium.

The pericardium derives its blood from the internal mammary, bronchial and œsophageal arteries; its nerve supply from the phrenic nerves.

On separating the left pulmonary artery and pulmonary vein, you will notice a fold of serous membrane about three-quarters of an inch long and about one inch in depth; this is the *vestigial fold of the pericardium*, described by Marshall.[1] It passes from the side of the left auricle, curving round the lower left pulmonary vein, to the left superior intercostal vein. It is a vestige of the left v. c. superior (duct of Cuvier) which exists in fœtal life.

Open the pericardium, and observe that the heart is conical in form, and convex everywhere except upon its lower surface, which is flat, and rests upon the tendinous centre of the diaphragm. When the pericardium is thus laid open, the following objects are exposed: viz. 1. Part of the right ventricle; 2. Part of the left ventricle; 3. Part of the right auricle with its appendix overlapping the root of the aorta; 4. The appendix of the left auricle overlapping the root of the pulmonary artery; 5. The aorta; 6. The pulmonary artery; 7. The vena cava superior; 8. The right and left coronary arteries.

Position of the Heart, continued. The heart, then, placed behind the lower half of the sternum, occupies more of the left than the right half of the chest, and rests upon the tendinous centre of the diaphragm, which is a little below the lowest part of the fifth rib. At each contraction the apex of the heart may be

[1] "On the Development of the Great Anterior Veins in Man and Mammalia," *Philosoph. Transactions*, 1850.

felt beating between the cartilages of the fifth and sixth ribs, about two inches below the nipple and an inch to its sternal side, or about three and a half inches to the left of the middle of the sternum. Speaking broadly, the base corresponds with a line drawn across the sternum along the upper borders of the third costal cartilages. The right border of the heart is formed almost entirely by the free margin of the right auricle, and, when distended, bulges nearly an inch to the right of the sternum. The left border of the heart is formed by the round border of the left ventricle, and reaches from a point commencing at the second left intercostal space, to a point placed two inches below the nipple and an inch to its sternal side. The horizontal border is formed by a sharp margin of the right ventricle, and extends from the sternal attachment of the fifth right costal cartilage, to meet the lowest point of the left margin.

The normal position which the cardiac valves hold to the thoracic walls is difficult to define with precision, and this probably accounts for the discrepancies noticed in anatomical works on this subject. The following relations are the results of carefully made observations in the post-mortem room: The *right auriculo-ventricular valves* are situated behind the sternum on the level of the fourth costal cartilage; the *left auriculo-ventricular valves* are opposite the third intercostal space, about one inch to the left of the sternum; the cusps of these valves extend as low as the fifth costal cartilage. The *pulmonary valves* lie immediately behind the junction of the third left costal cartilage with the sternum; the *aortic valves* are on a level with the upper border of the third intercostal space, just at the left of the middle line of the sternum.[1]

The position of the heart varies a little with the position of the body. Of this any one may convince himself by leaning alternately forward and backward, by lying on this side and on that, placing at the same time his hand upon the præcordial region. He will find that he can, in a slight degree, alter the place and the extent of the impulse of the heart. Inspiration and expiration

[1] Anatomists differ much in the description they give of the relations of the valves to the thoracic walls; in fact, no two agree in all the details.

also alter the position of the heart. In inspiration the heart descends with the tendinous centre of the diaphragm about half an inch.

The student should now make out the large vessels in connection with the base of the heart, leaving the consideration of this organ to a later stage of the dissection.

Before we can display the brachio-cephalic veins, the layer of the deep cervical fascia must be removed which descends over them from the neck and is lost upon the pericardium. Their coats are intimately connected with this fascia; and one of its functions appears to be to keep the veins permanently open for the free return of blood to the heart.

Brachio-Cephalic Veins. The *right* and *left brachio-cephalic* (innominate) *veins* are formed, near the sternal end of the clavicle, by the confluence of the internal jugular and subclavian veins. They differ in their course and relations, and must, therefore, be described separately.

The *left brachio-cephalic vein* passes from the left side obliquely behind the first bone of the sternum, the sterno-hyoid and thyroid muscles, the remains of the thymus gland, toward the right side, to form, with the right innominate vein, the vena cava superior (Fig. 42). It is about three inches in length, and its direction inclines a little downward. It is larger than the right brachio-cephalic, and crosses over the trachea and the origins of the three primary branches of the arch of the aorta. We are reminded of this fact in some cases of aneurism of these vessels—for what happens? The vein becomes compressed between the aneurism and the sternum; hence the swelling and venous congestion of the parts from which it returns its blood; namely, of the left arm, and the left side of the neck. The upper border of the vein is not far from the upper border of the sternum; in some cases it lies even higher, and we have seen it crossing in front of the trachea fully an inch above the sternum. This occasional deviation should be borne in mind in the performance of tracheotomy.

The *right brachio-cephalic vein* descends nearly vertically to join the superior vena cava, opposite the first right intercostal space. It is about an inch and a half in length, and is situated

about one inch from the mesial line of the sternum. On its l[illegible] side, but on a posterior plane, runs the arteria innominata; on [illegible] right side is the pleura. Between the vein and the pleura is t[illegible] phrenic nerve. The brachio-cephalic veins are not provided w[illegible] valves. The veins which generally empty themselves into t[illegible] right and left brachio-cephalic are as follows:—

The RIGHT B.-C. Vein receives:—
- The vertebral.
- The internal mammary.
- The inferior thyroid.

The LEFT B.-C. Vein receives—
- The vertebral.
- The internal mammary.
- The inferior thyroid.
- The superior intercostal.
- The pericardiac.
- The thymic.

FIG. 42.



SUPERIOR VENA CAVA AND ITS TRIBUTARIES.

Opening into the point of junction of the internal jugular and subclavian veins, on the right side is the right lymphatic duct; on the left side is the thoracic duct.

Vena Cava Superior. This is the great vein through which the impure blood from the head, upper extremities and chest, returns into the right auricle. It is formed by the junction of the right and left brachio-cephalic veins, which unite at nearly a right angle opposite the upper part of the first intercostal space on the right border of the sternum; that is, about the level of the highest point of the arch of the aorta. The vena cava descends

vertically, with a slight inclination backward, to the upper and anterior part of the right auricle. It is from two and a half to three inches long, and has no valves. The lower half of it is covered by the pericardium; you must, therefore, open this sac to see how the serous layer of the pericardium is reflected over the front and sides of the vein. In respect to its relations, notice that the vein lies in front of the right bronchus and the right pulmonary vessels; and that it is overlapped by the ascending aorta, which lies to its left side. In the upper half of its course, that is, above the pericardium, it is covered on its right side by the pleura; on this side, in contact with it, descends the phrenic nerve.

Before it is covered by the pericardium, the vena cava receives the right vena azygos, which opens into it after hooking over the right bronchus; also some pericardiac and mediastinal veins.

Course of the Aorta. The aorta is the great trunk from which all the arteries of the body carrying arterial blood are derived. It commences at the upper and back part of the left ventricle of the heart. It ascends forward and to the right as high as the lower border of the first intercostal space on the right side; it then arches backward toward the left side of the body of the second dorsal vertebra, and turning downward over the left side of the third, completes the arch at the fifth dorsal vertebra. The aorta descends through the thorax on the left side of the bodies of the remaining dorsal vertebræ as far as the diaphragm; it enters the abdomen through the aortic opening of the diaphragm, and descends as far as the left side of the body of the fourth lumbar vertebra, where it bifurcates into the right and left common iliac arteries. The aorta has received different names in the various parts of its course: thus, the arched portion extending from its origin at the left ventricle to the fifth dorsal vertebra is called the *arch of the aorta;* the portion between this vertebra and the diaphragm is the *descending thoracic aorta;* and the remainder of its course to its division at the fourth lumbar vertebra is known as the *abdominal aorta.*

Course and Relations of the Arch of the Aorta. The arch of the aorta, as before stated, commences at the upper part of the left ventricle, and describes an arch which terminates at the

fifth dorsal vertebra. Its origin is situated behind the pulmonary artery, and on the left side of the middle of the sternum, about the level of the lower border of the third costal cartilage. The direction of the arch, therefore, is from the sternum to the spine and rather obliquely from right to left.

FIG. 43.



For convenience of description, the arch of the aorta is divided into an ascending, a transverse, and a descending portion.

Ascending portion.—To see this portion of the aorta, the pericardium must be opened. You then observe that this part of the

artery is enclosed all round by the serous layer of the pericardium, except where it is in contact with the pulmonary artery. It is about two inches in length, and ascends with a slight curve, the convexity looking forward and to the right side, as far as the upper border of the second costal cartilage of the right side, where it lies almost in contact with the sternum. Its commencement is *covered* by the pulmonary artery, and is overlapped by the appendix of the right auricle, and higher up by the remains of the thymus gland. On its *right side*, but on a posterior plane, is the superior vena cava and the right auricle; on its *left side* is the pulmonary artery; *behind it* are part of the right auricle, the right pulmonary artery and vein, and the root of the right lung. This part of the arch gives off the right and left coronary arteries for the supply of the heart.

The right border of the ascending portion of the arch bulges to the right of the sternum to the extent of a quarter of an inch, and may be seen at the sternal end of the second right intercostal space.

The arch of the aorta presents partial dilatations in certain situations. One of these, called the *great sinus of the aorta*, is observed on the right side of the arch, about the junction of the ascending with the transverse portion; it is little marked in the infant, but increases with age. Three other dilatations (the *sinuses of Valsalva*), one corresponding to each of the valves at the commencement of the aorta, will be examined hereafter.

Transverse portion.—This portion of the aorta arches from the front to the back of the thorax, and extends from the upper border of the second right costal cartilage to the left side of the third dorsal vertebra. Its highest convex portion ascends usually to about an inch below the upper border of the sternum, and its concavity corresponds with the articulation of the first and second bones of the sternum. *In front*, it is covered by the left pleura and lung, and is crossed by the left phrenic, the left pneumogastric, the superficial cardiac nerves, the pericardiac and the left superior intercostal veins. Near its *summit* runs the left brachio-cephalic vein. Within its *concavity* are the left bronchus, the bifurcation of the pulmonary artery, the left recurrent laryngeal nerve, and

the remains of the ductus arteriosus. The artery *rests* upon the trachea (a little above its bifurcation), the deep cardiac plexus, the œsophagus, the thoracic duct, and the left recurrent laryngeal nerve. From the upper part of the transverse portion of the arch arise the arteria innominata, the left carotid, and the left subclavian arteries; and lying in front of these arteries is the left-brachio-cephalic vein.

Descending portion.—This part of the arch lies upon the left side of the body of the fourth dorsal vertebra, and at the lower border of the body of the fourth, or the upper part of the fifth dorsal it takes the name of the descending thoracic aorta. On its *right side* are the œsophagus and thoracic duct; on its *left* is the pleura; *in front* are the pleura and the root of the left lung; *behind*, it lies on the anterior common ligament, corresponding to the fourth dorsal vertebra.

What parts are contained within the arch of the aorta? The left bronchus, the right pulmonary artery, the left recurrent nerve, the remains of the ductus arteriosus, and the superficial cardiac plexus of nerves.

Relations of the Arch of the Aorta to the Sternum.

These relations vary according to the size of the heart, the obliquity of the ribs, and the general development of the chest. In a well-formed adult the ascending aorta is, at the most prominent part of its bulge, about half an inch behind the first bone of the sternum. The highest part of the arch is about one inch below the upper edge of the sternum.[1]

The branches given off from the ascending portion of the arch are the right and left coronary arteries, which pass, one in front of and the other behind the heart, to supply its muscular tissue.

The *right coronary artery* arises from the anterior sinus of Val-

[1] The relations of the arch of the aorta to the sternum vary even in adults, more especially if there be any hypertrophy of the heart. As an instance among many, we may mention that of a young female who died of phthisis. The position of the aortic valves was opposite the middle of the sternum, on a level with the middle of the second costal articulation. The highest part of the arch was on a level with the upper border of the sternum; the arteria innominata was situated entirely in front of the trachea; and the left brachio-cephalic vein crossed the trachea so much above the sternum that it would have been directly exposed to injury in tracheotomy.

salva, and passes to the right between the pulmonary artery and the right auricular appendix, running in the auriculo-ventricular groove.

The *left coronary artery*, larger than the preceding, is given off from the left posterior sinus of Valsalva, and passes between the pulmonary artery and left auricular appendix; it runs down in the anterior interventricular sulcus toward the apex of the heart.

The further description of these vessels will be considered in the dissection of the heart.

From the highest part of the arch arise three large arteries for the head, neck and upper limbs; namely, the brachio-cephalic or innominate artery, the left carotid and the left subclavian.

Brachio-cephalic or Innominate Artery. This, the largest of the three, arises from the commencement of the transverse part of the arch. It ascends obliquely toward the right and after a course of about one inch and a half to two inches, divides behind the right sterno-clavicular joint into two arteries of nearly equal size—the right subclavian and the right common carotid.

The relations of the innominate artery are as follow: *In front* it has the manubrium sterni, the right sterno-clavicular joint, the origins of the sterno-hyoid and thyroid muscles, the remains of the thymus gland, the left brachio-cephalic vein, the right inferior thyroid vein, and the right inferior cervical cardiac branch of the pneumogastric nerve. *Behind*, it rests upon the trachea. On its *left side* are the left common carotid and the remains of the thymus. On its *right side* are the lung and pleura, the right brachio-cephalic vein and the pneumogastric nerve.[1]

With the anatomy of the parts before you, you can understand that an aneurism of the innominate artery might be distinguished from an aneurism of the aorta—1. By a pulsation in the neck between the sterno-mastoid muscles, *i. e.*, in the fossa above the

[1] In some cases the innominate artery ascends for a short distance above the clavicle before it divides, lying close to the right of the trachea. We have already alluded to the fact that it occasionally gives off a middle thyroid artery (p. 108), which ascends in front of the trachea to the thyroid body, and is therefore directly in the way in tracheotomy.

sternum; 2. By occasional dyspnœa, owing to pressure on the trachea; 3. By venous congestion in the *left* arm; 4. By the aneurismal thrill being confined to the *right* arm.[1]

Left Common Carotid Artery. This artery arises from the arch of the aorta, close to and to the left of the arteria innominata. It ascends obliquely to the left sterno-clavicular joint, and thence to the neck, where its course nearly corresponds with the right common carotid (p. 98). *In front* it has the sternum, the left sterno-hyoid and thyroid muscles, the left brachio-cephalic vein, and the remains of the thymus gland; *behind*, it has at first the trachea, and higher up the œsophagus and thoracic duct; to the *right side* is the innominate artery; to the *left side* are the left subclavian artery and left pneumogastric nerve.

Left Subclavian Artery. This is the third branch of the transverse part of the arch, and arises from it opposite the third dorsal vertebra. It ascends nearly vertically out of the chest to the inner border of the first rib, and then curves outward behind the scalenus anticus. *In front* it has the lung covered with pleura, the pneumogastric, phrenic and cardiac nerves, the left common carotid, the left internal jugular and the left innominate veins, the sterno-hyoid, sterno-thyroid, and sterno-mastoid muscles. To its *right side* are the left carotid, œsophagus and trachea; between the artery and œsophagus is the thoracic duct; to its *left side* is the lung, covered with pleura; *behind it* are the longus colli muscle covering the vertebræ, the œsophagus, thoracic duct, and the inferior cervical ganglion of the sympathetic. The upper part of its course, where the vessel passes in front of the apex of the lung, has been described with the anatomy of the neck (p. 133).

[1] If the innominate artery be ligatured, the circulation would be maintained by the following collateral branches:—1. Between the branches of the two external carotids, which anastomose across the middle line. 2. Between the aortic intercostal and the superior intercostal. 3. Between the aortic intercostals and the internal mammary, long thoracic, alar thoracic, and subscapular arteries. 4. Between the internal mammary and deep epigastric. 5. Between the inferior thyroid arteries. 6. Between the two vertebrals. 7. Between the two internal carotid arteries.

Course of the Phrenic Nerves through the Chest.

The *phrenic nerve* comes from the third, fourth and fifth cervical nerves, but chiefly from the fourth. It descends on the scalenus anticus, gradually inclining to its inner border, and enters the chest between the subclavian vein and artery. It then crosses over the internal mammary artery and runs in front of the root of the lung, between the pleura and the pericardium, to the diaphragm (Fig. 30), to the under surface of which it is distributed.[1]

The phrenic nerve is joined on the scalenus anticus by an offset from the fifth cervical branch of the brachial plexus; by another filament from the sympathetic nerve; and very frequently by a small loop from the nerve to the subclavius muscle; occasionally also by a branch from the descendens noni.

In what respects do the phrenic nerves differ from each other in their course? The right phrenic runs along the outer side of the brachio-cephalic vein and superior vena cava; the left crosses in front of the transverse part of the arch of the aorta; besides which, the left is rather longer than the right, since it curves over the apex of the heart.

Before the phrenic nerve divides into branches to supply the diaphragm, it sends off minute filaments to the pleura and the pericardium; after it has pierced the diaphragm it distributes branches to the peritoneum. The *right* phrenic gives off one or two filaments, which unite with some filaments from the solar plexus and form a small ganglion, from which branches are distributed to the supra-renal capsule, the hepatic plexus, and the inferior vena cava. The *left* phrenic gives off a branch which joins a twig from the sympathetic near the œsophageal opening of the diaphragm, but there is no appearance of a ganglion.

Having studied these anatomical details, consider for a moment what symptoms are likely to be produced by an aneurism of the arch of the aorta, or any of the primary branches. A glance at the important parts in the neighborhood helps to answer the question. The effects will vary according to the part of the artery which is

[1] In the Museum of the College of Surgeons there is a dissection showing that the right phrenic nerve enters the diaphragm close to the right side of the vena cava inferior, while the left phrenic enters the left muscle of the diaphragm.

the seat of the aneurism, and according to the size, the form and the position of the tumor. One can understand that compression of the vena cava superior, or either of the brachio-cephalic veins, would occasion congestion and œdema of the parts from which they return the blood; that compression of the trachea or one of the bronchi might occasion dyspnœa, and thus simulate disease of the larynx;[1] that compression of the œsophagus would give rise to symptoms of obstruction. Nor must we forget the immediate vicinity of the thoracic duct and the recurrent nerve,[2] and the effects which would be produced by their compression. Can one, then, be surprised that a disease which may give rise to so many different symptoms should be a fertile source of fallacy in diagnosis?

Thus you can understand how aneurisms of the aorta may prove fatal, by bursting into the contiguous tubes or cavities; for instance, into the trachea, the œsophagus, the pleura, or the pericardium. You will see, too, why an aneurism of the first part of the arch is so much more dangerous than elsewhere. The reason is, that in this part of its course the aorta is covered only by a thin layer of serous membrane. If an aneurism take place here, the coats of the vessel soon become distended, give way, and allow the blood to escape into the pericardium; an occurrence which is speedily fatal, because, the pericardium being filled with blood, the heart is prevented from acting.

Posterior Mediastinum and its Contents. The posterior mediastinum (p. 182) is formed by the reflection of the pleural sac on each side, from the root of the lung to the sides of the bodies of the dorsal vertebræ. It is bounded in front by the pericardium and the roots of the lungs. To obtain a view of it, cut away the ribs nearly as far as their angles, draw out the right lung toward the left side, and fasten it firmly to the left side of the thorax. Remove the pleura of the right side from the ribs, and the posterior aspect of the root of the right lung, and then

[1] In the Museum of Guy's Hospital there is a preparation, No. 1487, in which laryngotomy was performed under the circumstances described in the text.

[2] See *Med. Gaz.* Dec. 22d, 1843: a case in which loss of voice was produced by the pressure of an aneurismal tumor upon the left recurrent nerve.

by a little careful dissection the space and the structures contained in it will be displayed. This mediastinum contains the descending thoracic aorta with some of the right aortic intercostal arteries; in front of the aorta, the œsophagus, with the pneumogastric nerves, the left in front and the right behind; on the right of the aorta is the vena azygos major, between this vein and the aorta is the thoracic duct; superiorly is the trachea; inferiorly are the splanchnic nerves and some lymphatic glands. To expose these last, we must remove the pleura, and a layer of dense fascia which lines the chest outside of it.

Descending Thoracic Aorta. We have already traced the arch of the aorta to the body of the fifth dorsal vertebra (p. 196). From this point, the aorta descends on the left side of the spine, gradually approaching toward the middle line. The artery, moreover, following the dorsal spinal curve, is not vertical, but concave forward. Opposite the last dorsal vertebra it passes between the crura of the diaphragm and enters the abdomen. It is contained in the posterior mediastinum; on its *left side* it is covered with pleura enclosing the left lung, and below it has the œsophagus to the left; on its *right* run the vena azygos, the œsophagus and thoracic duct; *in front* of it are the root of the left lung and the pericardium. Lower down the œsophagus is in front of the artery, and subsequently lies a little to its left side; *behind* are the vertebral column and the vena azygos minor. Its branches will be described presently.

Vena Azygos Major. This vein commences in the abdomen opposite the first or second lumbar vertebra, by small branches from one of the lumbar veins of the right side, and generally communicates with the renal, or the vena cava itself. This, indeed, is the main point about the origin of the vena azygos, that it communicates directly or indirectly with the vena cava inferior. It enters the chest through the aortic opening of the diaphragm, and ascends on the right side of the aorta through the posterior mediastinum, in front of the bodies of the lower dorsal vertebræ, and over the right intercostal arteries. When the vein reaches the level of the third dorsal vertebra, it arches forward over the right bronchus, and terminates in the superior vena cava,

just before this vessel is covered by pericardium. In its course it receives nine or ten of the lower intercostal veins of the right side, the spinal veins, the posterior mediastinal, the œsophageal and the right bronchial veins. Opposite the sixth or seventh dorsal vertebra it is joined by the left vena azygos. It is occasionally connected with the right superior intercostal vein.

FIG. 44.



DIAGRAM TO SHOW THE COURSE OF THE VENA AZYGOS AND THE THORACIC DUCT.

The left vena azygos, *vena azygos minor*, runs up the *left* side of the spine. This vein commences in the abdomen from one of the lumbar veins of the left side, or from the left renal. It then ascends on the left side of the aorta, through the left crus of the diaphragm. On a level with the sixth or seventh dorsal vertebra, it passes beneath the aorta and thoracic duct to join the azygos major. Before passing beneath the aorta it usually communicates with the left superior intercostal vein. It generally receives six or seven of the lower intercostal veins of the left side, the œsophageal and mediastinal veins. These azygos veins are provided with imperfect valves, and are supplemental to the inferior vena cava.

The *left upper azygos vein* receives the intercostal veins of the left side, usually from the fourth to the sixth; it communicates above with the left superior intercostal vein, and opens below, either directly into the vena azygos major, or indirectly into it through the vena azygos minor.

Thoracic Duct and Receptaculum Chyli.

The *thoracic duct* (Fig. 44) is a canal, from fifteen to eighteen inches long, through which the contents of the lacteal vessels from the intestines and the lymphatics from the lower limbs are conveyed into the blood. These vessels converge to an oval dilatation, termed *receptaculum chyli* (cistern of Pecquet), situated a little to the right side of the front of the body of the second lumbar vertebra, behind the aorta and close to the right crus of the diaphragm. Then, getting to the right side of the aorta, it ascends through the aortic opening of the diaphragm into the chest, and runs up the posterior mediastinum, still along the right side of the aorta, between this vessel and the vena azygos major, and opposite the sixth dorsal vertebra crosses over the vena azygos minor. Near the *third* dorsal vertebra it inclines to the left side, and then passes behind the arch of the aorta and the œsophagus, and ascends on the left side of this tube, between it and the left pleura; subsequently the duct passes up between the œsophagus and the left subclavian artery, as high as the seventh cervical vertebra, resting on the longus colli. It then emerges from beneath the carotid sheath, curves downward over the subclavian artery, in front of the scalenus anticus, and opens into the back part of the confluence of the left internal jugular and subclavian veins. The orifice of the duct is guarded by two valves which permit fluid to pass from the duct into the vein, but not *vice versâ*. Valves, disposed like those in the venous system, are placed at short intervals along the duct, more numerous in its upper part, so that its contents can only pass upward.[1] The diameter of the duct varies in different parts of its course; at its commencement it is about three lines in diameter, at the sixth dorsal it is about two lines, and it enlarges again toward the termination. It receives the lymphatics from the lower extremities, and from all the abdominal viscera (except

[1] The thoracic duct varies in size in different individuals. It may divide in its course into two branches, which subsequently reunite; instead of one there may be several terminal orifices. Instances have been observed in which the duct has terminated on the right instead of the left side (Fleischmann, *Leichenöffnungen*, 1815; also Morrison, *Journal of Anat.*, vol. vi, p. 427). It has been seen to terminate in the vena azygos (Müller's *Archiv*, 1834).

the convex surface of the liver and the abdominal walls); above these it receives the lymphatics from the left side of the thorax, the left lung, the left side of the heart, the left upper extremity, and the left side of the head and neck.

Œsophagus.

The *œsophagus* is that part of the alimentary canal which conveys the food from the pharynx to the stomach. It commences at the lower border of the fifth cervical vertebra, at the back of the cricoid cartilage; runs down in front of the spine, to the right side of the transverse portion of the arch of the aorta, then through the posterior mediastinum in front of the descending aorta, and passes through the œsophageal opening in the diaphragm to end in the stomach, opposite the ninth dorsal vertebra. It is from nine to ten inches long. Its course is not exactly straight, for it describes three curves—one an antero-posterior, the other two lateral curves. In the neck, at its commencement, it lies at first in the middle line; it then gets behind and a little to the left of the trachea; in the chest, *i. e.*, about the fourth dorsal vertebra, it inclines toward the right side to make way for the aorta; but it again inclines to the left before it passes through the diaphragm. Its anterior-posterior curve corresponds to the curve of the spinal column.

The œsophagus, in the neck, rests *behind*, upon the front of the spine, covered by the longus colli muscle; in *front* it has the trachea; on *each side* it is in relation with the thyroid body, the common carotid (chiefly the left) and inferior thyroid arteries, and the recurrent laryngeal nerves; to the *left* of it is the thoracic duct.

In the thorax, the œsophagus has, *in front*, the trachea, the left bronchus, the arch of the aorta, the left carotid and left subclavian arteries; and, lastly, for about two inches, the posterior surface of the pericardium (behind the left auricle); this accounts for the pain which is sometimes experienced, in cases of pericarditis, during the passage of food; *behind*, it rests upon the spinal column, the longus colli, the thoracic duct, the third, fourth, and fifth intercostal arteries of the right side; and, lastly, it lies in front of and slightly to the left side of the aorta; *laterally*, the aorta and pleura are to the *left*, and the vena azygos major to the

right of the tube. As it passes down in the interpleural space, it is in connection with both pleuræ. The œsophagus is surrounded by a plexus of nerves formed by the pneumogastric nerves, the left being in front of, the right behind it.

The œsophagus is supplied with blood by the inferior thyroid, the œsophageal branches of the aorta, the coronaria ventriculi, and the left phrenic artery. It is supplied with nerves by the pneumogastric and the sympathetic, which ramify between the two muscular layers. The œsophagus is composed of three coats, an external or muscular, a middle or areolar, and an internal or mucous. The muscular coat consists of an outer longitudinal, and an inner circular layer of fibres. The longitudinal layer is particularly strong, and arranged in the upper part mainly in three fasciculi, an anterior attached to the vertical ridge on the cricoid cartilage, and two lateral, which are continuous with the inferior constrictor; these, lower down, spread out and form a continuous layer round the œsophagus and support the circular fibres. Under the microscope the muscular fibres composing the upper part are seen to consist entirely of the striped variety; at the lower part, almost exclusively of the non-striped variety. The middle coat is composed of areolar tissue, and connects very loosely the muscular and mucous coats. The mucous membrane is of a pale color and considerable thickness, and in the contracted state of the œsophagus is arranged in longitudinal folds within the tube, which lies flattened in front of the spine. On the surface of the mucous membrane there are numerous minute papillæ placed obliquely. It is lined by a very thick layer of scaly epithelium. In the submucous tissue are many small compound racemose glands—*œsophageal glands*—especially toward the lower end of the œsophagus.

Course and Branches of the Pneumogastric Nerves.

The course of the pneumogastric nerves in the chest is not the same on both sides. The *right* pneumogastric nerve enters the chest between the subclavian artery and vein, descends behind the right innominate vein by the side of the trachea to the back of the root of the lung, where it breaks up into a plexus forming the posterior pulmonary plexus. From this plexus two cords descend

to the posterior surface of the œsophagus, upon which they divide into numerous branches; forming, with corresponding branches of the left pneumogastric nerve, the œsophageal plexus (plexus gulæ). The plexus then reunites into a single trunk, consisting also of some fibres from the left pneumogastric, and passes into the abdomen through the œsophageal opening in the diaphragm. The *left* pneumogastric descends into the chest between the left subclavian and carotid arteries, and behind the left brachio-cephalic vein. It then crosses in front of the arch of the aorta, and passes behind the root of the left lung to the anterior surface of the œsophagus, upon which it also assists to form a plexus with the nerve of the right side. The branches of the pneumogastric nerve in the chest are as follow:—

a. The inferior laryngeal or recurrent.—This nerve on the right side turns under the subclavian and the common carotid arteries (p. 132); on the left, under the arch of the aorta, below the ductus arteriosus, and ascends to the larynx. It passes beneath the inferior thyroid artery, and lying in the groove between the trachea and œsophagus, it enters the larynx beneath the lower border of the inferior constrictor of the pharynx. It supplies with motor nerves all the muscles which act upon the rima glottidis, except the crico-thyroid (supplied by the external laryngeal nerve). As they turn beneath their respective arteries, they give off branches to the deep cardiac plexus; also some small filaments to the inferior cervical ganglion of the sympathetic. In the neck it distributes small branches to the trachea, œsophagus, and inferior constrictor muscle.

b. Cardiac branches.—These are very small, and join the cardiac plexuses; the right arise from the right pneumogastric, and the right recurrent laryngeal, close to the trachea; the left come from the left recurrent laryngeal nerve. On both sides these branches pass to the deep cardiac plexus.

c. Pulmonary branches.—These accompany the bronchial tubes. The greater number run behind the root of the lung, and constitute the *posterior* pulmonary plexus. A few, forming the *anterior* pulmonary plexus, supply the front part of the root of the lung. Both these plexuses are joined by filaments from the second, third

and fourth thoracic ganglia of the sympathetic. The nerves of the lungs are, however, very small, and cannot be traced far into their substance.[1]

d. Œsophageal plexus.—Below the root of the lung each pneumogastric nerve is subdivided so as to form an interlacement of nerves round the œsophagus (plexus gulæ). From this plexus numerous filaments supply the coats of the tube; but the majority of them are collected into two nerves—the one, chiefly the continuation of the left pneumogastric nerve, lying in front of the œsophagus; the other, chiefly that of the right, lying behind it. Both nerves pass through the œsophageal opening in the diaphragm for the supply of the stomach; the left also sending filaments to join the hepatic plexus; the right sending branches to the cœliac, splenic and left renal plexuses.

Having examined the contents of the posterior mediastinum from the right side, now do so from the left. The left lung should be turned out of its cavity and fastened by hooks toward the right side. After removing the pleura, we see the descending thoracic aorta, the pneumogastric nerve crossing the arch and sending the recurrent branch under it; also the first part of the left subclavian, covered externally by the pleura. The pneumogastric nerve must be traced behind the root of the left lung to the œsophagus, and the œsophageal plexus of this side dissected. Lastly, notice the lesser vena azygos which crosses under the aorta about the sixth or seventh dorsal vertebra, to join the vena azygos major.

Thoracic Portion of the Sympathetic. This portion of the sympathetic system is generally composed of twelve ganglia covered by the pleura; one ganglion being found over the head of each rib, except the last two, which lie on the side of the bodies of the vertebræ. Often there are only ten ganglia, in consequence of two of them being fused here and there. The first thoracic ganglion is the largest.

The ganglia are connected together by thick branches, and each ganglion is connected *externally* by two branches with the corresponding intercostal nerve. The nerves proceeding from the

[1] Upon this subject, see the beautiful plates of Scarpa.

ganglia pass inward to supply the thoracic and part of the abdominal viscera. The *internal* branches which proceed from the six upper ganglia are small, and are distributed as follows (see the diagram):—

FIG. 45.

DIAGRAM OF THE THORACIC PORTION OF THE SYMPATHETIC.

a. Minute nerves from the first and second ganglia to the deep *cardiac plexus.*

b. Minute nerves from the third and fourth ganglia to the *posterior pulmonary plexus.*

The branches arising from the six lower ganglia unite to form three nerves—the *great splanchnic*, the *lesser*, and the *smallest splanchnic nerves.*

a. The *great splanchnic* nerve is generally formed by branches from the fifth or sixth to the tenth ganglion, and also receiving filaments, according to Beck, from all the thoracic ganglia above the sixth. They descend obliquely by the sides of the bodies of the dorsal vertebræ, along the posterior mediastinum, and unite into a single nerve, which passes through the corresponding crus of the diaphragm, and joins the semilunar ganglion of the abdomen, sending also branches to the renal and supra-renal plexuses.

b. The *lesser splanchnic* nerve is commonly formed by branches from the tenth and eleventh ganglia. It passes through the crus of the diaphragm to the cœliac plexus, and occasionally to the renal plexus.[1]

[1] In a few instances we have traced a minute filament from one of the ganglia into the body of a vertebra. According to Cruveilhier each vertebra receives one.

c. The *smallest splanchnic* nerve comes from the twelfth ganglion, passes through the crus of the diaphragm, and terminates in the lower part of the cœliac and renal plexuses. (This is not represented in the diagram.)

Intercostal Muscles. The intercostal muscles fill in the intervals between the ribs and are arranged in each interval in two layers, an external and an internal, which cross each other like the letter X. The *external intercostals*, eleven on each side, run obliquely from behind forward, like the external oblique muscle of the abdomen. They connect the contiguous borders of the ribs, passing from the outer lip of the rib above to the upper border of the rib below; they extend from the tubercles of the ribs behind to the costal cartilages in front, and are continued forward to the sternum as a thin membrane. The *internal* run from before backward, like the internal oblique, and pass from the inner lip of the groove in the rib above and from the costal cartilage, and are inserted into the upper border of the rib below. Observe that a few fibres of the inner layer pass over one or even two ribs, chiefly near the angles (especially of the lower ribs), and terminate upon a rib lower down.[1]

Neither of these layers of intercostal muscles extends all the way between the sternum and the spine; the outer layer, beginning at the spine, ceases at the cartilages of the ribs; the inner, commencing at the sternum, ceases at the angles of the ribs.

The intercostal muscles present an intermixture of tendinous and fleshy fibres; and they are covered inside and outside the chest by a glistening fascia, to give greater protection to the intercostal spaces.

The external intercostal muscles elevate the ribs, and are therefore muscles of inspiration. The internal intercostal muscles depress the ribs, and are therefore muscles of expiration.

Intercostal Arteries. There are eleven intercostal arteries on each side which lie between the internal and external intercostal muscles. The *two upper* arteries are derived from the superior intercostal branch of the subclavian; the remaining *nine* are furnished by the thoracic aorta; and since this vessel lies

[1] These irregular muscular bundles are called the *subcostal muscles*.

rather on the left side of the spine, the right intercostal arteries are longer than the left. The upper intercostal arteries from the aorta ascend obliquely to reach their intercostal spaces; the lower run more transversely. They are given off from the back of the descending aorta, and as they pass outward across the bodies of the vertebræ they are covered by the pleura and the sympathetic nerves; the right, in addition, pass behind the œsophagus, thoracic duct, and the vena azygos major; the left behind the left superior intercostal vein and the vena azygos minor. Having reached the intercostal space, each artery divides into an *anterior* and a *posterior* branch. The *anterior* branch in direction and size appears to be the continuation of the common trunk. At first it runs *along the middle of the intercostal space*, lying upon the external intercostal muscle, and separated from the cavity of the chest by the pleura and intercostal fascia. Here, therefore, it is liable to be injured by a wound in the back. But near the angle of the rib it passes between the intercostal muscles, and occupies the groove in the lower border of the rib above. Here it gives off a small branch, the *collateral intercostal*, which runs for some distance along the upper border of the rib below. After supplying the muscles, the main trunk anastomoses with the anterior intercostal branch of the internal mammary artery. In some cases this branch is as large as the intercostal itself, and situated so as to be directly exposed to injury in the operation of tapping the chest.

In its course along the intercostal space, each artery sends branches to the intercostal muscles and the ribs. About midway between the sternum and the spine, each gives off a small branch, which accompanies the lateral cutaneous branch of the intercostal nerve. The continued trunk, gradually decreasing in size, becomes very small toward the anterior part of the space, and is placed more in the middle of it. Those of the true intercostal spaces inosculate with branches of the internal mammary, and thoracic branches of the axillary; those of the false run between the layers of the abdominal muscles, and anastomose with the epigastric and lumbar arteries.

The *posterior* or *dorsal* branch passes backward between the

transverse processes of the vertebræ, on the inner side of the anterior costo-transverse ligament, and is distributed to the muscles and skin of the back. Each sends an artery through the intervertebral foramen to the spinal cord and its membranes.

On the right side the intercostal veins terminate in the vena azygos major; on the left, the seven or eight lower terminate in the vena azygos minor, the remainder in the left superior intercostal vein.

The usual relation which the intercostal vessels and nerve bear to each other in the intercostal space, is, that the vein lies uppermost, the nerve lowest, and the artery between them.

Dorsal Nerves. The *dorsal nerves* are twelve in number, the first emerging between the first and second dorsal vertebræ, and do not form a plexus as in the cervical, lumbar and sacral regions. Each dorsal nerve (like all the spinal nerves) arises from the spinal cord by two roots, an anterior or motor, and a posterior or sensory. The sensory root has a ganglion upon it. The two roots unite in the intervertebral foramen and form a *compound* nerve. After passing through the foramen, it is connected by two filaments with the sympathetic nerve, and then divides into an *anterior* and a *posterior* branch. The *posterior* or *dorsal* branches pass backward between the transverse processes of the dorsal vertebræ and divide into internal and external branches: the *internal* branches pass between the multifidus spinæ and semispinalis dorsi, pierce the rhomboidei and trapezius muscles; the six upper branches become cutaneous at the spinous processes of the vertebræ; the six lower supply only the multifidus spinæ, not giving off any cutaneous filaments; the *external* branches pass through the longissimus dorsi and supply this muscle, the ilio-costalis and

FIG. 46.

DIAGRAM OF A SPINAL NERVE.

their continuations and the levatores costarum; the six lower branches, in addition, distribute cutaneous filaments to the skin. These branches will be described more fully later on in the dissection of the back.

Intercostal Nerves.

The *intercostal nerves* are the *anterior* divisions of the dorsal nerves, and are twelve in number. Each nerve receives a filament from the sympathetic, and then proceeds between the intercostal muscles in company with, and immediately below, the corresponding artery. Midway between the spine and the sternum they give off *lateral cutaneous branches*, which supply the skin over the scapula and the thorax. The intercostal nerves terminate in front in the *anterior cutaneous nerves*. In the anterior part of the intercostal space the nerves lie in the substance of the internal intercostal muscles, and at the costal cartilages get to the inner side of the muscles, passing between them and the pleura.

The intercostal nerves are divided into two sets; the six upper are called the pectoral intercostals, because they supply the structures of the pectoral region; the six lower, the abdominal intercostals, because they supply the chest and abdominal walls.

The *upper* or *pectoral intercostal nerves* pass between the external and internal intercostal muscles, run forward in the substance of the latter muscle, and at the sternal end of the intercostal spaces pierce the internal intercostal muscles and the pectoralis major, to be ultimately distributed to the skin of the chest. The upper intercostal nerves supply the levatores costarum, serratus posticus superior, the intercostals, and the triangularis sterni.

The *lower* or *abdominal intercostal nerves* pass, like the upper nerves, between the intercostal muscles, as far forward as the costal cartilages. They pass behind these, and then run between the tranversalis and internal oblique, as far as the outer border of the rectus. Piercing the sheath of the muscle, they supply it, and subsequently end as the anterior cutaneous nerves of the abdomen. They supply the intercostal muscles, the serratus posticus inferior, and the abdominal parietal muscles.

Notice that the first dorsal nerve ascends nearly perpendicularly

over the neck of the first rib, to form part of the brachial plexus. This nerve, however, gives off a small branch, the first intercostal nerve, to supply the first intercostal space. This, as a rule, has no lateral cutaneous branch.

Intercostal lymphatic glands.—These are situated near the heads of the ribs; there are some between the layers of the intercostal muscles. They are of small size, and their efferent vessels go into the thoracic duct. Some of the upper ones on the right side pass into the right lymphatic duct. We have seen these intercostal glands enlarged and diseased in phthisis.

Bronchial and Œsophageal Arteries.

Small *bronchial arteries*, arising on the right side most frequently from the first aortic intercostal (third intercostal) artery, and on the left from the thoracic aorta, accompany the bronchial tube on its posterior aspect into the substance of the lung.[1] Their distribution and office will be considered with the anatomy of the lung. *Œsophageal arteries*, four or five in number, proceed from the front of the thoracic aorta to ramify on the œsophagus, where they inosculate above with the œsophageal branches of the inferior thyroid, and below with the œsophageal branches of the coronaria ventriculi and phrenic arteries. Small *posterior mediastinal arteries* are given off from the posterior part of the aorta, and supply the lymphatic glands and tissues of the posterior mediastinum.

Having finished the posterior mediastinum, replace the lung, and turn your attention once more to the great vessels at the root of the heart.

Pulmonary Artery.

This vessel is about two inches in length, and conveys the venous blood from the heart to the lungs. It proceeds from the upper part of the base of the right ventricle, and passes upward and backward along the left side of the aorta to the concavity of the arch of the aorta, where it divides into two branches, a right and a left, one for each lung. At its origin it has on each side an auricular appendix and a

[1] On the left side there are usually two bronchial arteries—a superior, arising from the highest part of the thoracic aorta, and an inferior, arising about an inch lower down.

coronary artery, and lies in front of the root of the aorta. The pulmonary artery and the aorta are surrounded for two inches by a common sheath of pericardium. The right branch, the larger and longer, passes horizontally below the arch of the aorta, behind the ascending aorta and the superior vena cava, to the root of its lung; the left is easily followed to its lung by removing the layer of pericardium investing it, when it will be found to pass horizontally in front of the descending aorta and the left bronchus to the root of the lung.

Search should be made for a short fibrous cord which connects the commencement of the left pulmonary artery with the concavity of the arch of the aorta. This cord is the remains of the *ductus arteriosus*, a canal which in fœtal life conveyed blood from the pulmonary artery to the aorta.

Draw toward the left side the first part of the arch of the aorta, and dissect the pericardium from the great vessels at the base of the heart. Thus a good view will be obtained of the trachea and its bifurcation into the two bronchi. Below the division of the trachea the right pulmonary artery is seen passing in front of the right bronchus. The superior vena cava and aorta are seen in front of, and nearly at right angles to, the right pulmonary artery. The vena azygos major is seen arching over the right bronchus and terminating in the vena cava superior, just before this vein pierces the pericardium. Notice, especially, a number of lymphatic glands called *bronchial*, at the angle of bifurcation of the trachea. The situation of these glands in the midst of so many tubes explains the variety of symptoms which may be produced by their enlargement.

Nerves of the Heart and Cardiac Plexuses. The nerves of the heart come from the pneumogastric and its recurrent branch, and the three cervical ganglia of the sympathetic. The pneumogastric gives off (generally) two or more filaments (cardiac) which proceed from the main trunk in the neck, or from its recurrent branch. The sympathetic sends three (cardiac) filaments; one from the upper cervical ganglion, a second from the middle, and a third from the lower; and they are called, respectively, the upper, middle and lower cardiac nerves of the sympathetic.

The minute and delicate nerves from these several sources on each side, pass downward to the base of the heart. They vary very much in their precise relations to the great vessels upon which they run; but speaking generally, it may be said that the nerves on the right side run chiefly behind the arch of the aorta; those on the left, in front of it. Eventually they form, by their mutual subdivisions and interlacement, an intricate network of nerves, termed, according to their position, the superficial and the deep cardiac plexus.

The *superficial* and *smaller cardiac plexus* lies in the concavity of the arch of the aorta, in front of the right pulmonary artery. It is closely connected with the deep plexus. It receives the upper cardiac branch of the left sympathetic, the lower cervical cardiac branch from the left pneumogastric, and filaments from the deep plexus. In it is usually found a small ganglion, *ganglion of Wrisberg*, placed beneath the arch of the aorta on the right side of the ductus arteriosus. This plexus distributes branches to the anterior coronary and the anterior pulmonary plexuses.

The *deeper* and *larger cardiac plexus* is situated behind the arch of the aorta in front of the bifurcation of the trachea and immediately above the right pulmonary artery. To see it the pericardial covering of the aorta must be carefully removed and the vessel hooked forward and to the left. This plexus is formed by all the cardiac branches of the right and left sympathetic ganglia, and by the cardiac branches of the pneumogastric and recurrent laryngeal nerves, except the left superior cardiac branch of the sympathetic and the left cervical cardiac branch of the pneumogastric, both of which pass to the superficial cardiac plexus. The branches from the *right* side of this plexus descend chiefly in front of the pulmonary artery and pass to the anterior pulmonary plexus and to the anterior coronary plexus; a few branches which pass behind the pulmonary artery are distributed to the right auricle and to the posterior coronary plexus. The branches from the *left* side of the plexus go to the left auricle, the anterior pulmonary plexus, but chiefly to the posterior coronary plexus.

From the cardiac plexuses, as a common centre, the nerves pass off to the heart, forming plexuses around the coronary

arteries. Thus, the *anterior coronary plexus* (derived chiefly from the superficial cardiac) accompanies the anterior coronary artery. The *posterior coronary plexus* (derived chiefly from the left side of the deep cardiac) runs with the posterior coronary artery. The two plexuses communicate at the apex of the heart, and in the ventricular septum.

FIG. 47.



DIAGRAM SHOWING THE CONSTITUENTS OF THE ROOT OF EACH LUNG, AND THEIR RELATIVE POSITION; ALSO THE POSITION OF THE VALVES OF THE HEART. THE ARROWS INDICATE THE DIRECTIONS IN WHICH AORTIC AND MITRAL MURMURS ARE PROPAGATED.

It is not an easy matter to trace the nerves into the substance of the heart. For this purpose a horse's heart is the best, and previous maceration in water is desirable. The nerves in the substance of the heart are peculiar in this respect; that they present minute ganglia in their course, which are presumed to preside over the rhythmical contractions of the heart.

Constituents of the Root of Each Lung. Draw aside the margin of the right lung; divide the superior vena cava above the vena azygos, and turn down the lower part. Remove the layer of pericardium which covers the pulmonary veins, and the constituent parts of the roots of the right lung will be exposed. It is composed of the pulmonary artery, the pulmonary veins, bronchus, bronchial vessels, anterior and posterior pulmonary plexuses, and some lymphatics. The following is the disposition of the large vessels forming the root of the lung. In front are the two pulmonary veins; behind the veins are the subdivisions of the pulmonary artery; behind the artery are the divisions of the bronchus. From above downward they are disposed thus: On the right side we find—1st, the bronchus; 2d, the artery; 3d, the veins. On the left we find—1st, the artery; 2d, the bronchus; 3d, the veins—as shown in Fig. 47.

DISSECTION OF THE HEART.

Position. The heart is conical in form, and more or less convex on its external aspect, with the exception of that portion lying on the tendinous centre of the diaphragm, which is flattened. It is situated obliquely in the thorax between the two lungs, and is completely surrounded by the pericardium. It extends from the fifth to the eighth dorsal vertebra, with its base directed upward and to the right, its apex downward and to the left, where during life it beats in the fifth intercostal space, two inches below the nipple and an inch to its sternal side. The position which the heart bears to the thoracic walls has been already described (p. 186); it varies, however, in different subjects, and, as a rule, is higher in the dead body than in the living, owing to the shrinking of the lungs.

The anterior surface of the heart is convex and looks upward and forward; the posterior surface is flattened and rests upon the diaphragm; the former is chiefly formed by the right ventricle, the latter by the left ventricle. The right border is sharp (*margo acutus*), while the left border is thick and rounded (*margo obtusus*).

Size and Weight. The size of the heart is dependent upon so many conditions, that the following measurements

must be received with more or less limitation. An average heart will measure, in its transverse direction at the base, three and a half inches; in its length, about five inches; in its thickness, two and a half inches. The weight is from ten to twelve ounces in the male, and from eight to ten in the female; but much depends upon the size and condition of the body generally. As a rule, the heart gradually increases in length, breadth and thickness, from childhood to old age.[1]

Notice two longitudinal grooves (*sulci*) on the front and back surfaces of the heart, which extend from the base of the ventricles to the apex, and which indicate the septum between the two ventricles; the anterior groove lies nearer to the left side, the posterior to the right side of the heart.

A circular groove, nearer the base, marks the separation between the auricles and ventricles. In the circular and longitudinal furrows, surrounded by more or less fat, run the coronary vessels, the nerves and the lymphatics.

The heart is a double, hollow, muscular organ; that is, it is composed of two hearts, a right and a left, separated by a septum, and not communicating with each other except during uterine, and rarely in adult life. Each half consists of two cavities, an *auricle* and *ventricle*, which communicate by a wide orifice, the auriculo-ventricular opening. The right half of the heart propels venous blood to the lungs, and is called the *pulmonary;* the left propels arterial blood from the lungs throughout the body, and is called the *systemic*. These two hearts are not placed apart, because important advantages result from their union. By being enclosed in a single bag they occupy less room in the chest; and the action of their corresponding cavities being precisely synchronous, their fibres, mutually intermixing, contribute to their mutual support.

The cavities of the heart should now be examined in the order in which the blood circulates through them.

Right Auricle. This is situated at the right side of the base of the heart, and forms a quadrangular cavity, the

[1] T. B. Peacock, *London and Edinb. Monthly Journal of Medical Science*, 1846 and 1854; Clendinning, *Med.-Chir. Trans.*, 1838; Beneke, *Marburger Schriften*, xi, 1879.

atrium or *sinus venosus*, between the two venæ cavæ, from which it receives the blood. From its front, a small pouch projects toward the left, and overlaps the root of the aorta; this part is termed the *appendix auriculæ*, and resembles a dog's ear in shape.

To see the interior, make a horizontal incision through the anterior wall from the apex of the appendix transversely across the cavity; from this make another upward at right angles into the superior vena cava. The interior is lined by a polished membrane called the *endocardium*, and is everywhere smooth except in the appendix, where the muscular fibres are collected into bundles, called, from their resemblance to the teeth of a comb, *musculi pectinati*. They radiate from the auricle to the edge of the auriculo-ventricular opening.

The following objects are seen on opening the auricle:—

Superior vena cava.	Eustachian valve.
Inferior vena cava.	Coronary valve.
Coronary sinus.	Annulus ovalis.
Auriculo-ventricular opening.	Fossa ovalis.
Foramina Thebesii.	Tubercle of Lower.

Musculi pectinati.

Examine carefully the openings of the two *venæ cavæ;* they are not directly opposite to each other; the *superior cava* opens into the auricle on a plane rather in front, and a little to the left, of the inferior, so that its orifice is opposite to the auriculo-ventricular opening. The *inferior cava*, after passing through the tendinous centre of the diaphragm, makes a slight curve to the left before it opens into the lowest part of the auricle; its direction is upward and inward, so that the stream of blood is directed toward the auricular septum. The orifice of each vena cava is nearly circular, and surrounded by circular muscular fibres continuous with those of the auricle.

The posterior wall of the auricle is formed by the partition between the auricles, the *septum auricularum*. Upon this septum, above, and to the left of the orifice of the vena cava inferior, is an oval depression (*fossa ovalis*), bounded by a prominent border (*annulus ovalis*). This depression indicates the remains of the opening (*foramen ovale*) through which the blood in fœtal life

passed from the right into the left auricle. After birth this opening closes; but if the closure is imperfect, the stream of dark blood in the right auricle mixes with the florid blood in the left, and occasions what is called *cyanosis*. A valvular communication, however, not infrequently exists between the auricles in this situation, which is not attended with indications of this disease.

A more or less noticeable fold of the lining membrane, the *Eustachian valve*,[1] may be seen projecting from the front margin of the v. c. inferior to the front border of the fossa ovalis. It is placed between the inferior vena cava and the lower margin of the

FIG. 48.



DIAGRAM OF THE INTERIOR OF THE RIGHT AURICLE.

annulus ovalis. Curved in shape, it passes forward and ends in two cornua, of which, one is attached to the annulus ovalis, the other is lost on the wall of the auricle. It consists of a reduplication of the endocardium, and contains some muscular tissue. It is the remnant of a valve, which was of considerable size in fœtal life, and served to direct the current of blood from the v. c. inferior, through the foramen ovale, into the left auricle.

To the left of the Eustachian valve, that is, between its remains and the auriculo-ventricular opening, is the orifice of the *coronary sinus*. The sinus is about an inch in length and receives the great cardiac vein, the posterior cardiac vein, and the oblique vein (of

[1] Eustachius, *Libell. de Vena sine Pari.*

Marshall), and will nearly admit the end of the little finger. It is surrounded by muscular fibres, and is guarded by a semi-circular fold of the endocardium, called the *valve of Thebesius*,[1] to prevent regurgitation of the blood during the auricular contraction.

Here and there upon the posterior wall of the auricle may be observed minute openings, called *foramina Thebesii;* some being the orifices of small veins returning blood from the substance of the heart; others being simple depressions in the muscular tissue. To the left, and rather in front of the orifice of the vena cava inferior, is the *auriculo-ventricular* opening, guarded by the tricuspid valve. It is oval in form, and will admit the passage of three fingers. Lastly, between the orifices of the superior and inferior venæ cavæ is a rounded elevation, the *tubercle of Lower*,[2] (not seen in the diagram), which is supposed to direct the current of blood, in fœtal life, from the superior cava to the auriculo-ventricular opening.

The *musculi pectinati* are parallel muscular elevations running across the inner surface of the auricular appendix, and to a slight extent also of the sinus venosus.

Right Ventricle. This forms the right border and about two-thirds of the front surface of the heart. To examine its interior, a triangular flap should be raised from its anterior wall. The apex of this flap should be below; one cut along the right edge of the ventricle, the other along the line of the ventricular septum. Observe that the wall of the ventricle is much thicker than that of the auricle. The cavity of the ventricle is conical, with base upward and to the right. Its inner wall is convex, and is formed by the septum ventriculorum. The upper and front part presents a smooth passage, the *infundibulum* or *conus arteriosus*, which leads to the opening of the pulmonary artery. It is situated to the left and in front of the auriculo-ventricular opening, and about three-fourths of an inch higher.

The following objects are seen in the right ventricle :—

Columnæ carneæ. Auriculo-ventricular opening.
Chordæ tendineæ. Pulmonary opening—

guarded by the tricuspid and semilunar valves.

[1] This valve is occasionally double. [2] Most distinct in quadrupeds.

From its walls project bands of muscular fibres, *columnæ carneæ*, of various length and thickness, which cross each other in every direction; this muscular network is generally filled with coagulated blood. Of these columnæ carneæ there are three kinds; one stands out in relief from the ventricle; another is attached to the ventricle by its extremities only, the intermediate portion being free; a third, and by far the most important set, called *musculi papillares*, is fixed by one extremity to the wall of the ventricle, while the other extremity gives attachment to the fine tendinous cords, *chordæ tendineæ*, which regulate the action of the tricuspid valve. The number of these musculi papillares is equal to the number of the chief segments of the valve; hence there are three in the right, and two in the left ventricle. Of those in the right ventricle, one is attached to the septum.

There are two openings in the right ventricle. One, the *auriculo-ventricular*, through which the blood passes from the auricle, is oval in form and placed at the base of the ventricle. It is surrounded by a ring of fibrous tissue, to which is attached the tricuspid valve.

Tricuspid Valve. This is situated at the right auriculo-ventricular opening, and consists of three triangular flaps. Like all the valves of the heart, it is formed by a fold of the lining membrane (*endocardium*) of the heart, strengthened by fibrous tissue, in which a few muscular fibres may be demonstrated. The bases of the valves are continuous with one another, so that they form a membranous ring between the auricle and the ventricle, while the segments project into the cavity of the right ventricle. Of its three flaps, the largest or anterior is so placed that, when not in action, it partially covers the orifice of the pulmonary artery; another, the internal, corresponds with the inferior wall of the ventricle; the third, or posterior, rests upon the septum ventriculorum.

Observe the arrangement of the tendinous cords which regulate the action of the valve. First, they are all attached to the ventricular surface of the valve. Secondly, the tendinous cords proceeding from a given papillary muscle are attached to the adjacent halves of two of the flaps; consequently, when the

ventricle contracts, and the papillary muscle also, the adjacent borders of the flaps will be approximated. Thirdly, to insure the strength of every part of the valve, the tendinous cords are inserted at three different points of it, in straight lines; accordingly, they are divisible into three sets. Those of the first, which are three or four in number, are attached to the base of the valve; those of the second, from four to six, proceed to the middle of its ventricular surface; those of the third, which are the smallest and most numerous, are attached to its free margin.[1]

Pulmonary or Semilunar Valves. These are three semicircular membranous folds, like watch-pockets, situated at the orifice of the pulmonary artery. They are attached by their convex borders to the root of the artery; their free edges look upward, and present a festooned border, in the centre of which is a small cartilaginous body called the *nodulus* or *corpus Arantii*.[2] The use of these bodies is plain. Since the valves are semilunar, when they fall together they would not exactly close the artery; there would be a space of a triangular form left between them in the centre, just as there is when we put the thumb, fore, and middle fingers together. This space is filled up by these nodules, so that the closure becomes complete.

The valves, two anterior and one posterior, are composed of folds of the *endocardium*, or lining membrane of the heart. Between the folds is a thin layer of fibrous tissue, which is prolonged from the fibrous ring at the orifice of the artery. This layer of fibrous

[1] The best mode of showing the action of the valve is to introduce a glass tube into the pulmonary artery, and then to pour water through it into the ventricle until the cavity is quite distended. By gently squeezing the ventricle in the hand, so as artificially to imitate its natural contraction, the tricuspid valve will flap back like a flood-gate, and close the auriculo-ventricular opening. In this way one can understand how, when the ventricle contracts, the blood catches the margin of the valve, and by its pressure gives it the proper distention and figure requisite to block up the aperture into the auricle. It is obvious that the tendinous cords will prevent the valve from flapping back into the auricle; and this purpose is assisted by the papillary muscles, which nicely adjust the degree of tension of the cords at a time when they would otherwise be too much slackened by the contraction of the ventricle.

[2] So called after Arantius, an Italian anatomist, who lived toward the close of the sixteenth century.

tissue, however, reaches the free edge of the valve at three points only, namely, at the centre, or corpus Arantii, and at each extremity. Between these points it stops short, and leaves a crescent-shaped portion of the valve, which is thinner than the rest, and consists of the endocardial membrane. This crescent-shaped portion, called the *lunula*, is not wholly without fibrous tissue; a thin, tendinous cord runs along its free edge, to give it additional strength to resist the pressure of the blood. Behind each of the valves the artery bulges and forms three slight dilatations, called the *sinuses of Valsalva*.[1] These, we shall presently see, are more marked at the orifice of the aorta.

The action of these valves is evident. During the contraction of the ventricle the valves lie against the side of the artery, and offer no impediment to the current of blood; during its dilatation, the elasticity of the distended artery would force back the column of blood, but that the valves, being caught by the refluent blood, bag, and fall together so as to close the tube. The greater the pressure, the more complete is the closure. The coats of the artery are very elastic and yielding, while the valve, like the circumference to which it is attached, is quite unyielding; consequently, when the artery is distended by the impulse of the blood, its wall is removed from the contact of the free margin of the valves, and these are the more readily caught by the regurgitating motion of the blood. The force of the reflux is sustained by the tendinous part of the valves, and by the muscular wall of the ventricle (probably in a state of contraction). The valves are capable of sustaining a weight of sixty-three pounds before they give way.[2] The thinner portions (*lunulæ*) become placed so as to lie side by side, each one with that of the adjacent valve. This may be demonstrated by filling the artery with water.

Left Auricle. This is situated at the left side and posterior part of the base of the heart, and is somewhat smaller than the right auricle. It consists, like the right auricle, of a cavity—the sinus venosus—and the auricular appendix. It is quadrilateral, and receives the four pulmonary veins, two on either side, which return the oxygenated blood from the lungs.

[1] An Italian anatomist, b. 1666, d. 1723. [2] Haller.

From its upper and left side, the *auricular appendix* projects toward the right, curling over the root of the pulmonary artery. The auricle should be opened by a horizontal incision along the ventricular border of the auricle, and another should be made upward from the centre of the first incision.

The interior of the *atrium* is smooth and flat, but in the appendix there are numerous raised muscular bands, the musculi pectinati. The interior presents the following objects for examination :—

The orifices of the four pulmonary veins.
The auriculo-ventricular opening.
The musculi pectinati.

The openings of the pulmonary veins are seen in the posterior wall, two on the right side (sometimes three), and two on the left side. They are not guarded by valves. Upon the septum between the auricles is a semilunar depression, indicating the remains of the foramen ovale. The *auriculo-ventricular opening*, situated at the lower and front part of the auricle, is smaller than that of the right side, and somewhat oval. Its long axis is nearly transverse, and, in the adult, will admit the passage of two fingers. The *musculi pectinati* are also smaller and fewer than in the right auricle.

Left Ventricle. This occupies the left border, and forms the apex of the heart. One-third of it only is seen on the anterior surface, the rest being on the posterior. To examine the interior, raise a triangular flap, with the apex below, from its front wall. Observe that its wall is about three times as thick as that of the right ventricle, and that this thickness gradually diminishes toward the apex. The interior of the left ventricle presents the following objects for examination :—

Auriculo-ventricular opening. Auriculo-ventricular or mitral valves.
Aortic opening. Semilunar valves.
Columnæ carneæ.

These parts so closely resemble that of the right that there is no necessity to describe them in detail. The *auriculo-ventricular valve* consists of two flaps; hence its name *mitral* or *bicuspid*. The larger of these flaps is placed between the aortic and auriculo-

ventricular orifices. There are only two *musculi papillares:* one attached to the anterior, the other to the posterior wall of the ventricle. They are thicker, and their *chordæ tendineæ* stronger than those of the right ventricle, but their arrangement is precisely similar. From the upper and back part of the ventricle, a smooth passage leads to the orifice of the aorta. This orifice is placed in the groove between the two auricles, and somewhat in front and to the right side of the left auriculo-ventricular opening. The two orifices are close together, and only separated by the larger flap of the mitral valve. The aortic orifice is guarded by three semilunar valves, of which the arrangement, structure and mode of action are similar to those of the pulmonary artery. Their framework is proportionately stronger, consistently with the greater strength of the left ventricle and the greater impulse of the blood. In the *sinuses of Valsalva* are observed the orifices of the two coronary arteries; the left arising from the sinus behind the left posterior segment; the right from behind the anterior segment.

Size of the Auriculo-ventricular and Arterial Openings.

The circumferences of the four orifices are as follows: that of the tricuspid orifice, 4.74 inches; that of the mitral, 4 inches; that of the pulmonary, 3.55 inches; and that of the aortic, 3.14 inches.[1]

Coronary Arteries.

The heart is supplied with blood by the two coronary arteries, a right or posterior, and a left or anterior. They are about the size of a crow's quill. Both arise from the aorta just above the free margins of the two semilunar valves, and thus always allow the passage of blood; both run in the furrows on the surface of the heart; both are accompanied by the cardiac nerves and by lymphatics.

The *anterior* or *left coronary* artery, the smaller of the two, arises from behind the left posterior valve of the aortic orifice. It appears between the pulmonary artery and the appendix of the left auricle, and then divides into two branches: one which seems the continuation of the main trunk and runs down the interventricular furrow on the anterior surface of the heart to the apex

[1] Dr. Peacock, *Croonian Lectures*, 1865.

the other passes transversely to the left, in the left auriculo-ventricular groove, to the back of the heart.

The *posterior* or *right coronary* artery arises from behind the anterior cusp of the aortic opening, and descends obliquely between the pulmonary artery and the appendix of the right auricle. It then turns to the right in the groove between the right ventricle and auricle to the back of the heart, where it divides into two branches; one of which descends in the posterior inter-ventricular furrow toward the apex of the heart; the other, which appears to be the continuation of the main trunk, runs in the left auriculo-ventricular groove. Besides these branches, the right coronary gives off a large branch which runs along the free border of the right ventricle.

Thus, the leading trunks of the coronary arteries run in the furrows of the heart, usually surrounded by fat. Their numerous branches supply the walls of the auricles and ventricles, and their terminations communicate with each other.

Coronary Veins and Sinus. The vein which corresponds with the anterior coronary artery ascends in the anterior inter-ventricular sulcus, and then curves round the left side of the heart in the left auriculo-ventricular groove, where it takes the name of the *great cardiac vein.* This vein soon dilates into a large trunk, the *coronary sinus*, which opens at the back of the right auricle below the Eustachian valve. Another vein, known as the *posterior cardiac*, ascends along the posterior inter-ventricular groove, to open by valved orifices into the coronary sinus; while others, the *anterior cardiac veins*, three or four in number, are seen running up on the anterior surface of the right ventricle to terminate directly in the right auricle. The *venæ Thebesii* transmit the blood directly from the muscular structure into the right auricle by small apertures, the foramina Thebesii. The *coronary sinus* is about an inch in length, and receives the great cardiac vein, the posterior cardiac vein, and the oblique vein of Marshall, placed on the posterior surface of the left auricle. Its orifice in the right auricle is guarded by a semilunar valve (*valve of Thebesius*) to prevent regurgitation of the blood. It is covered

and more or less supported in its course by muscular fibres passing from one auricle to the other.

The *lymphatics* of the heart pass mainly into a trunk which runs in the anterior inter-ventricular groove, and then, passing into the glands between the aorta and trachea, opens into the right lymphatic duct; other smaller lymphatics pass into the thoracic duct.

The *nerves* are derived from the cardiac plexuses, which have been already described, p. 215.

FIG. 49.



DIAGRAM OF THE RELATIVE POSITION OF THE VALVES OF THE HEART, SEEN FROM ABOVE.

A is placed on the triangular interval where the fibrous skeleton is the thickest.

Fibrous Rings of the Heart. What may be termed the fibrous skeleton of the heart, consists of four rings, which surround, respectively, the four orifices at its base: namely, the two auriculo-ventricular, the aortic, and the pulmonary. These rings give attachment by their external circumference to the muscular fibres of the heart, and from their internal circumference send fibrous prolongations to form the framework of the several valves. The skeleton is strongest just in the triangular interspace between the aortic and the two auriculo-ventricular orifices (letter A in Fig. 49).

In some animals, as in the ox and the elephant, there is here an irregularly triangular bone, known as the os cordis.

The relative position of these rings is best seen by removing the auricles and the great vessels at the base of the heart—leaving the several valves, and looking at them from above, as shown in the diagram. The pulmonary ring is on the highest level, and nearest to the sternum; below it is the aortic ring lying between and in front of the auriculo-ventricular rings, which are on the lowest level.

Attachment of the Large Arteries to the Ventricles. The fibrous rings at the arterial orifices present three festoons with their concavities directed upward. These give attachment, above, to the middle coat of the artery; below, to the muscular fibres of the ventricles; and, internally, to the fibrous tissue of the valves. The vessels are also connected to the heart by the serous layer of the pericardium, and by a continuation of the lining membrane of the ventricle.

Epicardium. This, the visceral layer of the pericardium, closely invests the external surface of the heart, and presents the usual appearances of a visceral serous membrane; having externally a layer of polygonal epithelial cells which rest upon a connective tissue layer, intermingled with elastic tissue. Beneath this layer there exists the sub-serous areolar tissue stratum, which is closely connected with the muscular structure of the heart (myocardium), and in which the vessels, nerves and lymphatics lie imbedded in more or less fat.

Endocardium. This smooth membrane lining the cavities of the heart resembles the visceral layer of the pericardium, and is continuous with the inner coat of the blood vessels. It may be easily stripped off, and is thin and semi-transparent, thicker in the left than in the right cavities, thickest of all in the left auricle. It consists of three layers: 1, a layer of flattened polygonal cells, resting upon, 2, a layer of connective tissue in which are some elastic fibres resembling the fenestrated coat of an artery; and, 3, a thin layer of connective tissue.

The muscular fibres of the heart are of the striped variety, although they are less well marked and regular than in voluntary

muscular fibres; the striæ are both transverse and longitudina The fibres are smaller than in the voluntary kind, have a we marked single nucleus, are branched, and are destitute of sarco lemma.

Arrangement of the Muscular Fibres of the Auricles.

The fibres of the auricles are distinct from those of the ventricles. They consist of a *superficia* layer common to both cavities, and a *deeper* layer proper to each. The *superficial* fibres run transversely across the auricles, and are most marked on the anterior surface; some pass into the inter-auricular septum. Of the *deeper* fibres, some are *annular* and surround the auricular appendages and the entrance of the great veins, upon which a few may be traced for a short distance; others, *looped*, run over the auricles, and are attached in front and behind to the auriculo-ventricular rings.

Arrangement of the Muscular Fibres of the Ventricles.

Speaking generally, it may be said that the right and left ventricles of the heart are two conical muscular sacs, enclosed in a third, which not only envelops them, but is reflected into the interior of both, at their apices, so as to line their cavities. All the muscular fibres are attached by one end to the fibrous rings of the orifices, and, by the other end, after a more or less spiral course, they reach the rings again, either directly or through the medium of the chordæ tendineæ and valves.

The external or superficial fibres pass from the base, where they are attached to the auriculo-ventricular rings, to the apex. This layer is thin in front, but behind it is better marked, and here the fibres do not pass into the septum, but over it, while in front they pass over the anterior septum, only at the base and apex of the ventricle. The fibres run more or less spirally toward the apex, where they form a whorl and pass into the left ventricle, so as to form, in part, the innermost vertical muscular layer, in part, the fleshy columns of its cavity. The superficial anterior fibres pass backward to the left, and form, behind, the posterior papillary muscle; and, on the other hand, the superficial posterior fibres pass over the right side of the heart and constitute the anterior papillary muscle.

The remaining fibres of the left ventricle, which constitute its chief thickness, are attached to the fibrous rings at the base of the heart. They pass, more or less obliquely, in the posterior and anterior walls, and entering the lower end of the septum pass in three different directions: one set upward in the septum, to be attached to the fibrous tissue in the triangular interspace; a second set pass through the septum, to form the posterior wall of the right ventricle and its posterior papillary muscle; while the third set take a transverse circular course in the left ventricle, some of its fibres being continuous with those of the right ventricle.

The fibres of the right ventricle are arranged on a plan similar to that of the left ventricle, of which it may be considered an appendage. The fibres which correspond to those forming the chief thickness of the left ventricle are similarly arranged into an anterior, middle and posterior set; the anterior pass backward into the septum, to reach the posterior wall of the left ventricle and interlace in the septum with the posterior set which pass forward in the septum to the front wall of the left ventricle; the middle set come chiefly from the outer wall of the right ventricle, deep down at the lower part of the septum, and then ascend to be attached to the fibro-cartilage. Besides these there are more or less numerous annular fibres encircling the right ventricle.[1]

Thickness of the Cavities. The average thickness of the right auricle is about one line; that of the left, one and a half. The average thickness of the right ventricle at its thickest part—*i. e.* the base—is about two lines. That of the left ventricle at its thickest part—*i. e.* the middle—is about half an inch. In the female the average is less.

Peculiarities of the Fœtal Circulation. The heart and the circulation of the fœtus differ from that of the adult in the following points:—

1. The *Eustachian valve* is well developed as a crescentic fold

[1] There are other accounts given of the arrangements of the muscular structure of the heart, and that given by Pettigrew is one which is adopted by many of the best anatomists. For further information on this subject consult Pettigrew, *Philosoph. Transactions*, 1864; Dr. Sibson, *Medical Anatomy*, 1869; Winckler, *Müller's Archiv*, 1865; Quain's *Anatomy*, vol. ii, p. 495, 1882.

which guides the current of blood from the inferior vena cava through the right auricle into the foramen ovale.

2. The *foramen ovale* is widely open up to the fourth month after which a septum grows up from the lower border of the left side, so that at the sixth month the blood can only pass in the onward direction into the left auricle.

3. The *right* and *left pulmonary arteries* are very small and ill developed, so as to admit very little blood to the lungs.

4. The *ductus arteriosus*, from the commencement of the left pulmonary artery to the aorta, is widely open.

5. The *hypogastric* or *umbilical arteries*, branches of the anterior division of the internal iliac, emerge through the umbilicus and pass to the placenta, so that the impure blood may be oxygenated.

6. The *umbilical vein* returns the pure blood partly to the liver, and partly through the—

7. *Ductus venosus* into the inferior vena cava.

8. The *right* and *left ventricles* are of equal thickness, because they have equal work to perform.

FŒTAL CIRCULATION.

Circulation of the Blood in the Fœtus.

Arterial blood is brought from the placenta by the umbilical vein (Fig. 50), and enters at the umbilicus, whence it passes to the under surface of the liver. Here it gives off some branches to the left lobe, and others to the lobulus Spigelii and lobulus quadratus, which eventually return their blood into the inferior vena cava. At the transverse fissure it divides into two branches—one, the smaller, termed the *ductus venosus*, passes straight to enter into the inferior vena cava, having previously joined the left hepatic vein; the other, or right division, joins the vena portæ, and after ramifying in the right lobe of the liver, returns its blood through the hepatic veins into the inferior vena cava.

From the inferior vena cava, which thus receives its blood from three sources, the blood enters the right auricle, and the stream (directed by the Eustachian valve) flows through the foramen ovale into the left auricle, where it becomes mingled with a little blood, which is returned from the pulmonary veins. From

the left auricle it runs through the left auriculo-ventricular opening into the left ventricle, and thence through the aorta into the great vessels of the head and upper extremities (only a small quantity passing into the descending thoracic aorta), which are

FIG. 50.



SCHEME OF THE FŒTAL CIRCULATION.

thus supplied by almost pure blood. From the head and upper limbs, the blood returns (impure) through the superior vena cava into the right auricle, whence, mixed with a small quantity derived from the inferior cava, it passes into the right ventricle. From the right ventricle the blood passes through the pulmonary artery

and the ductus arteriosus into the commencement of the descending aorta, only a very small quantity of it being distributed to the lungs; the lungs are, in the fœtus, almost solid organs, and the blood distributed to them is returned by the pulmonary veins into the left auricle. The blood which passes into the descending aorta, through the ductus arteriosus, is mingled with the small amount coming through the arch, and is then conveyed through the abdominal aorta into the iliac arteries; part is transmitted through the umbilical arteries (branches of the internal iliac arteries) to the placenta, to become re-oxygenated; part passes into the lower extremities through the external iliac and femoral arteries.

Changes in the Circulation at Birth.

The following changes take place in the circulation after birth:[1]—

1. The *umbilical vein* becomes obliterated from the second to the fifth day after birth, and subsequently forms the round ligament of the liver.

2. The *ductus venosus* also becomes closed about the same period, and may be traced as a thickened cord in the fissure of the ductus venosus.

3. The *foramen ovale* becomes closed from the sixth to the tenth day; but not infrequently a small indirect valvular communication may be found forming a communication between the two auricles.

4. The *ductus arteriosus* contracts immediately after birth, and becomes closed from the sixth to the tenth day. It eventually forms a fibrous cord connecting the left pulmonary artery with the aorta, the left recurrent laryngeal nerve winding round its left border.

5. The *pulmonary arteries* enlarge and convey venous blood to the lungs. These organs during fœtal life receive only a small quantity of blood from these arteries.

6. The *hypogastric arteries* become obliterated on the fourth or fifth day after birth.

[1] It is well to bear in mind that these important vascular changes do not take place suddenly at birth, but that they are the result of a gradual development which is completed at, or soon after, birth, mainly by the act of inspiration, whereby the blood passes through the lungs, the placental circulation at the same time being interrupted.

STRUCTURE OF THE LUNGS.

The lungs are very vascular, spongy organs, in which the blood is oxygenated by exposure to atmospheric air. Their situation and shape have been described (p. 185). We must now examine the trachea, the common air passage to both lungs, and then trace this tube downward to its bifurcation into the two bronchi, which, with their minute subdivisions, form the main structure of the lungs.

Trachea. This is a partly cartilaginous, partly membranous tube, and is situated in the middle line. It extends from the cricoid cartilage, *i. e.* opposite the upper border of the sixth cervical vertebra, to the third dorsal vertebra, where it divides into two tubes, the right and the left bronchus; one for each lung. Its length is from four to four and a half inches, and its width from eight to ten lines; but these measurements vary according to the age and sex of the patient and the capacity of the lungs. The trachea is surrounded by a quantity of loose connective tissue, so as to allow of its free mobility. It is kept permanently open by a series of incomplete cartilaginous rings, from sixteen to twenty in number, which extend round the anterior two-thirds of its circumference. These rings are deficient at the posterior part of the tube, where it is completed by a fibro-muscular membrane. This deficiency allows the trachea to enlarge or diminish its calibre; and for this purpose the membranous part of the tube is provided with unstriped muscular fibres which can approximate the ends of the rings.

The relations of the trachea to the surrounding parts should be considered, first, in the neck, and then within the thorax.

In the neck, it has, in *front* of it, the isthmus of the thyroid body, the sterno-hyoid and sterno-thyroid muscles, the inferior thyroid veins, two layers of the deep cervical fascia, the arteria thyroidea ima, if present, and (at the root of the neck) the innominate and left common carotid arteries. *Laterally* it is in relation with the lobes of the thyroid body, the common carotid arteries, the recurrent laryngeal nerves, and the inferior thyroid arteries. *Behind* it is the œsophagus, inclining slightly to the left.

In the chest, the trachea is contained in the superior mediastinum, and has, in *front* of it, the manubrium sterni, the origins of the sterno-hyoid and thyroid muscles, the left brachio-cephalic vein, the first parts of the innominate and left common carotid arteries, the transverse portion of the arch of the aorta, and the deep cardiac plexus. On the *right* side are the pleura and right pneumogastric nerve; on the *left*, the pleura, the left carotid, the left pneumogastric, cardiac and recurrent laryngeal nerves.

Bronchi, Right and Left.

The two *bronchi* differ in length, direction and diameter. The *right*, wider but shorter than the left, is about an inch long, and passes more horizontally to the root of its lung, on a level with the fourth dorsal vertebra. It is larger in all its diameters than the left; hence, foreign bodies which have accidentally dropped into the trachea are more likely to be carried into the right bronchus by the current of the air. The vena azygos major arches over the right bronchus to terminate in the superior vena cava. The *left* is about two inches in length, and, descending more obliquely to its lung than the right, enters it on a level with the fifth dorsal vertebra. The left bronchus passes under the arch of the aorta, in front of the œsophagus and the thoracic duct, and subsequently crosses in front of the descending aorta.

The *cartilages* of the trachea vary in number from sixteen to twenty, of the right bronchus from six to eight, and of the left from nine to twelve. Those of the trachea form about two-thirds of a circle, somewhat like a horseshoe in shape, are about $\frac{1}{4}$ of an inch in their vertical direction, and $\frac{1}{24}$ in thickness, and thicker in the middle than at the upper and lower borders. The cartilages are connected and covered on their outer and inner surfaces by a tough membrane, consisting of connective and elastic tissues. This membrane is attached above to the circumference of the cricoid cartilage, and is continued through the whole extent of the trachea and bronchial tubes. Posteriorly, where the cartilages are deficient, it contemplates the integrity of the air tube. In this tissue, which is of a pale reddish color, is a layer of unstriped muscular fibres, arranged in a transverse and a longitudinal direction.

The first cartilage is the broadest, and is frequently divided at one end; the last cartilage is placed at the bifurcation of the trachea, and is shaped like the letter **v**; its angle projects into the centre of the main tube, and its sides belong one to each bronchus.

Muscular Fibres. This thin stratum of unstriped muscular fibres consists of two layers, and is brought into view when the fibrous membrane and tracheal glands have been removed. The longitudinal fibres are the more external, and are attached by minute tendons to the extremities of the cartilages; the transverse fibres (*trachealis muscle*) extend transversely between the posterior free ends of the cartilages. By their contraction they approximate the ends of the cartilages and diminish the calibre of the trachea.

Elastic Tissue. This lines the whole tube, but is most abundant at the posterior or membranous part of the trachea, and its fibres run in a longitudinal direction. It is this layer which raises the mucous membrane into folds, and its elasticity admits of the elongation and the recoil of the tube.

Tracheal Glands. Upon the outer surface of the fibrous layer of the trachea are a number of small mucous glands, most numerous on the posterior part of the tube. They are compound racemose glands lined with columnar epithelium, and their excretory ducts pierce the fibrous and muscular layers, and terminate on the free surface of the mucous membrane. In health their secretion is clear, and just sufficient to lubricate the air passages. In bronchitis they are the sources of the abundant viscid expectoration.

Mucous Membrane. The *mucous membrane* lining the air passages is a continuation of that of the larynx. Its color in the natural state is nearly white, but in catarrhal affections it becomes bright red, in consequence of the accumulation of blood in the capillary vessels. It is continued into the ultimate air cells, where it becomes thinner and more transparent. In its deeper layer is found a considerable amount of elastic tissue; in its superficial layer a quantity of lymphoid tissue. Its surface is lined with a layer of columnar ciliated epithelial cells. The vibratile

movement of the cilia is directed in such a way as to favor the expectoration of the mucus. The ciliated epithelium lining the mucous membrane ceases at the commencement of the air cells, where it is replaced by the squamous variety.

At the root of the lung each bronchus divides into two branches, an upper and a lower, corresponding to the lobes of the lung; on the right side, the lower branch sends a small division to the third lobe of the lung. The tubes diverge through the lung, and divide into branches, successively smaller and smaller, until they lead to the air cells. These ramifications do not communicate with each other; hence, when a bronchial tube is obstructed, all supply of air is cut off from those cells to which it leads.

The several tissues—cartilaginous, fibrous, muscular, mucous, and glandular—which compose the air passages, are not present in equal proportions throughout all their ramifications, but each is placed in greater or less amount where it is required. The cartilaginous rings necessary to keep the larger tubes permanently open become, in the smaller tubes, fewer and less regular in form. As the subdivisions of the tubes multiply, the cartilages consist of small pieces placed here and there; they become less and less firm, and finally disappear when the tube is reduced to one-fortieth of an inch in diameter. The smallest air passages are entirely membranous, being formed of fibrous, elastic, and muscular tissues.

The Lungs. The lungs are two in number, and occupy the lateral cavities of the chest. Each is conical in shape, its apex extending into the neck, the base resting on the upper or convex surface of the diaphragm. The lung presents, for examination, an apex, a base, two surfaces, and two borders.

The *apex* extends upward about an inch and a half above the first rib, and is generally marked by a slight groove for the subclavian artery. The *base* is concave, and slopes downward at its posterior part. Its *outer surface*, in contact with the chest wall, is smooth and convex, and is deeper behind than in front. Its *inner surface* is concave, and hollowed out to accommodate the heart and its large vessels. The *anterior border* is sharp and

overlaps the large vessels and the pericardium. The *posterior border* is rounded and rests in the broad groove on the side of the bodies of the dorsal vertebræ. On the inner concave surface, a little above the middle and nearer the posterior than the anterior border, is the *root*, where the large vessels and bronchi pass to and from the lungs.

Each lung is traversed on its external surface by an oblique fissure which passes deeply into its interior. It extends from the upper part of the posterior border, downward and forward to the anterior border, and on the right side there is a second fissure, passing forward and upward from the oblique fissure to the middle of the anterior margin. The left lung presents a deep notch in the anterior border in which the pericardium is seen as far as the apex of the heart.

Contractibility of the Lung. When an opening is made into the chest, the lung, which was in contact with the ribs, immediately recedes from them, and, provided there be no adhesions, gradually contracts. If the lungs be artificially inflated, either in or out of the chest, we observe that they spontaneously expel a part of the air. This disposition to contract, in the living and the dead lung, is due to the elastic tissue in the bronchial tubes and the air cells; but more especially to a layer of delicate elastic tissue on the surface of the lung, which has been described by some anatomists as a distinct coat, under the name of the second or inner layer of the pleura.[1]

Color. The lungs are of a livid red or violet color; they often present a mixture of tints, giving them a marble-like appearance. This is not the natural color of the organ, since it is produced in the act of dying. It depends upon the stagnation of the venous blood, which the right ventricle still propels into the lungs, though respiration is failing. The tint varies in particular situations in proportion to the amount of blood, and is always deepest at the back of the lung. But the color of the proper tissue of the lung, apart from the blood which it contains, is pale and light gray. This color is seldom seen

[1] In some animals, the seal especially, the elasticity of this tissue is very strongly marked.

except in the lungs of infants who have never breathed, or after death from profuse hemorrhage.

Upon or near the surface of the lungs, numerous dark spots are observed which do not depend upon the blood, since they are seen in the palest lungs. They vary in number and size, and increase with age. The source of these discolorations is not exactly known; but they are probably deposits of minute particles of carbonaceous matter which have been inhaled with the air.

The lungs are composed of cartilaginous and membranous tubes, of which the successive subdivisions convey the air into closely-packed minute cells, called the *air vesicles;* of the ramifications of the pulmonary artery and veins; of the bronchial vessels concerned in their nutrition; of lymphatics and nerves. These component parts are united by connective tissue, and covered externally by pleura. The part at which they respectively pass in and out is called the *root of the lung*.

The lungs are the lightest organs in the body, and float in water, their specific gravity varying from .345 to .746. When entirely deprived of air they sink. This is observed in certain pathological conditions; *e. g.*, when one lung is compressed by effusion into the chest, or rendered solid by inflammation.

In the male the average *weight* of the right lung is 22 oz., that of the left 20 oz.; in the female the average is about 17 oz. on the right, and 15 oz. on the left side.

The total capacity of the lungs in an adult male of ordinary height is 282 cubic inches; and the amount of air still contained in the lungs after a forced expiration has been estimated at 57 cubic inches. The difference between these volumes—viz. 225 cubic inches—indicates the amount of air which can be inhaled, from the deepest expiration to the fullest inspiration, and has been termed the *vital capacity of the lungs*.[1]

The surface of the lung is closely invested by a thin, transparent layer of *serous membrane*, immediately beneath which is a fine areolar tissue, called *sub-serous*, which is very soft and elastic, so as to allow of the free expansion of the organ. This tissue sends inward prolongations, called *interlobular*, which map out the

[1] Hutchinson, *Med.-Chir. Trans.*, vol. xxix, 1846.

lungs into a number of angular spaces of various sizes, termed *lobules;* those on the surface, indicated by faint white lines, are larger than those in the interior of the lung. Each lobule is a lung in miniature, and consists of a small bronchial tube and its termination in dilated extremities, called *infundibula*, of ramifications of the pulmonary vessels, lymphatics and nerves, and lastly, of the bronchial vessels. The cells of the interlobular tissue have no communication with the air vesicles, unless the latter be ruptured by excessive straining, and then this connective tissue becomes inflated with air and is called *interlobular emphysema.* When infiltrated with serum it constitutes *œdema* of the lung.

Each bronchial tube divides and subdivides into smaller and smaller divergent tubes, until each has reached a reduced size of about 1 mm.; it then enters a pulmonary lobule, when it is termed a *lobular bronchial tube*, and presents on its walls numerous dilatations, called air cells or *alveoli*, which vary from $\frac{1}{75}$ to $\frac{1}{75}$ of an inch in diameter (Fig. 51). Thus reduced in size, the walls of the tubes no longer present traces of cartilaginous tissue, but are composed of a delicate elastic membrane upon which the capillaries ramify in a very minute network.[1] Each tube finally terminates in an enlarged irregular passage—*alveolar passage*—from which proceed on all sides numerous blind dilatations, named *infundibula.*

FIG. 51.

ULTIMATE AIR CELLS OF THE LUNG (FROM KÖLLIKER). MAGNIFIED TWENTY-FIVE TIMES.

The smaller bronchial tubes are encircled by more or less complete rings of cartilage; but as the tubes lessen in calibre the rings become less perfect; so that when the tubes are reduced to $\frac{1}{75}$ of an inch in diameter the rings

[1] In phthisis the expectoration contains some of the *débris* of this elastic framework of the air vesicles; it can be seen under the microscope, and is a test of the character of the sputa.

entirely disappear. The continuation of the air tubes consists simply of fibrous tissue, which becomes gradually thinner, so that in the smallest tubes they are reduced to simply membranous tubes, and are continued on as irregular passages—*intercellular passages*[1]—which are studded with numerous small saccules, termed *air cells* or *alveoli*.

The *air cells* are small, shallow, polyhedral depressions, from $\frac{1}{70}$ to $\frac{1}{200}$ of an inch in diameter, separated by thin partitions or septa which communicate freely with the intercellular passages, but not with each other.[2]

The *mucous membrane* which invests the divisions of the bronchi as far as the intercellular passages is lined with epithelium of the columnar ciliated variety. At this situation the character of the epithelium changes to that of a squamous kind, consisting of a single layer of flat polygonal nucleated cells.

The structure of the air cells differs in some important features from that of the smaller bronchial tubes; the muscular tissue disappears, the elastic tissue is no longer arranged in bundles, but becomes frayed out and intermingled with the connective tissue.

Pulmonary Vessels.

The *pulmonary artery* conveying venous blood to the lungs divides and subdivides with the bronchial tubes, and terminates in a fine, dense capillary plexus on the walls of the intercellular passages and air cells, beneath the epithelium. These plexuses—the *pulmonary capillaries*—form a single layer of capillaries which is so close that the interstices are even narrower than the blood vessels, which average about $\frac{1}{3000}$ of an inch in diameter. The plexus which ramifies over one air cell does not communicate with the plexus covering another air cell. The blood and air are not in actual contact. Nothing, however, intervenes but the wall of the cell and the capillary vessels, which are such delicate structures that they oppose no obstacle to the free interchange of gases by which the blood is purified. This purification is effected by the taking in of oxygen,

[1] Rainey, *Med.-Chir. Trans.*, vol. xxviii, 1845.

[2] The structure of the minute air cells of the human lung is in all respects similar to the large respiratory sac of the reptile.

and the elimination of carbonic acid and watery vapor. The most complete purification takes place in the single layer of capillaries between the folds of membrane projecting into the cell; for in this situation both sides of these vessels are exposed to the action of the air. The blood, circulating in steady streams through this capillary plexus, returns through the pulmonary veins. These, at first extremely minute, gradually coalesce into larger and larger branches, which anastomose very freely, and accompany the arteries. They finally emerge from the root of the lung by two large trunks which carry the oxygenated blood to the left auricle of the heart. The pulmonary veins are not provided with valves.

Bronchial Arteries. These small arteries, two or more in number, are the nutrient vessels of the lungs. The right arises either from the first aortic intercostal, or, conjointly with the left bronchial, from the thoracic aorta. The left, usually two in number, come from the thoracic aorta. They enter the lung behind the divisions of the bronchi, which they accompany. The bronchial vessels are distributed in various ways; some of their branches supply the coats of the air passages, the large blood vessels and the lymphatic glands; others the interlobular tissue; a few reach the surface of the lung, and ramify beneath the pleura. The right *bronchial veins* terminate in the vena azygos; the left, in the superior intercostal vein.

The *nerves* of the lung are derived from the pneumogastric and the sympathetic. They enter with the bronchial tubes, forming a plexus in front and behind them, *anterior* and *posterior pulmonary plexus*, in which are found minute ganglia.

The *lymphatics* of the lungs consist of a superficial and deep set; some commence in the lymphatic capillaries in the interlobular tissue, and thence pass to the surface, forming a network which communicates with the sub-pleural lymphatic plexus; others take their origin in the mucous membrane of the bronchial tubes; and all eventually enter the bronchial glands. Of these, the larger are situated about the bronchi near the root of the lung, particularly under the bifurcation of the trachea.

DISSECTION OF THE PHARYNX.

Dissection. To obtain a view of the pharynx, cut through the trachea, the œsophagus, the large vessels and nerves of the neck, a short distance above the first rib, and then separate them from the prevertebral muscles which lie immediately in front of the bodies of the cervical vertebræ, and to which they are but loosely connected. The sawn surface of the skull should now be allowed to rest horizontally upon the table, and the base should then be sawn or chiseled through transversely between the vertebral column and the styloid processes of the temporal bone. It is not always easy to keep the saw so well behind the pharynx and the vessels and nerves as to detach them without injury, in which case it is well to use the chisel in preference. When this is accomplished the student will find that the pharynx and larynx are left attached to the anterior half of the section; the spinal column and the prevertebral muscles to the posterior half. Tow should then be introduced through the mouth and œsophagus, to distend the walls of the pharynx. The front section is now to be fastened to a block by means of hooks, so that the pharyngeal muscles are toward the dissector, and the œsophagus downward.

One side of the pharynx should be dissected to show the constrictor muscles, the other should be reserved for the vessels and nerves in immediate relation with the pharynx.

General Description of Pharynx. The term *pharynx* is applied to that part of the alimentary canal which receives the food after it has been masticated, and propels it downward into the œsophagus. It is a funnel-shaped, muscular bag, about four and a half inches in length, and broader in its transverse than in its antero-posterior diameter. Its broadest portion is situated opposite the os hyoides, and it then gradually tapers as far as the cricoid cartilage, where it is continuous with the œsophagus, which is its narrowest portion. Its *upper part* is attached to the basilar process of the occipital bone and the petrous portions of the temporal bones; *behind*, it is loosely connected, by deep cervical fascia, with the prevertebral muscles;[1] in *front*, it is

[1] It is in this tissue (which never contains fat) that post-pharyngeal abscesses are seated.

attached to the internal pterygoid plates and hamular processes of the sphenoid, to the pterygo-maxillary ligaments, the lower jaw, the tongue, the hyoid bone, and the stylo-hyoid ligaments, and to the thyroid and cricoid cartilages; *laterally*, it is loosely connected to the styloid muscles, and it has in close relation with it the common and internal carotid arteries, the glosso-pharyngeal, pneumogastric, spinal accessory, hypoglossal and sympathetic nerves; the internal pterygoid, tensor palati and stylo-pharyngeus muscles; the lingual and ascending pharyngeal arteries, the superior laryngeal and external laryngeal nerves, the ascending palatine artery, and the internal jugular vein. Its dimensions are not equal throughout. Its breadth at the upper part is equal to that of the posterior openings of the nose; here it is only required to convey air, but it becomes much wider in the situation where it transmits the food—that is, at the back of the mouth; thence it gradually contracts to the œsophagus. The pharynx, therefore, may be compared to a funnel communicating in front, by wide apertures, with the nose, the mouth and the larynx; while the œsophagus represents the tube leading from its lower end. The upper part of the funnel forms a cul-de-sac at the basilar process of the occipital bone. At this part there is, on each side, the opening of a narrow canal, called the Eustachian tube, through which air passes to the tympanum of the ear.[1]

Before the muscles of the pharynx can be examined, we must remove a layer of thin fascia, termed the *pharyngeal fascia*. It is the layer of deep cervical fascia behind the pharynx, and must not be confounded with the proper *pharyngeal aponeurosis*, which intervenes between its muscular and mucous walls.

[1] Observe that the pharynx conducts to the œsophagus by a gradual contraction of its channel. This transition, however, is in some cases sufficiently abrupt to detain a foreign body, such as a morsel of food more bulky than usual, at the top of the œsophagus. If such a substance become firmly impacted in this situation, one can readily understand that it will not only prevent the descent of food into the stomach, but that it may occasion, by its pressure on the trachea, alarming symptoms of suffocation. Supposing that the obstacle can neither be removed by the forceps, nor pushed into the stomach by the probang, it may then become necessary to extract it by making an incision into the œsophagus on the left side of the neck.

At the back of the pharynx, near the base of the skull, are few *lymphatic glands.* They sometimes enlarge, and form a perceptible tumor in the pharynx.

In removing the fascia from the pharyngeal muscles, notice that a number of veins ramify and communicate in all directions.

FIG. 52.



SIDE VIEW OF THE MUSCLES OF THE PHARYNX.

They constitute the *pharyngeal venous plexus,* and terminate in the internal jugular veins.

Constrictor Muscles of the Pharynx.

They are three in number, and arranged so that they overlap each other—*i. e.* the inferior overlaps the middle, and the middle the superior (Fig. 52).

They have the same attachments on both sides of the body; and the fibres from the right and left meet together, and are inserted in the mesial line, the insertion being marked by a white longitudinal line, called the *raphé*.

The *inferior constrictor*, the most superficial and thickest of the thin constrictors, *arises* from the side of the cricoid cartilage behind the crico-thyroid muscle, from the surface behind the oblique ridge and the lower cornu of the thyroid cartilage. Its fibres expand over the lower part of the pharynx. The superior fibres ascend; the middle run transversely; the inferior descend slightly, and are *inserted* into the posterior median raphé. The lower fibres are continuous with those of the œsophagus. Beneath its lower border the recurrent laryngeal nerve enters the larynx. Its nerve supply is from the pharyngeal plexus, the external laryngeal and the recurrent laryngeal nerves.

In order to completely expose the next muscle, the right half of the inferior constrictor should be reflected from the middle line.

The *middle constrictor arises* from the upper edge of the greater cornu of the os hyoides, from its lesser cornu, and part of the stylo-hyoid ligament, and is *inserted* into the posterior median raphé. Its fibres take different directions, so that, with those of the opposite muscle, they form a lozenge. The lower angle of the lozenge is covered by the inferior constrictor; the upper angle ascends nearly to the basilar process of the occipital bone, and terminates upon the pharyngeal aponeurosis. The external surface of the muscle is covered at its origin by the hyo-glossus, from which it is separated by the lingual artery; while beneath it are the superior constrictor, the stylo-pharyngeus and palato-pharyngeus muscles and the pharyngeal aponeurosis. Its nerve comes from the pharyngeal plexus.

Between the middle and inferior constrictors, the superior laryngeal artery and nerve perforate the thyro-hyoid membrane to supply the larynx.

The *superior constrictor* consists of pale muscular fibres, and *arises* from the hamular process of the sphenoid bone, and from the lower part of its internal pterygoid plate; from the tuberosity of

the palate bone and the reflected tendon of the tensor pal
from the pterygo-maxillary ligament (which connects it with
buccinator); from the back part of the mylo-hyoid ridge of
lower jaw, and from the side of the tongue. The fibres pass ba
ward to the mesial raphé; some of them are *inserted* throu

FIG. 53.



VIEW OF THE CONSTRICTOR MUSCLES FROM BEHIND.

the medium of the pharyngeal aponeurosis into the basilar process. Its nerve comes from the pharyngeal plexus.

The upper border of the superior constrictor presents, on either side, a free semilunar edge with its concavity upward, so that, between it and the base of the skull, a space is left in which the

muscle is deficient (Fig. 53). Here the pharynx is strengthened and walled in by its own aponeurosis. The space is called the *sinus of Morgagni;* and in it, with a little dissection, we expose the muscles which raise and tighten the soft palate, *i. e.*, the levator palati and the tensor palati. The Eustachian tube opens into the pharynx just here. The fibres of the stylo-pharyngeus pass in between the superior and middle constrictors, and expand upon the side of the pharynx; some of them mingle with those of the constrictors, so as to be able to lift up the pharynx in deglutition; but most of them are inserted into the superior and posterior margins of the thyroid cartilage.

Pharyngeal Aponeurosis. The *pharyngeal aponeurosis* intervenes between the muscles and the mucous membrane of the pharynx. It is attached to the basilar process of the occipital bone, and to the points of the petrous portions of the temporal bones. It maintains the strength and integrity of the pharynx at its upper part, where the muscular fibres are deficient; but it gradually diminishes in thickness as it descends, and is finally lost on the œsophagus. Notice the number of mucous glands upon this aponeurosis, especially near the base of the skull and the Eustachian tube. These glands sometimes enlarge and occasion deafness, from the pressure on the tube.

Openings into the Pharynx. Lay open the pharynx by a longitudinal incision in the middle line, up to the pharyngeal tubercle; then divide transversely, for a short distance, that part of the pharyngeal aponeurosis which is attached to the basilar process, so as the better to view the cavity of the pharynx. Observe the seven openings leading into it (Fig. 54):—1. The two posterior nares; below the nares is the soft palate, with the uvula. 2. On either side of them, near the lower turbinated bones, are the openings of the Eustachian tube. 3. Below the soft palate is the communication with the mouth, called the isthmus faucium. On either side of this are two folds of mucous membrane, constituting the anterior and posterior half-arches of the palate; between them are the tonsils. Below the isthmus faucium is the epiglottis, which is connected to the base of the tongue by three folds of mucous membrane. 4. Below the epiglottis is the

aperture of the larynx. 5. Lastly, is the opening into the œsophagus.[1]

The pharynx consists of three coats, viz., muscular, fibrous and mucous. The two former have been already described.

FIG. 54.



DIAGRAMMATIC VIEW OF THE PHARYNX LAID OPEN FROM BEHIND.

Mucous Membrane.

The *mucous membrane* is common to the entire tract of the respiratory passages and the aliment-

[1] On reflecting the mucous membrane at the pharyngeal termination of the Eustachian tube, a thin, pale muscle, the *scalpingo-pharyngeus*, can be made out. It arises by a thin tendon from the Eustachian tube, and joins the palato-pharyngeus. It is lost among the fibres of the constrictor muscles.

nal. This membrane, however, presents varieties in the nt parts of these channels, according as they are intended ssages for air or for food. The mucous membrane of the ynx above the velum palati, being intended to transmit air , is very delicate in its texture, and lined by columnar ciliated helium like the rest of the air passages. But opposite the es, the mucous membrane resembles that of the mouth, and rovided with squamous epithelium. At the back of the larynx membrane is corrugated into folds, to allow the expansion of pharynx during the passage of the food.

The membrane is lubricated by a secretion from the numerous ucous glands which are situated in the submucous tissue roughout the whole extent of the pharynx, particularly in the eighborhood of the Eustachian tubes.[1]

Posterior Openings of the Nasal Fossæ. These are two oval openings, each of which is about an inch in the long, and half an inch in the short diameter. They are bounded above by the body of the sphenoid bone, externally by its pterygoid plate, below by the horizontal portion of the palate bone; they are separated from each other by the vomer.

On removing the mucous membrane from the posterior part of the roof of the nose and the top of the pharynx, you will find beneath it much fibrous tissue. Hence polypi growing from these parts are, generally, of a fibrous nature.

Isthmus Faucium. This name is given to the opening by which the mouth communicates with the pharynx. It is bounded above by the soft palate and uvula, below by the root of the tongue, and on either side by the arches of the palate, enclosing the tonsils between them.

Soft Palate, or Velum Pendulum Palati. This movable prolongation of the roof of the mouth is attached to the border of the hard palate, and laterally to the side of the pharynx. Posteriorly it has a free edge, with a pendulous, conical projection in the centre, called the *uvula*. It constitutes an imperfect partition between the mouth and the posterior nares. Its upper or nasal surface is convex and continuous with the floor of the nose;

[1] This aggregation of mucous glands is called the pharyngeal tonsil.

its lower surface is concave, in adaptation to the back of the tongue, and is marked in the middle by a ridge or raphé, indicating its original formation by two lateral halves. The soft palate, when at rest, hangs obliquely downward and backward; but in swallowing, it is raised to the horizontal position by the levatores palati, comes into apposition with the back of the pharynx, and thus prevents the food from passing through the nose.

On making a perpendicular section through the soft palate, you come first upon the oral mucous membrane; then you see that the great bulk of it is made up of *muciparous glands*, which lie thick on its under surface to lubricate the passage of the food. Above these glands is the thin layer of the palato-glossus, then the insertion of the tensor palati forming the broad aponeurosis of the palate; still higher, are the two portions of the palato-pharyngeus, separated by the fibres of the levator palati, the azygos uvulæ, and, lastly, the nasal mucous membrane. The soft palate is supplied with blood by the descending palatine branch of the internal maxillary, the ascending palatine branch of the facial, the ascending pharyngeal and the dorsales linguæ of the lingual artery. Its nerves are derived from the palatine branches of the superior maxillary division of the fifth and from the glosso-pharyngeal.

Uvula. The uvula projects from the middle of the soft palate, and gives the free edge of it the appearance of a double arch. It contains a number of muciparous glands, and a small muscle, the *azygos uvulæ*. Its length varies according to the state of its muscle. It occasionally becomes permanently elongated, and causes considerable irritation, a tickle in the throat, and harassing cough. When you have to remove a portion of it, cut off only the redundant mucous membrane.

Arches or Pillars of the Palate. The *soft palate* is connected with the tongue and pharynx by two folds of mucous membrane on each side, enclosing muscular fibres. These are the *anterior* and *posterior arches* or *pillars* of the palate. The *anterior arch* describes a curve downward and forward, from the base of the uvula to the side of the tongue. It is well seen when the tongue is extruded. The *posterior arch*, commencing at the side of the

uvula, curves downward and backward, along the free margin of the palate, and terminates on the side of the pharynx. The posterior arches, when the tongue is depressed, can be seen through the span of the anterior. The pillars of each side diverge from their origin, and in the triangular space thus formed is situated the tonsil. The chief use of the arches of the palate is to assist in deglutition. The anterior, enclosing the *palato-glossi* muscles, contract so as to prevent the food from coming back into the mouth; the posterior, enclosing the *palato-pharyngei*, contract like side curtains, and coöperate in preventing the food from passing into the nose. In vomiting, food does sometimes escape through the nostrils, but one cannot wonder at this, considering the violence with which it is driven into the pharynx.

Muscles of the Soft Palate. The muscles of the soft palate lie immediately beneath the mucous membrane. There are five pairs—namely, the levatores palati, the circumflexi or tensores palati, the palato-glossi, the palato-pharyngei, and the azygos uvulæ. This last pair is sometimes described as a single muscle. To clean the muscles, the soft palate should be made tense by means of hooks, as they are severally dissected.

Levator Palati. This muscle *arises* from the under aspect of the apex of the petrous portion of the temporal bone, and from the under part of the cartilage of the Eustachian tube. It descends obliquely inward, and then passes over the concave border of the superior constrictor into the pharynx, where its fibres spread out and are *inserted* along the upper surface of the soft palate below the azygos uvulæ, meeting those of its fellow in the middle line (Fig. 54). Its action is to raise the soft palate, so as to make it horizontal in deglutition. It is supplied by the descending palatine branch from the spheno-palatine ganglion.

Circumflexus or Tensor Palati. This muscle is situated between the internal pterygoid m. and the internal pterygoid plate of the sphenoid bone. It *arises* by a flattened muscular belly from the scaphoid fossa, and from the spine of the sphenoid; from the vaginal process of the temporal bone and from the outer and

anterior side of the cartilage of the Eustachian tube. Thence it descends perpendicularly, and ends in a tendon which winds round the hamular process, where there is a synovial bursa. Now changing its direction, the tendon passes horizontally inward, and expands into a broad aponeurosis, which is *inserted* into the horizontal plate of the palate bone, and is also connected to its fellow of the opposite side. It gives strength to the soft palate. A synovial membrane facilitates the play of the tendon round the hamular process. Its *action* is to draw down and tighten the soft palate, and, owing to its insertion into the palate bone, also to keep the Eustachian tube open. Its nerve is derived from the otic ganglion, and enters the muscle on its inner aspect.

Axygos or Levator Uvulæ. This consists of two thin bundles of parallel muscular fibres situated one on each side of the middle line. It *arises* from the aponeurosis of the palate and descends along the uvula nearly down to its extremity. It receives its nerve from the descending palatine branch of the spheno-palatine ganglion.

Palato-glossus and Palato-pharyngeus. These muscles are contained within the arches of the soft palate, and the mucous membrane must be removed in order to expose them. The *palato-glossus*, within the anterior arch, proceeds from the anterior surface of the soft palate to the side of the tongue, and is lost in the stylo-glossus muscle. The *palato-pharyngeus*, within the posterior arch, arises from the posterior border of the soft palate by two origins, separated by the levator palati. As it descends its fibres spread out and, passing along the side of the pharynx, blend with the fibres of the inferior constrictor and the stylo-pharyngeus. Both these muscles are supplied by the descending palatine branches of the spheno-palatine ganglion.

Tonsils. The tonsils are two glandular bodies, situated at the entrance of the fauces, between the arches of the soft palate. They are rounded in shape, and their use is to lubricate the fauces during the passage of the food. On their inner surface are visible from twelve to fifteen orifices leading into crypts, which make the tonsil appear like the shell of an almond. Hence, as well as from their oval figure, they are called the *amygdalæ*.

These openings lead into small follicles in the substance of the tonsil, lined by mucous membrane. Their walls are thick, and around them is a layer of closed cells (like Peyer's glands) situated in the submucous tissue. The fluid secreted by these cells is viscid and transparent, in the healthy state; but it is apt to become white and opaque in inflammatory affections of the tonsils, and occasionally accumulates in these superficial depressions, giving rise to the deceptive appearance of a small ulcer, or a slough, or even a false membrane on the part.

The tonsil lies close to the inner side of the internal carotid artery. It is only separated from this vessel by the ascending pharyngeal artery, the superior constrictor, and the aponeurosis of the pharynx. Therefore, in removing a portion of the tonsil, or in opening an abscess near it, the point of the instrument should never be directed outward, but *inward*, toward the mesial line.[1] The tonsil is supplied with blood by the tonsillar and palatine branches of the facial, and by the descending palatine branch of the internal maxillary. Nerves are furnished to it from the glosso-pharyngeal and from Meckel's ganglion.

Eustachian Tube. This canal conveys air from the pharynx to the tympanum. Its orifice is situated opposite the back part of the inferior spongy bone. The direction of the tube from the pharynx is upward, backward, and outward; it is an inch and a half long. The narrowest part is about the middle, and here its walls are in contact. Near the tympanum its walls are osseous, but toward the pharynx they are composed of fibro-cartilage and fibrous membrane. The cartilaginous end, about an inch in length, projects between the origins of the levator and the tensor palati, and gives attachment to some of their fibres. It is situated at the base of the skull, in the furrow between the petrous portion of the temporal and the great wing of the sphenoid bone.

[1] Cases are related by Portal and Béclard, in which the carotid artery was punctured in opening an abscess in the tonsil. The result was immediately fatal hemorrhage. It should, however, be borne in mind that the artery usually injured is the tonsillar branch of the facial artery, and not the internal carotid. The surgical treatment of this accident is, therefore, ligature of the external carotid artery between its superior thyroid and lingual branches, and not ligature of the common carotid artery, as is often recommended.

It adheres closely to the bony furrow, as well as to the fibro-cartilage filling up the foramen lacerum medium. The orifice is not trumpet-shaped, as usually described, but an elliptical slit about half an inch long, and nearly perpendicular. The fibro-cartilage bounds it only on the inner and the upper part of the circumference; the integrity of the canal below is maintained by tough fibrous membrane.

The Eustachian tube is lined by a continuation of the mucous membrane of the pharynx, and covered by ciliated epithelium. That which lines the cartilaginous portion of the tube is thick and vascular, and gradually becomes thinner toward the tympanum. Hence, inflammatory affections of the throat or tonsils are liable to be attended with deafness, from temporary obstruction of the tube.

Mucous glands surround the orifice of the tube, and are similar in nature and function to the glands beneath the mucous membrane of the mouth, the palate and the pharynx.

Hard Palate. The hard palate, formed by the superior maxillary and palate bones, is a resisting surface for the tongue in tasting, in mastication, in deglutition, and in the articulation of sounds. The tissue covering the bones is thick and close in texture, and firmly united to the asperities on the bones. But it is not everywhere of equal thickness. Along the raphé in the mesian line, it is much thinner than at the sides; for this reason, the hard palate is in this situation more prone to be perforated in syphilitic disease.

A thick layer of glands (*glandulæ palatinæ*) is arranged in rows on either side of the hard palate. These glands become more numerous and larger toward the soft palate. Their orifices are visible to the naked eye. The mucous membrane has a very thick epithelial coat, which gives the white color to the palate. The descending palatine branch of the internal maxillary artery, and the palatine nerves from the superior maxillary, may be traced along each side of the roof of the mouth. The ramifications of these arteries and nerves supply the soft as well as the hard palate.

Mechanism of Deglutition. With the anatomy of the parts fresh in your mind, consider for a moment the mechanism of deglutition. The food, duly masticated, is collected into a mass upon the back of the tongue; the lower jaw is then closed, to give a fixed point for the action of the muscles which raise the os hyoides and larynx, and the food is carried back into the pharynx by the pressure of the tongue against the palate, at the same time that the pharynx is elevated and expanded to receive it (by the stylo-pharyngei on each side).[1] The food, having reached the pharynx, is prevented from ascending into the nasal passages by the approximation of the posterior palatine arches and the elevation of the soft palate, which thus forms a horizontal temporary roof to the pharynx; it is prevented from returning into the mouth by the pressure of the retracted tongue and the contraction of the anterior palatine arches; it cannot enter the larynx, because its upper opening is closed and protected by the falling of the epiglottis;[2] consequently, being forcibly compressed by the constrictors of the pharynx, the food passes into the œsophagus, through which it is conveyed into the stomach by the undulatory contraction of that tube.

The food passes with different degrees of rapidity through the different parts of its course; but most rapidly through the pharynx. The necessity of this is obvious, as the air tube must be closed while the food passes over it, and the closure produces a temporary interruption to respiration. The progress of the food through the œsophagus is slow and gradual.

DISSECTION OF THE LARYNX.

Situation and Relations. The larynx is the upper dilated part of the windpipe, in which phonation takes place. It consists of numerous cartilages articulated together to form an

[1] The larynx being also elevated and drawn forward, a greater space is thus left between it and the vertebræ for the distention of the pharynx.

[2] This falling of the epiglottis is effected, not by special muscular agency, but by the simultaneous elevation of the larynx and the retraction of the tongue. A perpendicular section through all the parts concerned is necessary to show the working of this mechanism.

open tube, and to protect the delicate structures concerned in vocalization.

It forms a prominence in the middle line of the neck, covered in front by the integument and cervical fasciæ, the sterno-hyoid, sterno-thyroid and thyro-hyoid muscles, and the thyroid body. It has the large vessels of the neck on each side. Above, it is attached to the hyoid bone; below, it is continuous with the trachea; behind it is the pharynx, into the anterior part of which it opens.

Before commencing the dissection of the larynx the student should make himself acquainted with the cartilages which compose it, and the ligaments which connect them, as seen in a dry preparation.

Os Hyoides. This bone, named from its resemblance to the Greek Upsilon, is situated between the larynx and the tongue, and serves for the attachment of the muscles of the tongue. It may be felt immediately below and one inch and a half behind the symphysis of the jaw. It is arched in shape, and consists of a body, two greater and two lesser cornua. The *body* (*basi-hyal*) is the thick central portion. Its anterior surface is convex, and has a median vertical ridge, on each side of which are depressions for the attachments of muscles; its posterior surface is smooth, deeply concave, and corresponds to the epiglottis. The *greater cornua* (*thyro-hyals*), right and left, project backward for about an inch and a half, with a slight inclination upward, and terminate in blunt ends tipped with cartilage. In young subjects they are connected to the body of the bone by fibro cartilage; this, in process of years, becomes ossified. The *lesser cornua* (*cerato-hyals*) are connected, one on each side, to the point of junction between the body and the greater cornua, by means of a little joint lined with synovial membrane, which admits of free motion. They are of the size of a barleycorn, and give attachment to the stylo-hyoid ligaments.

Ligaments. The os hyoides is connected with the thyroid cartilage by several ligaments, which contain a quantity of elastic tissue. There is:—1. The *thyro-hyoid membrane,* a broad, fibrous membrane, which proceeds from the supe-

rior border of the thyroid cartilage to the upper and posterior part of the hyoid bone. In front of this membrane there is a bursa, of which the use is to facilitate the play of the thyroid cartilage behind the os hyoides. The central portion is stronger than the lateral, and is called the *anterior thyro-hyoid ligament.* Through the lateral part of this membrane the superior laryngeal nerve and artery enter the larynx. 2. The right and left *lateral thyro-hyoid ligaments* extend between the extremities of the greater cornua of the os hyoides, and the ascending cornua of the thyroid cartilage. They contain a small nodule of cartilage (*cartilago triticea*).

Cartilages of the Larynx. The framework of the larynx is composed of nine cartilages—viz., the thyroid, the cricoid, the two arytenoid, the two cornicula laryngis, the two cuneiform cartilages and the epiglottis. These are connected by joints and elastic ligaments, so that they can be moved upon each other by their respective muscles; the object of this motion being to act upon two elastic ligaments, called the *vocal cords*, upon the vibration of which phonation depends.

Thyroid Cartilage. This cartilage, so called because it shields the mechanism behind it,[1] consists of two lateral halves (*alæ*), united at an acute angle in front, which forms the prominence termed the *pomum Adami.* This prominence presents a notch at its upper part, to allow it to play behind the os hyoides in deglutition. There is a bursa in front of it. Each ala is somewhat quadrilateral in form, and presents for examination two surfaces and four borders. The *outer surface* of each ala is marked by an oblique line passing downward and forward from the base of the upper cornu, which gives attachment to the sterno-thyroid and thyro-hyoid muscles. The smooth surface behind the ridge gives attachment to the inferior constrictor. The *inner surface* is smooth, slightly concave, and is covered with mucous membrane. In the acute angle in front there are attached, from above downward, the epiglottis, the false and true vocal cords, the thyro-arytenoidei and thyro-epiglottidei muscles. The *inferior* border is slightly arched in the middle, affording attachment to the crico-

[1] θυρεός, a shield.

thyroid membrane, and on either side presents a convex prominence, which gives attachment to the crico-thyroid muscle and the crico-thyroid membrane. The *superior* border is nearly horizontal, and affords attachment to the thyro-hyoid membrane. The *posterior* border is thick, rounded and nearly vertical, and gives insertion to the stylo-pharyngeus and palato-pharyngeus muscles. This border terminates, above and below, in round projections called the *upper* and *lower cornua*. The upper is the longer; the lower articulates with the side of the cricoid cartilage.

Cricoid Cartilage. This cartilage, named from its resemblance to a ring,[1] is situated below the thyroid. It is not of equal depth all round. It is narrow in front, where it may be felt about a quarter of an inch below the thyroid; from this part, the upper border gradually rises, so that, posteriorly, the ring is an inch in vertical depth, and occupies part of the interval left between the alæ of the thyroid. In the middle of this broad posterior surface is a vertical ridge, on either side of which observe a superficial excavation for the origin of the crico-arytenoidei postici; to the lower part of the vertical ridge are attached some of the longitudinal fibres of the œsophagus. On its *upper* part are two oval slightly convex surfaces for the articulation of the arytenoid cartilages, between which is a concavity for the attachment of the arytenoideus. In front, its upper border presents a broad excavation to which the crico-thyroid membrane (on which is seen the crico-thyroid artery), is attached. On its *outer* surface, external to the depression for the crico-arytenoideus posticus, is an elevated facet which articulates with the inferior cornu of the thyroid cartilage. In front of this articular surface it gives attachment to the inferior constrictor of the pharynx. The *lower* border is straight, and is connected by fibrous membrane to the first ring of the trachea. The *inner* surface is smooth, and the upper border is elliptical; its lower being nearly circular.

Ligaments. The thyroid cartilage is connected to the cricoid by a membrane—the *crico-thyroid*. It consists of a median triangular portion, composed mainly of

[1] *Κρίκος*, a ring.

elastic tissue, with its base directed downward. The lateral portions are thin and membranous, extending as far backward as the articular facets for the thyroid cartilage, and are intimately connected with the inferior vocal cords. Between the inferior cornu of the thyroid cartilage and the cricoid there is a distinct joint, having a synovial membrane, and strengthened by a *capsular ligament*. The articulation allows of a movement revolving upon its own axis, and, consequently, permits the approximation of the two cartilages.

Arytenoid Cartilages. These cartilages are situated, one on each side, at the back of the upper border of the cricoid cartilage. In the recent state, before the membranes and muscles have been removed, the space between them resembles the lip of a pitcher;[1] hence their name. Each is pyramidal, with the apex upward, is about five or six lines in height, and three lines in diameter at its base, and presents for examination three surfaces (marked off by three borders), a base and an apex. The *posterior* surface of each is triangular and concave, and gives attachment to the arytenoideus muscle; the *anterior* surface is irregular and convex, affording attachment to the thyro-arytenoideus, and to the superior or false vocal cord; the *internal* surface, the narrowest and nearly flat, faces the corresponding surface of the opposite cartilage, and is covered with mucous membrane. The *base* is broad, and presents a smooth, somewhat concave, triangular surface, which articulates with the cricoid cartilage; in front of the base is the pointed *anterior angle*, which gives attachment to the true vocal cord, and contributes to form part of the boundary of the rima glottidis; at the outer and back part of the base is the *external angle*, into which certain muscles moving the cartilage are inserted, viz., the crico-arytenoideus posticus and crico-arytenoideus lateralis. The base is articulated with the cricoid by a joint which has a loose capsular ligament and a synovial membrane, permitting motion in all directions, like the first joint of the thumb. The *apex* is truncated and points backward and inward. It is surmounted by a cartilaginous nodule, called the *corniculum laryngis*.

[1] Ἀρύταινα, a pitcher.

Cornicula Laryngis. Are two small conical cartilaginous nodules, and continue the direction of the arytenoid cartilages upward and inward.

Cuneiform Cartilages. These cartilages, sometimes called the cartilages of Wrisberg, are conical in form, and somewhat curved, with their broader part directed upward and forward. They are contained in the aryteno-epiglottidean fold.

Epiglottis. This piece of yellow fibro-cartilage is situated in the middle line, and projects over the larynx like a valve. It is like a leaf with its stalk directed downward. Its ordinary position is perpendicular, leaving the upper opening of the larynx free, for respiration; but during the elevation of the larynx in deglutition it becomes horizontal, falls downward and backward over the larynx, and prevents the entrance of food into it. This descent of the epiglottis is accomplished, not by special muscular agency, but by the simultaneous elevation of the larynx and the retraction of the tongue. Its *apex* or lower part is attached by the *thyro-epiglottic ligament* to the angle of the thyroid cartilage; it is also connected by an elastic ligament, *hyo-epiglottic*, to the posterior surface of the os hyoides. *Laterally*, its borders are rather turned backward, and to them are attached two folds of mucous membrane, which pass to the arytenoid cartilages, called the *aryteno-epiglottic folds*. Its *anterior surface* is only free at its base, where it is connected with the base of the tongue by the three *glosso-epiglottic folds*. Its *posterior* or *laryngeal surface* is smooth, concavo-convex and free, and looks toward the larynx. The surface of the epiglottis is closely invested by mucous membrane; this being removed, the yellow cartilage of the epiglottis is seen pitted and often perforated by the small mucous glands.

The cartilages of the larynx resemble those of the ribs in structure. In the young they are dense and elastic, but some have a tendency to ossify with age. In very old subjects, the thyroid and cricoid cartilages are often completely ossified, and their interior presents an areolar tissue, containing oily matter, analogous to the spongy texture of the bones. The epiglottis, cornicula laryngis, and cuneiform cartilages are rarely ossified, on account

of their consisting of yellow fibro-cartilage resembling that of the ear and nose.

The larynx must now be examined in its perfect condition.

Mucous Membrane of the Larynx.

The mucous membrane lines the whole of the interior of the larynx, being continuous above with that of the pharynx and mouth, below with that of the trachea. It is intimately adherent to the posterior part of the epiglottis and to the true vocal cords; elsewhere it is loosely connected to the subjacent structures by an abundance of areolar tissue, which admits of its being elevated into large folds. This is chiefly found about the upper opening of the larynx, and it deserves notice from the rapidity with which it becomes the seat of serous effusion in acute inflammation of the larynx, and thus produces symptoms of suffocation. In the remaining part of the interior of the larynx the mucous membrane is moderately adherent to the subjacent tissues, and at the upper or false vocal cord it reduplicates upon itself and then lines the sacculus laryngis. Naturally, the mucous membrane is of pale rose color, except where it covers the cushion of the epiglottis, where it is bright pink. It is covered by columnar ciliated epithelium below the false vocal cords, and this variety is continued up the epiglottis as high as its middle; above this, by squamous epithelium. From the root of the tongue to the anterior surface of the epiglottis, the membrane forms three folds, *glosso-epiglottic*, one median, and two lateral, containing elastic tissue. From the epiglottis, to which it is intimately adherent, it is continued backward on either side to the apices of the arytenoid cartilages, forming the *aryteno-epiglottic folds* which bound the upper entrance into the larynx.

The mucous membrane of the larynx is remarkable for its acute sensibility. This is requisite to guard the upper opening of the larynx during the passage of the food over it. The larynx is closed during the act of deglutition; but if, during this process, any one attempt to speak or laugh, the epiglottis is raised, and allows the food to pass, as it is termed, the wrong way. As soon as the foreign body touches the mucous membrane of the larynx, a spasmodic fit of coughing expels it.

The submucous tissue of the larynx is studded with mucous glands. An oblong mass of them lies in the aryteno-epiglottic fold, and they are particularly numerous about the ventricles of the larynx. The surface of the epiglottis toward the tongue is abundantly provided with them. Their ducts pass through the epiglottis, and may be recognized as minute openings on the laryngeal aspect.

Superior Opening of the Larynx. This is the opening through which the larynx communicates with the pharynx. Its outline is triangular, with its base directed forward, and it slopes from before backward. Anteriorly it is bounded by the epiglottis, laterally by the aryteno-epiglottic folds and cuneiform cartilages, posteriorly by the arytenoid cartilages and the cornicula laryngis. The apex presents the funnel-shaped appearance from which the arytenoid cartilages derive their name.

On looking down through this superior opening you see the *cavity of the larynx*, which is divided into an upper and a lower part of the narrow triangular fissure, called the *glottis* or *rima glottidis;* so that the upper part gradually narrows to this chink, while the lower part gradually widens and becomes continuous at the lower border of the cricoid cartilage with the trachea.

The objects seen above the rima glottidis are, in the middle line, below the base of the epiglottis, a round elevation covered with mucous membrane of bright pink color, termed the *cushion* of the epiglottis; on each side is an arched fold, the *false vocal cords*, with their concavity looking downward, and forming the upper boundary of a small recess, the *ventricle of the larynx*, leading into a pouch, called the *sacculus laryngis;* below this are the two white bands, the *true vocal cords*, which form the boundaries of the glottis. The larynx below the true vocal cords gradually enlarges and presents nothing calling for special description.

Glottis, or Rima Glottidis. The *rima glottidis* is the triangular horizontal opening between the inferior or true vocal cords. Its apex is directed forward, its base backward. The anterior two-thirds of this opening is bounded by the true vocal cords, the posterior third by the arytenoid cartilages. The length in the male is eleven lines, its width at rest from three to four lines; in

the female its length is eight lines, its width two lines. Before the age of puberty these dimensions are much less.

Superior or False Vocal Cords. These are the prominent crescentic folds of mucous membrane which form the upper boundaries of the ventricles and enclose within them thin ligamentous fibres, called the *superior thyro-arytenoid ligaments*. They are called the *false vocal cords*, because they have little or nothing to do with the production of the voice. They are composed of elastic tissue like the true vocal cords; but they also contain fatty tissue, which the true ones do not.

Inferior or True Vocal Cords. These two cords, called also the *inferior thyro-arytenoid ligaments*, are composed of yellow elastic tissue, and extend horizontally from the angle of the thyroid cartilage to the anterior angles of the base of the arytenoid cartilages. Their inner or free edges are thin and sharp, and look upward; their outer borders are continuous with the crico-thyroid membrane, and are in contact with the thyro-arytenoidei muscles. They diverge as they pass backward, and are covered with very thin and closely adherent mucous membrane, having columnar ciliated epithelium. We shall presently see that, by the muscles which act upon the arytenoid cartilages, these cords can be approximated or separated from each other; in other words, the rima glottidis can be closed or dilated. When sufficiently tightened, and brought parallel by means of certain muscles, the cords are made to vibrate by the current of the expired air, and thus is produced the voice.

FIG. 55.



SHAPE OF THE GLOTTIS WHEN AT REST.

In the adult male the true vocal cords measure about seven lines; in the female, about five lines. In boys they are shorter; hence their peculiar voice. At puberty, the cords lengthen and the voice breaks.

The glottis admits of being dilated, contracted, and even com-

pletely closed by its appropriate muscles. When at rest, its shape is triangular, as shown in Fig. 55, where the arytenoid cartilages are cut through on a level with the vocal cords. During every inspiration, the glottis is dilated by the crico-arytenoidei postici; it then becomes pear-shaped (Fig. 57). During expiration, it resumes its triangular shape; and this return to a state of rest is effected, not by muscular agency, but by two elastic ligaments shown in Fig. 55, which draw the arytenoid cartilages together. Thus, then, the glottis, like the chest, is dilated by *muscular* tissue; like the chest, also, it is contracted by *elastic* tissue. In speaking or singing, the glottis assumes what is called the vocalizing posi-

FIG. 56.



DIAGRAM SHOWING THE ACTION OF THE CRICO-THYROID MUSCLE.

tion—that is, the opening becomes narrower, and its edges nearly parallel.

Ventricles of the Larynx. These are the recesses between the upper and lower vocal cords, and each leads to a small conical pouch, the *sacculus laryngis*. Each ascends for about half an inch, as high as the upper border of the thyroid cartilage, which bounds it on its outer side, while on the inner side is the upper vocal cord. It contains from sixty to seventy muciparous glands. Over its inner and upper part is a layer of muscular tissue, *compressor sacculi laryngis* of Hilton (aryteno-epiglottideus inferior), which connects it with the aryteno-epiglottic fold; on its outer side is the upper part of the thyro-arytenoideus.

Intrinsic Muscles of the Larynx. There are eleven muscles which act upon the larynx, five on each side and one in the middle. The five pairs are—the crico-thyroidei, the crico-arytenoidei postici, the crico-arytenoidei laterales, the thyro-arytenoidei, and the aryteno-epiglottidei. The single one is the arytenoideus.

M. Crico-Thyroideus. This muscle is situated on the front of the larynx. It *arises* from the front and side of the cricoid cartilage, ascends obliquely outward, and is *inserted* into the inferior border and lower cornu of the thyroid. Its *action* is to tighten the vocal cords. It does this by raising the anterior part of the cricoid cartilage, since this cartilage cannot be raised

FIG. 57.



GLOTTIS DILATED: MUSCLES DILATING IT REPRESENTED WAVY.

without lengthening these cords, as shown by the dotted line, Fig. 56. Its nerve is the *external laryngeal* branch of the superior laryngeal. Between the anterior borders of the two muscles is seen the crico-thyroid membrane, which is divided in laryngotomy.

M. Crico-Arytenoideus Posticus. This muscle *arises* from the broad depression on the posterior part of the cricoid cartilage; its fibres converge, and pass outward and upward, to be *inserted* into the outer angle of the base of the arytenoid. Its *action* is to dilate the glottis. It does this by drawing the posterior tubercle of the arytenoid cartilage *toward* the mesial line, and therefore the anterior angle (to which the vocal cord is

attached) *from* the mesial line (Fig. 57). In this movement th arytenoid cartilage rotates as upon a pivot, and acts as a leve of the first order; the fulcrum or ideal pivot being interme diate between the power and th weight. This muscle dilates th glottis at each inspiration. It nerve comes from the inferior la ryngeal.[1]

FIG. 58.



SIDE VIEW OF THE MUSCLES OF THE LARYNX.

1. Thyro-epiglottideus.
2. Thyro-arytenoideus, upper and lower portions.
3. Crico-arytenoideus lateralis.
4. Crico-arytenoideus posticus.
5. Arytenoideus.

M. Arytenoideus. This singl muscle is situate immediately at the back of th arytenoid cartilages. The fibre pass across from one cartilage t the other, running in a transvers direction. *Action.*—By approxima ing the arytenoid cartilages, the assist in contracting the glotti It is supplied by the inferior l ryngeal nerve.

M. Aryteno-Epiglottideus. This muscle *aris* from the inferi and outer angle of the aryteno cartilage, and, crossing its fello like the letter X, is *inserted*, part into the apex of the opposite ar tenoid cartilage, and partly into th aryteno-epiglottic fold.

M. Crico-Arytenoideus Lateralis. To expose th muscle, reflect t crico-thyroid muscle, the crico-thyroid membrane, and then c away the ala of the thyroid cartilage. It *arises* from the upp border of the side of the cricoid cartilage, and the fibres, passi

[1] A slip called the "kerato-cricoid" is occasionally present. It is a short fasci lus passing from the cricoid to be inserted into the inferior cornu of the thyroid ca lage. It is in connection with the crico-arytenoideus posticus, and is usually fou on one side only, being present in about one in five subjects. (Merkel, *Anat. u Phys. des mensch. Stimme- und Sprach-Organs*, 1857; Turner, *Month. Med. Jour* Feb. 1860).

backward and upward, converge to be *inserted* into the external angle of the base of the arytenoid, in front of the crico-arytenoideus posticus. *Action.*—By drawing the arytenoid cartilages inward, the muscles of opposite sides contract the glottis (Fig. 59). Its nerve comes from the inferior laryngeal.

M. Thyro-Arytenoideus. This muscle *arises* from the side of the angle of the thyroid cartilage and the crico-thyroid membrane, runs horizontally backward, and is *inserted* into the base and anterior surface of the arytenoid. Its fibres run parallel with the true vocal cord, and some of them are directly inserted into it. It consists of two fasciculi—an upper and a lower. The *lower* and larger portion is inserted into the anterior

FIG. 59.

GLOTTIS CLOSED; MUSCLES CLOSING IT REPRESENTED WAVY.

angle and the anterior surface of the arytenoid; the *upper* is inserted into the upper part of the anterior surface and the anterior border of the arytenoid. Its nerve comes from the inferior laryngeal.

This muscle relaxes the vocal cord. More than this, it puts the lip of the glottis in the vocalizing position; in this position the margins of the glottis are parallel, and the chink is reduced to the breadth of a shilling.

The following table shows the action of the several muscles which act upon the glottis:—

Crico-thyroidei . . .	Stretch the vocal cords.
Thyro-arytenoidei . .	Relax the vocal cords, and place them in the vocalizing position.

Crico-arytenoidei postici	Dilate the glottis.	
Crico-arytenoidei laterales	Draw together the arytenoid cartilages	close the glottis.
Arytenoideus	" " "	close the glottis.
Aryteno-epiglottidei	Contract the upper opening of the larynx.	

The epiglottis is connected by muscles with the arytenoid and thyroid cartilages; they are the thyro-epiglottideus, the aryteno-epiglottideus superior and inferior.

The *thyro-epiglottideus* is a thin muscle, *arising* from the angle of the thyroid cartilage just above the thyro-arytenoideus, and *inserted* by diverging fibres into the border of the epiglottis and into the aryteno-epiglottic fold.

The *aryteno-epiglottideus superior* passes as thin muscular fibres from the tip of the arytenoid cartilage to the mucous membrane attached to the side of the epiglottis.

The *aryteno-epiglottideus inferior*, separated from the preceding by a distant interval, *arises* from the anterior surface of the arytenoid cartilage, and is *inserted* into the upper and inner part of the epiglottis. This muscle is also called the *compressor sacculi laryngis* of Hilton.[1]

The *blood vessels of the larynx* are derived from the *superior* and *inferior thyroid arteries*. The laryngeal branch of the superior thyroid passes through the thyro-hyoid membrane with the corresponding nerve, and divides into branches, which supply the muscles and the mucous membrane. The laryngeal branches of the inferior thyroid ascends behind the cricoid cartilage. A constant little artery passes through the crico-thyroid membrane.

The *nerves* of the larynx are the *superior* and *inferior* (*recurrent*) *laryngeal* branches of the pneumogastric.

The *superior laryngeal*, having passed through the thyro-hyoid membrane, divides into branches, distributed to the mucous membrane of the larynx. Its filaments spread out in various directions; some to the anterior and posterior surfaces of the epiglottis, and to the aryteno-epiglottidean folds, others to the interior

[1] The *triticeo-glossus* is a small muscle frequently present; it arises from the cartilago triticea, and passing forward and upward joins the cerato-glossus, to be inserted into the tongue.

of the larynx and the vocal cords. A constant filament descends behind the ala of the thyroid cartilage, and communicates with the inferior laryngeal, and another communication with the same nerve is found behind the larynx beneath the pharyngeal mucous membrane. Its external laryngeal branch supplies the crico-thyroid muscle.

The *inferior* (or *recurrent*) *laryngeal nerve* enters the larynx beneath the inferior constrictor, and ascends behind the joint between the thyroid and cricoid cartilages. It supplies all the intrinsic muscles of the larynx except the crico-thyroid. If the recurrent nerve be divided, or in any way injured, the muscles moving the glottis become paralyzed, but its sensibility remains unimpaired. When the nerve is compressed by a tumor—for instance, an aneurism of the arch of the aorta—the voice is changed to a whisper,[1] or even lost.

Difference between the Male and the Female Larynx. Until the approach of puberty, there is no great difference in the relative size of the male and female larynx. The larynx of the male, within two years after this time, becomes nearly doubled in size; that of the female grows, but to a less extent.

The larynx of the adult male is in all proportions about one-third larger than that of the adult female.

The alæ of the thyroid cartilage form a more acute angle in the male; hence the greater projection of the pomum Adami and the greater length of the vocal cords in the male.

The average length of the vocal cords is in the	Male	7 lines.
	Female	5 "
The average length of the glottis is in the	Male	11 "
	Female	8 "

The size of the larynx does not necessarily follow the proportions of the general stature; it may be as large in a little person as in a tall one; this corresponds with what we know of the voice.

[1] *Medical Gazette*, Dec. 1843.

DISSECTION OF THE TONGUE.

The tongue is a complex muscular organ, subservient to taste, speech, suction, mastication and deglutition. It is situated in the space formed by the lower dental arch; its upper surface is convex and free, as is also its anterior part or tip, which lies behind the lower incisor teeth; its posterior and inferior part is connected to the os hyoides by the hyo-glossi, to the styloid process of the temporal bone by the stylo-glossi, and to the symphysis of the lower jaw by the genio-hyo-glossi muscles.

The upper surface or *dorsum* is convex, and slopes on all sides from the centre; it is divided into two symmetrical halves by a median groove—*raphé*—running along the middle, and terminates posteriorly in a depression—the *foramen cæcum*—into which open several mucous glands. The posterior third of the dorsum is comparatively smooth; the anterior two-thirds is rough, and covered with small eminences called *papillæ*.

Mucous Membrane. The surface of the tongue is covered with mucous membrane, which is composed of structures similar to those of the skin generally—that is to say, it consists of a cutis vera with numerous elevations called papillæ, and of a thick layer of squamous epithelium. The cutis is much thinner than that of the skin of the body, and affords insertion to some muscular fibres of the tongue.

The mucous membrane on the *under aspect* of the tongue is smooth and comparatively thin, and, in the middle line in front, forms a fold—the *frænum linguæ*—which connects it to the mucous membrane of the floor of the mouth. On each side of the frænum are the elevated orifices of the submaxillary ducts; and further back, in the furrow between the tongue and gums, are the openings of the sublingual ducts. *Laterally*, the mucous membrane is reflected from the under part of the tongue to the lower jaw, and forms the floor of the mouth.

From the posterior part of the tongue the mucous membrane passes to the soft palate on each side, forming the folds termed the *anterior palatine arches*, which enclose the palato-glossi; there are also three folds to the epiglottis, termed the *glosso-epiglottic—two*

lateral and *one median*—the latter enclosing a layer of elastic tissue called the *glosso-epiglottic ligament.* This ligament raises the epiglottis when the tongue is protruded from the mouth; hence the rule of never pulling the tongue forward when passing a tube into the œsophagus, otherwise the tube might pass into the larynx.

Papillæ of the Tongue. The anterior two-thirds of the tongue is studded with numerous small eminences called *papillæ;* these, according to their size and form, are distinguished into three kinds, viz., *papillæ circumvallatæ, papillæ fungiformes* and *papillæ filiformes* (Fig. 60).

FIG. 60.

UPPER SURFACE OF THE TONGUE, WITH THE FAUCES AND TONSILS.
1. Papillæ circumvallatæ.
2. Papillæ fungiformes.

The *papillæ circumvallatæ* vary in number from eight to twelve, and are arranged at the back of the tongue in two rows, which converge like the branches of the letter V, with the apex backward, toward the foramen cæcum. Each of these papillæ is circular, from the $\frac{1}{16}$ to $\frac{1}{12}$ of an inch wide, and slightly broader above than below. Each is surrounded by a circular fossa, which itself is bounded by an elevated ring (*vallum*). The papillæ are covered with a thick stratum of scaly epithelium, beneath which are numerous secondary papillæ. Buried in the epithelium surrounding the sides (but not on the upper surface) of these papillæ, numerous flask-shaped bodies, called the *taste-buds*, have been discovered. Their bases rest upon the corium, and their apices open upon the surface. Each consists of a cortical and a medullary portion; the cortical is made up of one or more layers of long, flattened cells; the central consists of numerous spindle-shaped cells, whose free extremity projects from the orifice of the taste-bud, while the deeper extremity rests in the corium, and is in close connection with a fine plexus of nerve filaments derived from the glosso-pharyngeal nerve.[1]

[1] For further information about these bodies the student is referred to Engelmann, in Stricker's *Handbook*. He states that each taste-bud consists of from fifteen to thirty cells

[illegible] smaller and more numerous than the [illegible] are scattered [illegible] and in front of the [illegible] are [illegible]. They vary in [illegible] the long epithelial [illegible] rounded heads like [illegible]. Near the apex of the tongue [illegible] to distinguish during life from the other papillæ by [illegible]. In [illegible] and some cruciform [illegible] vascular, elongated, and of a bright red color; on [illegible] these points acquire a brownish tint, giving [illegible] to what is called the strawberry tongue.

The papillæ filiformes (conicæ), are the smallest and most numerous. They are so closely aggregated that they give the tongue a velvet-like appearance. Their points are directed backwards, so that the tongue feels smooth if the finger be passed [illegible] from apex to base, but rough if in the contrary direction. These papillæ consist of small conical processes arranged for the most part in a series of lines running parallel to the two rows of the papillæ circumvallatæ. Each papilla is covered with a thick layer of epithelium, which is prolonged into a number of free, hair like processes.

If the papillæ be injected, and examined under the microscope, it is found that they are not simple elevations, like those of the skin, but that from them arise secondary papillæ. The papillæ circumvallatæ consist of an aggregation of smaller papillæ arranged parallel to each other; and the papillæ fungiformes consist of central stems from which minute secondary papillæ shoot off. This elaborate structure escapes observation because it is buried beneath the epithelium.[1] Each secondary papilla receives a blood vessel, which passes nearly to its apex, and returns in a loop like manner.

The papillæ are covered with one or more layers of squamous epithelium. That which covers the filiform is supernumerary to such a degree as to give it sometimes the appearance of a [illegible] when viewed by the microscope. The various forms of [illegible] consist of thick and sodden epithelium.

Regarding the use of the papillæ, it is probable that they

[1] See Kölliker and Todd's Physiological Anatomy.

enable the tongue to detect impressions with greater delicacy. From the density and arrangement of their epithelial coat, the filiform papillæ give the surface of the tongue a roughness, which is useful in its action upon the food. An apparatus of this kind, proportionately stronger and more developed, makes the tongue of ruminant animals an instrument by which they lay hold of their food. In the feline tribe—*e. g.* the lion and tiger—these papillæ are so sharp and strong that they act like rasps, and enable the animal to lick the periosteum from the bones by a single stroke of his tongue. In some mammalia, they act like combs for cleaning the skin and the hair.

Glands. Numerous small racemose and acino-tubular glands, *lingual glands*, are found in the sub-mucous tissue at the root of the tongue. They are similar in structure and secretion to the tonsillar and palatine glands, so that there is a complete ring of glands round the isthmus faucium. Small, round orifices upon their surface indicate the termination of their ducts. Other mucous glands, with ducts from one-quarter to half an inch long, are situated in the muscular substance of the tongue.

Lymphoid Tissue. A considerable amount of *lymphoid tissue* is situated at the back of the tongue, which in some parts is collected into definite masses called *follicles*. Small depressions also occur in this situation, whose walls are studded with lymphoid tissue, and into which some of the mucous glands open.

Glands Beneath the Apex of the Tongue. On the under surface of the apex of the tongue is placed, on either side, a group of glands presumed to be salivary. Considering each group as one gland, observe that it is oblong, with the long diameter from seven to ten lines, parallel with the axis of the tongue. It lies near the mesial line, a little below the ranine artery, on the outer side of the branches of the gustatory nerve, under some of the fibres of the stylo-glossus. Four or five ducts proceed from each group, and terminate by separate orifices on the under surface of the tongue.

Muscular Fibres of the Tongue. The substance of the tongue is composed of muscular fibres and of a small quantity of fat.

The *extrinsic* muscles of the tongue have been described in the dissection of the submaxillary region (p. 120). We have now to examine its *intrinsic* muscles. For this purpose the mucous membrane must be removed from the dorsum of the tongue. On dissection it will be found that the great bulk of the organ consists of fibres which proceed in a longitudinal direction, constituting the *linguales* muscles.

The *superficial lingualis* runs longitudinally beneath the mucous membrane of the dorsum; its fibres are attached posteriorly to the hyoid bone and run forward to the front and margin of the tongue. Posteriorly the muscle is thin and is covered by the fibres of the palato-glossus and hyo-glossus.

The *inferior lingualis* is larger than the preceding, and is situated on the under aspect of the tongue between the genio-hyo-glossus and the hyo-glossus. It may be readily exposed by dissecting the under surface of the tongue immediately on the outer aspect of the genio-hyo-glossus. It *arises* posteriorly from the hyoid bone and the substance of the tongue, and its fibres pass forward to the tip of the tongue after being reinforced by fibres from the stylo-glossus. On its under aspect it is in relation with the ranine artery.

The *transverse fibres* form a considerable part of the thickness of the tongue, and arise from the fibrous septum. They pass outward between the superficial and inferior linguales, ascending as they near the sides of the tongue, where the fibres become continuous with those of the palato-glossus. A considerable amount of fat is found among these fibres.

The *vertical fibres* run in a curved direction, descending from the dorsum to the under aspect of the tongue, with the concavity outward. They interlace with the transverse fibres and with the genio-hyo-glossus.

On tracing the genio-hyo-glossi into the tongue, we find that some of their fibres ascend directly to the surface; others cross in the middle line, intersect the longitudinal fibres, and finally terminate upon the sides of the tongue. Lastly, the fibres of the stylo-glossi should be traced along the side of the tongue to the apex.

Fibrous Septum of the Tongue.

The *fibrous septum* of the tongue is a vertical plane of fibrous tissue which extends, in the mesial line, from the base to the apex. It is thick posteriorly, where it is connected behind with the hyoid bone, and is lost in front between the muscles. In it is sometimes found a piece of fibro-cartilage, called *nucleus fibrosus linguæ*, a representative of the lingual bone in some of the lower animals.

The *arteries* supplying the tongue are the dorsal and ranine branches of the lingual artery. It is important to bear in mind that the arteries do not anastomose across the middle line, and only very slightly at the apex, so that it is possible to cut along the septum of the tongue from the apex to the base with very little hemorrhage—a fact of much importance in the removal of the tongue or cancer of that organ.

The *nerves* to the tongue should now be followed to their termination. The *hypoglossal* supplies with motor power all the muscles. The *gustatory* or *lingual* branch of the inferior division of the fifth is distributed to the mucous membrane and papillæ of the apex and sides of the tongue, supplying the anterior two-thirds with common sensation. Upon this nerve depends the sensation of all ordinary impressions, such as that of hardness, softness, heat, cold, and the like.

The *glosso-pharyngeal* nerve supplies the mucous membrane at the back and the sides of the tongue, and the papillæ circumvallatæ. Under the microscope small ganglia may be distinguished on the terminal fibres of this nerve.

DISSECTION OF THE SUPERIOR MAXILLARY NERVE.

To trace this nerve and its branches we must remove the outer wall of the orbit as far as the foramen rotundum, so as to expose the spheno-maxillary fossa.

The superior maxillary nerve is a sensory nerve, and is the second division of the fifth cranial nerve. Proceeding from the Gasserian ganglion (Fig. 61), it leaves the skull through the foramen rotundum, and passes horizontally forward across the spheno-maxillary fossa. It then passes into the orbit through the spheno-maxillary fissure, enters the infra-orbital canal with the

corresponding artery, and finally emerges upon the face, through the infra-orbital foramen, beneath the levator labii superioris, where it divides into a number of spreading branches, distributed to the lower eyelid, the nose and the upper lip. The branches given off are:—

1. Within the skull.

a. A *recurrent branch*, to the dura mater and the middle meningeal artery, is given off near the Gasserian ganglion.

FIG. 61.



DIAGRAM OF THE SUPERIOR MAXILLARY NERVE.

1. Spheno-palatine ganglion. 2. Otic ganglion.

2. In the spheno-maxillary fossa.

b. The *orbital branch*, already described (p. 80).

c. Two *spheno-palatine branches*, which descend to the spheno-palatine ganglion (Meckel's), situated in the spheno-maxillary fossa (Fig. 61).

d. The *dental branches* are three in number: the two posterior are given off immediately before the nerve enters the infra-orbital canal, and descend along the tuberosity of the superior maxillary bone; the anterior is given off in the infra-orbital canal. The *posterior* branch divides into two branches, which send small fila-

ments to the gums and the mucous membrane of cheek, and then run in bony canals in company with small arteries to supply the molar teeth and the antrum; the *middle* branch passes down in a special canal in front of the antrum to be distributed to the bicuspid teeth; the *anterior* branch, the largest, is given off before the nerve emerges from the infra-orbital foramen, and enters the canal in the front wall of the antrum; it divides into *dental* branches for the supply of the canine and incisor teeth, and into a *nasal* branch for the mucous membrane of the floor of the nasal fossa. The anterior branch while in its bony canal gives off some filaments which join with the posterior dental branches, and above the canine fossa it also forms a communication with a nasal branch from the spheno-palatine ganglion, to form the *ganglion of Bochdalek*.

e. The terminal branch of the superior maxillary nerve is called the *infra-orbital*, which divides on the face into palpebral, nasal, and labial branches. These have been already dissected and described (p. 65).

Dissection. At this stage the student should make the dissection to expose the spheno-palatine ganglion and its branches. To do this, it is necessary to saw through the skull rather on one side of its middle line, so as to expose the cavity of the nose. Search must now be made for the spheno-palatine foramen (just external to which is the spheno-palatine ganglion), which is situated immediately above the posterior extremity of the middle turbinated bone. Remove the mucous membrane at this point, when the terminal branch of the internal maxillary artery, which comes through this foramen into the nose, may be readily made out. The student should next cut away the thin plate of bone which forms the inner boundary of the posterior palatine canal. Then, by tracing upward the branches contained within the canal, he will find the ganglion.

Spheno-palatine Ganglion. This ganglion is called, after its discoverer, *Meckel's ganglion*. It is the largest of the ganglia in connection with the branches of the fifth cerebral nerve, is triangular, convex on its outer surface, of reddish-gray color, about one-fifth of an inch in diameter, and is placed immediately below

the superior maxillary nerve, as it crosses the spheno-maxillary fossa. Like other ganglia, it has three roots—a *sensory* from the superior maxillary; a *motor*, from the great petrosal branch of the facial; and a *sympathetic*, from the carotid plexus.

Its branches pass *upward* to the orbit, *downward* to the palate, *inward* to the nose, and *backward* to the pharynx, as follows:—

FIG. 62.



DEEP VIEW OF THE SPHENO-PALATINE GANGLION, AND ITS CONNECTION WITH OTHER NERVES.
(After Hirschfeld.)

1. Superior maxillary n.
2. Spheno-palatine ganglion, from the lower part of which are seen proceeding the palatine nerves.
3. Posterior superior dental brs.
4. Sixth n. receiving two filaments from the carotid plexus of the sympathetic n.
5. The carotid br. of the Vidian.
6. The great petrosal br. of the Vidian.
7. Lesser petrosal nerve.
8. External deep petrosal n., uniting with lesser petrosal n.
9. The internal deep petrosal nerve joining the great petrosal nerve.
10. Filament to fenestra ovalis.
11. Filament to Eustachian tube.
12. Filament to fenestra rotunda.
13. Chorda tympani.
14. Infra-orbital nerve.
15. Anterior dental n.
16. Junction of posterior and anterior dental filaments.
17. Glosso-pharyngeal n., giving off tympanic branch.

a. Ascending branches.—These are very small, and run through the spheno-maxillary fissure to be distributed to the periosteum of the orbit.[1]

[1] Anatomists describe several branches ascending from the ganglion; one to join the sixth nerve (Bŏck); another to join the ophthalmic ganglion (Tiedemann); and, lastly, some to join the optic nerve through the cilary branches (Hirzel).

b. Descending branches.—To see these the mucous membrane must be removed from the back part of the nose; we shall then be able to trace the nerves through their bony canals. Their course is indicated by their accompanying arteries. They descend through the palatine canals, and are three in number. The *anterior palatine* nerve, the largest, descends through the posterior palatine canal to the roof of the mouth, and then divides into branches, which run in grooves in the hard palate nearly to the gums of the incisor teeth, where it communicates with the naso-palatine nerve. Within its canal it sends two *inferior nasal branches* which enter the nose through foramina in the palate bone to supply the membrane on the middle and lower spongy bones. The *posterior or smaller palatine* descends in the same canal with the anterior, or in a smaller one of its own, and supplies the mucous membrane of the soft palate, the tonsil, and (according to Meckel) the levator palati muscle.[1] The *external palatine* may be traced in a special canal down to the soft palate, where it terminates in branches to the uvula, the palate and tonsil. The two last branches communicate with the tonsillar filaments of the glosso-pharyngeal to form the tonsillar plexus of nerves.

c. Internal branches.—These, three or four in number, pass through the spheno-palatine foramen to the mucous membrane of the nose. To see them clearly, the parts should have been steeped in dilute nitric acid; afterward, when well washed, these minute filaments may be recognized beneath the mucous membrane covering the spongy bones. The *upper nasal* branches, four or five in number, pass inward, and are distributed on the two upper spongy bones, the upper and back part of the septum and the posterior ethmoidal cells. The *naso-palatine* (nerve of Cotunnius) traverses the roof of the nose, distributes branches to the back part of the septum narium, and then proceeds obliquely

[1] According to Longet (*Anat. et Physiol. du Système Nerveux*, Paris, 1842), the posterior palatine nerve supplies the levator palati and the azygos uvulæ with motor power. In this view of the subject the nerve is considered to be the continuation or terminal branch of the motor root of the ganglion; that, namely, derived from the facial. This opinion is supported by cases in which the uvula is stated to have been drawn on one side in consequence of paralysis of the opposite facial nerve.

forward, along the septum, to the foramen incisivum, through which it passes, and finally terminates in the palate behind the incisor teeth, communicating here with the anterior palatine nerve.

d. Posterior branches.—The *pharyngeal nerve* (pterygo-palatine), very small, comes off from the back of the ganglion, and, after passing through the pterygo-palatine canal with its corresponding artery, supplies the mucous membrane of the back of the pharynx and the Eustachian tube. The *Vidian nerve* is the principal branch. It proceeds backward from the posterior part of the ganglion, through the Vidian canal, where it distributes small branches to the back part of the roof of the nose and septum. It then traverses the fibro-cartilage of the foramen lacerum medium, and divides into two branches. Of these two branches, one, the larger—the *carotid*—joins the sympathetic plexus on the outer side of the internal carotid artery; the other, the *great petrosal*, enters the cranium, and runs beneath the Gasserian ganglion and the dura mater in a small groove on the anterior surface of the petrous bone; it then enters the hiatus Fallopii, and joins the facial nerve in the aquæductus Fallopii.

It would seem to be more in accordance with modern views to regard the Vidian nerve, not as dividing to form the carotid and great superficial petrosal branches, but rather as formed by the junction of these branches. In this view, the Vidian runs, not *from*, but *to*, the spheno-palatine ganglion.

Otic Ganglion. The otic ganglion (Arnold's[1]) is situated on the inner side of the inferior maxillary division of the fifth nerve, immediately below its exit through the foramen ovale (Fig. 62). It is oval, of reddish-gray color, and always small. Its *inner* surface is in contact with the circumflexus palati muscle and the cartilage of the Eustachian tube; *behind* it is the middle meningeal artery; *externally* it is in relation with the inferior maxillary nerve, where the motor root joins the sensory root.

This ganglion has branches of connection with other nerves; namely, a *sensory*, from the auriculo-temporal nerve; a *motor*, from the branch of the inferior maxillary, which goes to the internal

[1] J. Arnold, *Diss. Inaug. Med., etc.* Heidelbergæ, 1826.

pterygoid muscle; and a *sympathetic*, from the plexus around the arteria meningea media. It communicates also with the facial and the glosso-pharyngeal nerves by the *lesser petrosal* nerve. This branch passes backward, either through the foramen ovale or the

FIG. 63.

AGRAM OF THE COMMUNICATIONS OF THE FACIAL, GLOSSO-PHARYNGEAL, PNEUMOGASTRIC, SPINAL ACCESSORY, HYPOGLOSSAL, SYMPATHETIC, AND THE TWO UPPER CERVICAL NERVES.

1. Great petrosal nerve.
2. Lesser "
3. External "
4. Nerve to stapedius muscle.
5. Spheno-palatine ganglion.
6. Otic ganglion.

ramen spinosum, or through a small hole between them, and runs neath the dura mater in a minute groove on the petrous bone, cternal to that for the *great petrosal* nerve. Here it divides into

two filaments, one of which joins the facial nerve in the aquæductus Fallopii; the other joins the tympanic branch of the glosso-pharyngeal. These nerves are difficult to trace, not only on account of their minuteness, but because they frequently run in canals in the temporal bone.

The otic ganglion sends a branch forward to the tensor palati, and one backward to the tensor tympani, on the outer side of the Eustachian tube.

DISSECTION OF THE NINTH, TENTH AND ELEVENTH CRANIAL NERVES AT THE BASE OF THE SKULL.

In this dissection we propose to examine the glosso-pharyngeal, pneumogastric and spinal accessory nerves in the jugular fossa, and the ganglia and nerves belonging to them in this part of their course. These are difficult to trace, and cannot be followed unless the nerves have been previously hardened by spirit, and the bones softened in acid. The next thing to be done is to remove the outer wall of the jugular fossa.

Glosso-pharyngeal Nerve. This nerve emerges from the cranium through a separate tube of dura mater, in front of that for the tenth and eleventh cranial nerves. Looking at it from the interior of the skull, we notice that it is situated in front, and rather to the inner side of the jugular fossa, where it lies in a groove.

In its passage through the foramen, the nerve presents two enlargements, termed the *jugular* and the *petrous* ganglia.

The *jugular ganglion*[1] is found upon the nerve immediately after its entrance into the canal of the dura mater, and averages about the $\frac{1}{16}$ of an inch in length and breadth. It is situated on the outer side of the nerve, and does not implicate all its fibres. According to our observation, this ganglion is not infrequently absent (Fig. 63).

The *petrous ganglion*[2] is lodged in a groove in the lower part of the jugular fossa. It is oval, about a quarter of an inch long, and involves all the filaments of the nerve. From it are given off

[1] Müller, *Medicin. Zeitung*, Berlin, 1833. No. 52.

[2] Andersch, *Fragm. Descript. Nerv. Cardiac.* 1791.

branches of communication with other nerves and the tympanic nerve[1] (Fig. 63).

The *communicating branches* which connect this ganglion with the pneumogastric are, one to its auricular branch, and a second to the ganglion of the root. It is also connected with the sympathetic by a small filament from the superior cervical ganglion. Another communicating branch pierces the posterior belly of the digastricus to join the facial just external to the stylo-mastoid foramen.

The *tympanic nerve* (Jacobson's) ascends, through a minute canal in the bony ridge which separates the carotid from the jugular fossa, to the inner wall of the tympanum, grooving the surface of the promontory, where it terminates in six filaments. Of these three are branches of distribution, and three of communication with other nerves. The *branches of distribution* are, one each to the fenestra rotunda and the fenestra ovalis, which pass backward, and one to the Eustachian tube, which is directed forward. The *branches of communication* are four small filaments; one or two traverse a bony canal in the anterior wall of the tympanum, and arching forward, join the plexus on the outer side of the carotid artery; another, the *small deep petrosal nerve*, runs in a canal in the processus cochleariformis, passes through the foramen lacerum medium to join the carotid plexus; a third ascends in front of the fenestra ovalis, and, passing forward, joins the great petrosal nerve in the hiatus Fallopii; the fourth leaves the front of the tympanum, under the name of the *small superficial petrosal nerve*, through a canal, where it is joined by a filament from the geniculate ganglion of the facial nerve; then passing beneath the canal for the tensor tympani, it emerges through a foramen on the anterior surface of the pars petrosa, external to the hiatus Fallopii; it proceeds along the anterior surface of pars petrosa, and emerges from the skull between the great wing of the sphenoid and the petrous bones, to join the otic ganglion. Thus the tympanic branch is distributed to the mucous membrane of the tympanum and the Eustachian tube,

[1] This nerve, though commonly called Jacobson's, was fully described by Andersch.

and communicates with the spheno-palatine ganglion through the great petrosal nerve, and with the otic ganglion through the lesser petrosal (Fig. 62).

Pneumogastric Nerve. This nerve leaves the cranium with the nervus accessorius through a common canal in the dura mater, behind that for the glosso-pharyngeal. At its entrance into the canal, it is composed of a number of separate filaments, which are soon collected into a single trunk. In the jugular foramen the nerve presents a ganglionic enlargement, called the *ganglion of the root;* and after the nerve has emerged from the jugular foramen, it presents a second ganglion—the *ganglion of the trunk* of the nerve—where it is joined by the accessory portion of the spinal accessory nerve.[1] The *ganglion of the root* is about ¼ of an inch in length. It is connected by filaments with the sympathetic through the superior cervical ganglion, with the petrous ganglion of the glosso-pharyngeal, with the auricular branch of the facial, and with the spinal accessory by one or two branches. It gives off the *auricular branch*,[2] which is distributed to the pinna of the ear. This branch, shortly after its origin, is joined by a branch from the petrous ganglion of the glosso-pharyngeal, and, passing outward behind the internal jugular vein, it enters a minute foramen in the jugular fossa near the styloid process. It then proceeds through a canal in the bone, crosses the aquæductus Fallopii, where it communicates with the facial nerve, and passes to the outside of the skull through the fissure between the mastoid process and the meatus auditorius externus. It here divides into two branches, one being distributed to the skin of the auricle, and communicating with the great auricular nerve; the other communicating with the posterior auricular branch of the facial over the mastoid process. This ganglion also sends backward a *meningeal branch*, which passes through the jugular foramen to be distributed to the dura mater of the posterior fossa (Fig. 63).

The *ganglion of the trunk* has communications with the hypoglossal nerve, with the loop formed between the first and second

[1] Arnold, *Der Kopftheil des Veget. Nerven Systems.* Heidelberg, 1831.

[2] Arnold's nerve.

cervical nerves, and with the superior cervical ganglion of the sympathetic. It gives off, as a branch of distribution, the pharyngeal and superior laryngeal nerves. This has been previously described (p. 166).

Facial Nerve in the Temporal Bone. The facial nerve is contained within the meatus auditorius internus, together with the auditory nerve. At the bottom of the meatus the two nerves are connected by one or more filaments. The facial nerve then enters the aquæductus Fallopii. This is a tortuous canal in the substance of the temporal bone, and terminates at the stylo-mastoid foramen. The nerve proceeds from the meatus auditorius

FIG. 64.

1. The chorda tympani.
2. The geniculate ganglion of the facial nerve.
3. The great petrosal nerve.
4. The lesser petrosal nerve lying over the tensor tympani.
5. The external petrosal nerve communicating with the sympathetic plexus on the arteria meningea media (6).
7. The Gasserian ganglion.

THE GENICULATE GANGLION OF THE FACIAL NERVE, AND ITS CONNECTIONS WITH THE OTHER NERVES. (From Bidder.)

internus for a short distance outward toward the hiatus Fallopii, where it presents a ganglionic enlargement—the *intumescentia gangliformis*, or *geniculate ganglion*—where it is joined by several nerves; it then makes a sudden bend backward along the inner wall of the tympanum above the fenestra ovalis, and lastly, curving downward along the back of the tympanum, it leaves the skull through the stylo-mastoid foramen.

Its *branches* of *communication* in the temporal bone are:—

Those in the meatus auditorius internus—

a. With the auditory nerve.

Those in the aquæductus Fallopii—

b. With Meckel's ganglion through the large petrosal nerve

c. With the otic ganglion through the small superficial petro nerve.

d. With the sympathetic around the middle meningeal arte through the external superficial petrosal nerve.

Its *branches* of *distribution* are :—

e. The tympanic branch.

f. The chorda tympani.

a. The *communicating branches* with the auditory are by seve filaments, in the meatus auditorius internus.

b. The *large petrosal nerve* joins the carotid branch from t sympathetic to form the Vidian nerve, which joins the sphen palatine ganglion (Fig. 64 [3]).

c. The *small superficial petrosal nerve* passes along the anteri surface of the pars petrosa to join the otic ganglion below t foramen ovale (Fig. 64 [4]).

d. The *external superficial petrosal nerve* passes from the ge iculate ganglion to the sympathetic plexus around the midd meningeal artery (Fig. 64 [5]).

e. The *tympanic branch* passes through a foramen in the ba of the posterior pyramid to supply the stapedius and the laxat tympani [1] (Fig. 63 [4]).

f. The *chorda tympani* is given off from the facial nerve befo its exit from the stylo-mastoid foramen.[2] It ascends a sh distance in a bony canal at the back of the tympanum, and ente that cavity through a small foramen—foramen chordæ posterius below and external to the pyramid, close to the membrana ty pani. It runs forward, ensheathed in mucous membrane, throu the tympanum, between the handle of the malleus and the lo process of the incus, to the anterior part of that cavity. It emerg through a small aperture—foramen chordæ anterius—then tra erses a special bony canal—canal of Huguier—and makes

[1] This is often not muscular, but ligamentous in structure.

[2] In the fœtus, this nerve is given off outside the foramen, but subsequently bone grows downward so as to enclose more of the facial nerve, and with it chorda tympani.

exit close to the fissura Glaseri. It passes downward and forward between the two pterygoid muscles, behind the arteria meningea media, the auriculo-temporal and inferior dental nerves, to join, at an acute angle, the lower border of the gustatory nerve. It then proceeds in part to the submaxillary ganglion, and in part to the lingualis muscle.

External to the stylo-mastoid foramen, the facial nerve communicates with the pneumogastric, the glosso-pharyngeal, the great auricular, the auriculo-temporal nerves, and with the carotid plexus; and on the face, with the numerous branches of the three divisions of the fifth nerve. Its branches of distribution, close to the stylo-mastoid foramen, are the posterior auricular, digastric and stylo-hyoid branches; and on the face, branches to all the facial muscles and the platysma myoides.

Course of the Internal Carotid through Base of Skull.

The cervical portion of the internal carotid has been already described (p. 164). Its subsequent course may be divided into the petrous, cavernous and cerebral portions.

In the *petrous portion*, the artery takes a very tortuous course; at first it ascends for a short distance; it then curves forward and inward; and lastly, it again ascends to reach the side of the body of the sphenoid. It is situated in front of the tympanum, from which it is separated by a thin lamella of bone, which is frequently absorbed in advanced age. It gives off a *tympanic branch* to the tympanum and membrana tympani.

In the *cavernous portion*, the artery again makes a series of curves; at first it ascends forward on the side of the body of the sphenoid, and then curves upward on the inner side of the anterior clinoid process. The artery in this part of its course lies in the inner wall of the cavernous sinus, having the sixth nerve below and to its outer side. From this portion are given off *arteriæ receptaculi* to supply the pituitary body, Gasserian ganglion, and neighboring structures; the *anterior meningeal* to supply the dura mater; and the *ophthalmic artery* already described (p. 74).

In the *cerebral portion*, it pierces the dura mater on the inner side of the anterior clinoid process, and is surrounded by a sheath

of the arachnoid membrane. It gives off the *anterior cerebral*, the *middle cerebral*, the *anterior choroid*, and the *posterior communicating arteries*.

The internal carotid is accompanied in the carotid canal by the cranial branch of the superior cervical ganglion of the sympathetic, described p. 170. Its position on the inner wall of the cavernous sinus, and the nervous plexuses upon it, are described at p. 40.

At this stage of the dissection we may conveniently trace the anterior divisions of the two upper cervical nerves.

Suboccipital Nerve. The anterior division of the *first cervical* or *suboccipital nerve* descends in front of the transverse process of the atlas to form a loop with the ascending branch of the second cervical nerve. It lies beneath the vertebral artery, on the inner side of the rectus capitis lateralis, to which it gives a branch; as also, one to the occipito-atloid joint, one to the rectus capitis anticus minor, and one to the sympathetic around the vertebral artery. From its loop of communication with the second nerve it gives filaments of communication to the superior cervical ganglion, to the hypoglossal and pneumogastric nerves; and muscular branches to the longus colli and rectus capitis anticus major.

Second Cervical Nerve. The anterior division of this nerve emerges between the arches of the atlas and axis, and passes between the vertebral artery and the inter-transversalis muscle, in front of which it subdivides into an ascending branch which joins the first cervical nerve, and into a descending which joins the third cervical nerve.

DISSECTION OF THE NOSE.

Presuming that the dissector is familiar with the bones composing the skeleton of the nose, we shall now describe: 1. The nasal cartilages; 2. The general figure and arrangement of the nasal cavities; 3. The membrane which lines them; and, 4. The distribution of the olfactory nerves.

Cartilages of the Nose. The framework of the external nose is formed by five cartilages; on each side by two lateral

cartilages; and by one in the centre, which completes the septum between the nasal fossæ.

The *lateral cartilages* are termed, respectively, upper and lower, which are covered externally by integument, and are lined internally by mucous membrane. The *upper*, triangular in shape, is connected superiorly to the margin of the nasal and superior maxillary bones; anteriorly, which is its thickest part, to the cartilage of the septum; and inferiorly, to the lower cartilage, by means of a tough fibrous membrane. The *lower* is elongated, and curved upon itself in such a way as to form not only half the apex, but the outer and inner boundaries of the external opening of the nostrils. Superiorly, it is connected by fibrous membrane

FIG. 65.



CARTILAGES OF THE NOSE.

to the upper cartilage; internally, it is in contact with its fellow of the opposite side, forming the upper part of the columna nasi; posteriorly, it is attached by fibrous tissue to the superior maxillary bone; in this tissue are usually found two or three nodules of cartilage, called *cartilagines sesamoideæ;* below, it is firmly connected to dense connective tissue. By their elasticity these several cartilages keep the nostrils continually open, and restore them to their ordinary size whenever they have been expanded by muscular action.

The *cartilage of the septum* is placed perpendicularly in the middle line; it may lean a little, however, to one side or the other, and in some instances it is perforated, so that the two

nasal cavities communicate with each other. The cartilage smooth and flat, and its outline is nearly triangular. The poster border is received into a groove in the perpendicular plate of t ethmoid; the anterior border is much thicker than the rest the septum, and is connected, superiorly, with the nasal bon and on either side with the lateral cartilages. The inferior bord is attached to the vomer and the median ridge at the junction the palatine processes of the superior maxillæ.

The nose receives its blood supply from the lateralis nasi, t artery of the septum, the facial, the nasal branch of the ophth mic, and the infra-orbital arteries. The veins are returned to t facial and ophthalmic veins. The nerves are derived from t nasal branch of the ophthalmic, the infra-orbital, and infra-troc lear nerves. Its muscles are supplied by branches from the faci nerve.

The muscles moving the nasal cartilages have been describe with the dissection of the face (p. 51).

Interior of the Nose. A vertical section should be made through th right nasal cavity, a little on the same side of th middle line,[1] to expose the partly bony and partly cartilaginou partition of the nasal cavities (*septum narium*). Each nasal foss is narrower above than below. The greatest perpendicular dept of each fossa is about the centre; from this point the depth grad ually lessens toward the anterior and the posterior openings o the nose. Laterally, each fossa is very narrow, in consequence of the projection of the spongy bones toward the septum; this narrowness in the transverse direction explains the rapidity with which swelling of the lining membrane from a simple cold obstructs the passage of air.

Boundaries of Nasal Fossæ. The nasal fossæ are bounded by the following bones:—*superiorly*, by the nasal, the nasal spine of the frontal, the cribriform plate of the ethmoid, the body of the sphenoid, and the sphenoidal turbinated bones; *inferiorly*, by the horizontal plates of the superior maxillary and palate bones; *internally*, is the smooth and flat septum formed by the perpendicular plate of the ethmoid, the ridge formed by the two nasal

[1] This has already been done in order to dissect the spheno-palatine ganglion.

bones, the vomer, the septal cartilage, also by the nasal spine of the frontal, the rostrum of the sphenoid, and the crest of the superior maxillary and palate bones; *externally*, by the nasal process and the inner surface of the superior maxillary, the lachrymal, the ethmoid, the palate, the inferior turbinated bones, and the internal pterygoid plate of the sphenoid.

Meatuses of the Nose. The outer wall of each nasal cavity is divided by the turbinated bones into three compartments —*meatuses*—of unequal size; and in these are orifices leading to air-cells—*sinuses*—in the sphenoid, ethmoid, frontal, and superior maxillary bones. Each of these compartments should be separately examined.

a. The *superior meatus* is the smallest of the three, and does not extend beyond the posterior half of the wall of the nose. The posterior ethmoidal and sphenoidal cells open into it. The spheno-palatine foramen is covered by the mucous membrane, and is posterior to the meatus.

b. The *middle meatus* is larger than the superior. At its anterior part a long narrow passage (*infundibulum*), nearly hidden by a fold of membrane, leads upward to the frontal and the anterior ethmoidal cells. About the middle a small opening leads into the antrum of the superior maxilla; this opening in the dry bone is large and irregular, but in the recent state it is reduced nearly to the size of a crow-quill by mucous membrane, so that a very little swelling of the membrane is sufficient to close the orifice entirely.

Notice that the orifices of the frontal and ethmoid cells are so disposed that their secretion will pass easily into the nose. But that is not the case with the maxillary cells, to empty which the head must be inclined on one side. To see all these openings the respective turbinated bones must be raised.

c. The *inferior meatus* extends nearly along the whole length of the outer wall of the nose. By raising the lower turbinated bone, we observe, toward the front of the meatus, the termination of the nasal duct, through which the tears pass down from the lachrymal sac into the nose. This sac and duct can now be conveniently examined.

Lachrymal Sac and Nasal Duct. The lachrymal sac and nasal duct constitute the passage through which the tears are conveyed from the canaliculi into the nose (p. 49). The *lachrymal sac* occupies the groove formed by the lachrymal bone and the nasal process of the superior maxilla. The upper end is round and closed; the lower gradually contracts into the nasal duct, and opens into the inferior meatus. The sac is composed of a strong fibrous and elastic tissue, which adheres very closely to the bone, and is lined by mucous membrane, continuous above with that lining the canaliculi, and below with that of the nasal duct. Its front surface is covered by the tendo oculi and the fascia proceeding from it, and by the tensor tarsi muscle.

The *nasal duct* is from half to three-quarters of an inch in length, and is directed downward, backward, and a little outward. Its termination is rather dilated, and is guarded by a valvular fold of mucous membrane—*valve of Hasner;* consequently, when air is blown into the nasal passages while the nostrils are closed, the lachrymal sac does not become distended. The lachrymal sac and the nasal duct are lined with ciliated epithelium, and the canaliculi with the squamous variety.

Behind the inferior turbinated bone is the opening of the Eustachian tube (p. 255). Into this, as well as into the nasal duct, we ought to practice the introduction of a probe. The chief difficulty is to prevent the probe from slipping into the cul-de-sac between the tube and the back of the pharynx.

Mucous or Schneiderian Membrane.[1] This membrane lines the cavities of the nose and the air cells communicating with it, and adheres very firmly to the periosteum. Its continuity may be traced into the pharynx, into the orbits through the nasal ducts and canaliculi, into the various air sinuses, viz., the frontal, ethmoidal, sphenoidal sinuses, and the antra of Highmore, and into the tympana and mastoid cells through the Eustachian tubes. At the lower border of the turbinated bones it is disposed in thick and loose folds. The membrane varies in thickness and vascularity in different parts of the nasal cavities. Upon the lower half of the septum and the inferior turbinated bones it

[1] Schneider, *De Catarrhis.* Wittenberg, 1660.

is much thicker than elsewhere, owing to a fine plexus of arteries and veins in the submucous tissue. In the sinuses the mucous membrane is thinner, less vascular, and closely adherent to the periosteum.[1]

The great vascularity of the mucous membrane raises the temperature of the inspired air, and pours out a copious secretion which prevents the membrane from becoming too dry.

The mucous membrane of the nasal cavities is not lined throughout by the same kind of epithelium. Near the nostrils the mucous membrane is furnished with papillæ, with a squamous epithelium like the skin, and a few small hairs (*vibrissæ*). In the lower part of the nose—namely, along the respiratory tract and in the sinuses—the epithelium is columnar and ciliated; but in the true olfactory region—that is, upon the superior and middle turbinated bones and the upper half of the septum—the epithelium is columnar, but not ciliated. In this region the mucous membrane is extremely vascular, thick, and studded with branched mucous glands. The columnar epithelial cells taper off at their deep ends into fine processes. Lying between these processes are fusiform cells, with central well-defined nuclei, to which the name of *olfactory cells*[2] has been given; and it is probable that the attenuated processes which pass inward from these cells are in direct connection with the terminal fibrils of the olfactory nerves.

The *arteries* of the nasal cavities are derived from the anterior and posterior ethmoidal branches of the ophthalmic, which supply the roof of the nose, the anterior and posterior ethmoidal cells, and the frontal sinuses; from the nasal artery of the internal maxillary, which supplies the septum, the meatuses, and the turbinated bones; from the posterior dental branch of the internal

[1] In the mucous membrane covering the lower part of the septum, in front, may be observed a small orifice which leads into a narrow blind channel, a few millimetres in length. This canal has numerous glands opening into it, and is the representative of a much larger tubular organ (*organ of Jacobson*) in some quadrupeds, in whom it is surrounded by a curved plate of cartilage (*cartilage of Jacobson*). In man this cartilage is very narrow, and is situated below the rudimentary organ. For further information, see Klein, *Quart. Journ. of Micros. Science*, 1881 and 1882.

[2] Max Schultze, *Med. Centralblatt*, 1864.

maxillary, which supplies the antrum. The external nose is supplied by the nasal branch of the ophthalmic (p. 76), the arteria lateralis nasi, the angular and the artery of the septum.

The *veins* of the nose correspond with the arteries, and, like them, form close plexuses beneath the mucous membrane. They communicate with the veins within the cranium, through the foramina in the cribriform plate of the ethmoid bone; also through the ophthalmic vein and the cavernous sinus. These communications explain the relief frequently afforded by hemorrhage from the nose in cases of cerebral congestion.

The mucous membrane of the nose is supplied with sensory nerves by the fifth pair. Thus, its *roof* is supplied by filaments from the external division of the nasal branch of the ophthalmic, and from the Vidian; its *outer wall*, by filaments from the superior nasal branches of the spheno-palatine ganglion, from the nasal, from the inner branch of the anterior dental, and from the inferior nasal branches of the large palatine nerve; its *septum*, by the septal branch of the nasal nerve, by the nasal branches of the spheno-palatine ganglion, by the naso-palatine, and by the Vidian; its *floor*, by the naso-palatine, and the inferior nasal branches of the large palatine nerve.

Olfactory Nerves. The olfactory nerves, proceeding from each olfactory bulb, in number about twenty on each side, pass through the foramina in the cribriform plate of the ethmoid bone. In its passage each nerve is invested with a coat derived from the dura mater. They are arranged into an inner, a middle and an outer set. The *septal*, which are the largest, traverse the grooves in the upper third of the septum. The *middle* ramify on the roof of the nose. The *outer* pass through grooves, and are divided into an anterior and a posterior group; the anterior being distributed over the superior turbinated bone, the posterior over the os planum of the ethmoid and the middle turbinated bone.

The nerves descend obliquely between the mucous membrane and the periosteum, and break up into filaments, which communicate freely with one another, and form minute plexuses with small elongated intervals. Microscopically, the filaments differ

from the other cerebral nerves, in containing no white substance of Schwann, and in their axis-cylinders being provided with a very distinct nucleated sheath with fewer nuclei and at longer intervals.

DISSECTION OF THE MUSCLES OF THE BACK.

Dissection to Expose the Third Layer of Muscles. Those muscles of the back—namely, the trapezius, latissimus dorsi, levator anguli scapulæ, and rhomboidei—which are concerned in the movements of the upper extremity, will be examined in the dissection of the arm. These must be reflected near to their insertions, together with the cutaneous vessels and nerves. We now proceed to examine the three muscles forming the *third layer* of muscles, named, from their appearance, *serrati postici, superior* and *inferior*, and the *splenius*. The nerves and arteries will be described after the dissection of the suboccipital triangle.

Serratus Posticus, Superior. This muscle is situated beneath the rhomboidei. It is a thin, flat muscle, and *arises* from the lower part of the ligamentum nuchæ,[1] from the spinous processes of the last cervical and two or three upper dorsal vertebræ, by a sheet-like aponeurosis which makes up nearly half the muscle; the fibres run obliquely downward and outward, and are *inserted* by four fleshy slips into the second, third, fourth and fifth ribs, beyond their angles. Its *action* is to raise these ribs, and therefore to assist in inspiration.

Serratus Posticus, Inferior. This muscle is situated in the upper lumbar region, beneath the latissimus dorsi. It *arises*, by means of the *lumbar aponeurosis*, from the spinous processes of the two last dorsal and two upper lumbar vertebræ and their supra-spinous ligament. It ascends obliquely outward, and is *inserted* by four fleshy slips into the four lower ribs, external to their angles. Its *action* is to pull down these ribs, and therefore to assist in expiration. The posterior serrati muscles are supplied,

[1] The ligamentum nuchæ is a rudiment of the great elastic ligament of quadrupeds (termed the *pack-wax*) which supports the weight of the head. It proceeds from the spine of the occiput to the spines of all the cervical vertebræ except the atlas; otherwise it would interfere with the free rotation of the head.

respectively, by the *external* branches of the posterior divisions of the cervical and dorsal nerves.

Vertebral Aponeurosis. The thin aponeurosis which in the posterior aspect of the thoracic region separates the muscles of the upper extremity from those of the back, is called the *vertebral aponeurosis*. *Superiorly*, it is continued beneath the splenius, and is continuous with the deep cervical fascia; *inferiorly* it binds down the muscles contained in the vertebral groove, and is attached to the upper border of the serratus posticus inferior, and the tendon of the latissimus dorsi; *internally*, it is attached

FIG. 66.



TRANSVERSE SECTION THROUGH THE ABDOMEN, TO SHOW THE ATTACHMENT OF THE THREE LAYERS OF THE LUMBAR FASCIA TO THE TRANSVERSE AND SPINOUS PROCESSES OF THE LUMBAR VERTEBRÆ.

to the spinous processes of the dorsal vertebræ, and *externally* to the angles of the ribs.

Lumbar Fascia. This aponeurosis consists of three layers, of which only the posterior layer can now be seen; the other two being demonstrated in the dissection of the abdominal muscles. The posterior or superficial layer is attached to the crest of the ilium, to the spinous processes of all the lower dorsal, lumbar and sacral vertebræ; it forms a sheath for the erector spinæ, and serves for the attachment of the latissimus dorsi, the serratus posticus inferior, and the internal oblique.

FIG. 67.



THE SUPERFICIAL MUSCLES OF THE BACK.

The serratus posticus superior must now be reflected from its origin, and turned outward, to expose the following muscle.

Splenius. This muscle, so called from its resemblance to a strap, *arises* from the spinous processes of the five or six upper dorsal and the last cervical vertebræ, from the supra-spinous ligament, and from the lower half of the ligamentum nuchæ. The fleshy fibres pass upward and outward and divide into two portions, named, according to their respective insertions, *splenius capitis* and *splenius colli*.

a. The *splenius capitis*, the inner of the two portions, is *inserted* into the mastoid process, and into the outer part of the superior curved line of the occipital bone, beneath the sterno-mastoid.

b. The *splenius colli*, the outer of the two portions, is *inserted* by tendinous slips into the posterior tubercles of the transverse processes of the upper three cervical vertebræ. The splenius is supplied by the external branches of the posterior divisions of the cervical nerves.

The *action* of the splenius, taken as a whole, is to draw the head and the upper cervical vertebræ toward its own side; so far, it coöperates with the opposite sterno-mastoid muscle. When the splenii of opposite sides contract, they extend the cervical portion of the spine, and keep the head erect. The permanent contraction of a single splenius may occasion wry-neck. It is necessary to be aware of this, otherwise one might suppose the opposite sterno-mastoid to be affected, considering that the appearance of the distortion is alike in either case.

Dissection to Expose the Fourth Layer. To lay bare the *fourth layer* of muscles, the splenius and serratus posticus inferior are to be detached from their origins. After reflecting the vertebral aponeurosis and the lumbar fascia from its internal attachment, the erector spinæ and its prolongations are exposed.

Erector Spinæ. The mass of muscle which occupies the vertebral groove on each side of the spine is, collectively, called *erector spinæ*, since it counteracts the tendency of the trunk to fall forward. It is pointed at its lower tendinous extremity, where it arises from the sacral region; in the lumbar region it is broad, thick and muscular; in the lower dorsal region

it divides into two portions, which are continued upward, with additional muscles, into the cervical vertebræ and the head. Observe that it is thickest and strongest at that part of the spine where it has the greatest weight to support, namely, in the lumbar region; and that its thickness gradually decreases toward the top of the spine.

It *arises* by thick tendinous fibres from the spinous processes of the two or three lowest dorsal and of all the lumbar vertebræ, from the spines of the sacrum, from the supra-spinous ligament, from the posterior fifth of the inner lip of the crest of the ilium, from the lower and back part of the sacrum, and from the posterior sacro-iliac ligament. From this extensive origin the muscular fibres ascend, at first as a single mass. Near the last rib, this mass divides into two—an outer, called the *ilio-costalis* or *sacro-lumbalis;* an inner, the *longissimus dorsi*. These two portions should be followed up the back; and there is no difficulty in doing so, because the division is indicated by a longitudinal groove, in which we observe the external cutaneous branches of the intercostal vessels and nerves.



ARRANGEMENT OF THE ERECTOR SPINÆ AND ITS PROLONGATIONS INTO THE POSTERIOR THORACIC AND CERVICAL REGIONS.

Ilio-Costalis or Sacro-Lumbalis. Tracing the *ilio-costalis* or *sacro-lumbalis* upward, we find that it terminates in a series of tendons which are *inserted* into the angles of the six lower ribs.

Musculus Accessorius. By turning outward the ilio-costalis, we o serve that it is continued upward under the nar of *musculus accessorius ad ilio-costalem.* This *arises* by a seri of tendons from the angles of the six lower ribs, internal to t preceding, and is *inserted* by muscular slips into the angles the six upper ribs.

Cervicalis Ascendens. This is the cervical continuation of the mu culus accessorius. It *arises* by tendinous sli from the angles of the four or five upper ribs, internal to t musculus accessorius, and is *inserted* into the posterior tubercl of the transverse processes of the fourth, fifth and sixth cervic vertebræ.

Longissimus Dorsi. The longissimus dorsi (the inner portion of t erector spinæ) terminates in tendons which a *inserted*, internally, into the tubercles[1] at the root of the transve processes of the lumbar vertebræ, into the tubercles of the articu processes of the same vertebræ, into the middle layer of the fas lumborum, also into the transverse processes of all the dor vertebræ, and, externally, into the greater number of the r (varying from eight to eleven) between their tubercles and angl

Transversalis Colli. This is the cervical continuation of the lo issimus dorsi. It *arises* by long tendinous sl from the tips of the transverse processes of the five or six up dorsal vertebræ, and is *inserted* into the posterior tubercles of t transverse processes of the four or five lower cervical verteb except the last.

Trachelo-mastoid. This muscle, situated on the inner side of t preceding, and external to the complexus, is internal continuation of the longissimus dorsi to the cranium. *arises* from the transverse processes of the three or four up dorsal, and the articular processes of the three or four lo cervical vertebræ, and is *inserted* by a flat tendon into the ba part of the mastoid process beneath the splenius.[2]

[1] Called *anapophyses* by Professor Owen.

[2] Those who are familiar with the transcendental nomenclature of the verteb skeleton will understand from the following quotation the plan upon which muscles of the back are arranged:—

"The muscles of the back are either longitudinal or oblique; that is, they ei pass vertically downward from spinous process to spinous process, from diapoph

Spinalis Dorsi.

This is a long, narrow muscle, situated close to the spines of the dorsal vertebræ, and apparently the inner part of the longissimus dorsi; it is by some considered the innermost column of the erector spinæ. It *arises* by tendinous slips from the spinous processes of the two lower dorsal and two upper lumbar vertebræ, and is *inserted* by little tendons into the spinous processes of the six or eight upper dorsal vertebræ. Beneath it is the semi-spinalis dorsi, which is closely connected with the spinalis dorsi.

Spinalis Colli.

This small but not constant muscle corresponds in the cervical region to the spinalis dorsi in the dorsal region. It *arises* by tendinous slips from the spinous processes of the two or three lower cervical vertebræ (sometimes also from the two upper dorsal), and is *inserted* into the spine of the axis, and occasionally into the spinous processes of the third and fourth cervical.

to diapophysis, from rib to rib (pleurapophysis), etc., or they extend obliquely from diapophysis to spine, or from diapophysis to pleurapophysis, etc.

"The erector spinæ is composed of two planes of longitudinal fibres aggregated together, below, to form one mass at their point of origin, from the spines and posterior surface of the sacrum, from the sacro-iliac ligament, and from the posterior third of the iliac crest. It divides into two portions, the sacro-lumbalis and the longissimus dorsi.

"The former, arising from the iliac crest, or from the pleurapophysis (rib) of the first sacral vertebra, is inserted by short flat tendons into (1) the apices of the stunted lumbar ribs, close to the tendinous origins of the transversalis abdominis; (2) the angles of the eight or nine inferior dorsal ribs; (3) it is inserted, through the medium of the musculus accessorius, into the angles of the remaining superior ribs, and into the long and occasionally distinct pleurapophysial element of the seventh cervical vertebra; and (4) through the medium of the cervicalis ascendens, into the pleurapophysial elements of the third, fourth, fifth and sixth cervical vertebræ. In other words, the muscular fibres extend from rib to rib, from the sacrum to the third cervical vertebra.

"The longissimus dorsi, situated nearer the spine than the sacro-lumbalis, is inserted (1) into the metapophysial spine of the lumbar diapophyses; (2) into the diapophyses of all the dorsal vertebræ, near the origin of the levatores costarum; (3) through the medium of the transversalis colli into the diapophyses of the second, third, fourth, fifth and sixth cervical vertebræ; and (4) through the medium of the trachelo-mastoid into the mastoid process, or the only element of a transverse process possessed by the parietal vertebra. In other words, its fibres extend from diapophysis to diapophysis; from the sacrum, upward, to the parietal vertebra."—*Homologies of the Human Skeleton*, by H. Coote, p. 75.

The muscles of the spine hitherto examined are all longitudina in their direction. We now come to a series which run obliquel from the transverse to the spinous processes of the vertebræ And first, of the complexus.

Complexus. This powerful muscle *arises* by tendinous slip from the transverse processes of the three or fou upper dorsal and the last cervical vertebræ, also from the articu lar processes of four or five cervical vertebræ and their capsula ligaments. It is *inserted* between the two curved lines of the occiput, near the vertical crest. In the centre of the muscle there is generally a transverse tendinous intersection. The muscle is perforated by the posterior branches of the second (the great occipital), third and fourth cervical nerves. It is chiefly supplied by the great occipital nerve. Its *action* is to maintain the head erect.

Biventer Cervicis. Is placed in the inner side of the preceding muscle and frequently forms part of it. It has an intermediate tendon, and *arises* from the transverse processes of two or three upper dorsal vertebræ, and ascends between the ligamentum nuchæ and the complexus, to be *inserted* into the innermost depression between two curved lines of the occipital bone.

Cut transversely through the middle of the complexus, and reflect it to see the arteria cervicalis profunda (p. 141), and the posterior branches of the cervical nerves.

Dissection to Expose the Fifth Layer. Remove the complexus, and then turn aside the erector spinæ and its prolongations, when the *fifth layer* of muscles will be seen occupying the interval between the spinous and transverse processes.

Transverso-spinalis. This is the mass of muscle which lies in the vertebral groove after the reflection of the complexus and the erector spinæ. It consists of a series of fibres which extend from the transverse and articular processes to the spinous processes of the dorsal and cervical vertebræ, and is, for convenience, divided into the *semispinalis dorsi* and *semispinalis colli*.

a. The *semispinalis dorsi arises* by long, thin, tendinous slip

from the transverse processes of the dorsal vertebræ, from the sixth to the tenth, and is *inserted* into the spinous processes of the four upper dorsal and the two or three lower cervical vertebræ. Its nerves are derived from the internal posterior branches of the dorsal nerves.

b. The *semispinalis colli* lies beneath the complexus, and *arises* from the transverse processes of the five or six upper dorsal vertebræ, and the articular processes of the four lower cervical, and is *inserted* into the spinous processes of the axis and the three or four succeeding vertebræ, that into the axis being the most fleshy fasciculus. It is supplied by the internal posterior branches of the cervical nerves.

Now reflect part of the semispinalis dorsi in order to expose the multifidus spinæ.

Multifidus Spinæ. This may be considered a part of the preceding muscle, since its fixed points and the direction of its fibres are the same. It consists of a series of little muscles which extend between the spinous and transverse processes of the vertebræ, from the sacrum to the second cervical vertebra. Those in the lumbar region are the largest. In the sacral region the fibres *arise* from the back of the sacrum as low down as the fourth foramen, from the deep surface of the aponeurosis of the erector spinæ, from the inner part of the posterior superior iliac spine, and from the posterior sacro-iliac ligament; in the lumbar region, from the mammillary processes on the superior articular processes; in the dorsal region, from the transverse processes, and in the cervical region from the articular processes of the four lower cervical vertebræ. They all ascend obliquely, and each fasciculus is *inserted* into the lamina and spinous process of the vertebra above, except the atlas. It should be observed that their fibres are not of uniform length; some extend only from vertebra to vertebra, while others extend between one, two, or even three vertebræ. It is supplied by the internal posterior branches of the sacral, lumbar, dorsal and cervical nerves.

Rotatores Spinæ. Beneath the multifidus spinæ, in the dorsal region of the spine only, are eleven flat and somewhat square muscles, called *rotatores spinæ*. They *arise* from the

upper and back part of the transverse processes, and are *inserte* into the lower border of the laminæ of the vertebra above. Thes muscles form but a part of the multifidus spinæ, and are supplie by the internal posterior branches of the dorsal nerves.

The action of the preceding muscles is, not only to assist i maintaining the trunk erect, but to incline and rotate the spin to one or the other side. They are all supplied by the posteric branches of the spinal nerves.

Levatores Costarum. These small muscles, twelve in number, o each side, *arise* from the apices of the transvers processes of the seventh cervical and the eleven upper dors vertebræ, and are *inserted* into the rib below. The direction their fibres corresponds with that of the outer layer of the inte costal muscles, and they are supplied by the internal posteric branches of the dorsal nerves. They are muscles of inspiratio

Supra-spinales. These are formed by a series of small muscul slips lying over the spinous processes of the ce vical vertebræ. Their nerves are derived from the internal post rior branches of the cervical nerves.

Inter-spinales. These muscles extend between the spino processes of the contiguous vertebræ. They a arranged in pairs, and only exist in those parts of the vertebr column which are most movable. In the cervical region the are the most distinct, and pass between the spinous processes the six lower cervical vertebræ. In the dorsal, they are foun between the spinous processes of the first and second, an between those of the eleventh and twelfth dorsal vertebræ. The are also found more or less distinctly between the spinous pr cesses of the lumbar vertebræ. They are supplied by the intern posterior branches of the spinal nerves.

Inter-transversales. These muscles extend between the transver processes of the vertebræ. In the *cervical regi* they are seven in number, and are most marked, being arrange in pairs, and extend between the anterior and posterior tubercl of contiguous vertebræ. The anterior branch of the correspondi cervical nerve separates the two fasciculi. In the *dorsal regi* these muscles in the upper part are represented by small roun

tendons, but in the three lower dorsal vertebræ they again become muscular in structure. In the *lumbar region* the muscular fasciculi are four in number, and are also arranged in pairs between the transverse processes. Their nerve supply is derived from the internal posterior branches of the cervical, dorsal and lumbar nerves.

We have next to examine the muscles concerned in the movements of the head upon the first and second cervical vertebræ (Fig. 68).

FIG. 68.



DRAWING, FROM NATURE, OF THE SUBOCCIPITAL TRIANGLE.

1 and 7. Complexus. 2. Rectus cap. posticus minor. 3. Rectus cap. posticus major. 4. Obliquus inferior. 5. Sterno-mastoid. 6. Semispinalis colli. 8. Obliquus superior. 10. Splenius. 11 Trachelo-mastoid. 12. Great occipital nerve. 13. Occipital artery giving off its descending branch—the *princeps cervicis*. 14. Suboccipital nerve. 15. Third cervical nerve (posterior branch).

Rectus Capitis Posticus Major. This is a largely developed interspinal muscle. It *arises* by a small tendon from the well-marked spinous process of the second cervical vertebra, and, expanding considerably, is *inserted* into the inferior curved ridge of the occipital bone, and into the surface of bone below it. These recti muscles, as they ascend, one on each side, to their insertions,

diverge and leave an interval between them, in which are f
the recti capitis postici minores.

Rectus Capitis Posticus Minor. This is an interspinal muscle, but smaller the preceding. *Arising* from the posterio bercle of the first vertebra, it expands as it ascends, and is *in* into the occipital bone between the inferior curved ridge an foramen magnum. The *action* of the two preceding musc to raise the head. They are supplied with nerves from the terior branch of the suboccipital.

Obliquus Inferior. This *arises* from the spinous process o second cervical vertebra, and is *inserted* int transverse process of the first. Its *action* is to rotate th upon the second vertebra; in other words, to turn the head r to the same side. It is supplied with a nerve by the great c ital (posterior division of the second cervical), which curv under its lower border.

Obliquus Superior. This muscle *arises* from the transverse pr of the atlas, and, ascending obliquely inwa *inserted* in the interval between the curved ridges of the occ bone. Its *action* is to draw the occiput toward the spine.

Suboccipital Triangle. Observe that the obliqui (superior and inf and the rectus capitis posticus major form is called the *suboccipital triangle*. The outer side is formed b obliquus superior; the inner, by the rectus capitis posticus m the lower, by the obliquus inferior. Within this triangle m seen the arch of the atlas, the vertebral artery lying in a groo its upper surface, and the posterior occipito-atloid ligament. tween the artery and the bone appears the posterior division suboccipital nerve, which here sends branches to the recti p the obliqui, and the complexus; that is to say, it supplie muscles which form the triangle, and the complexus that cov

Rectus Capitis Lateralis. This small muscle extends between the verse process of the first vertebra and the nentia jugularis of the occiput; but, since this eminence transverse process of the occipital vertebra, the muscle shou considered as an intertransverse one. Its nerve comes fro anterior division of the suboccipital.

Nerves of the Back. The posterior branches of the spinal nerves supply the muscles and skin of the back. They pass backward between the transverse processes of the vertebræ, and divide into *external* and *internal* branches. The general plan upon which these nerves are arranged is the same throughout the whole length of the spine; but, since there are certain peculiarities deserving of notice in particular situations, we must examine each region separately.

Cervical Region. The posterior division of the *first cervical nerve* (the suboccipital) passes between the arch of the atlas and the vertebral artery; it then enters the suboccipital triangle, and divides into branches which supply the muscles; one, which passes downward to supply the inferior oblique, and also sends downward a branch to communicate with the second cervical nerve; another passes upward to supply the recti capitis major and minor; another supplies the obliquus superior; another enters the complexus; and lastly, a cutaneous branch is sometimes given off, which accompanies the occipital artery, and is distributed to the back of the scalp.

The posterior branch (the great occipital) of the *second cervical nerve* is the largest in the series, and emerges between the arches of the atlas and axis. It turns upward beneath the inferior oblique muscle, passes through the complexus, and runs with the occipital artery to the back of the scalp.

The posterior divisions of the six lower cervical nerves divide into *external* and *internal* branches. The *external* are small, and terminate in the splenius, and the continuation of the erector spinæ—viz.: the trachelo-mastoid, the transversalis colli, and the cervicalis ascendens. The *internal*, by far the larger, proceed toward the spinous processes of the vertebræ; those of the third, fourth and fifth lie between the complexus and the semispinalis,[1] and after supplying the muscles terminate in the skin over the trapezius; those of the sixth, seventh and eighth lie between the semispinalis and the multifidus spinæ, to which

[1] The posterior branches of the second, third and fourth nerves are generally connected, beneath the complexus, by branches in the form of loops. This constitutes the *posterior cervical plexus* of some anatomists.

they are distributed, and do not, as a rule, give off any cutaneous branches.

Dorsal Region. The posterior divisions of the spinal nerves in this region come out between the transverse processes and the tendons attached to them. They soon divide into *external* and *internal* branches. The *external* pass obliquely over the levatores costarum, between the ilio-costalis and the longissimus dorsi; and successively increase in size from above downward. The upper six terminate in the erector spinæ and the levatores costarum; the lower six, after supplying these muscles, pass through the latissimus dorsi, and become the cutaneous nerves of the back. The *internal* successively decrease in size from above downward. They run toward the spine between the semispinalis dorsi and the multifidus spinæ. The upper six, after giving branches to the muscles, perforate the trapezius and become cutaneous nerves. The lower ones terminate in the muscles of the vertebral groove.

Lumbar Region. The general arrangement of the nerves in this region resembles that of the dorsal. Their *external* branches, after supplying the erector spinæ, become cutaneous and terminate in the skin over the buttock. The *internal* branches supply the multifidus spinæ.

Sacral Region. The posterior divisions of the spinal nerves in this region are small. With the exception of the last, they come out of the spinal canal through the foramina in the back of the sacrum. The upper two or three divide into *external* and *internal* branches. The *internal* terminate in the multifidus spinæ; the *external* become cutaneous and supply the skin of the gluteal region. The last two sacral nerves proceed, without dividing, to the integument.

The *coccygeal* nerve is exceedingly small, and, after joining a small branch from the last sacral, terminates in the skin over the coccyx.

Arteries of the Back. The arteries which supply the back are:—1. Small branches from the occipital; 2. Small branches from the vertebral; 3. The deep cervical; 4. The posterior branches of the intercostal and lumbar arteries.

FIG. 69.

DIAGRAM OF THE CUTANEOUS NERVES OF THE BACK. (Quain.)

a, *a*, Small occipital nerve from the cervical plexus; 1, external muscular branches of the first cervical nerve and union by a loop with the second; 2, the rectus capitis posticus major, with the great occipital nerve passing round the short muscles and piercing the complexus; the external branch is seen to the outside; 2′, the great occipital; 3, external branch of the posterior primary division of the third nerve; 3′, its internal branch, or third occipital nerve; 4′, 5′, 6′, 7′, 8′, internal branches of the several corresponding nerves on the left side; the external branches of these nerves proceeding to muscles are displayed on the right side: *d* 1 to *d* 6, and thence to *d* 12, external muscular branches of the posterior primary divisions of the twelve dorsal nerves on the right side; *d* 1′, to *d* 6′, the internal cutaneous branches of the six upper dorsal nerves on the left side; *d* 7′ to *d* 12′, cutaneous branches of the six lower dorsal nerves from the external branches; *l*, *l*, external branches of the posterior primary branches of several lumbar nerves on the right side piercing the muscles, the lower descending over the gluteal region; *l*′, *l*′, the same more superficially on the left side; *s*, *s*, on the right side, the issue and union by loops of the posterior primary divisions of four sacral nerves; *s*′, *s*′, some of these distributed to the skin on the left side.

The *occipital artery* furnishes several small branches to the muscles at the back of the neck; one, larger than the rest, the *arteria princeps cervicis*, descends beneath the complexus, and generally inosculates with the deep cervical artery and with small branches from the vertebral.

The *vertebral artery* runs along the groove in the arch of the atlas, and before perforating the posterior occipito-atloid ligament to enter the skull, distributes small branches to the adjacent muscles.

The *deep cervical artery* is the posterior branch of the first intercostal artery (from the subclavian). It passes backward between the transverse process of the last cervical vertebra and the first rib; it then ascends between the complexus and the semispinalis colli, and anostomoses with the princeps cervicis.

The *posterior* branches of the intercostal and lumbar arteries accompany the corresponding nerves, and are in all respects similar to them in distribution. Each sends a small branch into the spinal canal (*intraspinal*), and small branches to the vertebræ.

The *veins* correspond to the arteries.

Prevertebral Muscles. We have, lastly, to examine three muscles situated in front of the spine; namely, the longus colli, the rectus capitis anticus major, and the rectus capitis anticus minor. In order to have a complete view of the two latter, a special dissection should be made, before the head is removed from the first vertebra.

Longus Colli. This muscle is situated in front of the spine and extends from the third dorsal vertebra to the atlas. For convenience of description it is divided into three sets of fibres, of which one extends *longitudinally* from the body of one vertebra to that of another; the two others extend *obliquely* between the transverse processes and the bodies of the vertebræ.

The *longitudinal* portion of the muscle *arises* from the bodies of the three upper dorsal and the three lower cervical vertebræ and is *inserted* into the bodies of the second, third and fourth cervical vertebræ.

The *superior oblique* portion, *arising* from the anterior tubercles of the transverse processes of the third, fourth and fifth cervical

vertebræ, ascends inward, and is *inserted* into the front part or body of the atlas. The *inferior oblique* portion proceeds from the bodies of the three upper dorsal vertebræ, and passing upward and outward, is *inserted* into the transverse processes of the fifth and sixth cervical vertebræ. The *action* of this muscle, taken as a whole, must be to bend the cervical region of the spine. Its nerves come from the lower cervical nerves.

FIG. 70.



DIAGRAM OF THE PREVERTEBRAL MUSCLES.

1-7. The bodies of the cervical vertebræ; below are the bodies of the three upper dorsal vertebræ.
a. Rectus capitis lateralis.
b. Rectus capitis anticus major.
c. Rectus capitis anticus minor.
d. Intertransverse muscle.
e. Scalenus anticus.
f. Scalenus medius.
g. Longus colli.
h. Scalenus posticus.

Rectus Capitis Anticus Major. This muscle *arises* by tendinous slips from the anterior tubercles of the transverse processes of the third, fourth, fifth and sixth cervical vertebræ, and, ascending

obliquely inward, is *inserted* into the basilar process of the occipital bone, in front of the foramen magnum.

Rectus Capitis Anticus Minor. This muscle *arises* from the front of the root of the transverse process of the atlas, and is *inserted* into the basilar process of the occipital bone, nearer to the foramen magnum than the preceding muscle. The *action* of the rectus muscles is to bend the head forward. They are supplied with nerves from the anterior division of the suboccipital, and from the deep cervical plexus.

LIGAMENTS OF THE SPINE.

The vertebræ are connected by their intervertebral fibro-cartilages, by ligaments in front of and behind their bodies, and by ligaments which extend between their arches and their spines. Their articular processes have capsular ligaments and synovial membranes.

Anterior Common Ligament. This is a strong, broad band of longitudinal fibres which extends along the front of the bodies of the vertebræ, from the axis to the sacrum. The ligament is broader below than above, thickest in the dorsal region, and its fibres are more firmly adherent to the intervertebral cartilages and to the borders of the vertebræ, than to the middle of the bones. The fibres are not all of equal length; the more superficial extend from one vertebra to the fourth or fifth below it; those a little deeper pass from one vertebra to the second or third below it; while the deepest of all proceed from vertebra to vertebra. Above, it is attached to the axis by a pointed process where it is connected with the longus colli, and it is thicker over the bodies of the vertebra than over the intervertebral cartilages thus filling up the concavities of the bodies and rendering the surface more smooth and even.

Posterior Common Ligament. This extends longitudinally, in a similar manner to the anterior common ligament, within the spinal canal, along the posterior surface of the bodies of the vertebræ, from the axis to the sacrum. It is broader above than below, and, like the anterior ligament, is thickest in the dorsal region, and is more intimately connected with the intervertebral

fibro-cartilages than with the bodies of the vertebræ. It sends up a prolongation to the anterior border of the foramen magnum continuous with the apparatus ligamentosus.

Interspinous Ligaments. These bands of ligamentous fibres fill up the intervals between the spines of the dorsal and lumbar vertebræ. They are the most marked in the lumbar region. Those fibres which connect the apices of the spines, being stronger than the rest, are described as a separate ligament under the name of *supraspinous*.

Supraspinous Ligament. It extends from the spinous process of the seventh cervical to the spine of the sacrum; and is strongest in the lumbar region. Their use is to limit the flexion of the spine.

Ligaments Between the Arches of the Vertebræ. These are called, on account of their color, *ligamenta subflava*. To obtain a good view of them, the arches of the vertebræ should be removed with a saw, and the ligaments should be seen from within, since viewed from without they are to a large extent hidden by the overlapping laminæ. They pass between the laminæ of the contiguous vertebræ, from the axis to the sacrum; none existing between the occiput and the atlas, or between the atlas and the axis. Each ligament consists of two halves which are attached to the corresponding half laminæ above and below on each side. They are composed of yellow elastic tissue, the fibres being arranged vertically, and their strength increases with the size of the vertebræ. This elasticity answers a double purpose; it not only permits the spine to bend forward, but materially assists in restoring it to its *curve of rest*. They economize muscular force, like the ligamentum nuchæ in animals.

Intervertebral Fibro-Cartilage. This substance, placed between the bodies of the vertebræ, is by far the strongest bond of connection between them, and fulfills most important purposes in the mechanism of the spine. Its peculiar structure is adapted to break shocks, and to render the spine flexible and resilient. To see the structure of an intervertebral fibro-cartilage, a horizontal section must be made through it. It is firm and resisting near the circumference, but soft and pulpy toward the centre. The circumferential portion is composed of concentric

layers of fibro-cartilage, placed vertically. These layers are attached by their edges to the vertebræ; they gradually decrease in number from the circumference toward the centre; and the interstices between them are filled by soft pulpy tissue. The central portion is composed almost entirely of this pulpy tissue and it bulges when no longer under pressure. Thus the bodies of the vertebræ, in their motions upon each other, revolve upon an elastic cushion tightly girt all round by bands of fibrous tissue. These motions are regulated by the articular processes.

Dissect an intervertebral substance, layer after layer, in front and you will find that the circumferential fibres extend *obliquely* between the vertebræ, crossing each other like the branches of the letter X (Fig. 72).

The thickness of the intervertebral cartilages is not the same in front and behind. It is this difference in their thickness, more than that in the bodies of the vertebræ, which produces the several curves of the spine. In the lumbar and cervical regions they are thicker in front; in the dorsal region, behind.

The structure of the intervertebral cartilages explains the well-known fact that a man becomes shorter after standing for some hours; and that he regains his usual height after rest. The difference between the morning and evening stature amounts to more than half an inch.

It also explains the fact that a permanent lateral curvature of the spine may be produced (especially in the young) by the habitual practice of leaning to this or that side. Experience proves that the cause of lateral curvature depends more frequently upon some alteration in the structure of the fibro-cartilages than upon the bones. From an examination of the bodies of one hundred and thirty-four individuals with crooked spines, it was concluded that, in two-thirds, the bones were perfectly healthy; that the most frequent cause of curvature resided in the intervertebral substances, these being, on the concave side of the curve, almost absorbed, and, on the convex side, preternaturally developed. As might be expected in these cases, the muscles on the convex side become lengthened, and degenerate in structure.[1]

[1] On this subject see Hildebrandt's *Anatomie*, B. ii, s. 155.

Ligamentum Nuchæ. This ligament is a thin, fibrous septum, intermingled with elastic tissue, situated in the middle line, and extends from the spinous processes of the cervical vertebræ to the external occipital protuberance. It forms an intermuscular septum down the back of the neck, and may be regarded as the continuation upward of the supraspinous ligament.

Capsular Ligaments. Each joint between the articular processes has a synovial membrane surrounded by loose ligamentous fibres, forming a capsular ligament, which is longest in the cervical vertebræ, thus allowing free movement in this region. The surfaces of the bones are crusted with cartilage.

Intertransverse Ligaments. These are thin bands of fibres which pass between the transverse processes of the vertebræ. They are rudimentary in the cervical region, and are sometimes absent.

Movements of the Spine. Though but little movement is permitted between any two vertebræ (the atlas and axis excepted), yet the collective motion between them all is considerable. The spine can be bent forward, backward, or on either side; it also admits of slight rotation. In consequence of the elasticity of the intervertebral cartilages and the ligamenta subflava, it returns spontaneously to its natural curve of rest, like an elastic bow. Its mobility is greatest in the cervical region, on account of the thickness of the fibro-cartilages, the small size of the vertebræ, the oblique direction of their articulations, and, above all, the horizontal position and the shortness of their spines. In the dorsal region there is very little mobility, on account of the vertical direction of the articular processes, and the manner in which the arches and the spines overlap each other. In the lumbar region, the spine again becomes more movable, on account of the thickness of the intervertebral cartilages, and the horizontal direction of the spinous processes.

Ligaments between the Occipital Bone and the Atlas. The occiput is connected to the atlas by the following ligaments: viz.: two anterior occipito-atloid, a posterior occipito-atloid, two lateral occipito-atloid, and two capsular ligaments.

The *two anterior ligaments* are composed of a superficial and a

deep portion; the *superficial* part is a strong, rounded cord which passes from the basilar process above, to the tubercle on the anterior arch of the atlas below; the *deep* portion is membranous, and passes from the anterior margin of the foramen magnum to the front arch of the atlas.

The *posterior ligament* extends in a similar manner from the posterior border of the foramen magnum to the posterior arch of the atlas. It is thin, and superiorly becomes blended with the dura mater, and is pierced by the vertebral artery and the suboccipital nerve.

The *two lateral ligaments* pass from the jugular eminences of the occiput, downward and outward to the transverse processes of the atlas.

FIG. 71.



DIAGRAM OF THE ODONTOID AND TRANSVERSE LIGAMENTS.

The *capsular ligaments* extend from the margin of the condyles of the occipital bone to the upper articular borders of the atlas.

The movements which take place between the occipital bone and the atlas are flexion and extension, as in nodding forward and backward; and lateral movement, as in inclining the head sideways

Ligaments between the Occipital Bone and the Axis.

These are the most important; and to see them, the spinal canal must be exposed by removing the posterior arches of the upper cervical vertebræ, and the posterior common ligament, which

is here very thick and strong. It ascends from the posterior surface of the axis, then passes over the odontoid and transverse ligaments, and is attached to the basilar process of the occipital bone. It is called the *occipito-axial ligament*, or the *apparatus ligamentosus colli.*

Odontoid or Check Ligaments. The *odontoid* or *check ligaments* (Fig. 71) are two very strong ligaments, which proceed from the sides of the odontoid process to the tubercles on the inner sides of the condyles of the occiput. Their use is to limit the rotation of the head. A *third* or *middle odontoid ligament* passes from the apex of the odontoid process to the margin of the foramen magnum. It is sometimes called the *ligamentum suspensorium.*

Articulation between the Atlas and the Axis. This joint forms a lateral ginglymus or diarthrosis rotatoria, and is maintained by the following ligaments: two anterior atlo-axial, a posterior atlo-axial, two capsular, and a transverse.

The *two anterior ligaments* consist of a superficial and a deep portion; the *superficial* is a rounded ligament passing from the tubercle of the atlas to the base of the odontoid process; the *deep* passes as a membranous layer from the anterior arch of the atlas to the body of the axis.

The *posterior ligament* extends from the posterior arch of the atlas to the upper border of the lamina of the axis.

The *capsular ligaments* are thin, loose, ligamentous sacs connecting the borders of the articular surfaces.

The *transverse ligament* (Fig. 71) passes transversely behind the odontoid process, and is attached to the tubercles on the inner sides of the articular processes of the atlas. From the centre of this ligament a few fibres pass upward, to be attached to the basilar process, and some downward to the body of the axis, giving it a cruciform appearance. Thus it forms with the atlas a ring, into which the odontoid process is received. If this transverse ligament be divided, we observe that the odontoid process is covered with cartilage in front and behind, and is provided with two synovial membranes.

The ribs articulate by their heads with the bodies of the dorsal vertebræ; by their necks and tubercles with the transverse pro-

cesses of the vertebræ, and by their cartilages with the ste
in front.

Articulations between the Heads of the Ribs with the bodies of the Vertebræ.

The head of each rib presents two art
surfaces, corresponding to the bodies o
vertebræ. There are two distinct articul
each provided with a separate synovial
brane. The ligaments are—

1. An *anterior costo-central* or *stellate*, which connects the
of the head of the rib with the sides of the bodies of two
bræ and the intervening fibro-cartilage (Fig. 72). It is com
of three fasciculi of fibres which radiate from the rib,

FIG. 72.

1, 1, 1. Superior costo transverse ligaments.

2, 2, 2. Anterior co
tral or stellate liga

COSTO-VERTEBRAL LIGAMENTS.

which passes upward to be attached to the body of the v
above; the lower one passes to the body of the vertebra
while the intermediate one passes horizontally forward
intervertebral disc.

In the three lower ribs the fasciculi are not separately
guishable, although the fibres pass upward to the vertebr
downward to the vertebra with which the rib articulates.

Some anatomists describe a capsular ligament surroundi
articulation; the fibres are very thin, and form part of the
central ligament.

2. An *interarticular ligament*, which passes across the joi
the ridge on the head of the rib to the intervertebral ca
It divides the articulation into two joints which do not c

nicate with each other. It is absent in the three lower articulations.

Articulations of the Neck and Tubercle of the Ribs with the Transverse Processes.

The ligaments connecting these bones are the capsular, the anterior, middle and posterior costo-transverse.

The *capsular ligament* surrounds the articular surfaces of the tubercle of the rib and the transverse process of its corresponding vertebra, and has a synovial membrane. It is absent in the eleventh and twelfth ribs.

The *anterior* or *superior costo-transverse ligament* ascends from

FIG. 73.

DIAGRAM SHOWING THE LIGAMENTS CONNECTING THE RIB WITH THE VERTEBRA.

1. The anterior costo-central ligament.
2. The interosseous, or middle costo-transverse ligament.
3. The posterior costo-transverse ligament.
4. The synovial membrane between the rib and the body of the vertebra.

the upper border of the neck of the rib to the lower border of the transverse process above it. It is continuous externally with the aponeurosis covering the external intercostal muscle. The first and twelfth ribs have no anterior costo-transverse ligament (Fig. 72).

The *middle costo-transverse ligament* is an interosseous one, and connects the adjacent surfaces of the neck of the rib and the transverse process. It is badly developed in the eleventh and twelfth ribs (Fig. 73).

The *posterior costo-transverse ligament* passes from the apex of

the transverse process to the summit of the tubercle of
It is wanting in the eleventh and twelfth ribs (Fig. 73).

Connection between the Cartilages of the Ribs and Sternum.

The anterior extremities of the ribs a
cave, and receive the cartilages of the rib
junction is maintained by the periosteum
cartilages of all the true ribs are receiv
slight concavities on the side of the sternu
are secured by *anterior*, *posterior*, *upper* and *lower lig*
There is a synovial membrane between the cartilage of e
and the sternum, except that of the first, and usually
articulation the synovial membrane is separated into tw
interarticular ligament.

The *costal cartilages from the sixth to the tenth* are conne
ligamentous fibres. There are intercostal synovial membr
front, between the adjacent borders of the sixth, seventh,
and ninth costal cartilages.

Movements of the Ribs.

The movements permitted between th
of the ribs and the bodies of the verteb
those of elevation and depression, and those of rotation forw
backward; the centre of these movements being at the inte
lar ligament. Between the tubercles and the transverse pr
there is the movement of an arthrodial nature; and betw
costal cartilages and the sternum, that of elevation and dep

The movement of the first rib is very slight; that of the
is freer; and mobility of the ribs gradually increases from
downward.

Articulation of the Lower Jaw.

The condyle of the lower jaw articulat
the glenoid cavity of the temporal bone, an
an arthrodial joint. The joint is provided with an intera
fibro-cartilage, with external and internal lateral and c
ligaments, and two synovial membranes (Fig. 74).

The *external lateral ligament* extends from the zygoma
tubercle; its fibres pass downward and backward to th
surface and posterior border of the neck of the jaw.

The *internal lateral ligament*—a long, thin, flat band—e
from the spinous process of the sphenoid bone to the inner
of the dental foramen.

The *capsular ligament* consists of a few scattered fibres attached above to the margin of the glenoid cavity, below to the neck of the jaw.

The *interarticular fibro-cartilage* is a thin plate of an oval form, and thicker at the margin than at the centre. It is placed horizontally, and its upper surface is concavo-convex from before backward; its lower surface is concave. It is connected on the outer side to the external lateral ligament, and on the inner side

FIG. 74.



TRANSVERSE SECTION, TO SHOW THE LIGAMENTS AND THE FIBRO-CARTILAGE OF THE JOINT OF THE LOWER JAW. THE DOTTED LINES REPRESENT THE TWO SYNOVIAL MEMBRANES.

some of the fibres of the external pterygoid muscle are inserted into it.

There are two *synovial membranes*—an upper and a lower—for the joint. The larger and looser of the two is situated between the glenoid cavity and the fibro-cartilage. The lower is interposed between the fibro-cartilage and the condyle of the jaw. They sometimes communicate through a small aperture in the centre of the fibro-cartilage.

The form of the articulation of the lower jaw admits of move-

ment, upward and downward, forward, backward, and from sid to side. A combination of these movements takes place in masti cation; during this act the condyles of the jaw describe an oblique rotatory movement in the glenoid cavity. The purposes serve by the fibro-cartilage in this joint are: first, it follows the condyle and interposes a convenient socket for all its movements; second being elastic, it breaks shocks; for shocks here would be almos fatal, considering what a thin plate of bone the glenoid cavity is and that just above it is the brain. Its *nerves* are derived fro the auriculo-temporal and the masseteric branches of the inferio maxillary.

The stylo-hyoid and stylo-maxillary ligaments have bee previously described.

DISSECTION OF THE UPPER EXTREMITY.

The subject should be placed on its back, and, the thorax bei raised by a block placed under the shoulders, the arm is be extended to a right angle with the trunk and slightly rotat outward. A narrow board must be placed under the arm keep it in position, and the hand, with the palm upward, is be firmly fastened, by string, to the board.

Surface Marking. Before commencing the dissection of the ar the student should carefully examine with t eye and the finger the various inequalities of the surface of t skin, which are caused by or are landmarks of important st jacent structures.

Beginning in the middle line, we notice a broad, shallow groo in front of the sternum, between the sternal origins of the pect alis major; about two inches below the upper border of t sternum is a prominent transverse bony ridge (angulus ster which corresponds to the junction of the first and second portio of the sternum.

The clavicle may be easily traced, convex as to its sternal h and concave in its outer half; not placed quite horizontally, b inclined upward in the present position of the limb, and arti lating externally with the prominent acromion process. Extendi obliquely downward and outward, from the middle of the clavic

is a groove, marking the separation between the contiguous borders of the deltoid and pectoralis major, and in which may, by deep pressure, be felt the coracoid process. Another groove, passing outward from the sterno-clavicular joint, indicates the interval between the sternal and clavicular attachments of the pectoralis major. The upper arm below the acromion is rounded, the convexity being caused by the greater and lesser tuberosities of the humerus. In the more common forms of dislocations of the humerus this roundness is lost, and a depression takes its place. Between the thorax and the arm there is a deep hollow —the axilla—which varies according to the position of the arm to the side. Its front border is formed by the pectoralis major, and its hinder border by the latissimus dorsi; and if the fingers be pushed up into this space, the head of the humerus can be easily felt. The free border of the pectoralis major muscle corresponds with the fifth rib, and below this can be distinguished the lower digitations of the serratus magnus with the external oblique.

Dissection. The student must now make three incisions through the skin; the first, along the middle of the whole length of the sternum; the second, along the lower border of the clavicle, and down along the front of the upper arm for four inches; the third, from the ensiform cartilage, backward to the posterior border of the axilla.

The skin should now be taken up with the forceps, at the upper and inner angle, and when the skin has been so far reflected as to enable the fingers to take it up, lay aside the forceps and use the fingers in their place. The skin should be carefully dissected from the subjacent layer of subcutaneous fascia and fat. In doing so, notice the thin, pale fibres of the broad subcutaneous muscle of the neck—*platysma myoides* (Fig. 13).

Beneath this subcutaneous fascia and fat there is the strong deep fascia which closely invests the muscles, and in the axilla it forms a dense fascia, which passes from the pectoralis major to the latissimus dorsi.

Cutaneous Nerves. The numerous nerves which run through the subcutaneous tissue to the skin and mammary

gland must be carefully dissected out. They are derived from various sources; some, branches of the superficial cervical plexus, descend over the clavicle; others, branches of the intercostal nerves, come through the intercostal spaces close to the sternum, each with a small artery; a third series, also branches of the intercostal nerves, come out on the side of the chest, and run forward over the outer border of the pectoralis major.

The *supra-clavicular nerves*, which arise from the third and fourth cervical nerves, descend over the clavicle, and are subdivided, according to their direction, into *sternal*, *clavicular* and *acromial* branches (diagram, p. 83). The *inner* or *sternal* cross the inner end of the clavicle, to supply the skin over the upper part of the sternum. The *middle* or *clavicular* pass over the middle of the clavicle, and supply the integument over the chest and the mammary gland. The *outer* or *acromial* branches cross over the outer end of the clavicle, and distribute their filaments to the skin of the shoulder.

Near the sternum are found the *anterior cutaneous branches* or terminal filaments of the *intercostal nerves*. After piercing the internal intercostal and pectoralis major muscles, each nerve sends an inner filament to the skin over the sternum, and an outer larger one, which supplies the skin over the pectoral muscle. Those of the third and fourth intercostal supply also the mammary gland.

Branches of the internal mammary artery, for the supply of the mammary gland, accompany these nerves. During lactation they increase in size, ramifying tortuously over the surface of the gland. They are occasionally as large as the radial at the wrist.

The *lateral cutaneous branches of the intercostal nerves* come out between the digitations of the serratus magnus on the side of the chest, and divide into anterior and posterior branches. The *anterior* branches curve round the free border of the pectoralis major, and then supply the skin over that muscle and the mammary. The *posterior* branches supply the skin of the back of the chest.

Dissection. Dissect off the superficial fascia and fat with the mammary gland. Thus you will expose the strong *deep fascia* which is closely attached to the pectoral

major, and deltoid muscles. It is continuous, above, with the fascia of the neck; below, with that of the arm. At the axilla it becomes denser, where it passes from the pectoral to the latissimus dorsi muscles.

Reflect this fascia from the pectoralis major by dissecting parallel with the course of its fibres. The muscle having been fully exposed, observe its shape, the course of its fibres, their origin and insertion.[1]

Pectoralis Major. The pectoralis major is the large triangular muscle in front of the chest. It *arises* from the anterior surface of the sternal half of the clavicle, from the front of its own half of the sternum, from the cartilages of all the true ribs except the last, and from the aponeurosis of the external oblique muscle of the abdomen. From this extensive origin the fibres converge toward the arm, the upper ones passing downward and outward, the middle ones transversely outward, and the lower fibres upward and outward; they terminate in a flat tendon, about two inches in breadth, which is *inserted* into the anterior margin of the bicipital groove of the humerus. The arrangement of its fibres, as well as the structure of its tendon, is peculiar. The lower fibres, which form the boundary of the axilla, are folded beneath the rest, and terminate upon the upper part of the tendon—*i. e.* nearer to the shoulder-joint; the upper fibres, which arise from the clavicle, and are frequently separated from the main body of the muscle by a slight interval, descend in front of the lower, and terminate upon the lower part of the tendon. Consequently the upper and lower fibres of the muscle cross each other previously to their insertion.

The object of this arrangement is to enable all the fibres to act simultaneously when the arm is extended.

The upper part of the tendon sends off a fibrous prolongation, which binds down the long head of the biceps, and is attached to the great tuberosity of the humerus; another tendinous expansion

[1] Sometimes we find a thin little muscle running perpendicularly in front of the inner part of the pectoralis major. This is the *rectus sternalis* or *sternalis brutorum*. It arises inferiorly by a tendinous expansion from the rectus abdominis, and is connected above to the tendon of the sterno-mastoid.

is prolonged backward to the tendon of the deltoid muscle
third passes downward to be intimately connected with th
of the upper arm.

The chief *action* of the pectoralis major is to draw the h
toward the chest, as in placing the hand on the opposite sh
or in pulling an object toward the body. When the arm i
and made the fixed point, the muscle assists in raising th
as in climbing. Thus too, on emergency, it can act as an a
muscle of inspiration.

Between the pectoralis major and the deltoid, the great
covering the shoulder, is an interval varying in extent in d
subjects, but always more marked toward the clavicle. It c
a small artery—the *thoracica humeraria*—and the *cepha*
which ascends on the outer side of the arm, and emptie
into the axillary. This interval is the proper place to feel
coracoid process. In doubtful injuries about the shoulc
point of bone is a good landmark in helping the surgeon t
at a correct diagnosis.

The pectorialis major is supplied with nerves by the
thoracic branches of the brachial plexus; with blood, by t
and short thoracic branches of the axillary artery.

Dissection. Anatomy of the Infra-clavicular Region. Reflect the clavicular part of the pe
major by detaching it from the clavicle, a
it downward; in doing so, notice a smal
the external anterior thoracic, which en
under surface of this part of the muscle. Beneath the
thus reflected, part of the pectoralis minor will be expos
this triangle—bounded above by the clavicle; below by th
border of the sternal origin of the pectoralis major; and
outer side by the deltoid—is an important space in wh
relative position of the following objects must be carefi
amined:—

Costo-coracoid Membrane. *a.* A strong ligamentous expansion, ca
costo-coracoid membrane, extends from t
tilage of the first rib to the coracoid process. Betwee
points it is attached to the clavicle, and forms a complete
ment for the subclavius muscle. Its lower crescent-shap

arches over and protects the axillary vessels and nerves; from this edge is prolonged downward a funnel-shaped fascia, which covers the axillary vessels, forming the anterior portion of their sheath; the posterior being formed by a prolongation of the deep cervical fascia. The front portion of this sheath is perforated by the cephalic vein, the thoracica acromialis artery and vein, the anterior thoracic nerves, and the superior thoracic artery. This fascia must be removed.

b. The subclavius muscle enclosed in its fibrous sheath.

c. The axillary vein, artery, and brachial plexus of nerves.

d. Two arteries, the superior or short thoracic and the thoracica acromialis.

e. The termination of the cephalic vein in the axillary.

f. Two nerves, the anterior thoracic, which descend from the brachial plexus below the clavicle, and cross in front of the axillary vessels to supply the pectoral muscles.

Subclavius. This muscle lies between the clavicle and the first rib. It *arises* from the first rib by a short round tendon at the junction of the bone and cartilage in front of the costo-clavicular ligament, and is *inserted* into the groove on the under surface of the clavicle as far outward as the coraco-clavicular ligament. Its nerve comes from the fifth and sixth cervical nerves. Its *action* is to depress the clavicle, and prevent its too great elevation.

Relative position of the Axillary Vessels and Nerves. In the infra-clavicular space before us are the great vessels and nerves of the axilla in the first part of their course. They lie at a great depth from the surface. They are surrounded by a sheath of fascia, which descends with them beneath the clavicle. Their relations with regard to each other are as follows: The axillary vein lies in front of the artery, and rather to its thoracic side. The brachial plexus of nerves is situated above the artery, and on a posterior plane. The plexus consists of two, or sometimes three, large cords, which result from the union of the anterior branches of the four lower cervical and the first dorsal, nerves. The course and relations of the axillary artery will be examined subsequently.

Superior Thoracic and Acromio-thoracic Arteries. These are two branches which arise from th axillary artery in the first part of its course, abov the pectoralis minor. The *superior thoracic* fre quently arises in common with the acromio-thoracic, and passin along the upper border of the pectoralis minor, descends betwee this muscle and the pectoralis major, supplying both and anast mosing with the intercostal and internal mammary arteries. Th *thoracica acromialis* is given off just above the pectoralis mino and shortly divides into three sets of branches, viz: two or thr small *thoracic branches* to the serratus magnus and pectoral mu cles; the *thoracica humeraria*, which descends with the cepha vein, in the interval between the pectoralis major and delto and ramifies in both; and, lastly, the *acromial branch*, wh passes over the coracoid process to the under surface of t deltoid, which it supplies, and communicates with the poster circumflex, a branch of the axillary, and the suprascapular branch of the subclavian. A constant though small branch, t *clavicular*, runs along the anterior aspect of the subclavius. these arteries are accompanied by veins, which most frequen empty themselves into the cephalic, but occasionally into axillary vein.

Cephalic Vein. The cephalic vein is one of the principal cu neous veins of the arm. Commencing on back of the thumb and forefinger, it runs up the radial side of forearm, in front of the elbow joint; thence ascending along outer edge of the biceps, it runs up the interval between the p toralis major and deltoid, pierces the costo-coracoid membra crosses over the axillary artery, and finally empties itself into axillary vein.[1]

Anterior Thoracic Nerves. These nerves come from the brachial ple below the clavicle to supply the pectoral musc There are generally two—an external and an internal—one each pectoral muscle. The *external*, the more superficial, ari from the outer cord of the brachial plexus, passes over the a

[1] The cephalic vein, in some cases, runs over the clavicle to join the exte jugular; or there may be a communication (termed jugulo-cephalic) between th veins.

lary artery and vein, pierces the costo-coracoid membrane, and supplies the pectoralis major on its under aspect; it communicates with the next nerve by a filament which forms a loop on the inner side of the artery; the *internal* and smaller branch comes from the internal cord, and descends between the axillary artery and vein (occasionally through the vein) to supply the pectoralis minor on its under surface.

Difficulty of tying the first part of the Axillary Artery. From this view of the relations of the axillary artery in the first part of its course, some idea may be formed of the difficulty of passing a ligature round it in this situation. In addition to its great depth from the surface, variations sometimes occur in the position of the nerves and veins, which render the operation still more embarrassing. For instance, the anterior thoracic nerves may be more numerous than usual, and form by their mutual communication a plexus around the artery. A large nerve is often seen crossing obliquely over the artery, immediately below the clavicle, to form one of the roots of the median nerve. The cephalic vein may ascend higher than usual, and open into the subclavian; and as it receives large veins corresponding to the thoracic axis, a concourse of veins would be met with in front of the artery. Again, it is by no means uncommon to find a deep-seated vein, the supra-scapular, crossing over the artery to join the axillary vein.

DISSECTION OF THE AXILLA.

Sebaceous Glands. On the under surface of the skin of the axilla, near the roots of the hairs, are numerous sebaceous glands. They are of a reddish-brown color, and rather larger than a pin's head.

Axillary Fascia. This dense fascia, which lies immediately beneath the skin of the axilla, is a continuation of the general fascial investment of the muscles. It closes in and forms the floor of the cavity of the axilla. Externally, it is strengthened by fibres from the tendons of the pectoralis major and latissimus dorsi, and is continuous with the fascia of the arm; internally, it is prolonged on the side of the chest, over the serratus magnus muscle; in front and behind it divides, so as to

enclose between its layers the muscles which form the boundaries of the axilla. Thus the anterior layer encloses the two pectoral muscles, and is connected with the coracoid process, the costo-coracoid ligament and the clavicle; the posterior layer encloses the latissimus dorsi, and passes backward to the spine.

A subcutaneous artery, sometimes of considerable size, is often found in the substance of the axillary fascia. It generally arises from the brachial, or from the lower part of the axillary artery, and runs across the floor of the axilla toward the lower edge of the pectoralis major. It is not a named branch, but should be remembered, as it would occasion much hemorrhage if wounded in opening an abscess.

Dissection and Contents of the Axilla. Reflect the axillary fascia, to display the boundaries and the contents of the axilla. The dissection of this space is difficult, and must be done cautiously. Bear in mind that the trunk blood vessels and nerves run through the *upper* and *outer* part of the axilla; that the long thoracic artery runs along the *anterior* border, and the subscapular artery along the *posterior*. Commence dissecting, therefore, in the middle; break down with the handle of the scalpel the loose connective tissue, fat and lymphatic glands which occupy the cavity. You will soon discover some cutaneous nerves coming out between the ribs, and then crossing the axillary space. These nerves are the *posterior lateral cutaneous branches of the intercostal nerves;* they perforate the intercostal spaces between the digitations of the serratus magnus, midway between the sternum and the spine, and divide into anterior and posterior branches. The *anterior* turn over the pectoralis major, to supply the skin on the front of the chest and the mammary gland. The *posterior* pass backward over the latissimus dorsi, and are distributed to the skin covering this muscle and the scapula.

Intercosto-humeral Nerves. The posterior lateral branch of the second intercostal nerve requires a special description. It is larger than the others, and is called the *intercosto-humeral* because it supplies the integuments of the arm. It comes through the second intercostal space, traverses the upper part of the

axilla, where it receives a branch of the lesser internal cutaneous nerve (nerve of Wrisberg), and, piercing the fascia, terminates in filaments, which are distributed to the skin on the inner side and back of the arm, as low as the internal condyle. The corresponding branch of the third intercostal is also an *intercosto-humeral* nerve. It receives a branch from the second, and runs a similar course. The distribution of these nerves accounts for the pain down the arm which is sometimes experienced in pleurisy.

Boundaries of the Axilla. The axilla is a conical space, of which the *apex* is beneath the clavicle, and the *base* between the pectoralis major and the latissimus dorsi. Obviously it varies in capacity according to the position of the arm to the side. On the *inner* side, it is bounded by the four upper ribs, with their corresponding intercostal muscles and the serratus magnus; on the *outer*, by the humerus, covered by the coraco-brachialis and biceps; in *front*, by the pectoralis major and minor; *behind*, by the latissimus dorsi, teres major and subscapularis. Its anterior and posterior boundaries converge from the chest, so that the axilla becomes narrower toward the arm. With a full view of the axilla before you, bear in mind that pus may burrow under the pectoral muscles, or under the scapula, or that it may run up beneath the clavicle and point in the neck, if the abscess be allowed to remain unopened.

Axillary Lymphatic Glands. The axillary glands form a continuous chain, beneath the clavicle, with the cervical glands. They are from ten to twelve in number, of a reddish-brown color and variable size. Most of them lie near the axillary vessels; others are embedded in the loose tissue of the axilla; sometimes one or two small ones are observed along the lower border of the pectoralis major. They are supplied with blood by a branch—*thoracica alaris*—of the axillary artery, and by branches from the thoracic and subscapular arteries.

These glands receive the lymphatics from the arm, from the front and side of the chest, and from the outer half of the mammary gland. It is these glands which frequently become enlarged in cancer of the mammary gland. From these glands the

efferent lymphatics pass along with the subclavian artery and terminate, on the right side, in the right lymphatic duct; and on the left side, in the thoracic duct.

Dissection. Now cut through the pectoralis major, about the middle, and turn the inner part of the muscle toward the sternum, and the outer part toward the arm. The pectoralis minor is thus exposed, together with the ramifications of the short and long thoracic arteries. Preserve the arteries, as far as possible, in connection with the main trunks.

Pectoralis Minor. This triangular muscle *arises* from the third, fourth and fifth ribs, near the costal cartilages, and from the thick fascia over the intercostal spaces. The fibres run obliquely upward and outward, and converge to a strong tendon, which is *inserted* into the anterior surface of the coracoid process. The tendon is connected to that of the coraco-brachialis and biceps by a strong fascia, which forms a protection for the subjacent axillary vessels and nerves. The action of this muscle is to draw the scapula downward and forward. Its nerve is derived from the internal anterior thoracic.

Dissection. Having examined the muscles which form the anterior boundary of the axilla, we pass now to the course and relations of the axillary artery and its branches. To have a clear view, reflect the subclavius from its insertion, and cut the pectoralis minor through its middle.

Axillary Artery, its Course and Relations. This artery, the continuation of the subclavian, takes the name of *axillary* at the outer border of the first rib. It then passes downward and outward, through the upper part of the axilla, beneath the two pectoral muscles, and along the inner border of the coraco-brachialis, as far as the lower border of the tendon of the teres major, beyond which it is continued under the name of the brachial. Its course is divided, for convenience of description, into three parts; the first lies above the pectoralis minor; the second behind that muscle; and the third below it.

In the *first part* of its course the artery is *covered* by the pectoralis major and the costo-coracoid membrane, the subclavius, and is crossed by the cephalic and acromio-thoracic veins. On

its *inner side*, and slightly in front, is the axillary vein; on its *outer side* is the brachial plexus of nerves; *behind* it are the first intercostal space, the second digitation of the serratus magnus, and the posterior thoracic nerve (external respiratory of Bell).

FIG. 75.



DIAGRAM OF AXILLA.

1. Axillary artery.
2. Brachial artery.
3. Thoracica humeraria a.
4. Superior thoracic a.
5. Subscapular a.
6. Dorsalis scapulæ a.
7. Posterior circumflex a.
8. Superior profunda a.
9. Posterior thoracic nerve.
10. Long subscapular n.
11. Median n.
12. Cephalic vein.
13. Musculo-cutaneous n.
14. Teres major.

In the *second part* of its course it lies *behind* the pectoralis

major and minor; on its *inner side* is the axillary vein, still slightly anterior, but separated from the artery by the inner cord of the brachial plexus; on its *outer side* is the outer cord of the brachial plexus; and *behind* it is the posterior cord of the plexus, and also a quantity of loose connective tissue which separates it from the subcapularis muscle. The inner head of the median nerve is often in front of the artery in this part of its course.

In the *third part*, in *front* of the artery, are the pectoralis major, the two roots of the median nerve, converging like the letter V; and lower down is the skin and the fascia of the arm; on the *outer side* are the coraco-brachialis, the musculo-cutaneous and median nerves; on the *inner side* are the axillary vein, the ulnar, and the

FIG. 76.

PLAN OF THE BRANCHES OF THE AXILLARY ARTERY.

1. Thoracic axis, giving off—
2. Short thoracic.
3. Thoracica acromialis.
4. Thoracica humeraria.
5. Long thoracic.
6. Subscapular.
7. Dorsalis scapulæ.
8. Anterior circumflex.
9. Posterior circumflex.

two internal cutaneous nerves; *behind* it are, in succession, th[illegible]he subscapularis, the latissimus dorsi, the teres major, and th[illegible] musculo-spiral and circumflex nerves.

Branches of the Axillary Artery. The number and origin of these branches often[illegible]n vary, but their general course is in most case[illegible]s similar, and they usually arise in the following order:—

a. The *superior thoracic* arises above the pectoralis minor, and[illegible] divides into branches, which have already been described (p. 330).

b. The *acromial thoracic* also arises above the pectoralis minor, and gives off numerous branches already described (p. 330).

c. The *alar thoracic*, variable in its origin, supplies the lymphatic glands and the connective tissue of the axilla.

d. The *inferior* or *long thoracic artery* (external mammary) runs along the lower border of the pectoralis minor to the side of the chest. It supplies the mammary gland, the serratus magnus and pectoral muscles, and maintains a free anastomosis with the short thoracic, internal mammary and intercostal arteries.

e. The *subscapular* is the largest branch of the axillary; it arises opposite the lower border of the subscapularis, and after running a short course of about an inch and a half, divides into an anterior and posterior branch.

FIG. 77.

1. Subscapularis.
2. Teres major.
3. Long head of triceps.
4. Square space for post. circumflex a. and n.
5. Triangular space for dorsalis scapulæ a.
6. Space for musculo-spiral n., and superior profunda a.

DIAGRAM OF THE ORIGINS OF THE TRICEPS.

The *anterior branch* runs along the anterior edge of the subscapularis toward the lower angle of the scapula. Its numerous branches supply the subscapularis, latissimus dorsi, serratus magnus and teres major, and anastomose with the intercostal and thoracic arteries, and with the posterior scapular (a branch of the subclavian).

The *posterior branch* (dorsalis scapulæ) runs to the back of the scapula, through a triangular space, bounded in front by the long head of the triceps; below by the teres major; and above by the subscapularis and teres minor (Fig. 77). It gives off a small branch which enters the subscapular fossa beneath the

subscapularis, supplying it and anastomising with the supr scapular and posterior scapular arteries. On the back of t scapula it divides into two branches; one runs in the groove the axillary border of the scapula, lying beneath the teres min and ramifies in the infraspinous fossa between the bone and infraspinatus; the other runs down between the teres minor a major on their dorsal aspects, and passes to the inferior angle the scapula, anastomosing with the posterior and suprascapu arteries. The subscapular vein empties itself into the axillary v

f. The *posterior circumflex artery* arises from the back the axillary artery, and is as large as the subscapular, cl to which it is given off; or both may arise by a comm trunk from the axillary. It passes backward, with its c responding veins and nerve, through a quadrilateral spa bounded above by the subscapularis and teres minor, bel by the teres major, externally by the neck of the humerus, a internally by the long head of the triceps (Fig. 77). It then wi round the back of the neck of the humerus, and is chiefly tributed to the under surface of the deltoid.

Besides the deltoid, the posterior circumflex artery supp the long head of the triceps, the head of the humerus, and shoulder joint. It inosculates above with the acromio-thora and suprascapular arteries; below with the ascending branch the superior profunda (a branch of the brachial), and in front w the anterior circumflex artery. Should you not find the poste circumflex artery in its normal position, look for it (as a bra of the brachial) below the tendon of the teres major.

g. The *anterior circumflex artery*, much smaller than the p terior, runs in front of the neck of the humerus, above the ten of the latissimus dorsi. It passes directly outward beneath coraco-brachialis and short head of the biceps, close to the bo and terminates in the under surface of the deltoid, where it in culates with the posterior circumflex.

The anterior circumflex artery sends a small branch which r with the long tendon of the biceps up the groove of the hume and is called, on that account, the *bicipital* artery. It supp the shoulder joint and the neck of the humerus.

If the axillary were tied below the pectoralis minor, the collateral circulation would be established by the suprascapular, and its branches anastomosing with the subscapular, the dorsalis scapulæ and the posterior circumflex; the posterior scapular with the dorsalis scapulæ and subscapular arteries.[1]

FIG. 78.

THE BRACHIAL PLEXUS OF NERVES.

C 4-8. The five lower cervical nerves.
D 1. The first dorsal nerve.
9. The rhomboid nerve—to rhomboidei major and minor.
10. The supra-scapular nerve—to supra and infra-spinati.
11. The nerve to the subclavius.
12. Outer anterior thoracic nerve—to pectoralis major.
13. Inner anterior thoracic nerve—to pectoralis minor.
14, 15, 16. The subscapular nerves—to subscapularis, latissimus dorsi and teres major.
17. Lesser internal cutaneous nerve.
18. Musculo-cutaneous nerve.
19. Musculo-spiral nerve.
20. Median nerve.
21. Circumflex nerve—to deltoid and teres minor.
22. Ulnar nerve.
23. Internal cutaneous nerve.
24. External respiratory nerve of Bell, or posterior thoracic.

Axillary Vein. The *axillary vein* is formed by the continuation upward of the basilic vein, and extends from the

The axillary artery varies much as to the branches it gives off; occasionally (1 in gives off the radial artery; more rarely (1 in 72) it gives off the ulnar; and rarely still (1 in 506) it gives off the interosseous artery.

lower border of the teres major to the outer border of the first rib. It receives the venæ comites of the brachial artery near the lower border of the subscapularis. It receives the subscapular and the other veins corresponding to the branches of the axillary artery, with the exception of the circumflex, which usually join either the subscapular or one of the venæ comites. The axillary near its termination also receives the cephalic vein.

The axillary vein in the upper part of its course lies in front of the artery, and close to its sternal side; in the lower two-thirds of its course the vein lies still to the sternal side of the artery, but is separated from it by some of the nerves of the brachial plexus.

Axillary or Brachial Plexus of Nerves.

This plexus is formed by the anterior trunks of the four lower cervical and first dorsal nerves and receives also a small communicating branch from the fourth cervical nerve. The plexus is broad at the lower part of the neck, where it emerges between the anterior and middle scalene muscles; but it gradually contracts as it descends beneath the clavicle into the axilla, and on a level with the coracoid process distributes its large branches to the upper limb.

The arrangement of the cervical nerves in the formation of the plexus is variable, often not alike on both sides.[1] The most frequent disposition is this—the fifth and sixth cervical unite at the outer border of the scalenus medius to form an upper cord; the eighth and the first dorsal form between the scaleni muscles a lower cord; the seventh cervical runs alone, as a middle cord for a short distance. Each of these nerves divides, just external to the outer border of the scalenus medius, into an anterior and a posterior branch; the anterior branches given off from the fifth, sixth and seventh cervical form the *outer cord* of the plexus; the anterior branches given off from the eighth cervical and the first dorsal form the *inner cord;* while the posterior branches of all the nerves—viz., the fifth, sixth, seventh, eighth cervical and the first dorsal—unite to form the *posterior cord.*

At first these cords are placed on the outer side of the axillary artery, but behind the pectoralis minor they are situated one on

[1] Frequently the second dorsal nerve sends upward a communicating filament to the first dorsal nerve. (*Journal of Anatomy*, vol. xi, p. 539.)

the outer side of, one on the inner side of, and one behind, the axillary artery.

The brachial plexus gives off some branches above the clavicle, which were dissected with the neck (p. 143). Below the clavicle it gives off the following:—

From the *outer* cord proceed an anterior thoracic branch, the musculo-cutaneous, and the outer head of the median; from the *inner* cord proceed the inner anterior thoracic nerve, the inner head of the median, the ulnar, the internal cutaneous, and the lesser internal cutaneous; from the *posterior* cord proceed the three subscapular nerves, the circumflex and the musculo-spiral.

The *anterior thoracic* nerves have been described (p. 330).

Subscapular Nerves. The *three subscapular* nerves are found on the surface of the subscapularis. They come from the posterior cord of the brachial plexus, and supply, respectively, the latissimus dorsi, teres major and subscapularis. The nerve to the latissimus dorsi (*long subscapular nerve*) runs with the anterior branch of the subscapular artery to the lower border and inner surface of the muscle.

The *nerve to the teres major* is either a branch of the preceding, or comes separately from the posterior cord. It lies nearer to the humerus than the long subscapular. It gives off also a small branch to the anterior border of the subscapularis.

The *nerve to the subscapularis* arises from the posterior cord, higher than the others, and enters the muscle not far from its upper border, in company with a small artery.

Circumflex Nerve. The circumflex nerve accompanies the *posterior circumflex artery*. This large nerve comes from the posterior cord, and after giving a small filament to the shoulder joint, passes, with its companion artery, through the quadrilateral space (p. 338), to the under surface of the deltoid. Here the nerve divides into an upper and a lower branch. The *upper* supplies the anterior part of the deltoid and the skin over it; the *lower* supplies the back part of the deltoid, and gives the nerve to the teres minor,[1] upon which nerve sometimes a little

[1] This branch to the teres minor is said to be constant in all mammalia that have been examined in reference to this point.

gangliform swelling can be seen; it enters the under aspect of the middle of this muscle. After furnishing these muscular branches the nerve turns round the posterior border of the deltoid, and diverges in filaments which supply the skin over the back of this muscle and over the long head of the triceps.

Latissimus Dorsi. This broad, flat muscle forms the posterior margin of the axilla. It arises from the external l of the crest of the ilium, from an aponeurosis attached to the spino processes of the six lower dorsal, of all the lumbar, and of the sacr vertebræ and their supraspinous ligament, and by fleshy digitatio from the three or four lower ribs, interdigitating with those of t external oblique; in some cases, as it passes over the inferior an g of the scapula, it has an additional origin from the angle. Its fib converge from this large origin, and the muscle is *inserted* in the bottom of the bicipital groove of the humerus by a broad f tendon, which curves round the lower border of the teres maj The axillary vessels and nerves lie upon the tendon close to insertion. Its nerve is the long subscapular branch from t posterior cord of the brachial plexus, and it enters the musc close to its anterior border, in company with a large branch of t subscapular artery.

Teres Major. This muscle lies behind and to the inner si of the latissimus dorsi, is closely connected wi it, and assists in forming the posterior boundary of the axilla. *arises* from the oval surface on the dorsal aspect of the low angle of the scapula, and from the fibrous septa between it a the teres minor and infraspinatus, and is *inserted* by a broad fl tendon, about two inches in length, behind the latissimus dor into the posterior margin of the bicipital groove of the humeru The tendon extends below that of the latissimus dorsi, and bursa or sac, lubricated with serum, intervenes between the tw tendons. The action of this and the preceding muscle is to dra the humerus inward and backward. Its nerve is the midd subscapular, and lies along the dorsalis scapulæ artery.

Subscapularis. This muscle *arises* from the posterior two-thir of the subscapular fossa of the scapula, with th exception of the angles, neck, and the posterior border, and fro

the intermuscular septa attached to the bony ridges. Its fibres converge to a strong tendon, which passes under the axillary vessels and nerves, over the inner side of the shoulder joint, and is *inserted* into the lesser tuberosity of the humerus, and into the neck of the humerus for an inch below it. The tendon of the muscle is intimately connected with the capsular ligament of the shoulder joint, and between the base of the coracoid process and the tendon is a bursa, which communicates with the joint. Its action is to rotate the humerus inward, and when the arm is raised to draw it to the side. Its nerves come from the upper and middle subscapular nerves.

Serratus Magnus. This muscle covers the side of the chest like a girth. It *arises* from the front of the outer surfaces of the eight upper ribs by nine slips or digitations, the second rib having two. Its fibres converge, and are *inserted* into the posterior border of the scapula in the following manner: the first two digitations are attached into the upper angle of the scapula; the third and fourth digitations along nearly the whole length of the posterior border; the remainder are inserted into the inferior angle. Its action is to draw the scapula forward; but of this more hereafter. It is supplied by the following nerve, which is seen on its outer surface.

Posterior Thoracic or External Respiratory Nerve of Bell. This nerve supplies the serratus magnus only. It comes from the fifth and sixth cervical nerves, and after passing through the scalenus medius, runs behind the axillary vessels, along the outer surface of the serratus magnus, each digitation receiving a separate filament.[1]

DISSECTION OF THE UPPER ARM.

Surface Marking. In front of the upper arm may be distinguished the long prominence of the biceps muscle, and lower down at the bend of the elbow its tendon may be easily felt. The bend of the elbow, in muscular subjects, presents a triangu-

[1] It may be asked why this nerve is called the external respiratory. It was so named by Sir C. Bell, who considered the serratus magnus as the external respiratory muscle, co-operating with the diaphragm or internal respiratory muscle.

lar depression, with its boundaries formed on the inner side by the pronator teres, and on the outer side by the supinator longus. Superficially in this space the subcutaneous veins can be recognized, of which a fuller description will be entered into later on. On the inner side of the elbow, the internal condyle of the humerus is very prominent, and behind this is the olecranon; between these is a hollow in which may be felt the ulnar nerve. The olecranon is situated nearer the internal than the external condyle, which is visible on the outer side; below this, is a dimple which corresponds with the head of the radius.

Dissection. Continue the incision down the inner side of the arm as far as two inches below the elbow, and then make a transverse incision from the inner to the outer side of the forearm. Reflect the skin, and trace out the cutaneous nerves, and the numerous veins in front of the elbow.

Cutaneous Nerves. On the inner side of the arm are the intercosto-humeral, the internal cutaneous branch of the musculo-spiral, the internal cutaneous, and the lesser internal cutaneous (nerve of Wrisberg) nerves; on the outer side are the cutaneous branches of the circumflex, the external cutaneous branches of the musculo-spiral, and lower down is the musculo-cutaneous nerve.

The filaments of the *intercosto-humeral nerves* (p. 332) descend along the inner and posterior part of the arm as far as the olecranon, and communicate with the internal cutaneous branch of the musculo-spiral nerve.

The *internal cutaneous nerve* perforates the fascia, with the basilic vein about the middle of the arm, and divides into an anterior and a posterior branch; the *anterior* passes down in front of the arm (as a rule, beneath the median basilic vein), and supplies the skin as far as the wrist, communicating with a cutaneous branch of the ulnar nerve; the *posterior* winds round to the back of the forearm, behind the internal condyle, as far as the wrist, and communicates above the elbow with the nerve of Wrisberg, and above the wrist with the dorsal branch of the ulnar nerve.

The *lesser internal cutaneous* (*nerve of Wrisberg*) perforates the

fascia about the lower third of the arm, and supplies the skin over the internal condyle and the olecranon. This nerve, as it lies close to the axillary vein, communicates with the first or second intercosto-humeral nerve.

The *internal cutaneous branch of the musculo-spiral nerve* pierces the fascia, and supplies the skin of the inner and posterior aspect of the middle of the arm as far as the olecranon.

FIG. 79.

1. Acromial branches } of the superficial cervical plexus.
2. Clavicular branches }
3. Cutaneous branches of the circumflex nerve.
4. Branches of the internal cutaneous nerve.
5. External cutaneous branch of the musculo-spiral nerve.
6. Internal cutaneous nerve.
7. Its posterior cutaneous branch.
8. The cutaneous branch of the musculo-cutaneous nerve.

DISTRIBUTION OF CUTANEOUS NERVES TO THE FRONT OF THE SHOULDER AND ARM.

The *cutaneous branches of the circumflex nerve* pierce the fascia over the insertion of the deltoid, and supply the skin of the upper part of the arm on its outer side.

The *external cutaneous branches of the musculo-spiral nerve* are two in number; the *upper* and smaller accompanies the cephalic vein in the lower half of the arm; the *lower* may be traced down the outer and back part of the forearm nearly as far as the wrist, where it joins the posterior branch of the musculo-cutaneous nerve.

On the outer side of the tendon of the biceps, the *cutaneous branch of the musculo-cutaneous nerve* perforates the fascia, and divides into many filaments, which supply the skin of the outer part of the forearm.

Disposition of Veins in front of the Elbow.

Attention should now be directed to the disposition of the veins in front of the elbow. In cleaning these veins, take care not to divide the

FIG. 80.



SUPERFICIAL VEINS AND NERVES AT THE END OF THE LEFT ELBOW.

branches of the internal and external cutaneous nerves which pa over and under them.

The following is the ordinary arrangement of the superfic veins at the bend of the elbow (Fig. 80). On the outer side is t *radial;* on the inner side is the *ulnar vein*, formed by the ju tion of the anterior and posterior ulnar cutaneous veins; in t centre is the median, which divides into two branches, the exter of which, uniting with the radial to form the cephalic vein, called the *median cephalic;* the internal, uniting with the uln

to form the basilic, is named the *median basilic*. Near its bifurcation, the median vein communicates by a branch (mediana profunda) with the deep veins which accompany the arteries of the forearm.

Trace the *cephalic vein* up the arm. It runs along the outer border of the biceps to the groove between the pectoralis major and the deltoid, and dipping down between these two muscles, terminates in the axillary vein immediately above the pectoralis minor.

The *basilic vein*, the largest of the veins of the upper arm, ascends along the inner side of the arm with the internal cutaneous nerve. Near the middle of the arm it perforates the fascia, and empties itself either into the internal vena comes of the brachial artery, or into the axillary vein.

Relation of the Cutaneous Nerves and Veins at the Elbow.

The principal branches of the cutaneous nerves pass beneath the veins: that is to say, as a rule, the internal cutaneous passes behind the median basilic vein, and the external cutaneous behind the median cephalic; but it should be remembered that many small filaments cross in front which are exposed to injury in venesection.

Relation of Median Basilic Vein to Brachial Artery.

Since the median basilic vein is larger than the median cephalic, and, on account of the strong fascia beneath, more easily compressible, it is usually chosen for venesection; its position, therefore, in reference to the brachial artery, becomes important. The vein is only separated from the artery by the *semilunar fascia*, derived from the tendon of the biceps. This fascia is in some subjects remarkably thin. Sometimes the artery lies above the fascia, in contact with the vein. In choosing, therefore, this vein for venesection, there is a risk of wounding the artery; hence the practical rule, to bleed either from the median cephalic, or from the median basilic above the situation where it crosses the brachial artery.

Lymphatic Glands.

Immediately above the internal condyle, in the neighborhood of the basilic vein, we find one or two small *lymphatic glands*. Others may be found higher up

along the inner side of the arm. A gland is occasionally n with at the bend of the elbow; but never below this joint. Th little glands are the first which are liable to become tender a enlarged after a poisoned wound of the hand.

Muscular Fascia and its Connections. The fascia which invests the muscles of t upper arm is a continuation of the fascia of t trunk and the axilla. This membrane varies density; thus it is thin over the biceps, stronger on the inner si of the arm, to protect the brachial vessels and nerves, and strong over the triceps. At the upper part of the arm it is connect with the coracoid process and the clavicle; it is strengthened the axilla by an expansion from the tendons of the pectoralis ma and latissimus dorsi; posteriorly, it is attached to the spine the scapula. The fascia surrounds the brachial vessels witl sheath, and furnishes partitions which separate the muscles fr each other. Of these partitions, the most marked are, the *exter* and *internal intermuscular septa*, which divide the muscles the anterior from those on the posterior surface of the up arm. These septa are attached to the condyloid ridges of humerus and to the condyles. The *internal intermuscular tum*, the stronger of the two, begins at the insertion of teres major, and is connected with the tendinous insertion of coraco-brachialis; it separates the triceps extensor from brachialis anticus, to both of which it affords attachment to t muscular fibres. It is pierced by the ulnar nerve and inferior profunda and anastomotic arteries. The *external i muscular septum* commences from the insertion of the delt and separates the brachialis anticus, the supinator longus, the extensor carpi radialis longior in front, from the triceps tensor behind, to all of which muscles it affords attachm It is pierced by the musculo-spiral nerve and the supe profunda artery.

At the lower part of the upper arm the fascia is remark strong, especially where it covers the brachialis anticus and brachial vessels, and is continued over the muscles on the in side of the forearm. At the back of the elbow the fasci attached to the tendon of the triceps and the olecranon.

Dissection. Now remove the fascia corresponding to the incisions through the skin, in order to see the muscles on the front of the arm—namely, the biceps, the coraco-brachialis, and the brachialis anticus. The long rounded muscle in front is the biceps; the muscle attached with it to the coracoid process is the coraco-brachialis; and the broad flat muscle covering the lower end of the humerus is the brachialis anticus.

Biceps. The *biceps*, as its name implies, arises by two heads—a long and a short. The *short head arises* from the tip of the coracoid process of the scapula, by a thick, flat tendon in common with a slender muscle on its inner side, called the coraco-brachialis. The *long head arises* from the upper border of the glenoid fossa of the scapula and the glenoid ligament, by a long, rounded tendon, which, traversing the shoulder-joint, passes over the head of the humerus; there pierces the capsular ligament, and descends in the groove between the two tuberosities. The tendon is retained in the groove by a fibrous bridge derived from the capsule of the joint, and connected with the tendon of the pectoralis major. Divide this bridge, and see that the synovial membrane of the joint is reflected round the tendon, and accompanies it for about two inches down the groove, thus forming a synovial fold. The object of this is to facilitate the play of the tendon, and to carry little arteries (from the anterior circumflex) for its supply. The two heads unite about the middle of the arm, and form a single muscle, which terminates on a strong, flat tendon of considerable length; this dips down into the triangular space at the bend of the elbow, and after a slight twist upon itself, is *inserted* into the posterior part of the tubercle of the radius. The anterior part of the tubercle, over which the tendon plays, is crusted with cartilage, and a *bursa* intervenes to diminish friction. The most internal fibres of the muscle are inserted into a *strong, broad aponeurosis*, which is prolonged from the inner border of the tendon to the fascia on the inner side of the forearm. This aponeurosis, called the *semilunar fascia of the biceps*, protects the brachial vessels and the median nerve at the bend of the elbow.

The *action* of the biceps is twofold. 1. It is a flexor of the

forearm. 2. It is a powerful supinator of the forearm, in co
quence of its insertion into the *posterior* part of the tubercle of
radius. Its power of supination is greatest when the forearm
half bent, because its tendon is then inserted at a right an
Why does the long tendon pass through the shoulder joint?
acts like a strap, and confines the head of the humerus in its pro
centre of motion.[1] But for this tendon, the head of the bone, wh
the deltoid acts, would be pulled directly upward and strike agai
the under surface of the acromion. When the tendon is ruptu
or dislocated from its groove a man can move his arm backw
and forward, but he cannot raise the smallest weight.[2] T
biceps is supplied with blood by a branch from the brach
which runs into the middle of its inner side, and divides i
ascending and descending branches. Its nerve comes from
musculo-cutaneous.

Coraco-brachialis. This thin muscle is situated at the upper p
of the arm, and runs parallel to the inner bor
of the short head of the biceps. It *arises* by fleshy fibres fr
the point of the coracoid process, in common with the short h
of the biceps, and from a fibrous septum which lies betw
them. The muscle descends backward and outward, and ter
nates on a flat tendon, which is *inserted* into the inner side of
middle of the humerus, between the brachialis anticus and
inner head of the triceps. Its action is to draw the humerus
ward and inward—*e. g.*, in bringing the gun up to the shoul
It is supplied by a branch from the musculo-cutaneous ne
which pierces it.

Concerning the coraco-brachialis, remember: 1. That the n
culo-cutaneous nerve runs through it; 2. That its inner fle
border is the guide to the axillary artery in the last part o
course; 3. That the brachial artery lies upon its flat tendo
insertion, and can here be effectually compressed by the fi
or the tourniquet.

[1] Another action of the long tendon of the biceps would seem to be that of an
nal rotator of the humerus when that bone is rotated externally. The marked p
nence of the internal tuberosity and the groove on its outer aspect would favor this

[2] See a preparation in the Museum of St. Bartholomew's Hospital.

The coraco-brachialis and biceps are covered at their upper part by the deltoid and pectoralis major. The head of the humerus rolls beneath the coraco-brachialis and short origin of the biceps; and a large *bursa* is interposed between these muscles and the tendon of the subscapularis, which covers the head of the bone.

Brachialis-anticus. This broad muscle covers the lower half of the humerus, and is partially concealed by the biceps. Between the two muscles is the musculo-cutaneous nerve, which supplies them both.

It *arises* from the humerus by a fleshy digitation on either side of the tendon of the deltoid; from the lower half of the front and inner surfaces of the bone, and from the intermuscular septa. The muscle, becoming thicker and broader, covers the front of the capsule of the elbow-joint, to which it is more or less attached, and terminates on a tendon, which is *inserted* in a pointed manner into the anterior surface of the coronoid process of the ulna. Its *action* is to bend the forearm. Its nerves come from the musculo-cutaneous, and it usually receives in addition a small branch from the musculo-spiral.

Now examine the course and relations of the brachial vessels and nerves.

Course and Relations of the Brachial Artery. The *brachial artery*—the continnation of the axillary—takes its name at the lower border of the teres major. It runs down the anterior and the inner side of the arm, along the *inner border of the coraco-brachialis and biceps*, to about an inch below the elbow, where it divides, near the coronoid process of the ulna, into the radial and ulnar arteries.

Thus its direction corresponds with a line drawn from the deepest part of the axilla to the middle point between the condyles of the humerus.

In the upper part of its course it *lies on* the long and inner heads of the triceps (from the long head it is separated by the musculo-spiral nerve and superior profunda artery); in the middle, it lies on the tendon of the coraco-brachialis; in the lower part, on the brachialis anticus.

In *front* of the artery are the internal cutaneous nerve, t
median basilic and basilic veins; the median nerve, which cross
obliquely over the artery, being on its outer side near the axil
and on its inner side near the elbow; and lastly, the artery
more or less overlapped, in the first part of its course, by t
coraco-brachialis, lower down by the fleshy belly of the biceps; t
inner borders of these muscles, in their respective situations, bei
the best guides to the artery.

On the *outer side* of the artery are the median nerve, t
coraco-brachialis, and biceps.

On the *inner side* are, at first, the ulnar nerve, the intern
cutaneous nerves; and, below, the median nerve.

The artery is accompanied by two veins (*venæ comites*) and t
median nerve, all of which are invested in a common sheath
fascia.

The *ulnar nerve* runs along the inner side of the artery as
as the middle of the arm. Below this point, the nerve leaves t
artery, and passes through the internal intermuscular septum
get behind the internal condyle.

About the middle of the humerus, the artery lies for nea
two inches on the tendon of the coraco-brachialis, and is so clo
to the bone that it can be effectually compressed, provided t
pressure be made in the proper direction—namely, outwa
Here, too, it is crossed by the median nerve.

At the bend of the elbow the artery is crossed by the semilu
fascia from the biceps. It enters a triangular space, bounded
the pronator radii teres internally, and by the supinator ra
longus externally. It sinks into this space, with the tendon
the biceps to its outer side, and the median nerve to its inn
all three rest upon the brachialis anticus. To compress
artery here, pressure should be made directly backward. Op
site the coronoid process of the ulna it divides into the radial a
ulnar arteries.

Two veins, of which the internal is the larger, lie in close c
tact with the brachial artery, and communicate at frequent interv
by transverse branches. Near the axilla they join and form
axillary vein.

Branches of Brachial Artery. The brachial artery gives off four branches, all from its inner side, namely, the superior profunda, the inferior profunda, the nutrient artery, and the anastomotica magna. It also distributes muscular branches to the coraco-brachialis and biceps, which are given off from its outer side.

FIG. 81.



PLAN OF THE CHIEF BRANCHES OF THE BRACHIAL ARTERY AND THE ARTERIAL INOSCULATIONS ABOUT THE RIGHT ELBOW JOINT.

a. The *profunda superior* arises from the inner and back part of the brachial artery, immediately below the tendon of the teres major.[1] It winds round the back of the humerus, between the

[1] If the profunda be not in its usual place, look for it above the tendon of the latissimus dorsi, where it will probably be given off from a common trunk with the anterior circumflex.

outer and inner heads of the triceps, accompanied by the musculo spiral nerve, and a little above the middle of the arm divide into two branches, which run for some distance on either sid of the nerve. One of these runs in the substance of the tricep muscle, with the nerve to the anconeus, as far as the olecranor and anastomoses with the posterior ulnar recurrent, the inter osseous recurrent, and anastomotica magna arteries; the othe branch accompanies the musculo-spiral nerve to the outer sid of the arm, where it perforates the external intermuscular septum It then descends deep in the interval between the brachiali anticus and supinator radii longus, and terminates in numerou ramifications, some of which pass in front of the externa condyle, others behind it, to inosculate with the radial and inter osseous recurrent arteries.

Before its division, the superior profunda sends several branche to the deltoid, coraco-brachialis, and the triceps, some of whic inosculate with the circumflex. These assist in establishing collateral circulation when the brachial artery is ligatured abov the origin of the profunda.

b. The *profunda inferior* arises from the brachial, opposite to th insertion of the coraco-brachialis, or sometimes by a commc trunk with the superior profunda. It runs with the ulnar ner on the inner head of the triceps (which it supplies), pass through the internal intermuscular septum, and then descen to the interval between the internal condyle and the olecrano inosculating with the posterior ulnar recurrent and anastomoti magna arteries. It also sends a small branch down in fro of the internal condyle to anastomose with the anterior uln recurrent.

c. The *nutrient artery* of the humerus arises sometimes fro the brachial, sometimes from the inferior profunda. It pierc the tendon of the coraco-brachialis, runs obliquely downwa through the bone, and in the medullary canal divides into ascen ing and descending branches, which anastomose with the nutrie vessels of the bone derived from the periosteum.

d. The *anastomotica magna* arises from the inner side of th brachial, about two inches above the elbow, runs tortuous

inward, transversely across the brachialis anticus, and divides into branches, some of which pass in front of the internal condyle, anastomosing with the anterior ulnar recurrent artery; another passes behind the internal condyle by piercing the internal intermuscular septum, and anastomoses with the inferior profunda and posterior ulnar recurrent arteries; and one branch forms an arch, above the olecranon fossa, with the superior profunda.

c. Numerous *muscular branches* arise from the outer side of the brachial artery; one of these, the bicipital, more constant than the rest, supplies the biceps; another runs transversely beneath the coraco-brachialis and biceps, over the insertion of the deltoid, supplying this muscle and the brachialis anticus.

Venæ Comites. The two veins which accompany the brachial artery are continuations of the deep radial and ulnar veins. The internal is usually the larger, and generally receives the veins corresponding to the principal branches of the artery. In their course they are connected at intervals by transverse branches, either in front of or behind the artery. Near the subscapularis, the *vena comes externa* crosses obliquely in front of the axillary artery, to join the *vena comes interna*, which then takes the name of axillary.

Now trace the great nerves of the upper arm, which proceed from the brachial plexus near the tendon of the subscapularis; namely, the median, the musculo-cutaneous, the ulnar and the musculo-spiral nerves.

Median Nerve. The *median nerve*, so called from the course it takes along the front of the arm and the forearm, arises by two roots, which converge in front of the axillary artery (p. 341). The external root is derived from the outer cord, in common with the musculo-cutaneous; the internal from the inner cord, in common with the ulnar and internal cutaneous. In its course down the arm, the nerve is situated at first on the outer side of the brachial artery, between it and the coraco-brachialis; about the middle of the arm the nerve crosses obliquely over (in some cases under) the vessel, so that at the bend of the elbow it is found on the inner side of the artery, lying upon

outer and inner heads of the triceps, accompanied by the musculo-spiral nerve, and a little above the middle of the arm divides into two branches, which run for some distance on either side of the nerve. One of these runs in the substance of the triceps muscle, with the nerve to the anconeus, as far as the olecranon and anastomoses with the posterior ulnar recurrent, the inter-osseous recurrent, and anastomotica magna arteries; the other branch accompanies the musculo-spiral nerve to the outer side of the arm, where it perforates the external intermuscular septum. It then descends deep in the interval between the brachialis anticus and supinator radii longus, and terminates in numerous ramifications, some of which pass in front of the external condyle, others behind it, to inosculate with the radial and inter-osseous recurrent arteries.

Before its division, the superior profunda sends several branches to the deltoid, coraco-brachialis, and the triceps, some of which inosculate with the circumflex. These assist in establishing a collateral circulation when the brachial artery is ligatured above the origin of the profunda.

b. The *profunda inferior* arises from the brachial, opposite to the insertion of the coraco-brachialis, or sometimes by a common trunk with the superior profunda. It runs with the ulnar nerve on the inner head of the triceps (which it supplies), passes through the internal intermuscular septum, and then descends to the interval between the internal condyle and the olecranon inosculating with the posterior ulnar recurrent and anastomotica magna arteries. It also sends a small branch down in front of the internal condyle to anastomose with the anterior ulnar recurrent.

c. The *nutrient artery* of the humerus arises sometimes from the brachial, sometimes from the inferior profunda. It pierces the tendon of the coraco-brachialis, runs obliquely downward through the bone, and in the medullary canal divides into ascending and descending branches, which anastomose with the nutrient vessels of the bone derived from the periosteum.

d. The *anastomotica magna* arises from the inner side of the brachial, about two inches above the elbow, runs tortuously

inward, transversely across the brachialis anticus, and divides into branches, some of which pass in front of the internal condyle, anastomosing with the anterior ulnar recurrent artery; another passes behind the internal condyle by piercing the internal intermuscular septum, and anastomoses with the inferior profunda and posterior ulnar recurrent arteries; and one branch forms an arch, above the olecranon fossa, with the superior profunda.

e. Numerous *muscular branches* arise from the outer side of the brachial artery; one of these, the bicipital, more constant than the rest, supplies the biceps; another runs transversely beneath the coraco-brachialis and biceps, over the insertion of the deltoid, supplying this muscle and the brachialis anticus.

Venæ Comites. The two veins which accompany the brachial artery are continuations of the deep radial and ulnar veins. The internal is usually the larger, and generally receives the veins corresponding to the principal branches of the artery. In their course they are connected at intervals by transverse branches, either in front of or behind the artery. Near the subscapularis, the *vena comes externa* crosses obliquely in front of the axillary artery, to join the *vena comes interna*, which then takes the name of axillary.

Now trace the great nerves of the upper arm, which proceed from the brachial plexus near the tendon of the subscapularis; namely, the median, the musculo-cutaneous, the ulnar and the musculo-spiral nerves.

Median Nerve. The *median nerve*, so called from the course it takes along the front of the arm and the forearm, arises by two roots, which converge in front of the axillary artery (p. 341). The external root is derived from the outer cord, in common with the musculo-cutaneous; the internal from the inner cord, in common with the ulnar and internal cutaneous. In its course down the arm, the nerve is situated at first on the outer side of the brachial artery, between it and the coraco-brachialis; about the middle of the arm the nerve crosses obliquely over (in some cases under) the vessel, so that at the bend of the elbow it is found on the inner side of the artery, lying upon

the brachialis anticus and covered by the semilunar fascia the biceps.[1]

As a summary of the distribution of the median nerve, we say that it supplies the two pronators and all the flexors of forearm (except the flexor carpi ulnaris and the ulnar half of flexor profundus digitorum); the muscles of the ball of the th the two radial lumbricales, both sides of the thumb, fore middle fingers, and the radial side of the ring finger, on palmar aspect.

Musculo-Cutaneous Nerve. This nerve (often called the external cutaneous or perforans Casserii) arises in common with external root of the median, from the external cord of brachial plexus behind the pectoralis minor, and is situated the outer side of the axillary artery. It perforates the co brachialis obliquely, and then runs down between the bicep the brachialis anticus to the outer side of the arm. A little the elbow joint, between the tendon of the biceps and the supi

[1] I have observed the following *varieties* relating to the median nerve, course in regard to the artery.

a. The roots may be increased in number by one on either side of the arte the internal root may be deficient.

b. They may vary in their position with regard to the artery; both may be s behind the vessel; or one behind and the other in front of it.

c. The nerve, formed in the usual manner, may be joined lower down by branch from the external cutaneous; such a case presents a junction of two nerves in front of the brachial artery, in the middle of the arm.

d. The nerve in many cases crosses under instead of over the artery.

e. The nerve sometimes runs parallel and external to the artery; or it m parallel to and in front of the artery.

In one hundred arms the relative position of the nerve to the artery in its down the arm was as follows:—

In 72 the nerve took ordinary course.
" 20 the nerve crossed obliquely under the artery.
" 5 the nerve ran parallel and superficial to the artery.
" 3 the nerve ran parallel and external to the artery.

These varieties of the median nerve are of practical importance, for this whenever, in the operation of tying the brachial artery, we do not find the ne its normal position, we may expect to find some irregular distribution of the —*e. g.* a high division of the brachial, or even, which I have often seen, aberrans' coming from the upper part of the brachial, and joining either the ra ulnar arteries.

radii longus, the nerve pierces the deep fascia and becomes subcutaneous; then passing under the median cephalic vein, it divides into an anterior and a posterior branch, for the supply of the integuments of the forearm.

The musculo-cutaneous nerve, in the upper part of its course, sends branches to the coraco-brachialis and the short head of the biceps, and as it descends between the biceps and the brachialis anticus, it supplies both. Consequently, if the nerve were divided in the axilla, the result would be inability to bend the arm.[1] This nerve also sends small filaments to supply the elbow joint.

Ulnar Nerve. This nerve arises from the inner cord of the brachial plexus, in common with the internal cutaneous and the inner head of the median. It descends along the inner side of the brachial artery, as far as the insertion of the coraco-brachialis. The nerve then diverges from the artery, running obliquely over the inner head of the triceps, perforates the internal intermuscular septum, and runs with the inferior profunda artery, behind the internal condyle.

The distribution of the nerve is to the elbow joint, to the flexor carpi ulnaris, to half the flexor profundus digitorum, to all the interosseus muscles of the hand, to both sides of the little finger, to the ulnar side of the ring finger, on their dorsal and palmar aspects, to the muscles of the ball of the little finger, to the wrist joint, to the two ulnar lumbricales, and, lastly, to the adductor pollicis, and the inner head of the flexor brevis pollicis.

Previous to the examination of the musculo-spiral nerve, we should examine the great muscle which occupies the whole of the posterior part of the humerus—viz.: the triceps extensor cubiti.

[1] In some instances the musculo-cutaneous nerve descends on the inner side of the coraco-brachialis without perforating the muscle; in these cases it often sends a larger branch than usual to the median nerve.

The trunk of the musculo-cutaneous nerve may come from the median at any point between the axilla and the middle of the arm. In some subjects the nerve is absent; all its branches are then supplied by the median, which is larger than usual. Such anomalies are easily explained by the fact of the two nerves having a common origin.

Triceps Extensor Cubiti. This muscle has three distinct origins, named, from their positions, the *external*, the *internal*, and the *middle* or *long* heads (p. 337). The *middle* or *long* head *arises* by a flat tendon from the inferior border of the scapula, close to the glenoid cavity, and in connection with the glenoid and capsular ligaments. The *external* head *arises* from the humerus, beginning in a pointed form immediately below the insertion of the teres minor, from the posterior surface between this and the musculo-spiral groove, and from the external intermuscular septum. The *internal* head *arises* from the humerus below the insertion of the teres major, from the posterior surface of the bone below the musculo-spiral groove, and from the internal intermuscular septum. The three heads unite, near the middle of the arm, to form a single fleshy mass, which covers the posterior part of the elbow-joint, and is *inserted* by a thick tendon into the summit and sides of the olecranon. There is a bursa between the tendon and the olecranon, which is sometimes multilocular. Each head is supplied by a separate branch from the musculo-spiral nerve.[1]

Musculo-Spiral Nerve. This, the largest of the brachial nerves, arises, in common with the circumflex, from the posterior cord of the axillary plexus (p. 339). It descends at first behind the axillary artery, and then behind the brachial artery; it subsequently winds obliquely round the posterior part of the humerus, between the external and internal heads of the triceps, in company with the superior profunda artery. About the lower third of the outer side of the arm, the nerve perforates the external intermuscular septum, and then runs deeply embedded between the brachialis anticus and the supinator radii longus.

The nerve gives off branches on the *inner* side of the humerus, to the inner and long heads of the triceps, and the internal cutaneous branch; on the *back* of the humerus, to the external head of the triceps and the anconeus; on the *outer* side of the humerus, to the supinator radii longus, the extensor carpi radialis longior, and the brachialis anticus (usually); lastly, after perforating the

[1] The *subanconeus*, a small muscle situated beneath the triceps, will be described later on.

septum, it gives off the upper and lower external cutaneous branches.

A little above the elbow joint the nerve divides into its two principal branches—the *radial*, which accompanies the radial artery along the forearm, and the *posterior interosseous*, which perforates the supinator brevis, and supplies the muscles on the back of the forearm.

To sum up the muscular distribution of this nerve, we may say that it supplies *all* the extensors of the forearm, wrist, thumb, and fingers; and *all* the supinators except one—namely, the biceps (supplied by the musculo-cutaneous nerve).

DISSECTION OF THE FRONT OF THE FOREARM.

Surface Marking. The front of the forearm presents, at the bend of the elbow, a triangular depression, from which there extends down to the wrist a groove which corresponds to the radial artery; on the inner side is another groove, increasing in depth toward the wrist, indicating the course of the ulnar artery. The head of the radius can be easily felt on the outer side, below the external condyle of the humerus, and in the lower third the bone becomes again defined, terminating below in the styloid process, beyond which is the prominence of the tubercle of the scaphoid. The border of the ulna can be felt on the inner side of the forearm, in the lower half, and it ends at the wrist in an ill-defined styloid process, which does not descend as low as the corresponding process of the radius. The lower part of the forearm presents, an inch beyond the wrist-joint, a transverse furrow, which corresponds with the border of the annular ligament.

Dissection. Prolong the incision down to the wrist, and, at its termination, make another transversely. Reflect the skin, and dissect the subcutaneous veins and nerves.

Cutaneous Veins. On the inner side is the *anterior ulnar vein*, which commences on the front of the wrist, and is then continued upward on the inner side of the forearm as far as the elbow, where it is joined by the posterior ulnar vein to form the common ulnar vein. This vein communicates with the median vein by numerous branches (p. 346).

The veins on the back of the hand commence at the extremities of the fingers, run up *between* the knuckles, and unite on the back of the hand, forming an arch with its concavity upward. The *posterior ulnar vein* arises from this arch by a branch (vena salvatella) situated over the fourth interosseous space, and runs up on the back of the forearm, toward the inner condyle, to join the anterior ulnar vein.

The *radial vein*, situated on the outer side of the forearm, commences on the back of the hand from the venous arch, runs up the radial side of the front of the forearm to the elbow, where, after receiving the median cephalic, it becomes the cephalic vein.

Running up in front of the middle of the forearm is the *median vein;* it communicates in the forearm with the radial and anterior ulnar veins, and near the bend of the elbow it is joined by a deep branch—*mediana profunda*—after which it divides into two branches, an outer or *median cephalic*, which joins the cephalic, and an inner or *median basilic*, which joins the basilic (Fig. 80).

Cutaneous Nerves. On the radial side of the forearm, as low down as the wrist, are found the terminal filaments of the *anterior branch* of the musculo-cutaneous nerve, which, about the middle of the forearm, sends a *posterior branch* backward to supply the posterior and lower part of the forearm as low as the wrist, communicating with the radial and external cutaneous branch of the musculo-spiral. At the lower part of the front of the forearm one or more of these filaments are situated over the radial artery, and one branch passes to the palm, to supply the skin over the muscles of the ball of the thumb; it communicates with the palmar branch of the median and with the radial nerve.

In front of the upper part of the forearm are some filaments of the *external cutaneous branch* of the musculo-spiral nerve; on the outer and back part of the forearm, near the elbow, the lower *external cutaneous branch* of the musculo-spiral runs down as far as the wrist, to supply the skin.

At the lower third of the radial side of the forearm the *radial vein* becomes superficial, and turns over the radius, to supply the back of the hand and fingers.

On the ulnar side the *anterior division* of the internal cutaneous nerve descends as far as the wrist; its *posterior branch* passing to the back of the forearm, to supply it as far as the middle. Near the styloid process of the ulna the *dorsal branch* of the ulnar nerve perforates the fascia, to reach the back of the hand.

Deep Fascia of the Forearm. The muscles of the forearm are enveloped by a dense shining aponeurosis, continuous with that of the arm. Its thickness increases toward the wrist, that the tendons in this situation may be kept in their position. It is composed of fibres which cross each other obliquely, and is attached, above, to the condyles of the humerus and olecranon; internally, to the ridge on the posterior part of the ulna. At the back of the wrist it forms the posterior annular ligament, and in front it is continuous with the anterior annular ligament. Above, the fascia is strengthened by fibres from the tendons of the biceps and brachialis anticus. The aponeurotic expansion from the inner edge of the tendon of the biceps is exceedingly strong. It braces the muscles on the inner side of the forearm, and interlaces at right angles with the fibres of the fascia attached to the internal condyle. The under surface of the fascia gives origin to the muscular fibres in the upper part of the forearm, and furnishes septa which separate the muscles, and form surfaces for their origin. The fascia is perforated at various parts for the passage of the cutaneous vessels and nerves of the forearm.

Dissection. Remove the fascia from the muscles by incisions corresponding to those for reflecting the skin; taking care of the cutaneous branches of the median and ulnar nerves close to the wrist.

Triangle at the Elbow. At the bend of the elbow is a triangular space, with its base toward the humerus; on the inner side this space is bounded by the pronator teres; on the outer, by the supinator radii longus. In it are the following objects which must be carefully dissected:—1. In the centre is the brachial artery (with its companion veins) dividing into the radial as its outer, and into the ulnar as its inner branch; 2, on the outer side of the artery is the tendon of the biceps; 3, on the inner side is the median nerve; 4, the musculo-spiral nerve on the outer side

is partly concealed by the supinator longus; 5, the radial current artery; 6, the anterior ulnar recurrent; 7, the comm interosseous branch of the ulnar artery; 8, the vena media profunda.

FIG. 82.

1. Biceps.
2. Brachialis anticus.
3. Supinator longus.
4. Pronator radii teres.
5. Flexor carpi radialis.
6. Palmaris longus.
7. Flexor carpi ulnaris.
8. Flexor sublimis digitorum.
9. Flexor longus pollicis.
10. Extensor ossis metacarpi the primi internodii polli
11. Triceps.
12. Radial artery.
13. Ulnar artery.

DIAGRAM OF THE MUSCLES OF THE FRONT OF THE FOREARM.

Muscles of the Forearm. The muscles of the forearm are arranged two groups: one, consisting of supinators extensors, is attached to the outer condyloid ridge and condy the other, consisting of pronator and flexors, is attached to inner condyle. The inner group should be examined first. T

r a common tendon, and are arranged in the following pronator teres; flexor carpi radialis; palmaris longus; ublimis digitorum, and flexor carpi ulnaris.

nator Teres. This muscle forms the inner boundary of the triangular space at the elbow. It *arises* by two one, from the anterior surface of the internal condyle, ıe common tendon, from the fascia of the forearm, and ıe septum between it and the flexor carpi radialis; the ıy a small tendinous origin from the inner border of the d process of the ulna. From these two origins, between the median nerve passes, the muscle proceeds obliquely ard and outward across the forearm, and is *inserted* by a don into a rough surface on the outer and back part of the third of the radius. In amputating the forearm, it is very le to save the insertion of this muscle, that the stump may pronator. Its nerve comes from the median.

r Carpi lialis. This muscle, situated on the ulnar side of the preceding muscle, *arises* by the common tendon e internal condyle, from the intermuscular septa, and from cia of the forearm. The fleshy fibres terminate a little he middle of the forearm, in a flat tendon, which runs in a e sheath outside the anterior annular ligament of the wrist, through a groove in the os trapezium, bridged over by tissue and lined by a synovial membrane, and is *inserted* e base of the metacarpal bone of the index finger. The order of its tendon is the guide to the radial artery in the alf of the forearm. Its nerve comes from the median.[1]

maris gus. This slender muscle *arises* from the common tendon at the internal condyle, from the inter- ar septa, and from the fascia of the forearm. About the of the forearm it terminates in a flat tendon, which de- along the middle of the forearm to the wrist, lying upon or sublimis digitorum; it then passes over the anterior

[1] scle is not infrequently found beneath this muscle, called by Mr. Wood the pi radialis brevis, or profundus. It arises from the front of the radius above tor quadratus, and is inserted into the base of the metacarpal bone of the ger. (*Journ. of Anat. and Phys.*, p. 55, Nov. 1866.)

annular ligament, and is continued into the palmar fascia. Thi muscle is a tensor of the palmar fascia.[1] Its nerve comes fro the median.

Flexor Carpi Ulnaris. This muscle *arises* by two heads; one from the internal condyle, the common tendon, and the intermuscular septum; the other from the inner edge of the olecranon; these two origins form an arch, under which the ulnar nerve and the posterior ulnar recurrent artery pass. It also *arises* from the upper two-thirds of the posterior edge of the ulna, through the medium of the aponeurosis, which is common to this muscle, the flexor profundus digitorum, and the extensor carpi ulnaris. The tendon appears on the radial side of the muscle, about the lower third of the forearm, and receives fleshy fibres on its ulnar side as low as the wrist. It is *inserted* into the pisiform bone, and thence by a strong tendon into the unciform and the base of the fifth metacarpal bone. Its nerve comes from the ulnar.

The tendon of the flexor carpi ulnaris is the guide to the ulna artery, which lies close to its radial side, and is *overlapped* by it As it passes over the annular ligament, the tendon furnishes fibrous expansion to protect the ulnar artery and nerve.

Flexor Sublimis Digitorum. This muscle has three distinct origins, and situated beneath those previously mentioned, s that, in order to expose it fully, the preceding muscles should b reflected, by cutting them through the middle, and turning th ends upward and downward. The longer *origin* takes place fro the internal condyle, from the internal lateral ligament, the con mon tendon, and the intermuscular septa; the second *origin* tak place from the coronoid process of the ulna above the pronat teres; the third *origin*, by tendinous and fleshy fibres from t oblique ridge on the front of the radius, extending from t tubercle to about an inch below the insertion of the pronator ter This, called its *radial origin*, is partly concealed by the pronat

[1] The palmaris longus is absent in about one out of ten subjects. The situation its muscular portion is subject to variety; sometimes occupying the middle, sometin the lower third of the forearm. The tendon is in some instances wholly insert into the anterior annular ligament.

teres. The muscle, thus formed, passes down the middle of the forearm, and divides into four distinct muscular slips; from these four tendons arise, which pass beneath the annular ligament, arranged in two pairs; the tendons of the middle and ring fingers being placed over those of the fore and little fingers. The tendons pass through the palm to the fingers, where they diverge and split to allow the passage of the deep flexor tendons, and are *inserted* into the sides of the second phalanges, where they will be subsequently traced. Its *action* is, therefore, to bend the second joint of the fingers.

The muscles described as arising from the internal condyle are all supplied by the median nerve, except the flexor carpi ulnaris, which is supplied by the ulnar.

Having finished the superficial muscles on the inner side of the forearm, notice one of those on the outer side, named supinator radii longus, before tracing the vessels and nerves of the forearm.

Supinator Radii Longus. This muscle forms the external boundary of the triangular space at the bend of the elbow. It *arises* by fleshy fibres from the upper two-thirds of the external condyloid ridge of the humerus, commencing a little below the insertion of the deltoid, and from the external intermuscular septum. The muscular fibres terminate about the middle of the forearm, in a flat tendon, which is *inserted* into the outer side of the base of the styloid process of the radius. The inner border of the muscle is the guide to the radial artery which lies between this muscle and the flexor carpi radialis. It supinates the hand, but acts much more powerfully as a flexor of the forearm. It is supplied by the musculo-spiral nerve.

Radial Artery. The radial artery, the smaller division of the brachial, runs down the radial side of the forearm to the wrist, where it turns over the external lateral ligament of the carpus, beneath the extensor tendons of the thumb, and sinks into the angle between the first and second metacarpal bones to form the deep palmar arch. Thus, its course corresponds with a line drawn from the middle of the bend of the elbow to the front of the styloid process of the radius.

In the upper third of the forearm, the artery lies deep between the pronator teres on the inner and the supinator longus on the outer side; the fleshy border of the latter overlaps it in muscular subjects. In the lower two-thirds of the forearm the artery is more superficial, and is placed between the tendons of the supinator longus on the outer and the flexor carpi radialis on the inner side. In its course, it lies successively on the following: first, upon the tendon of the biceps; secondly, upon the supinator radii brevis; thirdly, upon the insertion of the pronator teres; fourthly, upon the radial origin of the flexor sublimis; fifthly, upon the flexor longus pollicis; sixthly, upon the pronator quadratus, and lastly, upon the lower end of the radius. The artery then turns round the outer side of the wrist-joint, lying upon the external lateral ligament, and covered by the tendons of the extensores ossis metacarpi, and primi internodii pollicis, some cutaneous veins and branches of the radial nerve; next, it lies upon the trapezium; it is then crossed by the extensor secundi internodii pollicis; and lastly, passing between the two heads of the first dorsal interosseous muscle, it enters the palm to form the deep palmar arch. It is accompanied by two veins, which communicate at frequent intervals, and join the venæ comites of the brachial artery at the bend of the elbow.

In the middle third of its course the artery is accompanied by the radial nerve (a branch of the musculo-spiral), which lies to its outer side. Below this point the nerve leaves the artery, and passes under the tendon of the supinator longus, to the back of the hand.

Thus, in the situation where the pulse is usually felt, the radial nerve no longer accompanies the artery; nevertheless, the vessel is accompanied by a branch of the musculo-cutaneous (or external cutaneous), which lies superficially to it.

The radial artery sends off in the forearm the following branches, besides offsets, which supply the muscles on the outside of the forearm—

a. The *radial recurrent* is given off just below the elbow; it ascends up the supinator brevis, between the supinator longus and the brachialis anticus, to supply the long and short supinator

tors and the two radial extensors. It runs up with the musculo-spiral nerve, and forms a delicate inosculation with the superior profunda (p. 353).

b. The *muscular branches* which are given off to the muscles on the outer side of the forearm.

c. The *arteria superficialis volæ* arises from the radial, about half an inch, or more, above the lower end of the radius; it runs over the anterior annular ligament, above, or through, the origin of the muscles of the ball of the thumb, into the palm of the hand, where it sometimes inosculates with the superficial branch of the ulnar, and completes the superficial palmar arch.[1]

d. The *anterior carpal* artery is a small branch of the radial, which arises close to the lower border of the pronator quadratus, and then runs beneath the tendons, and supplies the anterior surface of the synovial membrane and bones of the carpus, anastomosing with the anterior interosseous, the anterior carpal branch of the ulnar, and the recurrent carpal branch of the deep palmar arch.

At the wrist it gives off—

e. The *posterior carpal* artery, which runs beneath the extensor tendons, and joins the corresponding branch of the ulnar to form an arch; it also anastomoses with the anterior interosseous artery on the back of the wrist.

Radial Nerve. The *radial nerve*, a branch of the musculo-spiral, is given off above the bend of the elbow, deep between the supinator radii longus and brachialis anticus; it descends on the outer side of the radial artery, covered by the supinator radii longus. In the *upper* third of the forearm, the nerve is at some distance from the artery; in the *middle* third, it approaches nearer to it, lying to its outer side; but in the *lower* third the nerve leaves the artery, passes underneath the tendon of the supinator longus, perforates the deep fascia on the outer side of the forearm, and becomes subcutaneous. It then divides

[1] There is great variety in the size and origin of the superficialis volæ; sometimes it is very large, arises higher than usual, and runs to the wrist parallel with the radial; sometimes it is very small, terminating in the muscles of the thumb; or it may be absent.

into two branches: an *outer*, the smaller, which supplies the s of the ball of the thumb, and communicates with the ante branch of the musculo-cutaneous nerve; and an *inner*, wh generally supplies both sides of the dorsal aspects of the thu of the index and middle fingers, and of the radial side of ring finger.

Ulnar Artery. This artery, the larger of the two divisions the brachial, comes off below the elbow, r obliquely inward along the ulnar side of the forearm to the wr passes over the annular ligament near the pisiform bone, a entering the palm, forms the superficial palmar arch, by inos lating with the superficialis volæ.

In the upper half of its course the artery describes a ge curve with the concavity toward the radius, and lies deep bene the superficial layer of muscles, namely, the pronator teres, fle carpi radialis, palmaris longus, and flexor sublimis digitorum. is also crossed in its upper part by the median nerve. In lower part of its course it comes nearer the surface, and desce between the flexor sublimis and flexor carpi ulnaris, of which tendon partially overlaps it at the wrist. The artery lies fo short distance on the brachialis anticus; in the remainder of course it lies on the flexor profundus digitorum.

The ulnar nerve is at first separated from the artery by a c siderable interval; about the middle of the forearm it joins artery, and accompanies it in the rest of its course, lying clos its inner side. Both pass over the anterior annular ligament the carpus, lying close to the pisiform bone, the nerve be nearer to the ulnar side and a little behind the artery. A str expansion from the tendon of the flexor carpi ulnaris prot them in this exposed situation.

Observe that the ulnar artery, in the lower third of its cou lies under the radial border of the tendon of the flexor ca ulnaris, which is the *surgical guide* to the vessel. The artery accompanied by two veins, which join the venæ comites of brachial.

The ulnar artery gives off the following branches in the fo arm:—

a. The *anterior and posterior ulnar recurrent arteries* arise immediately below the elbow joint—sometimes by a common trunk. The *anterior* passes upward between the brachialis anticus and the pronator teres, and inosculates with the inferior profunda and anastomotica magna. The *posterior*, the larger, ascends between the flexor sublimis and the flexor profundus digitorum, to the space between the internal condyle and the olecranon; it then passes up between the two heads of the flexor carpi ulnaris, where it inosculates with the inferior profunda, the anastomotica magna, and, above the olecranon, with the posterior interosseous recurrent (p. 354).

b. The *common interosseous* artery is about half an inch long. It arises from the ulnar, just below the tubercle of the radius, and soon divides into the anterior and posterior interosseous, which we shall examine presently.

c. The *muscular branches* which supply the muscles on the ulnar side of the forearm.

d. The *carpal branches* are given off just above the pisiform bone; the *posterior carpal* runs beneath the tendon of the flexor carpi ulnaris and the extensor tendons, and forms, with the corresponding branch of the radial artery, an arch, from which are usually given off the second and third dorsal interosseous arteries; these anastomose with the perforating arteries. The *anterior carpal* runs in front of the carpus, beneath the flexor tendons, supplies the synovial membrane and the ligaments, and anastomoses with the anterior carpal from the radial.

Ulnar Nerve. This nerve runs behind the internal condyle, between two origins of the flexor carpi ulnaris. In its course down the ulnar side of the upper part of the forearm, the nerve is still covered by this muscle, and lies upon the flexor profundus digitorum. About the middle of the forearm, the nerve joins the ulnar artery, and runs along its inner side over the anterior annular ligament into the palm.

The ulnar nerve gives off the following branches:—

a. The *articular branches* to the joint are given off to it, immediately behind the elbow.

b. The *muscular branches* are distributed to the flexor carpi

ulnaris and the ulnar half of the flexor profundus digitorum, ar are given off from the ulnar a short distance below the elbow.

c. A *cutaneous branch* is given off about the middle of the fore arm, one filament of which, called the *palmar cutaneous branch* accompanies the ulnar artery to the palm, and communicates with branches from the median nerve.

d. The *dorsal cutaneous branch*, of considerable size, is given off from the ulnar about two inches below the styloid process of the ulna, to pass to the back of the hand. It crosses under the tendon of the flexor carpi ulnaris, pierces the deep fascia, and immediately below the styloid process of the ulna, appears on the back of the hand, where it divides into branches which supply the back of the little finger, and half the ring; here also it sends a branch which communicates with the corresponding branch of the radial nerve.

e. Articular branches are also distributed to the wrist joint.

Median Nerve This nerve, at the bend of the elbow, lies on the inner side of the brachial artery and beneath the bicipital fascia. It then passes between the two heads of origin of the pronator teres, and descends along the middle of the forearm, between the flexor sublimis and the flexor profundus digitorum. At the lower part of the forearm, it becomes more superficial, lying above the wrist between the outer tendon of the flexor sublimis and the inner border of the tendon of the flexor carpi radialis; beneath, or to the ulnar side of the palmaris longus, and having in front of it the skin and deep fascia; it then enters the palm beneath the anterior annular ligament, and divides into five branches for the supply of the thumb, both sides of the fore and middle fingers, and the radial side of the ring finger.

Immediately below the elbow, the median nerve sends off:

a. The *muscular branches* to the pronator teres, and to all the flexor muscles of the forearm, except the flexor carpi ulnaris and the ulnar half of the flexor profundus, which are supplied by the ulnar nerve.

b. The *anterior interosseous nerve*, also a branch of the median, runs with the anterior interosseous artery on the interosseous

membrane, lying on its radial side, between the flexor longus pollicis and flexor profundus digitorum; it supplies both these muscles and the pronator quadratus.

c. The *palmar cutaneous branch* is given off from the median before it passes beneath the annular ligament. This branch passes over the ligament, and divides into numerous filaments to supply the skin of the palm and the ball of the thumb, communicating with the cutaneous palmar branches of the ulnar, the external cutaneous, and the radial nerves.

Dissection. Now reflect the superficial layer of muscles, to see those more deeply seated. Preserve the principal vessels and nerves.

The deep-seated muscles are, on the ulnar side, the flexor digitorum profundus; and on the radial side, the flexor longus pollicis; beneath both, near the wrist, lies a transverse muscle, the pronator quadratus. On the interosseous membrane, between the first two named muscles, run the anterior interosseous artery and nerve.

Flexor Profundus Digitorum. This is the thickest muscle of the forearm. It *arises* from the upper two-thirds of the anterior surface of the ulna, surrounding the insertion of the brachialis anticus above, from the same extent of its internal surface, from the aponeurosis attached to the posterior edge of the ulna, and from the ulnar two-thirds of the interosseous membrane. About the middle of the forearm the muscle is *inserted* into four flat tendons, of which only that which goes to the index finger is separate from the others above the wrist. These tendons lie upon the same plane, and pass beneath the annular ligament, under those of the superficial flexor, into the palm, where they diverge to pass to their respective fingers. On the first phalanx of the fingers, the tendons of the deep flexor perforate those of the superficial, and are *inserted* into the bases of the third or ungual phalanges. It derives its nerves from the interosseous branch of the median and from the ulnar.

Flexor Longus Pollicis. This muscle is situated on the front surface of the radius, outside the preceding. It *arises* from the front surface of the radius, between the tubercle and the oblique ridge above, and the pronator quadratus below, and from the inter-

osseous membrane.[1] Its tendon, which begins on the ulnar of the muscle, proceeds beneath the annular ligament to the of the last phalanx of the thumb. Its nerve comes from the i osseous branch of the median.

Pronator Quadratus. This square muscle *arises* from the lower f of the ulna and from a strong aponeurosis w covers its anterior surface; its fibres pass, some transversely, obliquely outward, and are *inserted* into the lower fourth o anterior surface and the outer border of the radius. It pro the radius on the ulna. Its nerve proceeds from the interos branch of the median.

Anterior Interosseous Artery. Nearly on a level with the insertion o biceps, the ulnar artery gives off from its side the *common interosseous*, which runs ward for about an inch, and divides into the *anterior* and *pos interosseous*.

The *anterior interosseous artery* runs down on the interos membrane, lying deeply between the flexor profundus digit and flexor longus pollicis. At the upper edge of the pro quadratus it divides into two branches; one of which, the sm passes beneath the muscle, supplies it and the front of the c bones, communicating with the anterior carpal arteries fron radial and ulnar; the other, the more important, perforate interosseous membrane, and helps to supply the muscles o back of the forearm.

A branch, the *arteria comes nervi mediani*, proceeds fron anterior interosseous. It lies in close contact with the sometimes in its very centre; though usually of small size, it be as large as the ulnar artery itself, and, in such cases, it p under the annular ligament with the nerve to join the palmar This is interesting, because it helps to explain the recurren hemorrhage from a wound in the palm, even after the radia ulnar arteries have been tied.

The anterior interosseous artery gives off branches t muscles on each side; also the *nutrient arteries* which ente radius and ulna, near the centre of the forearm, to supply

[1] Sometimes by a slip from the coronoid process.

medullary membrane; these arteries pass upward toward the elbow.

Anterior Interosseous Nerve. This nerve is a branch of the median; it generally runs close to the radial side of the artery, and supplies the flexor longus pollicis, half the flexor profundus digitorum, and the pronator quadratus.

DISSECTION OF THE PALM OF THE HAND.

Surface Marking. On the ulnar side of the palm of the hand is a round, long eminence, *hypothenar*, which corresponds with the muscles of the ball of the little finger; and on the radial side, placed obliquely over the metacarpal bone of the thumb, is another eminence, *thenar*, which is caused by the muscles of the ball of the thumb. Between the two eminences, at the wrist, is a slight depression, corresponding with the middle of the annular ligament, and which broadens out toward the fingers. The palm of the hand, about an inch from the clefts of the fingers, presents a transverse furrow, which corresponds with the metacarpo-phalangeal articulations, with the distal limit of the synovial sheaths of the flexor tendons, with the divisions of the palmar fascia into its four processes, and with the transverse metacarpal ligament. The superficial palmar arch may be indicated by a line drawn from the cleft of the extended thumb across the palm; the deep palmar arch lies half an inch nearer the annular ligament.

Dissection. Make a vertical incision along the centre of the palm, and a transverse one along the bases of the fingers; from this transverse cut continue vertical incisions along the front of the fingers, and reflect the skin; taking care not to remove a small cutaneous muscle—the palmar brevis—situated over the ball of the little finger, and also two small cutaneous branches of the median and ulnar nerves, which are found in the fat of the palm.

Observe how closely, in the centre of the palm, the skin adheres to the palmar fascia beneath it. On the ball of the little finger and the distal ends of the metacarpal bones, the subcutaneous structure is composed of a dense filamentous tissue, which

contains numerous pellets of fat, forming an elastic pad. A similar padding protects the palmar surfaces of the fingers. These cushions on the ends of the fingers defend them in the powerful actions of the hand; they are also useful in subservience to the nerves of touch.

The palm is supplied with nerves by three small branches: the *palmar branch* of the median passes in front of the anterior annular ligament to the centre of the palm; the *palmar branch* of the ulnar supplies the inner aspect of the hand; and the *anterior branch* of the musculo-cutaneous nerve is distributed to the skin over the thenar eminence. The terminal branches of these cutaneous nerves communicate with each other.

Palmaris Brevis. This small cutaneous muscle is situated on the inner side of the palm. It *arises* from the inner edge of the central palmar fascia and the annular ligament, and *inserted* into the skin on the ulnar border of the palm. Its use to support the pad on the inner edge of the palm; it acts powerfully as we grasp; it raises the inner edge of the palm, and deepens the hollow of it, forming the so-called "cup of Diogenes." It is supplied by the ulnar nerve.

Palmar Fascia. This fascia has a silvery lustre, and, in the centre of the palm, is remarkably dense and strong. It is divided into three portions; a *central*, by far the strongest; an *external*, covering the muscles of the thumb; and an *internal*, covering the muscles of the little finger. From the deep surface of the fascia two septa dip down, and divide the palm into three separate compartments; one for the ball of the thumb, a second for that of the little finger, and a third for the centre of the palm.

The fascia is formed by a prolongation from the anterior annular ligament. It is also strengthened by the expanded tendon of the palmaris longus.

The central portion of the fascia is triangular, with the apex at the wrist. About the middle of the palm it splits into four portions, which are connected by transverse tendinous fibres extending completely across the palm, and corresponding pretty nearly to the transverse furrow of the skin in this situation.

Examine any one of these four portions of the fascia, and you will find that it splits into two strips which embrace the corresponding flexor tendons, and are intimately connected with the transverse metacarpal ligament. The effect of this is that the flexor tendons of each finger are kept in place in the palm by a fibrous ring. Between the four divisions of the palmar fascia the digital vessels and nerves emerge, and descend in a line with the clefts between the fingers.

In the hands of mechanics, in whom the palmar fascia is usually very strong, we find that slips of it are lost in the skin at the lower part of the palm, and also for a short distance along the sides of the fingers.

The chief use of the palmar fascia is, to protect the vessels and nerves from pressure, when anything is grasped in the hand. It also confines the flexor tendons in their proper place.

Beneath the interdigital folds of the skin, there are aponeurotic fibres to strengthen them, constituting what are called the *transverse ligaments* of the fingers. They form a continuous ligament across the lower part of the palm, in front of the digital vessels and nerves.

Dissection. Cut through the palmar fascia at its attachment to the anterior annular ligament, and reflect it toward the fingers, so as to expose the vessels, nerves and tendons in the palm. The vessels lie above the nerves, and the tendons still deeper. There is an abundance of loose connective tissue to allow the free play of the tendons. When suppuration takes place in the palm, it is seated in this tissue. Reflect for a moment what mischief is likely to ensue. The pus cannot come to the surface through the dense palmar fascia, or on the back of the hand; it will, therefore, run up into the carpal bursa under the annular ligament, and make its way deep among the tendons of the forearm.

Superficial Palmar Arch. The ulnar artery, having passed over the annular ligament, near the pisiform bone, describes a curve across the upper part of the palm, beneath the palmar fascia, toward the thumb, and, gradually diminishing in size, inosculates with the superficialis volæ, and very commonly with a branch from

the arteria radialis indicis, to form the *superficial palmar a* The curve of the arch is directed toward the fingers, its great convexity descending as low as a horizontal line drawn acr the junction of the upper with the middle third of the palm.

FIG. 83.



DIAGRAM OF THE SUPERFICIAL AND DEEP PALMAR ARCHES.
1, 2, 3, 4, Interosseous branches.

In its passage over the annular ligament the artery lies in furrows, between the pisiform and unciform bones, and is prote by an expansion from the tendon of the flexor carpi ulnaris tc palmaris longus. The ulnar nerve lies close to the inner sic the artery, both being covered by the palmaris brevis. In

palm, the artery rests for a short distance upon the muscles of the little finger, then it lies upon the superficial flexor tendons and the divisions of the ulnar and median nerves, and is covered by the palmar fascia.

Immediately below the pisiform bone the ulnar artery gives off the *ulnaris profunda*, which sinks deeply into the palm, between the origins of the abductor and flexor brevis minimi digiti, to form the deep palmar arch, by joining the terminal branch of the radial artery. It is accompanied by the deep branch of the ulnar nerve.

From the concavity of the arch small *recurrent branches* ascend to the carpus, and inosculate with the other carpal branches of the radial and ulnar arteries.

Four digital arteries arise from the convexity of the arch. They supply all the digits, except the thumb and the radial side of the index finger. The *first* descends over the muscles on the inner side of the palm, to the ulnar side of the little finger, along which it runs to the apex. The *second*, *third* and *fourth* descend nearly vertically between the tendons, *in a line with the clefts* between the fingers, and, about half an inch above the clefts, each divides into two branches, which proceed along the opposite sides of the fingers nearly to the end of the last phalanges, where they unite to form an arch with the convexity toward the end of the finger; from this arch numerous branches supply the papillæ at the tip of the finger.

In the palm of the hand the digital arteries, before they divide, are joined by branches from the corresponding palmar interosseous arteries (branches of the deep palmar arch) (Fig. 83).

The digital arteries freely communicate, on the palmar and dorsal aspect of the fingers, by transverse branches, which supply the joints and the sheaths of the tendons. Near the ungual phalanx a considerable branch passes to the back of the finger, and forms a network of vessels which supply the matrix of the nail.

Ulnar Nerve in the Palm. The *ulnar nerve* passes over the annular ligament into the palm, on the inner side of the ulnar artery, and a little behind it. It lies in the groove between the

pisiform and unciform bones, so that it is perfectly secure from pressure. Immediately below the pisiform bone the nerve divides into a superficial and a deep palmar branch. The *deep* branch supplies the muscle forming the ball of the little finger, and accompanies the ulnaris profunda artery into the palm, to supply *all* the interosseous muscles, the two ulnar lumbricales, and it ends in branches which are distributed to the first dorsal interosseous, the adductor pollicis, and the inner head of the flexor brevis pollicis; it moreover sends filaments which ascend to supply the wrist joint, and others which descend to the metacarpo-phalangeal joints. The *superficial branch* sends filaments to the palmaris brevis, to the skin on the inner side of the palm, and then divides into two digital nerves, one for the supply of the ulnar side of the little finger, the other for the contiguous sides of the little and ring fingers. This branch also communicates with the median nerve behind the superficial palmar arch. All the digital branches run along the sides of the fingers to their extremities, superficial to their corresponding arteries.

Anterior Annular Ligament of the Carpus.

This exceedingly strong and thick ligament confines the flexor tendons of the fingers and thumb, and fastens together the bones of the carpus. It is attached, externally, to the tuberosity of the scaphoid and the ridge on the trapezium; internally, to the pisiform and unciform. Its upper border is continuous with the aponeurosis in front of the wrist; its lower is connected with the palmar fascia; its anterior surface receives the expanded tendon of the palmaris longus, and gives origin to most of the muscles of the ball of the thumb and little finger.

Now proceed to the muscles composing the ball of the thumb and the little finger. The dissection of them requires considerable care.

Muscles of the Ball of the Thumb.

The great strength of the muscles of the ball of the thumb is one of the distinguishing features of the human hand. This strength is necessary in order to oppose that of all the fingers. In addition to its strength, the thumb enjoys perfect mobility. It has no less than

ht muscles—namely, an abductor, an opponens, two flexors, ree extensors, and an adductor.

Abductor Pollicis. This is the most superficial. It is a thin, flat muscle, and *arises* from the ridge of the os trapezium and the annular ligament. It passes forward and outward, and is *inserted* by a flat tendon into the radial side of the base of the first phalanx of the thumb. Its *action* is to draw the thumb away from the fingers. Its nerve comes from the median. Reflect it from its insertion to expose the following:—

Opponens Pollicis. This muscle *arises* from the front of the os trapezium beneath the abductor, and from the annular ligament, and, passing forward and outward, is *inserted* into the whole length of the radial side of the metacarpal bone of the thumb. The *action* of this powerful muscle is to oppose the thumb to all the fingers. Its nerve comes from the median. Reflect it from its insertion, to expose the following:—

Flexor Brevis Pollicis. This muscle has two origins, between which runs the tendon of the flexor longus pollicis; one, the superficial, from the annular ligament and trapezium; the other, the deep, from the trapezoid, os magnum, the bases of the second and third metacarpal bones, and the sheath of the tendon of the flexor carpi radialis. It is *inserted* by two strong tendons into the base of the first phalanx of the thumb; the superficial tendon being connected with the abductor pollicis, and the deep one with the adductor pollicis. A sesamoid bone is found in each of the tendons. The tendons of insertion of this muscle are separated by the long flexor tendon of the thumb and the arteria magna pollicis. Its *action* is to bend the first phalanx of the thumb. The superficial portion is supplied by the median nerve; the deep, by the ulnar.

Adductor Pollicis. This triangular muscle *arises* from the palmar aspect of the shaft of the metacarpal bone of the middle finger; its fibres converge and are *inserted*, along with the deep or inner portion of the flexor brevis pollicis, into the base of the first phalanx of the thumb and the internal sesamoid bone. Its *action* is to draw the thumb toward the palm, as when we bring the tips of the thumb and little finger into contact. It

is supplied by the deep branch of the ulnar nerve, which also supplies the deep head of the flexor brevis pollicis. The other muscles of the ball of the thumb are supplied by the median nerve.

Muscles of the Ball of the Little Finger. The muscles of the little finger correspond in some measure with those of the thumb. Thus there is an abductor, a flexor brevis, and an opponens minimi digiti. All derive their nerves from the deep branch of the ulnar.

Abductor Minimi Digiti. This, the most superficial of the muscles of the little finger, *arises* from the pisiform bone, and from the tendinous expansion of the flexor carpi ulnaris; it is *inserted* by a flat tendon into the inner side of the base of the first phalanx of the little finger. Its *action* is to draw this finger from the others. Its nerve comes from the deep branch of the ulnar.

Flexor Brevis Minimi Digiti. This slender muscle may be considered as a portion of the preceding, to the radial side of which it is situated. It *arises* from the apex of the unciform bone and annular ligament, and is *inserted* with the tendon of the abductor into the base of the first phalanx of the little finger. Its *action* is similar to that of the abductor. Nerve from deep branch of ulnar. Between the origins of the abductor and flexor brevis minimi digiti, the deep branch of the ulnar artery and nerve sink down to form the deep palmar arch.

Opponens Minimi Digiti. The last two muscles must be reflected from their insertion, to expose the *opponens minimi digiti.* It *arises* from the unciform process and the annular ligament, and is *inserted* along the ulnar side of the shaft of the metacarpal bone of the little finger. Its *action* is to draw this bone, the most movable of all the metacarpal bones of the fingers, toward the thumb. Thus it greatly strengthens the grasp of the palm. Nerve from deep branch of ulnar.

Dissection. Cut vertically through the anterior annular ligament, and observe that, with the carpal bones, it forms an elliptical canal, with the broad diameter transversely. This canal is lined by a synovial membrane which is reflected

loosely over the tendons. Superficial to the ligament pass the palmaris longus, the ulnar artery and nerve, the fibrous expansion from the flexor carpi ulnaris covering these vessels and nerve, and the palmar branch of the median and ulnar nerves; beneath it pass the superficial and deep flexor tendons of the fingers, the long flexor tendon of the thumb, and the median nerve. The tendon of the flexor carpi radialis does not run with the other tendons, but is contained in a distinct sheath, lined by a separate synovial membrane, formed, partly by the annular ligament, and partly by the groove in the trapezium.

Median Nerve in the Palm. In its passage under the annular ligament, the median nerve is enveloped in a fold of synovial membrane, and lies upon the flexor tendons. Here it becomes enlarged and flattened, and of a pinkish color, and divides into two nearly equal parts; the *external* gives a recurrent branch to the muscles of the ball of the thumb—namely, to the abductor pollicis, the opponens pollicis, and the outer head of the flexor brevis pollicis, and then terminates in three digital nerves, two of which are distributed to the thumb, and the third to the radial side of the index finger; the *internal* gives digital branches which supply the ulnar side of the forefinger, both sides of the middle finger, and the radial side of the ring finger.

The *two* nerves to the thumb proceed, one on each side of the long flexor tendon, to the last phalanx; the outer one being connected with a terminal filament of the radial.

The *third* digital nerve runs along the radial side of the index finger. The *fourth* descends toward the cleft between the index and middle fingers, and subdivides into two branches, which supply their opposite sides. The *fifth* is joined by a filament from one of the ulnar digital nerves, and then subdivides above the cleft between the middle and ring fingers, to supply their opposite sides.

Two small branches are given off from the third and fourth digital nerves, to supply the two radial lumbricales; the two ulnar being supplied by the ulnar nerve.

About an inch and a quarter above the clefts between the fingers, each digital nerve subdivides into two branches, between which

the digital artery passes and bifurcates lower down; therefore a vertical incision down the cleft would divide the artery before the nerve.

In their course along the fingers and thumb, the nerves lie superficial to the arteries, and nearer to the flexor tendons. About the base of the first phalanx each nerve sends a *dorsal branch*, which runs along the back of the finger nearly to the extremity, communicating with the dorsal branches derived from the radial and ulnar nerves.[1] Near the ungual phalanx another dorsal or ungual branch is distributed to the skin around and beneath the matrix of the nail. Each digital nerve terminates in the cushion at the end of the finger, in a brush of filaments, with their points directed into the papillæ of the skin.

Flexor Tendons and their Sheaths.

Immediately below the annular ligament the tendons separate from each other; near the metacarpal joints they pass in pairs, through strong fibrous rings (p. 375) formed by the divisions of the palmar fascia. Below the metacarpal joint the two tendons for each finger enter the sheath, *theca*, which confines them in their course along the phalanges. It is formed by a strong fibrous membrane, which is attached to the ridges on the phalanges, and converts the groove in front of these bones into a complete canal, exactly large enough to contain the tendons. The density of the sheath varies in particular situations, otherwise there would be an obstacle to the easy flexion of the fingers. To ascertain this, cut open one of the sheaths along its entire length; you will then see that it is much

[1] Upon the cutaneous nerves of the hands and feet are little bodies, termed, after their discoverer, corpuscles of Pacini. Some of them will be found, by carefully examining the trunk of a nerve, or one of its smaller branches, in the subcutaneous tissue at the root of a finger. Each corpuscle is about $\frac{1}{12}$ of an inch long, and is attached by a slender fibro-cellular pedicle to the nerve upon which it is situated; through the pedicle, a single primitive nerve-fibril passes into the corpuscle. The corpuscle itself is composed of a series of concentric capsules, varying from twenty to fifty in number, and separated by intervals containing fluid; and the nerve-fibril terminates by a dilated extremity in a central cavity, which exists in the axis of the corpuscle. Their function is unknown. These bodies are found in many other situations, viz., in the solar plexus, the pudic nerves, the intercostal nerves, the cutaneous nerves of the arm and neck, the infra-orbital nerve, the sacral plexus, and in nerves supplying the periosteum. They can be best examined in the mesentery of the cat.

stronger between the joints than over the joints themselves. Through these sheaths inflammation, commencing in the integuments of the finger, may readily extend to the synovial membrane of the tendon.

In cases of whitlow, when pus forms in the theca, the incision should be made deep enough to lay open this fibro-osseous canal, without which the incision will be of no use. It is obvious that the incision should be made down the *centre* of the finger, to avoid the digital nerves and arteries. If this opening be not timely made, the flexor tendons are likely to slough, and the finger becomes stiff.

But what protects the joints of the fingers where the flexor tendons play over them? Look into an open sheath, and you will see that in front of the joints the tendons glide over a smooth fibro-cartilaginous structure called the *palmar* ligament.

To facilitate the play of the tendons, the interior of the sheath, as well as the tendons, is lined by a synovial membrane, of the extent of which it is important to have a correct knowledge. With a probe you may ascertain that the synovial membrane is reflected from the sheath upon the tendons, a little above the metacarpal joints of the fingers—that is, nearly in a line with the transverse fold in the skin in the lower third of the palm. Toward the distal end of the finger, the synovial sheath stops short of the last joint, so that it is not injured in amputation of the ungual phalanx.

And now notice how the tendons are adapted to each other in their course along the finger. The superficial flexor, near the root of the finger, becomes slightly grooved to receive the deep flexor; about the middle of the first phalanx it splits into two

[1] On closer inspection it will be observed that the sheath is composed of bands of fibres, which take different directions, and have received distinct names. The strongest are called the *ligamenta vaginalia*. They constitute the sheath over the body of the phalanx, and extend transversely from one side of the bone to the other. The *ligamenta cruciata* are two slips, which cross obliquely over the tendons. The *ligamenta annularia* are situated immediately in front of the joints, and may be considered as thin continuations of the ligamenta vaginalia. They consist of fibres, which are attached on either side to the lateral ligaments of the joints, and pass transversely over the tendons.

portions, through which the deep flexor passes. The two portions reunite below the deep tendon so as to embrace it, and then divide a second time into two slips, which interlace with each other, and are inserted into the sides of the second phalanx. The *deep* flexor, having passed through the opening of the superficial one, is inserted into the base of the last phalanx.[1]

In what way are the tendons supplied with blood? Raise and separate the tendons, and you will see that slender but very vascular folds of synovial membrane (*vincula tendinum*) run up from the phalanges and convey blood vessels to the tendons.

The *tendon of the flexor longus pollicis* lies on the radial side of the other tendons beneath the annular ligament. It passes between the two portions of the flexor brevis pollicis and the two sesamoid bones of the thumb, enters its proper sheath, and is *inserted* into the base of the last phalanx. Its synovial sheath is prolonged from the large bursa of the flexor tendons beneath the annular ligament, and accompanies the tendon down to the last joint of the thumb; consequently the sheath is injured in amputation of the last phalanx.

Bursal Sac of the Carpus. A large and loose synovial sac (bursa of the carpus) facilitates the play of the tendons beneath the anterior annular ligament. It lines the under surface of the ligament and the groove of the carpus, and is reflected in loose folds over the tendons. It is prolonged up the tendons for an inch and a half, or two inches, and forms a cul-de-sac above the ligament. Below the ligament the bursa extends into the palm, and sends off prolongations for each of the flexor tendons, which accompany them down to the middle of the hand. You will understand that when the bursa is inflamed and distended by fluid, there will be a bulging above the annular ligament, and another in the palm, with perceptible

[1] In the Museum of the College of Surgeons a preparation is put up which shows a beautiful piece of animal mechanics concerning the flexor tendons; namely, that in its passage along the phalanges, the deep flexor forms, at the first phalanx, a kind of little patella for the superficial one; but, at the second phalanx, the superficial flexor lies deeper than the other, and forms a little patella for it. This increases the leverage in each case.

fluctuation between them; the unyielding ligament causing a constriction in the centre.[1]

Lumbricales. These four slender muscles, one for each finger, are attached to the deep flexor tendons in the palm. All of them *arise* by fleshy fibres from the radial side and palmar surface of the deep tendon of their corresponding finger; the third and fourth also arise from the adjacent sides of two tendons. Each terminates in a broad, thin tendon which passes over the radial side of the first joint of the finger, and is *inserted*, by a broad expanded aponeurosis, into the extensor tendon on the dorsal aspect of the first phalanx of the finger. Their *action* is to bend the metacarpo-phalangeal joint of the fingers. Being inserted near the centre of motion, they can move the fingers with great rapidity. As they produce the quick motions of the musician's fingers, they were called by the old anatomists *fidicinales*.

The two ulnar lumbricales are supplied by the deep branch of the ulnar nerve; the two radial by the third and fourth digital branches of the median nerve.

Dissection. Now cut through all the flexor tendons, and remove the deep fascia of the palm, to see the deep arch of arteries and its branches.[2]

Branches of the Radial Artery in the Palm. The radial artery, sinking into the space between the first and second metacarpal bones, and between the two heads of the abductor indicis, enters the palm between the inner head of the flexor brevis and the adductor pollicis, and gives off three branches—the arteria princeps pollicis, the radialis indicis, and the palmaris profunda, which unites with the deep ulnar artery to form the deep arch.

The *arteria princeps pollicis* runs behind the deep head of the

[1] In only one subject have we seen an instance in which this bursa communicated with the wrist joint. It communicates always with the synovial sheath of the long flexor of the thumb, in most cases with that of the flexors of the little finger, and but rarely with that of the index, middle and ring fingers. For this reason, inflammation of the theca of the thumb or little finger is more liable to be attended with serious consequences than either of the others.

[2] The course and relations of the radial artery as it winds round the wrist will be described in the dissection of the back of the hand.

flexor brevis pollicis and in front of the abductor indicis (*first* dorsal interosseus), close along the metacarpal bone of the thumb; in the interval between the lower portions of the flexor brevis pollicis the artery divides into two digital branches, which proceed one on either side of the thumb, and inosculate at the apex of the last phalanx. Their distribution and mode of termination are like those of the other digital arteries.

The *arteria radialis indicis* runs between the abductor indicis and adductor pollicis, along the radial side of the index finger to the end, where it forms an arch with the other digital artery, a branch of the ulnar. Near the lower margin of the adductor pollicis, the radialis indicis generally receives a branch from the princeps pollicis, and gives a branch to the superficial palmar arch.

The *palmaris profunda* may be considered as the continuation of the radial artery. It enters the palm between the inner head of the flexor brevis and the adductor pollicis, and, running upon the bases of the metacarpal bones, inosculates with the deep branch of the ulnar artery, thus completing the *deep palmar arch*. From the concavity of the arch small recurrent branches ascend to supply the bones and joints of the carpus, inosculating with the other carpal arteries.

From the convexity of the arch three or four small branches, called *palmar interosseous* (Fig. 83, p. 376), descend to supply the interosseous muscles, and near the clefts of the fingers communicate with the digital arteries. These palmar interosseous branches are sometimes of considerable size, and take the place of one or more of the digital arteries ordinarily derived from the superficial palmar arch. Three branches, called *perforating*, pass between the upper ends of the metacarpal bones to the back of the hand, and communicate with the carpal branches of the radial and ulnar.

Deep Branch of the Ulnar Nerve. This nerve sinks into the palm with the ulnar profunda artery, between the abductor and flexor brevis minimi digiti. It then runs with the deep palmar arch toward the radial side of the palm, and terminates in the adductor pollicis, in the inner or deep head of the flexor brevis pollicis,

and in the first dorsal interosseous. Between the pisiform and unciform bones, the nerve gives a branch to each of the muscles of the little finger. Subsequently it sends branches to each interosseous muscle and to the two inner lumbricales.

The tendon of the flexor carpi radialis in the palm must now be followed to its insertion into the base of the second metacarpal bone.

The dissection of the remaining muscles of the palm, called, from their position, *interossei*, must be, for the present, postponed.

MUSCLES OF THE BACK CONNECTED WITH THE ARM.

Dissection. Make an incision down the middle of the spine from the occiput to the sacrum; another, from the last dorsal vertebra upward and outward to the acromion; and a third, from the sacrum along the crest of the ilium; then reflect the skin outward from the dense subcutaneous tissue, in which will be found the following cutaneous nerves.

Cutaneous Nerves of the Back. These are derived from the posterior divisions of the spinal nerves, and correspond, generally, to the number of the vertebræ. The posterior primary branches, much smaller than the anterior, divide, between the transverse processes, into *external* and *internal* branches, with the exception of the suboccipital, the fourth and fifth sacral, and the coccygeal nerves.

Posterior Branches of the Cervical Nerves. The *posterior primary branches of the cervical nerves* (except the first[1]) divide into external and internal branches; the *external* are distributed solely to some of the muscles of the neck, and which will be dissected later on; the *internal*, larger than the external, are distributed in the following manner; the *second*, or the *great occipital nerve*, perforates the complexus, and ramifies on the back of the scalp with the occipital artery; the *third*, *fourth* and *fifth* nerves, after sending branches to the multifidus spinæ, semi-spinalis and the complexus, emerge through the trapezius close to the spinous processes, and there pass transversely across that muscle to supply

[1] This nerve has already been described in the dissection of the suboccipital triangle (p. 309).

the skin over it; the branch of the *third cervical nerve* sometimes sends a branch to the back of the scalp;[1] the branches of the *sixth*, *seventh* and *eighth* are small, and are situated beneath the semi-spinalis, to which they are distributed.

Posterior Branches of the Dorsal Nerves. The *external branches* become superficial between the longissimus dorsi and the ilio-costalis, and supply these muscles and the other divisions of the erector spinæ; the six lower supply cutaneous nerves in the line of the angles of the ribs. The *internal branches*, as to the upper six dorsal, emerge between the multifidus spinæ and semi-spinalis, and passing horizontally outward, end in branches to the skin close to the spinous processes; that from the second ramifies over the spine of the scapula; the six lower do not become cutaneous, but terminate in the multifidus spinæ.

Posterior Branches of the Lumbar Nerves. The *external branches* from the first, second and third lumbar nerves perforate the ilio-costalis and the latissimus dorsi, and then descend over the crest of the ilium, supplying cutaneous branches to the gluteal region; the fourth supplies the erector spinæ without becoming cutaneous; the fifth sends down a branch to communicate with the first sacral nerve. The *internal branches* are small, and end in the multifidus spinæ.

Posterior Branches of the Sacral Nerves. The *external branches* of the upper three sacral nerves form a series of loops with themselves, and also with the last lumbar above and the fourth sacral below; they pass to the superficial surface of the great sacro-sciatic ligament, where they form another series of loops, from which filaments are distributed to the skin after piercing the gluteus maximus. The *internal branches* of the three upper sacral nerves are distributed to the multifidus spinæ. The posterior branches of the fourth and fifth sacral nerves do not divide into external and internal branches, but form a loop, the lower one being joined with the coccygeal nerve.

[2] The internal branches of the first, second and third cervical nerves form a communication beneath the complexus, which is called by Cruveilhier the *posterior cervical plexus*.

FIG. 84.

, *a*, Small occipital
ve from the cervical
rm; 1, external mus-
ar branches of the first
vical nerve and union
a loop with the second;
ar rectus capitis posti-
major, with the great
pital nerve passing
nd the short muscles
piercing the com-
m; the external
ch is seen to the out-
: 2′, the great occi-
: 3, external branch
e posterior primary
ion of the third
: 3′, its internal
Ba, or third occipital
: 4′, 5′, 6′, 7′, 8′,
nal branches of the
al corresponding
es on the left side;
external branches of
nerves proceeding
scles are displayed
e right side: *d* 2 to *d*
d thence to *d* 12, ex-
l muscular branches
e posterior primary
ions of the twelve
al nerves on the right
d 1′, to *d* 6′, the in-
cutaneous branches
e six upper dorsal
es on the left side;
to *d* 12′, cutaneous
hes of the six lower
l nerves from the ex-
branches; *l*, *l*, ex-
branches of the pos-
primary branches
eral lumbar nerves
right side piercing
muscles, the lower
nding over the glu-
gion; *l*′, *l*′, the same
superficially on the
le; *x*, *x*, on the right
he tissue and union
ps of the posterior
ry divisions of four
nerves; *x*′, *x*′, some
ese distributed to
in on the left side.



DIAGRAM OF THE CUTANEOUS NERVES OF THE BACK. (Quain.)

Coccygeal Nerve. The posterior division of this nerve, after being joined by a branch from the last sacral, is distributed to the posterior aspect of the coccyx.

Dissection. The trapezius and latissimus dorsi, which form the first layer of muscles, must now be cleaned by putting them on the stretch, and reflecting the connective tissue which covers them; they should then be dissected in the course of their fibres.

Trapezius. Alone, this muscle is triangular; with its fellow, it presents a trapezoid form. It *arises* from the inner fourth, more or less, of the superior curved line of the occiput, from the ligamentum nuchæ,[1] from the spinous processes of the seventh cervical, and all the dorsal vertebræ, and from their supraspinous ligament. The fibres converge toward the shoulder. The upper pass downward and outward, and are *inserted* by fleshy fibres into the external third of the clavicle; the middle pass transversely outward into the inner border of the acromion and superior lip of the spine of the scapula; the lower pass upward and outward, and terminate in a thin tendon, which plays over the triangular surface at the back of the scapula, and is *inserted* into the beginning of the spine. The insertion of the trapezius exactly corresponds to the origin of the deltoid, and the two muscles are connected by a thin aponeurosis over the spine and acromion. If both the trapezius muscles be exposed, observe that, between the sixth cervical and the third dorsal vertebræ, their origin presents an aponeurotic space of elliptical form (Fig. 85).

The structures covered by the trapezius are, the splenius, the complexus, the levator anguli scapulæ, the rhomboidei minor and major, the supraspinatus, a small part of the infraspinatus, the serratus posticus superior, the vertebral aponeurosis, the

[1] The ligamentum nuchæ is, in man, only a rudiment of the great elastic ligament which supports the weight of the head in quadrupeds. It extends from the spine of the occiput to the spines of all the cervical vertebræ, except the atlas; otherwise it would impede the free rotation of the head. In the giraffe this ligament is six feet long, and as thick as a man's forearm. Professor Quekett states that when divided it shrinks at least two feet.

FIG. 85.



THE SUPERFICIAL MUSCLES OF THE BACK.

latissimus dorsi, the ilio-costalis, the spinal accessory nerve, and the superficialis colli artery.

The fixed point of the muscle being at the vertebral column, all the fibres tend to raise the shoulder. The deltoid cannot raise the humerus beyond an angle of ninety degrees; beyond this, the elevation of the arm is principally effected by the rotatory movement of the scapula. The trapezius is in strong action when a weight is borne upon the shoulders; again, its middle and inferior fibres act powerfully in drawing the scapula backward, as in preparing to strike a blow. If both muscles act, they draw the head backward; if only one acts, it draws the head to the same side. It is supplied by the nervus accessorius and the deep branches of the cervical plexus, and by the superficialis colli artery.

Latissimus Dorsi. This broad flat muscle occupies the lumbar and lower dorsal regions, and thence extends to the arm, where it forms part of the posterior boundary of the axilla. It *arises* from the posterior third of the external lip of the crest of the ilium, from the spinous processes of the two upper sacral, all the lumbar and the sixth lower dorsal vertebræ, and their supraspinous ligament, by a strong aponeurosis; and, lastly, from the three or four lower ribs by fleshy slips, which interdigitate with those of the external oblique muscle of the abdomen. All the fibres converge toward the axilla, were they form a thick muscle, which curves over the inferior angle of the scapula, and is *inserted* by a broad, flat tendon into the bottom of the bicipital groove of the humerus. The upper fibres are inserted into the lowest of the groove, the lower fibres into the upper part. The tendon is about two inches broad, and lies in front of and higher than that of the pectoralis major and of the teres major, from which it is separated by a large *bursa*.[1] It is supplied mainly by the long subscapular nerve, also by the posterior branches of the dorsal and lumbar nerves.

The latissimus dorsi draws the humerus inward and backward; rotating it also inward. It co-operates with the pectoralis major

[1] The latissimus dorsi frequently receives a distinct accessory slip from the inferior angle of the scapula.

pulling any object toward the body; if the humerus be the ted point, it raises the body, as in climbing. The object of the uscle arising so high up the back is, that the transverse fibres the muscle may strap down the inferior angle of the scapula. sometimes happens that the scapula slips above the muscle: is displacement is readily recognized by the unnatural projection f the lower angle of the bone, and the impaired movements of e arm.[1]

The muscle covering the latissimus dorsi is the trapezius above; ose lying beneath it are, a small part of the rhomboideus major, the infraspinatus, and of the teres major, the serratus posticus erior, the spinalis dorsi, the longissimus dorsi, the ilio-costalis, d the external intercostals. Between the base of the scapula, trapezius, and the upper border of the latissimus dorsi, a ngular space is observed when the arm is raised, in which the er fibres of the rhomboideus major and part of the sixth inter- tal space are exposed. Immediately above the crest of the m, between the free margins of the latissimus dorsi and ernal oblique, there is, also, an interval in which a little of the ernal oblique can be seen.

mbar or Verte- al Aponeurosis. This dense, shining, aponeurosis of the back (sometimes termed the aponeurosis of the latissi- s dorsi) forms the posterior part of the sheath of the erector inæ. It is pointed above, where it is continuous with the deep rvical fascia, broader and stronger below. It consists of tendi- us fibres, which are attached internally to the spines of the rsal, all the lumbar and sacral vertebræ; externally, to the gles of the ribs; and inferiorly it is blended with the tendons the serratus posticus inferior and latissimus dorsi. When puration takes place in the loins, constituting a lumbar abscess connection with spinal disease, the pus is seated beneath this neurosis, and is therefore tardy in coming to the surface.

[1] We have seen several instances of this displacement. There is great projection e inferior angle of the scapula, especially when the patient attempts to raise the . He cannot raise the arm beyond a right angle, unless firm pressure is made on lower angle of the scapula, so as to supply the place of the muscular strap. Whether scapula can be replaced or not, a firm bandage should be applied around the chest.

Dissection. Reflect the trapezius from its insertion. On its under surface see the ramifications of its nutrient artery, the *superficialis colli*, a branch of the posterior scapular. A large nerve, the *spinal accessory*, enters its under surface near the clavicle, and divides into filaments, which, reinforced by filaments from the third and fourth cervical nerves, are distributed to the muscle as far as its lower border.

Spinal Accessory Nerve. This nerve, the eleventh cerebral nerve, arises by two roots—the accessory and the spinal portions: the former from the medulla oblongata, the latter from the spinal cord. The *accessory portion*, the smaller, arises by four or five filaments from a gray nucleus in the floor of the fourth ventricle, below the origin of the pneumogastric nerve; the *spinal portion* arises from the lateral part of the cervical portion of the spinal cord by several filaments, some of which arise as low as the sixth cervical vertebra, and which may be traced into the gray matter of the anterior horn. Formed by the union of these roots, the nerve enters the skull through the foramen magnum, and leaves it again, with the accessory portion, through the foramen jugulare. These portions communicate external to the skull; but while the accessory root joins the vagus, the spinal portion, in the main, runs behind the internal jugular vein, traverses obliquely the upper third of the sterno-mastoid muscle, and crosses the posterior triangle of the neck to the trapezius, which it supplies (p. 92). In front of the trapezius it is joined by branches from the third, fourth and fifth cervical nerves, together with which it communicates with the posterior branches of the spinal nerves.

The trapezius should now be cut through the middle, and the inner half turned inward toward the spine, the outer half over the clavicle and the spine of the scapula.

Beneath the trapezius we have to examine the second layer, consisting of three muscles connected with the scapula; namely, the levator anguli scapulæ, the rhomboideus major and minor. The scapula should be adjusted so as to stretch their fibres.

Levator Anguli Scapulæ. This muscle is situated at the back and side of the neck. It *arises* by four tendons from the

posterior tubercles of the transverse processes of the four upper cervical vertebræ. The muscular slips to which the tendons give rise form a single muscle, which descends outward along the side of the neck, and is *inserted* into the posterior border of the scapula between its spine and superior angle. Its *action* is to raise the posterior angle of the scapula; as, for instance, in shrugging the shoulders. Its nerve comes from the fifth cervical, and by filaments from the external series of the deep cervical plexus, which comes from the third and fourth cervical nerves.

Rhomboideus Major and Minor. These flat muscles extend from the spinous processes of the vertebræ to the base of the scapula. They often appear like a single muscle. The rhomboideus minor, the higher of the two, *arises* by a thin aponeurosis from the spinous processes of the last cervical and the first dorsal vertebræ, and is *inserted* into the base of the scapula opposite its spine. The rhomboideus major *arises* by tendinous fibres from the spinous processes of the four or five upper dorsal vertebræ and the supraspinous ligament, and is *inserted* by fleshy fibres into the base of the scapula between its spine and inferior angle, the larger number of the fibres being inserted into a tendinous arch, which is chiefly attached to the inferior angle. The *action* of these muscles is to draw the scapula upward and backward. They are the antagonists of the serratus magnus.

The *nerve of the rhomboid muscles* (posterior scapular) is a branch of the fifth cervical. It passes outward beneath the lower part of the levator anguli scapulæ, to which it sends a branch, and is lost in the under surface of the rhomboidei.

Omo-hyoideus. This muscle extends from the scapula to the os hyoides, and consists of two long, narrow, muscular portions, connected by an intermediate tendon beneath the sterno-mastoid. The posterior portion only can be seen in the present dissection. It *arises* from the upper border of the scapula, close behind the notch, and from the transverse ligament above the notch. Thence the slender muscle passes forward across the lower part of the neck, beneath the sterno-mastoid, where it changes its direction and ascends nearly vertically, to be attached to the os hyoides at the junction of the body with the greater

cornu (p. 95). Thus the two portions of the muscle form, beneath the sterno-mastoid, an obtuse angle, of which the apex is tendinous, and of which the angular direction is maintained by a layer of fascia, proceeding from the tendon to the first rib and the clavicle. Its *action* is to depress the os hyoides. Its nerve comes from the descendens noni and the communicantes noni (p. 128).

Suprascapular Artery. This artery (transversalis humeri), a branch of the thyroid axis (p. 139), runs behind and parallel with the clavicle, over the lower end of the scalenus anticus and subclavian artery, and beneath the sterno-mastoid and omo-hyoid muscles, to the upper border of the scapula, where it usually passes above the ligament bridging over the notch. It ramifies in the supraspinous fossa, supplying the supraspinatus, and then passes under the acromion to the infraspinous fossa, where it inosculates freely with the dorsalis scapulæ, a branch of the subscapular. It sends off—

a. The *inferior sterno-mastoid* artery to the sterno-mastoid and contiguous muscles; *b*. the *supra-acromial* branch, which ramifies upon the acromion, anastomosing with the other acromial arteries derived from branches of the axillary; *c*. a small *subscapular* branch to the fossa of the same name; *d*. *articular* arteries to the shoulder joint; and, lastly, *e*. the *infraspinous* branch, which anastomoses with the dorsalis scapulæ. The suprascapular vein terminates either in the subclavian or in the external jugular.

The *suprascapular nerve*, a branch of the fifth and sixth cervical nerves, runs with the corresponding artery, and, after passing through the suprascapular notch, is distributed to the supraspinatus and infraspinatus. In the supraspinous fossa, this nerve sends a small *articular* branch to the shoulder joint; in the infraspinous fossa it gives off two branches to the infraspinatus, and some to the shoulder joint.

Posterior Scapular Artery. This artery is one of the divisions of the transversalis colli, but comes very frequently from the subclavian in the third part of its course (p. 140). It runs across the lower part of the neck, above or between the nerves of the brachial plexus, toward the posterior superior angle of the scapula. Here it pursues its course along the posterior border of the scapu

neath the levator anguli scapulæ and the rhomboidei, anasto-sing with branches of the suprascapular and subscapular eries, and with branches from the intercostal arteries. The rresponding vein joins the external jugular or the subclavian.

Dissection. Divide the rhomboid muscles near their insertion, and trace the artery to the inferior angle of e scapula, where it terminates in the rhomboidei, serratus gnus, and latissimus dorsi.

Numerous muscular branches arise from the posterior scapular. e *superficialis colli* (the other division of the transversalis colli) given off near the upper angle of the scapula, for the supply of trapezius, which it enters together with the spinal accessory ve.

Divide and reflect the latissimus dorsi before the inferior angle he scapula, and draw the scapula forcibly outward, to have a e perfect view of the extent of the serratus magnus than was in the axilla. The abundance of connective tissue in this ation is necessary for the play of the scapula on the chest.

atus Magnus. This broad, thin, flat muscle intervenes between the scapula and the ribs. It *arises* by nine fleshy itations from the eight upper ribs, each rib giving origin to , and the second to two, and from the fascia covering the responding intercostal spaces. The four lower digitations respond with those of the external oblique muscle of the omen. The fibres pass backward and outward and are nged in three fasciculi; the upper portion *arises* from the first second ribs and the fascia between them, and is *inserted* into triangular surface in front of the upper angle of the scapula; middle portion *arises* from the second, third and fourth ribs, is *inserted* into the inner lip of the vertebral border between first and third portions; the third portion *arises* from the fifth , seventh and eighth ribs, and is *inserted* into the smooth ce in front of the inferior angle; this last portion consists of serrations, and are those which interdigitate with the exter-blique.

his is the most important of the muscles which regulate the ements of the scapula. It draws the scapula forward, and

thus gives additional reach to the arm; it counteracts all forces which tend to push the scapula backward; for instance, when a man falls forward upon his hands, the serratus magnus sustains the shock, and prevents the scapula from being driven back to the spine. Supposing the fixed point to be at the scapula, some anatomists ascribe to it the power of raising the ribs; hence Sir Charles Bell called it the external respiratory muscle, the internal respiratory muscle being the diaphragm.

The nerve which supplies it is a branch of the fifth and sixth cervical nerve; it descends along its outer surface, distributing a filament to each digitation of the muscle (p. 145).

Dissection. Divide the serratus magnus near the scapul and remove the arm by sawing through the mi dle of the clavicle, cutting through the axillary vessels and nerve These should be tied to the coracoid process. After the remov of the arm, examine the precise insertions of the precedi muscles.

DISSECTION OF THE MUSCLES OF THE SHOULDER.

Dissection. The remainder of the skin over the should is to be reflected, and in the subcutaneous tiss are found the cutaneous vessels and nerves. Some pass do over the shoulder, others ascend over the deltoid, emerging fr beneath its lower border.

The *acromial branches* come from the third and fourth cervi nerves, and descend over the acromion (Fig. 86) in front and behind the deltoid. The *cutaneous branch* of the *circumf* nerve comes out beneath the posterior border of the deltoid, a supplies the skin over the posterior and outer two-thirds of t muscle; others perforate the muscle, each accompanied by small artery.

Notice the strong layer of fascia upon the surface of the deltoi which extends from the aponeurosis covering the muscles on th back of the scapula, and is continuous with the fascia of the arm It dips down between the fibres of the muscle, dividing it int large bundles. This fascia is to be removed, by putting th deltoid on the stretch, and reflecting it in the direction of it

fibres, beginning from the front. The fascia will be seen to be continuous in front with the fascia covering the pectoralis major; above, it is attached to the clavicle and spine of the scapula; behind, it is continuous with that over the infraspinatus.

Deltoid. The large muscle which covers the shoulder joint is named deltoid, from its resemblance to the Greek Δ reversed. It *arises* from the external third of the anterior border of the clavicle, from the apex and outer border of

FIG. 86.



1. Supra-acromial br. of the cervical nerves.
2. Ascending and descending brs. of the circumflex n.
3, 4. Cutaneous brs. of the musculo-cutaneous n.
5. Internal cutaneous br. of musculo-spiral n.
6. Intercosto-humeral brs.
7. Filaments of the lesser internal cutaneous n.
8. Posterior cutaneous br. of internal cutaneous n.
9. Branch of internal cutaneous n.

CUTANEOUS NERVES OF THE LEFT SHOULDER AND ARM. (POSTERIOR VIEW.)

the acromion, and from the lower border of the spine of the scapula down to the triangular surface at its root. This origin, which corresponds to the insertion of the trapezius, is tendinous and fleshy everywhere, except at the commencement of the spine of the scapula, where it is simply tendinous, and connected with the infraspinous aponeurosis. The muscular fibres descend, the anterior backward, the posterior forward, the middle perpen-

dicularly ; all converge to a tendon which is *inserted* into a rough surface on the outer side of the humerus, a little above the middle of the shaft. The insertion of the tendon extends one inch and a half along the humerus, and terminates in a V-shaped form, the origin of the brachialis anticus embracing it on either side. Sometimes a few fibres of the pectoralis major are connected with its front border.

The muscular bundles composing the deltoid have a peculiar arrangement; a peculiarity arising from its broad origin and its narrow insertion. It consists in the interposition of tendons between the bundles for the attachment of the muscular fibres. The annexed wood-cut shows this arrangement better than any description. The action of the muscle is not only concentrated upon one point, but its power is also greatly increased by this arrangement.

Action of the Deltoid. It raises the arm ; but it cannot do so beyond an angle of ninety degrees. The elevation of the arm beyond this angle is effected through the raising of the shoulder by the trapezius and serratus magnus. Its anterior fibres draw the arm forward ; its posterior, backward.

FIG. 87.

ANALYSIS OF THE DELTOID.

This powerful muscle is supplied with blood by the anterior and posterior circumflex, the thoracica humeraria, the thoracica acromialis, all from the axillary artery; also by the deltoid branch of the brachial. Its nerve is the circumflex.

The rotundity of the shoulder is due, not so much to the deltoid as to the upper end of the humerus. When the head of the humerus is dislocated into the axilla, the fibres of the muscle run vertically to their insertion ; hence the flattening of the deltoid, and the greater prominence of the acromion.

It is below the deltoid that a

ununited fracture of the humerus is most commonly met with, owing to the muscle displacing the upper fragment.

Dissection. Reflect the deltoid from its origin, and turn it downward. Observe the ramifications of the circumflex nerve and the anterior and posterior circumflex arteries on its under surface; notice also the large bursa between it and the tendons inserted into the great tuberosity of the humerus.

Parts Covered by the Deltoid. The structures seen on reflecting the deltoid are as follows: the bursa, already alluded to, the coracoid process, the coraco-acromial ligament, the origins of the biceps and coraco-brachialis, the insertions of the pectoralis minor and major, the long head of the biceps, the insertions of the supraspinatus, infraspinatus, and teres minor, the long and external heads of the triceps, the circumflex vessels and nerve, and the neck and upper part of the humerus.

Bursa under the Deltoid, or Sub-acromial. The large bursa under the deltoid extends for some distance beneath the acromion and the coraco-acromial ligament, and covers the tendons attached to the great tuberosity of the humerus. It communicates, very rarely, with the shoulder joint. Its use is to facilitate the movements of the head of the bone under the acromial arch.

Posterior Circumflex Artery. This artery is given off from the axillary in the third part of its course; it runs behind the surgical neck of the humerus, through a quadrilateral opening, bounded above by the subscapularis and teres minor; below, by the teres major; externally, by the neck of the humerus; and internally, by the long head of the triceps (p. 338). Its branches terminate in the under surface of the deltoid, anastomosing with the anterior circumflex, acromial thoracic, and suprascapular arteries.

From the posterior circumflex, a branch descends in the substance of the long head of the triceps, to inosculate with the superior profunda; this is one of the channels through which the circulation would be carried on if the axillary were tied in the last part of its course.

Circumflex Nerve. This nerve, a branch of the posterior cord of the axillary plexus, runs with the posterior

circumflex artery, through the same quadrilateral space, and then divides into two branches—an upper and a lower. The *upper branch* winds round the neck of the humerus, and supplies the anterior part of the deltoid, and gives off cutaneous branches to supply the skin over its lower part. The *lower branch* sends a filament to the teres minor, one or two to the integuments over the shoulder at its posterior part, and terminates in the substance of the deltoid. It also distributes an *articular* filament, which enters the shoulder joint in front, below the subscapularis.

The proximity of this nerve to the head of the humerus explains the occasional paralysis of the deltoid after dislocation or fracture of the humerus. The nerve is liable to be injured, if not actually lacerated, by the pressure of the bone. In the summer of 1840 a man was admitted into the hospital with a severe injury to the shoulder, and died of delirium tremens. On examination the humerus was found broken high up, the capsule of the joint opened, and the circumflex nerve torn completely across.[1]

A strong aponeurosis covers the muscles of the dorsum of the scapula, and is firmly attached to the spine and borders of the bone. At the posterior edge of the deltoid it divides into two layers, one of which passes over, the other under, the muscle. Remove the aponeurosis, so far as it can be done without injury to the muscular fibres which arise from its under surface.

Infraspinatus. This triangular muscle *arises* by fleshy fibres from the posterior two-thirds of the infraspinous fossa, by tendinous fibres from the ridges on the fossa, and from the aponeurosis which covers it. The fibres converge to a tendon, which is at first contained in the substance of the muscle, and then proceeds, over the capsular ligament of the shoulder joint, to be *inserted* into the middle depression on the greater tuberosity of the humerus. Its nerve comes from the suprascapular.

Teres Minor. This long narrow muscle is situated below the infraspinatus, along the inferior border of the scapula. It *arises* from the dorsum of the scapula close to the inferior border, and from the intermuscular septa between it and the infraspinatus above and the teres major below. The fibres

[1] See preparation in Museum of St. Bartholomew's Hospital.

ascend outward, parallel with those of the infraspinatus, and terminate in a tendon, which passes over the capsular ligament of the shoulder joint, and is *inserted* into the lowest depression on the great tuberosity of the humerus, and by muscular fibres into the bone below it. It is supplied by a branch of the circumflex nerve, which enters the muscle at its lower border, and it has (usually) a small ganglion-like enlargement upon it.

The *action* of the infraspinatus and teres minor is to rotate the humerus outward, and when the arm is raised it draws the humerus downward and backward.

Teres Major. This muscle is closely connected with the latissimus dorsi, and extends from the inferior angle the scapula to the humerus, contributing to form the posterior undary of the axilla. It *arises* from the flat surface on the dorsal pect of the inferior angle of the back of the scapula, from its ferior border, and the intermuscular septa, and terminates upon flat tendon, nearly two inches in breadth, which is *inserted* into e inner edge of the bicipital groove of the humerus, behind d a little lower than the tendon of the latissimus dorsi. Its *ion* is to draw the humerus backward and downward when e arm is raised, and to rotate it slightly inward. It is supplied the middle subscapular nerve, which enters it on its axillary pect.

A bursa is found in front of, and another behind, the tendon of he teres major; the former separates it from the latissimus dorsi, he latter from the bone.

Supraspinatus. This muscle *arises* from the posterior two-thirds of the supraspinous fossa, and from its aponeurotic overing. It passes under the acromion, over the capsular ligament of the shoulder joint, and is *inserted* by a strong tendon into he superior depression on the greater tuberosity of the humerus. o see its insertion, the acromion should be sawn off near the eck of the scapula. Its *action* is to assist the deltoid in raising he arm. It is supplied by two branches derived from the supracapular nerve.

Subscapularis. This triangular fleshy muscle occupies the subscapular fossa. It *arises* from the posterior three-

fourths of the fossa, except the posterior border and angles which give attachment to the serratus magnus, and from three or four tendinous septa attached to the oblique bony ridges on its surface. The fibres passing upward and outward, converge toward the neck of the scapula, where they terminate upon three or four tendons, which are concealed among the muscular fibres, and are *inserted* into the lesser tuberosity of the humerus and into the bone for an inch below the tuberosity. Its broad insertion is

FIG. 88.



DIAGRAM OF ARTERIES OF SCAPULA.

1. Suprascapular artery.
2. Posterior circumflex a.
3. Infraspinous br. of suprascapular a.
4. Dorsalis scapulæ a.
5. Posterior scapular a.
6. Subclavian a.

closely connected with the capsule of the shoulder joint, which it completely protects upon its inner side. Its *action* is to rotate the humerus inward, and when the arm is raised, draws it downward. The nerves which supply it come from the long and middle subscapular nerves.

The coracoid process, with the coraco-brachialis and short head of the biceps, forms an arch, under which the tendon of the subscapularis plays. There are several *bursæ* about the tendon. One of considerable size, on the upper surface of the tendon, facilitates its motion beneath the coracoid process and the coraco-brachialis; this sometimes communicates with the large bursa under the deltoid. Another is situated between the tendon and the capsule of the joint, and almost invariably communicates with it.

Dissection. Now reflect the muscles from the surfaces of the scapula, to trace the arteries which ramify upon it.

Continuation of Suprascapular Artery and Nerve. This artery, a branch of the thyroid axis, runs under and parallel with the clavicle, and passes above the notch of the scapula, into the supraspinous fossa; it sends a branch to the supraspinatus, another to the shoulder joint, and then descends behind the neck of the scapula into the fossa below the spine, where it inosculates directly with the dorsalis scapulæ. Its branches ramify upon the bone, and supply the infraspinatus and teres minor (Fig. 88).

The suprascapular nerve passes most frequently through the notch of the scapula, accompanies the corresponding artery, supplies two branches to the supraspinatus and one to the shoulder joint; it then enters the infraspinous fossa, to terminate in the infraspinatus.

Dorsalis Scapulæ Artery. This artery, after passing through the triangular space (p. 337), curves round the inferior border of the scapula, which it grooves, to the infraspinous fossa, where it ascends close to the bone, and anastomoses with the supra- and posterior scapular arteries. Another branch of the subscapular artery runs between the teres minor and major, toward the inferior angle of the scapula, where it anastomoses with the posterior scapular artery (Fig. 88).

The several communications about the scapula between the branches of the subclavian and axillary arteries would furnish a large collateral supply of blood to the arm, if the subclavian were tied above the clavicle (p. 142).

Dissection. If the skin has not been reflected from the back of the arm, it should now be done. In the subcutaneous tissue will be seen the *internal cutaneous branch* of the *musculo-spiral* nerve, which supplies the skin as low down as the olecranon. On the inner side of this branch is the *intercosto-humeral nerve*, supplying the skin as far as the lower third of the arm. The *nerve of Wrisberg* also supplies the lower third of the arm; and on the outer side for the same distance is the *external cutaneous* branch of the *musculo-spiral* nerve.

The fascia is now to be removed, when the triceps will be exposed, forming the only muscle on the back of the arm.

Triceps Extensor Cubiti. This muscle, which arises by three distinct heads, and was only partially seen in the dissection of the upper arm (p. 358), should now be thoroughly examined. The *long head arises* immediately below the glenoid cavity of the scapula, by a strong, flat tendon, which is connected with the capsular and glenoid ligaments of the shoulder joint. The *external head arises* from the posterior part of the humerus, below the insertion of the teres minor, as far as the musculo-spiral groove, from the outer border of the humerus, and the external intermuscular septum. The *internal head arises* from the posterior part of the humerus, below the teres major and the musculo-spiral groove, as far as the olecranon fossa; it has an additional origin from the internal intermuscular septum, and from the internal border of the humerus. The precise origin of these heads from the humerus may be ascertained by following the superior profunda artery and musculo-spiral nerve, which separate them. The three portions of the muscle terminate upon a broad tendon, which covers the back of the elbow joint, and is *inserted* into the summit and sides of the olecranon; it is also connected with the fascia on the back of the forearm. The effect of this connection is that the same muscle which extends the forearm tightens the fascia which gives origin to the extensors of the wrist and fingers. The same holds good in the case of the biceps, and its semilunar expansion in the fascia of the forearm.

Between the tendon and the olecranon is a *bursa*, commonly of small size, but sometimes so large as to extend upward

nd the capsule of the joint. This bursa must not be mistaken he subcutaneous one, which is situated between the skin and olecranon, and is so often injured by a fall on the elbow.

Dissection. By dividing the triceps transversely a little above the elbow, and turning down the lower ion, it will be seen that some of the muscular fibres terminate n the capsular ligament of the joint. They have been de- ed as a distinct muscle, under the name of the *subanconeus*; r use is to draw up the capsule, so that it may not be injured ng extension of the arm. The subanconeus is in this respect ogous to the subcrureus muscle of the thigh. Observe the *a* under the tendon, and the arterial arch formed upon the k part of the capsule by the superior profunda and the anas- otica magna (Fig. 89, p. 416).

race the continuation of the superior profunda artery (p. 353) musculo-spiral nerve round the posterior part of the humerus. y lie in a slight groove on the bone,[1] between the external and nal heads of the triceps, and are protected by an aponeurotic thrown over them by the external head of the triceps. After lying the muscles, the artery continues its course along the r side of the arm between the brachialis anticus and supinator longus, and inosculates with the radial recurrent. It gives branch, which runs down between the triceps and the bone, inosculates, at the back of the elbow, with the anastomotica na and posterior interosseous recurrent. The musculo spiral e which accompanies the artery sends branches to supply the portions of the triceps, the supinator radii longus, and ex- or carpi radialis longior.[2] It then divides into the posterior osseous and radial nerves. The small nerve must be made which runs down in the substance of the triceps, accompanied branch from the superior profunda artery, to supply the neus. The cutaneous branches of the musculo-spiral nerve been already dissected (p. 345).

[1] It is worth remembering that the nerve may be injured by a fracture of the rus in this situation, and even by too tight bandaging; the result being paralysis extensor muscles of the forearm.

[2] The brachialis anticus usually receives a branch from the musculo-spiral nerve.

DISSECTION OF THE BACK OF THE FOREARM.

Subcutaneous Bursæ.

Remove the skin from the back of the forearm, hand and fingers, and make out the subcutaneous *bursa* over the olecranon. It is of considerable size, and, if distended, would appear nearly as large as a walnut. Another *bursa* is sometimes found a little lower down upon the ulna. A subcutaneous *bursa* is generally placed over the internal condyle, another over the external. A *bursa* is also situated over the styloid process of the ulna; this sometimes communicates with the sheath of the extensor carpi ulnaris. Small *bursæ* are sometimes developed in the cellular tissue over each of the knuckles.

The cutaneous veins from the back of the hand and forearm join the venous plexus at the bend of the elbow (see p. 346).

Cutaneous Nerves of the Back of the Forearm.

The cutaneous nerves of the back of the forearm are derived from the external cutaneous branches of the musculo-spiral, from branches of the internal cutaneous and of the external cutaneous nerves. The greater number of these nerves may be traced down to the back of the wrist.

Nerves on the Back of the Hand and Fingers.

The skin on the back of the hand is united to the subjacent tendons by an abundance of loose connective tissue, in which are large veins, and branches of the radial and ulnar nerves. The *dorsal branch* of the ulnar nerve passes beneath the tendon of the flexor carpi ulnaris, pierces the fascia just above the wrist joint, runs over the posterior annular ligament of the wrist, and divides upon the back of the hand into filaments, which supply both sides of the back of the little finger, the ring finger and the ulnar side of the middle finger. The *radial nerve* passes obliquely beneath the tendon of the supinator longus, perforates the fascia about two inches above the wrist joint, and subdivides into filaments, which supply both sides of the back of the thumb and forefinger, and the radial side of the middle finger.[1]

[1] The relative share which the radial and ulnar nerves take in supplying the fingers varies. Under any arrangement the thumb and each finger has two dorsal nerves, one on either side, of which the terminal branches reach the root of the nail. They supply filaments to the skin on the back of the finger, and have frequent com-

The radial nerve commonly gives off, on the back of the hand, a branch which joins the nearest branch of the ulnar.

Fascia on Back of Forearm. The fascia on the back of the forearm is composed of fibres interlacing and stronger than that upon the front of the forearm. It is attached to the condyles of the humerus and to the olecranon, and is strengthened by an expansion from the tendon of the triceps. Along the forearm it is attached to the ridge on the posterior part of the ulna. Its upper third gives origin to the fibres of the muscles beneath it, and divides them by septa, to which their fibres are also attached.

Posterior Annular Ligament. This ligament should be considered as a part of the fascia of the forearm, specially strengthened by oblique aponeurotic fibres on the back of the wrist, to confine the extensor tendons. These fibres are attached to the outer margin of the radius, and thence pass obliquely inward to the inner side of the wrist, where they are connected with the pisiform and cuneiform bones. They pass below the styloid process of the ulna, to which they are in no way attached, otherwise the rotation of the radius would be impeded.

Separate Sheaths for Extensor Tendons. From the deep surface of the posterior annular ligament, processes are attached to the ridges on the back of the radius, so as to form six distinct fibro-osseous sheaths for the passage of the extensor tendons. Commencing from the radius, the first sheath contains the tendons of the extensor ossis metacarpi and the extensor primi internodii pollicis; the second, the tendons of the extensor carpi radialis longior and brevior; the third, the tendon of the extensor secundii internodii pollicis; the fourth, the tendons of the extensor indicis and the extensor communis digitorum; the fifth, the tendon of the extensor minimi digiti; and the sixth, the tendon of the extensor carpi ulnaris. All the sheaths are lined by synovial membranes, which extend nearly to the insertions of their tendons. Occasionally, but not often, one or more of them communicate with the wrist joint.

munications with the palmar digital nerves. In some instances one or more of the dorsal nerves do not extend beyond the first phalanx; their place is then supplied by a branch from the palmar nerve.

The *fascia of the metacarpus* consists of a thin fibrous layer, continued from the posterior annular ligament. It separates the extensor tendons from the subcutaneous veins and nerves, and is attached to the radial side of the second metacarpal bone, and the ulnar side of the fifth.

Dissection. The fascia must be removed from the muscles, without injuring the muscular fibres which arise from its under surface. Preserve the posterior annular ligament. The following superficial muscles are now exposed, and should be examined in the order in which they are placed, proceeding from the radial to the ulnar side:—1. The supinator radii longus (already described, p. 365). 2. The extensor carpi radialis longior. 3. The extensor carpi radialis brevior. 4. The extensor communis digitorum. 5. The extensor minimi digiti. 6. The extensor carpi ulnaris. 7. The anconeus.

Superficial Muscles on the Back of the Forearm.

A little below the middle of the forearm, the extensors of the wrist and fingers diverge from each other, leaving an interval, in which are seen the three extensors of the thumb—namely, the extensor ossis metacarpi pollicis, the extensor primi internodii pollicis, and the extensor secundi internodii pollicis. The two former cross obliquely over the radial extensors of the wrist, and pass over the lower third of the radius; the latter emerges from under the radial border of the extensor communis digitorum, and then passes over the insertion of the tendons of the radial extensors of the wrist.

Between the second and third extensors of the thumb, we observe a part of the lower end of the radius, which is not covered either by muscle or tendon. This subcutaneous portion of the bone is immediately above the prominent tubercle in the middle of its lower extremity, and, since it can be easily felt through the skin, it presents a convenient place for examination in doubtful cases of fracture.

Extensor Carpi Radialis Longior. This muscle is partly covered by the supinator radii longus. It *arises* from the lower third of the ridge leading to the external condyle of the humerus, and from the intermuscular septum. It descends along the outer side

of the forearm, and terminates about the middle, in a flat tendon, which passes beneath the extensor ossis metacarpi and primi internodii pollicis, traverses a groove on the outer and back part of the radius, lined by a synovial membrane, and is *inserted* into the radial side of the carpal end of the metacarpal bone of the index finger. Previous to its insertion, the tendon is crossed by the extensor secundi internodii pollicis. It is supplied by a branch from the musculo-spiral nerve.

Extensor Carpi Radialis Brevior. This muscle *arises* from the external condyle by the tendon common to it and the other extensors, from the intermuscular septa, from the external lateral ligament of the elbow joint and the aponeurosis covering the muscle. The muscular fibres terminate near the lower third of the forearm, upon the under surface of a flat tendon, which descends, covered by that of the extensor carpi radialis longior, beneath the three extensors of the thumb. The tendon traverses a groove on the back of the radius, on the same plane with that of the long radial extensor, but lined by a separate synovial membrane, and is *inserted* into the radial side of the base of the metacarpal bone of the middle finger. A bursa is generally found between the tendon and the bone. Its nerve comes from the posterior interosseous.

Extensor Digitorum Communis. This muscle *arises* from the common tendon attached to the external condyle, from the septa between it and the contiguous muscles, and from its strong fascial covering. A little below the middle of the forearm, the muscle divides into three tendons, which pass, together with the extensor indicis, beneath the posterior annular ligament, through a groove on the back of the radius lined by synovial membrane. On the back of the hand the tendons become broader and flatter, and diverge from each other toward the metacarpal joints of the fingers, where they become thicker and narrower, and give off, on each side, a fibrous expansion, which covers the sides of the joint. Over the first phalanx of the finger, each tendon again spreads out, receives the expanded tendons of the lumbricales and interossei muscles, and divides at the second phalanx into three portions, of which the middle is *inserted* into

the upper end of the second phalanx; the two lateral, reuniting over the lower end of the second phalanx, are *inserted* into the upper end of the third.[1] Its nerve comes from the posterior interosseous.

The oblique aponeurotic slips which connect the tendons on the back of the hand are subject to great variety. The tendon of the index finger is commonly free; it is situated on the radial side of the proper indicator tendon, and becomes united with it at the metacarpal joint.

The tendon of the middle finger usually receives a slip from that of the ring. The tendon of the ring finger generally sends a slip to the tendons on either side of it, and in some cases, entirely furnishes the tendon of the little finger. Thus the ring finger does not admit of independent extension.

The muscle is not only a general extensor of the fingers, but can extend some of the phalanges independently of the rest: *e.g.* it can extend the first phalanges while the second and third are flexed; or it can extend the second and third phalanges during flexion of the first.

Extensor Minimi Digiti or Auricularis. This long, slender muscle, situated on the ulnar side of the common extensor, *arises* from the common tendon from the external condyle, and from the septa between it and the contiguous muscles. Its slender tendon runs separately beneath the annular ligament immediately behind the joint between the radius and ulna, in a special sheath lined by synovial membrane. On emerging from the annular ligament, the tendon splits into two, which pass obliquely to the little finger. At the first joint of the little finger the outer tendon is joined by that of the common extensor, and both expand upon the first and second phalanges, terminating in the same manner as the extensor tendons of the other fingers. Its nerve comes from the posterior interosseous.

[1] The extensor tendons are inserted into the periosteum; but the flexor tendons are inserted into the substance of the bone. This accounts for the facility with which the former will tear off the bones in cases of necrosis, while the latter will adhere so tightly as to require cutting before the phalanx can be removed. It probably also explains the great liability to necrosis which is so frequently observed in cases of thecal abscess.

Extensor Carpi Ulnaris. This muscle *arises* from the common tendon from the external condyle, from the septum between it and the extensor minimi digiti, from the fascia of the forearm, and from the aponeurosis attached to the posterior ridge of the ulna common to this muscle, the flexor carpi ulnaris, and the flexor profundus digitorum. The fibres terminate upon a strong, broad tendon, which traverses a distinct groove on the back of the ulna, close to the styloid process, and is *inserted* into the posterior aspect of the carpal end of the metacarpal bone of the little finger. Below the styloid process of the ulna, the tendon passes beneath the posterior annular ligament, over the back of the wrist, and is confined in a very strong fibrous canal, which is attached to the back of the cuneiform, pisiform and unciform bones, and is lined by a continuation from the synovial membrane in the groove of the ulna. The *action* of this muscle is to extend the hand, and incline it toward the ulnar side. It is supplied by the posterior interosseous nerve.

In pronation of the forearm, the lower articular end of the ulna projects between the tendons of the extensor carpi ulnaris and the extensor minimi digiti. A subcutaneous *bursa* is sometimes found above the bone in this situation.

Anconeus. This small triangular muscle is situated at the outer and back part of the elbow. It is covered by a strong layer of fascia, derived from the tendon of the triceps, and appears like a continuation of that muscle. It *arises* by a tendon from the posterior part of the external condyle of the humerus, and is *inserted* into the triangular surface on the upper fourth of the outer part of the ulna. Part of the under surface of the muscle is in contact with the capsule of the elbow joint. Its *action* is to assist in extending the forearm. Its nerve comes from the musculo-spiral.

Dissection. To expose the deep layer of muscles, detach from the external condyle the extensor carpi radialis brevior, the extensor communis digitorum, the extensor minimi digiti, and the extensor carpi ulnaris; and, after noticing the vessels and nerves which enter their under surface, turn them down. The deep-seated muscles, with the posterior interosseous

artery and nerve, must be dissected. The muscles exposed are:

Deep-seated Muscles on the Back of the Forearm.

1. The extensor ossis metacarpi pollicis. 2. Extensor primi internodii pollicis. 3. Extensor secundi internodii pollicis. 4. Extensor indicis or indicator. 5. The supinator radii brevis. They are all supplied by branches from the posterior interosseous nerve.

Extensor Ossis Metacarpi Pollicis.

This muscle lies immediately below the supinator brevis, and *arises* from the posterior surface of the ulna below the supinator brevis, from the posterior surface of the middle third of the radius, and from the interosseous membrane. The muscle passes obliquely downward and outward, crosses the radial extensors of the wrist about three inches above the carpus, and terminates in a tendon, which passes along a common groove with the extensor primi internodii pollicis lined by synovial membrane, on the outer part of the lower end of the radius, and is *inserted* into the base of the metacarpal bone of the thumb, and frequently also by a tendinous slip into the trapezium.

Extensor Primi Internodii Pollicis.

This, the smallest of the deep muscles, *arises* from the posterior surface of the radius, below the preceding, and from the interosseous membrane. It descends obliquely in company with the preceding muscle, turns over the radial extensors of the wrist, and terminates upon a tendon which passes beneath the annular ligament, through the groove in the outer part of the radius, and is *inserted* into the radial side of the base of the first phalanx of the thumb.

Extensor Secundi Internodii Pollicis.

This muscle covers part of the origin of the preceding muscle, and *arises* from the posterior surface of the ulna, below the extensor ossis metacarpi pollicis and from the interosseous membrane. The tendon receives fleshy fibres as low as the wrist, passes between the annular ligament in a distinct groove on the back of the radius, crosses the tendons of the radial extensors of the wrist, proceeds over the metacarpal bone and the first phalanx of the thumb, and is *inserted* into the base of the last phalanx.

The tendons of the three extensors of the thumb may be easily distinguished in one's own hand. The extensor ossis metacarpi

and primi internodii pollicis cross obliquely over the radial artery, where it lies on the external lateral ligament of the carpus; the extensor secundi internodii pollicis crosses the artery just before it sinks into the palm, between the first and second metacarpal bones, and is a good guide to the vessel. The *action* of the three extensors of the thumb is implied by their names.

Extensor Indicis, or Indicator. This muscle *arises* from the posterior surface of the ulna, below the extensor secundi internodii pollicis, and slightly from the interosseous membrane. The tendon passes beneath the posterior annular ligament, in the same groove, on the back of the radius, with the tendons of the extensor digitorum communis. It then proceeds over the back of the hand to the first phalanx of the index finger, where it is united to the ulnar border of the common extensor tendon. By the *action* of this muscle the index finger can be extended independently of the others.

Dissection. Reflect the anconeus from its origin, to expose the following muscle:—

Supinator Radii Brevis. This muscle embraces the upper third of the radius. It *arises* from the external condyle of the humerus, from the external lateral ligament of the elbow joint, from the orbicular ligament surrounding the head of the radius, from an oblique ridge on the outer surface of the ulna below the insertion of the anconeus, by fleshy fibres from the triangular excavation below the lesser sigmoid notch of the ulna, and from the aponeurosis covering the muscle. The muscular fibres turn over the neck and upper part of the shaft of the radius, and are *inserted* into the upper third of this bone, as far forward as the ridge leading from the tubercle to the insertion of the pronator teres. The muscle is traversed obliquely by the posterior interosseous nerve, which sends a branch to it, and its upper part is in contact with the capsule of the elbow joint. It is a powerful supinator of the forearm, some of its fibres acting at nearly a right angle to the axis of the radius.

Posterior Interosseous Artery. This artery comes from the ulnar by a common trunk with the anterior interosseous (p. 372), and supplies the muscles on the back of the forearm. It

passes between the oblique ligament and the interosseous membrane, and appears at the back between the supinator radii brevis and the extensor ossis metacarpi pollicis. After supplying branches to all the muscles in this situation, the artery descends, much diminished in size, between the superficial and deep layer of muscles to the wrist, where it inosculates with the carpal

FIG. 89.

1. The superior profunda.
2. The anastomotica magna.
3. The posterior ulnar recurrent.
4. The posterior interosseous { its ascending and descending branches.
5. The termination of the anterior interosseous.
6. The posterior carpal arch.

DIAGRAM SHOWING THE ANASTOMOSES OF ARTERIES AT THE BACK OF THE ELBOW AND WRIST JOINTS.

branches of the anterior interosseous and the posterior carpal branches of the radial and ulnar arteries.

The largest branch of this artery is the *interosseous recurrent*. It ascends beneath the supinator brevis and the anconeous to the space between the external condyle and the olecranon, where it inosculates with the branch of the superior profunda which

descends in the substance of the triceps, with the posterior ulnar recurrent artery, and with the anastomotica magna.

In the lower part of the back of the forearm, a branch of the anterior interosseous artery is seen passing through the interosseous membrane to reach the back of the wrist.

Posterior Interosseous Nerve. The nerve which supplies the muscles on the back of the forearm is the *posterior interosseous*, one of the divisions of the musculo-spiral. It passes obliquely through the supinator radii brevis, and descends, lying on the lower fibres of this muscle, the extensores ossis metacarpi and primi internodii pollicis, and beneath the superficial extensores. It then, much diminished in size, passes under the extensor secundi internodii pollicis, on the interosseous membrane, as far as the posterior annular ligament, where it presents a gangliform enlargement. Between the superficial and deep layer of muscles, it sends to each a filament, generally in company with a branch of the posterior interosseous artery. It sends a branch to the extensor carpi radialis brevior, and supplies the supinator brevis in passing through its substance. The supinator radii longus and the extensor carpi radialis longior are supplied by distinct branches from the musculo-spiral nerve.

After the posterior interosseous nerve descends beneath the extensor secundi internodii pollicis, it lies in the interosseous membrane, beneath the extensor digitorum communis and the indicator. At the back of the wrist, beneath the annular ligament, it forms the gangliform enlargement from which filaments are sent to the carpal and metacarpal joints.

Dissection. Radial Artery on the Back of the Wrist. The radial artery is continued over the external lateral ligament of the carpus, beneath some filaments of the radial nerve, cutaneous veins, and the extensor tendons of the thumb, to the proximal part of the interval between the first and second metacarpal bones, where it dips down between the two origins of the abductor indicis, and, entering the palm, forms the deep palmar arch. In this part of its course it is accompanied by a filament of the musculo-cutaneous nerve; observe also that the tendon of the extensor secundi internodii pollicis passes over it immediately

before it sinks into the palm. It supplies in this part of its course the following small branches to the back of the hand:—

a. Posterior carpal artery.—This branch passes across the carpal bones, beneath the extensor tendons. It inosculates with the termination of the anterior interosseous artery, and forms an arch beneath the extensor tendons, with a corresponding branch from the ulnar artery. The carpal arch sends off small branches, called the *dorsal interosseous*, which descend along the third and fourth interosseous spaces from the arch just mentioned, beneath the extensor tendons, and inosculate near the carpal ends of the metacarpal bones with the perforating branches from the deep palmar arch.

b. The *first dorsal interosseous artery* is generally larger than the others. It passes forward, beneath the extensor of the thumb, on the second interosseous space, to the cleft between the index and middle fingers, communicating here with a perforating branch of the deep palmar arch; and terminates in small branches, some of which proceed along the back of the fingers, others inosculate with the palmar digital arteries.

c. The *dorsalis indicis*, a branch of variable size, passes over the first interosseous muscle, along the radial side of the back of the index finger.

d. The *dorsales pollicis* are two small branches which arise from the radial opposite the head of the first metacarpal bone, and run along the back of the thumb, one on either side. They are often absent.

These dorsal interosseous arteries supply the extensor tendons and their sheaths, the interosseous muscles, and the skin on the back of the hand, and the first phalanges of the fingers.

Dissection. Remove the tendons from the back and from the palm of the hand; observe the deep palmar fascia which covers the interosseous muscles. It is attached to the ridges of the metacarpal bones, forms a distinct sheath for each interosseous muscle, and is continuous inferiorly with the transverse metacarpal ligament. On the back of the hand the interosseous muscles are covered by a thin fascia, which is attached to the adjacent borders of the metacarpal bones.

Transverse Metacarpal Ligament. This consists of strong bands of ligamentous fibres, which pass transversely between the distal extremities of the metacarpal bones. These bands are intimately united to the fibro-cartilaginous ligament of the metacarpal joints, and are of sufficient length to admit of a certain degree of movement between the ends of the metacarpal bones.

Dissection. Remove the fascia which covers the interosseous muscles, and separate the metacarpal bones by dividing the transverse metacarpal ligament. A *bursa* is frequently developed between their digital extremities.

Interosseous Muscles. These muscles, so named from their position, extend from the sides of the metacarpal bones to the bases of the first phalanges and the extensor tendons of the fingers. In each interosseous space (except the first, in which there is only an abductor) there are two muscles, one of which is an abductor, the other an adductor, of a finger. Thus there are *seven* in all; four of which, situated on the back of the hand, are called dorsal; the remainder, seen only in the palm, are called palmar.[1] They are all supplied by the ulnar nerve.

Dorsal Interossei. Each dorsal interosseous is a bipenniform muscle, and *arises* from the opposite sides of two contiguous metacarpal bones (Fig. 90). From this double origin the fibres converge to a tendon, which passes between the metacarpal joints of the finger, and is *inserted* into the side of the base of the first phalanx, and by a broad expansion into the extensor tendon on the back of the same finger.

The *first dorsal interosseous* muscle (*abductor indicis*) is larger than the others, and occupies the interval between the thumb and forefinger. It *arises* from the proximal half of the ulnar side of the first metacarpal bone, and from the entire length of the radial side of the second; between the two origins, the radial artery passes into the palm. Its fibres converge on either side to a tendon, which is *inserted* into the *radial* side of the first phalanx of the index finger and its extensor tendon.

[1] If we consider the adductor pollicis as a palmar interosseous muscle, there would be four palmar and four dorsal—all supplied by the ulnar nerve.

The *second dorsal interosseous* muscle occupies the second metacarpal space. It is *inserted* into the radial side of the first phalanx of the middle finger and its extensor tendon.

The *third* and *fourth*, occupying the corresponding metacarpal spaces, are *inserted*, the one into the ulnar side of the middle, the other into the ulnar side of the ring finger.

If a line be drawn longitudinally through the middle finger, as represented by the dotted line in Fig. 90, we find that all the dorsal interosseous muscles are abductors from that line; consequently, they separate the fingers from each other.

FIG. 90.

DIAGRAM OF THE FOUR DORSAL INTEROSSEI, DRAWING FROM THE MIDDLE LINE.

FIG. 91.

DIAGRAM OF THE THREE PALMAR INTEROSSEI AND THE ADDUCTOR POLLICIS, DRAWING TOWARD THE MIDDLE LINE.

Palmar Interosseous. It requires a careful examination to distinguish this set of muscles, because the dorsal muscles protrude with them into the palm. They are smaller than the dorsal, and each *arises* from the lateral surface of only one metacarpal bone—that, namely, connected with the finger into which the muscle is inserted (Fig. 91). They terminate in small tendons, which pass between the metacarpal joints of the fingers, and are *inserted*, like those of the dorsal muscles, into the sides of the first phalanges and the extensor tendons on the back of the fingers.

The *first palmar interosseous* muscle *arises* from the ulnar side of the second metacarpal bone, and is *inserted* into the ulnar side of the index finger. The *second* and *third* arise, the one from the radial side of the fourth, the other from the radial side of the fifth metacarpal bone, and are *inserted* into the same sides of the ring and little fingers.

The palmar interosseous muscles are all adductors to a line drawn through the middle finger (Fig. 91). They are, therefore, the opponents of the dorsal interosseous, and move the fingers toward each other.[1]

The palmar and dorsal interossei are supplied by filaments from the deep branch of the ulnar nerve.

DISSECTION OF THE LIGAMENTS.

Sterno-clavicular Joint. The inner end of the clavicle articulates with the comparatively small and shallow excavation on the upper and outer part of the sternum, and is an arthrodial joint. The security of the joint depends upon the great strength of its ligaments. There are two synovial membranes, and an intervening fibro-cartilage.

The *anterior sterno-clavicular ligament* (Fig. 92) consists of a strong, broad band of ligamentous fibres, which pass obliquely downward and inward over the front of the joint, from the inner end of the clavicle to the anterior surface of the sternum.

[1] The interossei, probably, also assist the flexors of the fingers when the latter are slightly flexed at their metacarpo-phalangeal joints. M. Duchenne believes that, in addition to their usually ascribed function of abduction, adduction and supplemental flexion at the metacarpo-phalangeal articulation, the interossei act as extensors of the second and third phalanges; the common extensor tendons acting only as extensors of the first phalanges. (*Physiologie des Mouvements etc.* 1867.) The action of the lumbricales in extending the second and third phalanges (even if they are not the chief factors of this movement) must not be lost sight of, for in a case, recorded in *St. Bartholomew's Hospital Reports*, 1881, in which the ulnar nerve had been divided a short distance above the wrist joint, the first phalanges of the ring and little fingers were bent (extended) upon their articulating metacarpal bones, the second and third phalanges being flexed at obtuse angles upon their proximal phalanges; the index and middle fingers being normal. I attribute this condition to paralysis of the two ulnar lumbricales and not to loss of power of the interossei. I have seen about a dozen instances of division of the ulnar nerve, and in all of them the same condition of the little and ring fingers has existed.

The *posterior sterno-clavicular ligament* extends over the back of the joint, its fibres passing downward and forward from the back of the clavicle to the back of the sternum in a similar manner to the anterior.

The *interclavicular ligament* connects the clavicles directly. It extends transversely along the notch of the sternum, and has a broad attachment to the upper border of each clavicle. Between the clavicles it is more or less attached to the sternum, so that it forms a curve with the concavity upward.

The three ligaments just described are so closely connected that, collectively, they form for the joint a complete fibrous capsule, of such strength that dislocation of it is rare.

The *costo-clavicular* or *rhomboid ligament* connects the clavicle to the cartilage of the first rib. It ascends obliquely outward and backward from the cartilage of the rib to a rough surface beneath the sternal end of the clavicle. Its use is to limit the elevation of the clavicle. There is such constant movement between the clavicle and the cartilage of the first rib that a well-marked *bursa* is commonly found between them.

Interarticular fibro-cartilage.—To see this, cut through the rhomboid, the anterior and posterior ligaments of the joint, and raise the clavicle. It is nearly circular in form, and thicker at the circumference than the centre, in which there is sometimes a perforation, and divides the articulation into two cavities. Inferiorly, it is attached to the cartilage of the first rib, close to the sternum; superiorly, to the upper part of the clavicle and the interclavicular ligament. Its circumference is inseparably connected with the anterior and posterior ligaments.[1]

The joint is provided with *two synovial membranes;* one between the articular surface of the sternum and the inner surface of the fibro-cartilage; the other between the articular surface of the clavicle and the outer surface of the fibro-cartilage.

[1] Interarticular fibro-cartilages (menisci) also exist in the following joints: acromio-clavicular, temporo-maxillary, knee, and wrist joints. Professor Humphry has shown that interarticular cartilages augment the variety of movements in a joint, permitting, for instance, that of rotation in the knee joint, in addition to that of extension and flexion, which otherwise would be the only possible ones.

interarticular fibro-cartilage is a structure highly elastic, t admitting of any stretching. It equalizes pressure, breaks , and also acts as a ligament, tending to prevent the clavicle eing driven inward toward the mesial line.

erve the relative form of the cartilaginous surfaces of the that of the sternum is slightly concave in the transverse nvex in the antero-posterior direction; that of the clavicle everse.

form of the articular surfaces and the ligaments of a joint known, it is easy to understand the movements of which it ble. The clavicle can be moved upon the sternum in a

FIG. 92.



DIAGRAM OF THE STERNO-CLAVICULAR LIGAMENTS.

nterclavicular ligament.
nterior sterno-clavicular ligament.
3. Costo-clavicular ligament.
4. Interarticular fibro-cartilage.

n either upward, downward, backward, or forward; it mits of circumduction. These movements, though limited sternum, are considerable at the apex of the shoulder.[1]

[1] ssor Humphry, in describing the movements of this joint, in his valuable n the Human Skeleton," says, "The movements attendant on elevation and n of the shoulder take place between the clavicle and the interarticular the bone rotating upon the ligament, on an axis drawn from before back- ough its own articular facet. When the shoulder is moved forward and , the clavicle, with the interarticular ligament, rolls to and fro on the surface of the sternum, revolving, with a slightly sliding movement, round rawn nearly vertically through the sternum. In the circumduction of the which is compounded of these two movements, the clavicle revolves upon rticular cartilage, and the latter, with the clavicle, rolls upon the sternum."

Scapulo-clavicular Joint. The outer end of the clavicle articulates with the acromion, and is connected by strong ligaments to the coracoid process of the scapula.

The clavicle and the acromion articulate with each other by two flat, oval cartilaginous surfaces, of which the planes slant inward, and the long diameters are in the antero-posterior direction. It is an arthrodial joint.

The *superior acromio-clavicular ligament*, a broad band of parallel ligamentous fibres, strengthened by the aponeurosis of the trapezius, extends from the upper surface of the acromion to the upper surface of the clavicle.

The *inferior acromio-clavicular ligament*, of less strength, extends along the under surface of the joint, from bone to bone.

FIG. 93.

ANTERIOR VIEW OF THE SCAPULO-CLAVICULAR LIGAMENTS, AND OF THE SHOULDER JOINT.

An *interarticular fibro-cartilage* is sometimes found in this joint; but it is incomplete, and seldom extends lower than the upper half. There is only one synovial membrane.

Coraco-clavicular ligament.—The clavicle is connected to the coracoid process of the scapula by two strong ligaments—the *conoid* and *trapezoid*, which, being continuous with each other, should be considered as one. The *trapezoid* ligament is the more anterior and external. Quadrilateral in shape, it arises from the back of the upper surface of the coracoid process, and ascends obliquely backward and outward to the oblique line on the under aspect of the clavicle, near its outer end. The *conoid* ligament,

riangular in form, is situated behind the trapezoid ligament, to the posterior border of which it is attached. It is fixed, at its apex, to the root of the coracoid process, ascends nearly vertically, and is attached by its base to the clavicle. The coraco-clavicular ligaments fix the scapula to the clavicle, and prevent undue rotation of the scapula. When the clavicle is fractured in the line of the attachment of the coraco-clavicular ligament, there is little or no displacement of the fractured ends, these being kept in place by the ligament.

Ligaments of the Scapula. These are two: the *coraco-acromial* or *triangular* ligament, attached by its apex to the tip of the acromion process, and by its base to the outer border of he coracoid process; it is separated from the upper part of the apsule of the shoulder joint by a large bursa; and the *transrse* or *coracoid* ligament, which passes across the supraapular notch, converting it into a foramen. The suprascapr vessels pass over the foramen, the suprascapular nerve ough it.

houlder Joint. The articular surface of the head of the humerus, forming rather more than one-third of phere, moves upon the shallow glenoid cavity of the scapula, ich is of an oval form, with the broader end downward, and long diameter nearly vertical. The security of the joint pends, not upon any mechanical contrivance of the bones, but on the great strength and number of the ligaments and tendons ich surround and are intimately connected with it. It is an arthrodial, or ball-and-socket joint.

To admit the free motion of the head of the humerus upon e glenoid cavity, it is requisite that the *capsular ligament* of the int be loose and capacious. Accordingly, the head of the bone, hen detached from its muscular connections, may be separated om the glenoid cavity to the extent of an inch or more, without ceration of the capsule. This explains the elongation of the m observed in some cases in which effusion takes place into e joint; also in cases of paralysis of the deltoid.

The *capsular ligament* is attached above, round the circumrence of the glenoid cavity; below, round the anatomical neck

of the humerus. It is strongest on its upper aspect, weakest and longest on its lower. It is strengthened on its upper and posterior part by the tendons of the supraspinatus, infraspinatus, and teres minor; its inner part is strengthened by the broad tendon of the subscapularis and the coraco-humeral ligament; its lower part, by the long head of the triceps.

Thus the circumference of the capsule is surrounded by tendons on every side, excepting a small space toward the axilla. If the humerus be raised, it will be found that the head of the bone rests upon this unprotected portion of the capsule, between the tendons of the subscapularis and the long head of the triceps; through this part of the capsule the head of the bone is first protruded in dislocations into the axilla.

At the upper and inner side of the joint, a small opening is observable in the capsular ligament, through which the tendon of the subscapularis passes, so that the synovial membrane of the joint communicates with the bursa under the tendon of this muscle. A second opening exists in the lower part of the front of the capsular ligament, where the tendon of the biceps emerges from the joint. A third opening occasionally exists between the joint, and a bursa under the tendon of the infraspinatus muscle.

The upper and inner surface of the capsule is strengthed by a strong band of ligamentous fibres, called the *coraco-humeral* or *accessory ligament.* It is attached to the root of the coracoid process, expands over the upper surface of the capsule, with which it is inseparably united, and, passing downward and outward, is attached to the greater tuberosity of the humerus.

Open the capsule to see the *tendon* of the *long head of the biceps.* It arises by a rounded tendon from the upper margin of the glenoid cavity, and is continuous with the glenoid ligament; becoming slightly flattened, it passes over the head of the humerus, descends through the groove between the two tuberosities, and, after piercing the capsular ligament of the shoulder joint, it passes along the bicipital groove, being retained in situ by an aponeurotic prolongation from the tendon of the pectoralis major. It is loose and movable within the joint. It acts like a strap, keeping down the head of the bone when the

m is raised by the deltoid, and then might be considered as king the part of a ligament of the joint.

The tendon of the biceps, strictly speaking, does not perforate e synovial membrane of the joint. It is enclosed in a tubular eath, which is reflected over it at its attachment to the glenoid ity, and accompanies it for two inches down the groove of the umerus. During the earlier part of fœtal life it is connected to e capsule by a fold of synovial membrane, which subsequently sappears.

The margin of the glenoid cavity of the scapula is surrounded a fibro-cartilaginous band of considerable thickness, called the *noid ligament.* This not only enlarges, but deepens the cavity. periorly, it is continuous on either side with the tendon of the ceps; inferiorly, with the tendon of the triceps; in the rest of circumference it is attached to the edge of the cavity.

The cartilage covering the head of the humerus is thicker at e centre than at the circumference. The reverse is the case in glenoid cavity.

The *synovial membrane* lining the under surface of the capsule reflected around the tendon of the biceps, and passes with it, in form of a cul-de-sac, down the bicipital groove. On the inner e of the joint it always communicates with the bursa beneath tendon of the subscapularis.

There is also a large bursa situated between the capsule and deltoid muscle, which does not communicate with the joint.

The muscles in relation with the joint are: above, the supra-natus; behind, the infraspinatus and teres minor; below, the g head of the triceps; internally, the subscapularis; and, inside joint, the long head of the biceps.

The shoulder joint is an enarthrodial joint, and has a more ex-sive range of motion than any other joint in the body; it is at mechanics call a universal joint. It is capable of motion ward and backward, of adduction, abduction, circumduction, l rotation. The various movements are limited chiefly by surrounding muscles and by atmospheric pressure, for the sule is so lax as to offer no obstacle to the freedom of move-nt in any direction. The amount of rotation which the head

of the humerus is capable of, is to the extent of a quarter of a circle.

The movements of which the shoulder joint is capable are effected by the following muscles: thus—

Extension is effected by the posterior fibres of the deltoid, latissimus dorsi, teres major, and (when the arm is raised) by the infraspinatus and teres minor.

Flexion, by the anterior fibres of the deltoid, coraco-brachialis, and the pectoralis major (slightly).

Abduction, by the deltoid and supraspinatus.

FIG. 94.

a. External lateral ligament.
b. Orbicular or annular ligament.
c. Part of internal lateral ligament.
d. Radius, removed from the annular ligament.

LIGAMENTS OF THE ELBOW JOINT.

Adduction, by the pectoralis major, latissimus dorsi, teres maj r, coraco-brachialis; and (when the arm is raised) by the su b- scapularis.

Rotation inward, by the subscapularis, latissimus dorsi, an d teres major.

Rotation outward, by the infraspinatus and the teres minor.

Elbow Joint. The elbow joint is a ginglymus or hinge join t. The larger sigmoid cavity of the ulna is adapte d to the trochlea upon the lower end of the humerus, admittin

of flexion and extension; while the shallow excavation upon
ead of the radius admits not only of flexion and extension,
f central rotation, upon the rounded articular eminence
ellum) of the humerus, and of peripheral rotation at the
ior radio-ulnar articulation.

e joint is secured in front and behind by anterior and
rior ligaments, and laterally by two strong lateral ligaments.
igament is attached to the head of the radius, otherwise its
ory movement would be impeded. The head is simply sur-
ded by a ligamentous collar, called the annular ligament,
n which it freely rolls in pronation and supination of the hand.

e *anterior ligament* consists of broad, thin ligamentous fibres,
hed above to the front of the humerus, above the coro-
fossa, below to the coronoid process of the ulna and to the
ular ligament, and continuous on each side with the lateral
ents. Some of the fibres cross each other at right angles.

e *posterior ligament* is composed of thin, loose fibres attached
to the margin of the olecranon fossa, below to the border of
lecranon, and spread over the posterior aspect of the joint.

e *internal lateral ligament* is triangular, and is divided into
portions, an anterior and a posterior. Its *anterior part* is
ed to the front of the internal condyle of the humerus; from
oint the fibres radiate, and are inserted along the inner mar-
f the coronoid process of the ulna. The *posterior part* is also
gular, and passes from the back part of the internal condyle
e inner border of the olecranon.

band of fibres extends transversely from the olecranon to
oronoid process, across a notch observable on the inner side
e sigmoid cavity; through this notch small vessels pass into
int.

e *external lateral ligament* is attached to the external condyle
humerus, and is in intimate connection with the common
n of the extensors. The fibres spread out as they descend,
re interwoven with the annular ligament surrounding the
of the radius.

e preceding ligaments, collectively, form a continuous cap-
or the joint.

Superior Radio-ulnar Articulation.

The *orbicular* or *annular ligament of the radius* forms about three-fourths of a ring. Its ends are attached to the anterior and posterior borders of the lesser sigmoid cavity of the ulna, and it is broader in the middle than at either end. Its lower border is straight; its upper border is convex, and connected with the anterior and external lateral ligaments. With this sigmoid cavity it forms a complete collar, which encircles the head, and part of the neck, of the radius. The lower part of the ring is narrower than the upper, the better to clasp the neck of the radius, and maintain it more accurately in position.

Synovial membrane of the elbow joint.—Open the joint by a transverse incision in front, and observe the relative adaptation of the cartilaginous surfaces of the bones. The synovial membrane lines the interior of the capsule, and forms a cul-de-sac between the head of the radius and its annular ligament. It is widest and loosest under the tendon of the triceps. Where the membrane is reflected from the bones upon the ligaments, there is more or less adipose tissue, particularly in the fossæ on the front and back part of the lower end of the humerus.

The only movements permitted between the humerus and the ulna are those of flexion and extension, both of which are limited by the ligaments and tendons in front of and behind the joint, and probably not by the coronoid and olecranon processes. The head of the radius is most in contact with the capitellum of the humerus during semiflexion and semipronation; and it is kept, by the strong orbicular ligament which surrounds the neck of the radius, from being dislocated forward by the biceps. The movement at the superior radio-ulnar articulation is that of rotation in the lesser sigmoid cavity of the ulna, forming an example of a lateral ginglymus or diarthrosis rotatoria. It is by this rotation of the head of the radius that the hand is carried through an extensive range of pronation and supination; for it is articulated only to the lower end of the radius, the ulna being excluded by the interarticular fibro-cartilage from taking any share in the movement at the wrist joint.

Interosseous Membrane.

This is an aponeurotic septum, stretched between the interosseous ridges of the radius and

which the chief purpose is to afford an increase of surface ttachment of muscles. The septum is deficient above, g about an inch below the tubercle of the radius, and thus free rotation of that bone. Its fibres extend obliquely rd from the radius to the ulna. It is perforated in its ird by the anterior interosseous vessels.

ame of *round* or *oblique ligament* is given to a thin band , which extends obliquely between the bones of the fore- direction contrary to those of the interosseous membrane. ached, superiorly, to the front surface of the ulna, near the de of the coronoid process; inferiorly, to the radius imme- below the tubercle. Between this ligament and the upper f the interosseous membrane is a triangular interval through he posterior interosseous artery passes to the back of the . A *bursa* intervenes between the oblique ligament and rtion of the tendon of the biceps. The use of this liga- to limit supination of the radius.

erior -ulnar ulation.

This joint is a lateral ginglymus, and is formed by the inner concave surface of the lower end of the radius rotating upon the convex head of the hich mechanism is essential to the pronation and supina- the hand. These corresponding surfaces are encrusted hin layer of cartilage, and are provided with a very loose l membrane. The surfaces are maintained in position by ior and a posterior radio-ulnar ligament, and a triangular rtilage.

nterior radio-ulnar ligament is a thin fasciculus extending ly inward from the anterior border of the sigmoid cavity adius to the head of the ulna.

posterior radio-ulnar ligament passes from the posterior of the sigmoid cavity to the posterior surface of the sty- cess of the ulna.

triangular fibro-cartilage between the radius and ulna is ncipal uniting medium between the bones. To see it, saw the bones of the forearm, and separate them by cutting the interosseous membrane, and opening the synovial ne of the joint between the lower ends. Thus a good

view is obtained of the fibro-cartilage which connects them (Fig. 95). It is triangular, and placed transversely at the lower end of the ulna, filling up the interval caused by the greater length of the radius. Its base is attached to the lower end of the radius; its apex to a depression at the root of the styloid process of the ulna. It is thin at the base and centre, thicker at the apex and sides. Its upper surface is in contact with the ulna, and covered by the synovial membrane of the inferior radio-ulnar joint; its

FIG. 95.



1. External lateral ligament.
2. Internal lateral ligament.
3. Interarticular fibro-cartilage between radius and ulna.
4. Interosseous ligaments.
5. Lateral ligaments of the intercarpal joint.

DIAGRAM OF THE LIGAMENTS AND SYNOVIAL MEMBRANES OF THE WRIST JOINT.

lower surface, forming a part of the wrist joint, is contiguous w the cuneiform bone. Its borders are connected with the anteri and posterior ligaments of the wrist. In some instances there an aperture in the centre.

When, from accident or disease, this fibro-cartilage gets d tached from the radius, the consequence is an abnormal proje tion of the lower end of the ulna.

The *synovial membrane* of this joint is distinct from that the wrist, except in the case of a perforation through the fibr

tilage. On account of its great looseness, necessary for the
: rotation of the radius, it is called *membrana sacciformis.*

The movement between the lower ends of the radius and ulna
lue to the rotation of the radius round the articular head of the
a, and is confined to rotation forward or *pronation*, and to rota-
n backward or *supination;* the extent of movement being limited
the anterior and posterior ligaments.

adio-carpal or Wrist Joint. This is an arthrodial joint, and is formed, above, by the lower end of the radius and the distal sur-
e of the triangular fibro-cartilage; below, by the scaphoid, semi-
ar and cuneiform bones; the two former articulate with the two
ets on the radius, the latter with the fibro-cartilage. The three
rpal bones form a convex surface which is received into the con-
vity formed by the radius and the cartilage. The joint is secured
an anterior, a posterior and two lateral ligaments.

The *external lateral ligament* extends from the tip of the styloid
ocess of the radius to the outer side of the scaphoid bone, to the
terior annular ligament, and to the trapezium.

The *internal lateral ligament* is round, and proceeds from the
remity of the styloid process of the ulna to the cuneiform
e. Another fasciculus is attached to the pisiform bone and
anterior annular ligament.

The *anterior ligament* consists of two or more broad bands of
mentous fibres, which extend from the lower end of the radius
he first row of carpal bones, except the pisiform.

The *posterior ligament*, weaker than the preceding, proceeds
n the posterior surface of the lower end of the radius, and
ttached to the posterior surfaces of the first row of carpal
es.

The *synovial membrane* lines the under surface of the triangular
o-cartilage at the end of the ulna, is reflected over the several
ments of the joint, and thence upon the first row of the carpal
es (Fig. 95).

This articulation allows of all the movements of enarthrodial
ts, except that of rotation; thus, it allows of flexion, extension,
uction, adduction and circumduction, so that it is, strictly
aking, only an arthrodial joint.

Carpal Joints. The bones of the carpus are arranged in two rows, an upper and a lower, adapted to each other so as to form between them a joint. The articulations may be best arranged in three sets, those between the carpal bones of the first row; between those of the second row; and the articulation of the two rows with each other; they are all examples of arthrodial joints.

a. The *first row of carpal bones* are connected together by two palmar, two dorsal and two interosseous ligaments.

The *dorsal* and *palmar transverse ligaments* proceed, on the dorsal and palmar aspects, from the scaphoid to the semilunar bone, and from the semilunar to the cuneiform bone; the dorsal being the stronger; the *interosseous* ligaments connect the semilunar with the bones on each side of it (Fig. 95).

The *pisiform bone* is articulated to the palmar surface of the cuneiform bone, to which it is united by a fibrous capsule. Inferiorly, it is attached by two strong ligaments, the one to the unciform bone, the other to the carpal end of the fifth metacarpal bone. This articulation has a distinct synovial membrane.

b. The *second row of carpal bones* is connected by three dorsal, three palmar and two interosseous ligaments.

The *dorsal* and *palmar ligaments* pass transversely from one to the other. There are usually *two interosseous ligaments*, one on either side of the os magnum; sometimes there is a third, between the trapezium and trapezoid bones; they are thicker and stronger than those of the upper row, and unite the bones more firmly together.

c. The first row of carpal bones is arranged in the form of an arch, so as to receive the corresponding surfaces of the os magnum and unciforme. External to the os magnum, the trapezium and trapezoid bones present a slightly concave surface, which articulates with the scaphoid. In this way a joint, admitting of flexion and extension only, is formed between the upper and lower row.

The *two rows of carpal bones* are connected together by palmar and dorsal ligaments, and by an external and an internal lateral ligament.

The *palmar ligaments* consist of strong ligamentous fibres,

which pass obliquely from the bones of the first to those of the second row.

The *dorsal ligaments* consist of oblique and transverse fibres which connect the dorsal surfaces of the bones of the upper with the lower row.

The *external lateral ligament*, very distinct, passes from the scaphoid to the trapezium; the *internal lateral ligament* from the cuneiform to the unciform.

Divide the ligaments to see the manner in which the carpal bones articulate with one another. Their surfaces are crusted with cartilage, and have a *common synovial membrane* which is very extensive and lines the distal surfaces of the scaphoid, semilunar and cuneiform bones; it then passes forward between the trapezium and trapezoid, the trapezoid and os magnum, the os magnum and the cuneiform to the articulations between the second row of carpal bones and the metacarpal bones of the four fingers (Fig. 95).

oint between the Trapezium and the First Meta-carpal Bone.

The trapezium presents a cartilaginous surface, convex in the transverse, and concave in the antero-posterior direction (*i. e.* saddle-shaped), which articulates with the cartilaginous surface the metacarpal bone of the thumb, concave and convex in the posite directions. This peculiar adaptation of the two surfaces mits the several movements of the thumb—viz., flexion, exten-n, abduction and adduction; consequently circumduction. It an arthrodial joint, but permits of such extensive movement, at it is described by some anatomists as one by "reciprocal ception." Thus we are enabled to oppose the thumb to all the gers, which is one of the great characteristics of the human nd. The joint is surrounded by a *capsular ligament* sufficitly loose to admit free motion, and stronger on the dorsal than the palmar aspect. The security of the joint is increased the muscles which surround it. It has a *separate synovial embrane.*

rpo-metacarpal Joints.

The metacarpal bones of the fingers are connected to the second row of the carpal bones by gaments upon their *palmar* and *dorsal* surfaces, and by interseous ligaments.

The *dorsal* ligaments are the stronger. The metacarpal bone of the forefinger has two, one from the trapezium, the other from the trapezoid bone. That of the middle finger has also two, proceeding from the os magnum and the os trapezoides. That of the ring finger has also two, proceeding from the os magnum and the unciform bone. That of the little finger has one only, from the unciform bone.

The *palmar* ligaments are arranged nearly upon a similar plan. The metacarpal bone of the forefinger has one from the trapezoid bone. That of the middle finger has three, proceeding from the trapezium, the os magnum and the unciform bone. Those of the ring and little fingers have each one, from the unciform bone.

Besides the preceding ligaments, there are some of considerable strength, called the *interosseous*. They proceed from the adjacent sides of the os magnum and the os unciforme, descend vertically, and are fixed into the radial side of the metacarpal bone of the middle and ring fingers (Fig. 95). This ligament occasionally isolates the synovial membrane of the two inner metacarpal bones from the common synovial membrane of the carpus.

Separate the metacarpal bones from the carpus, and observe the relative form of their continuous surfaces. The metacarpal bones of the fore and middle fingers are adapted to the carpus in such an angular manner as to be almost immovable. The metacarpal bone of the ring finger, having a plane articular surface with the unciform bone, admits of more motion. Still greater motion is permitted between the unciform and the metacarpal bone of the little finger, the articular surfaces of each being slightly concave and convex in opposite directions. The greater freedom of motion of the metacarpal bone of the little finger is essential to the expansion and contraction of the palm.

The *carpal extremities of the metacarpal bones of the fingers* are connected with each other by *palmar* and *dorsal transverse ligaments*. They are also connected by *interosseous* ligaments, which extend between the bones, immediately below their contiguous cartilaginous surfaces.

The *distal extremities* of these bones are loosely connected on their palmar aspect by the *transverse metacarpal* ligament.

Synovial Membranes of the Wrist. There are five, sometimes six, distinct synovial membranes, proper to the lower end of the radius, and the several bones of the carpus (see the diagram, p. 432) as follows:—

a. One between the lower end of the radius and the ulna.

b. One between the radius and the first row of carpal bones.

c. One between the trapezium and the metacarpal bone of the thumb.

d. One between the cuneiform and pisiform bones.

e. One between the first and second rows of carpal bones (the intercarpal joint). This extends to the metacarpal bones of the four inner fingers.

The interosseous ligament between the os magnum and ring finger occasionally shuts off the synovial membrane between the unciform and two inner metacarpal bones from the large intercarpal sac; thus making the sixth distinct synovial membrane.

First Joint of the Fingers. The first phalanx of the finger presents a shallow oval cavity, crusted with cartilage, with the broad diameter in the transverse direction, to articulate with the round cartilaginous head of the metacarpal bone, of which the articular surface is elongated in the antero-posterior direction, and of greater extent on its palmar than its dorsal aspect. This formation of parts permits flexion of the finger to a greater degree than extension; and also a slight lateral movement.

Each joint is provided with two strong *lateral* ligaments, and an anterior or *palmar* ligament.

The *lateral ligaments* arise from the tubercles on either side of each metacarpal bone, and, inclining slightly forward, are inserted into the sides of the base of the first phalanx of the finger.

The *anterior* (*glenoid*) *ligament* is a thick, compact, fibrous structure, which extends over the palmar surface of the joint between the lateral ligaments. Its distal end is firmly attached to the base of the first phalanx of the finger; its proximal end is loosely adherent to the rough surface above the head of the metacarpal bone. On either side it is inseparably connected with the lateral ligaments, so that with them it forms a strong capsule over the front and sides of the joint. Its superficial surface, firmly

connected with the transverse ligament, is slightly grooved for the play of the flexor tendons; its deep surface is adapted to cover the head of the metacarpal bone. Two sesamoid bones are found in the palmar ligament belonging to the joint between the metacarpal bone and the first phalanx of the thumb.

The palmar ligaments have a surgical importance for the following reason: In dislocation of the fingers, the facility of reduction mainly depends upon the extent to which the glenoid ligament is injured. If it be much torn, there is but little difficulty; if entire, the reduction may require much manipulation.

These joints are secured on their dorsal aspect by the extensor tendon, and the expansion proceeding from it on either side. Their synovial membranes are loose, especially beneath the extensor tendons.

Second and Last Joint of the Fingers. The corresponding articular surfaces of the phalanges of the fingers and thumb are so shaped as to form a hinge-joint, and, therefore, incapable of lateral movement. The ligaments connecting them are similar in every respect to those between the metacarpal bones and the first phalanges. The palmar ligament of the last joint of the thumb generally contains a sesamoid bone.

The wrist joint is a complex articulation, in which the seat of movement is partly in the radio-carpal, and partly in the inter-carpal articulation. Thus the hand at the radio-carpal joint is capable of extension (dorsi-flexion) and flexion, the latter being the most free; it is also capable of adduction (ulnar flexion) and of abduction (radial flexion) to a lesser extent. Between the carpal bones and carpo-metacarpal bones, the movement which takes place when the hand is pressed down so as to support the weight of the body, is that of separation of the anterior part of their apposed surfaces; undue separation being prevented by the interosseous and palmar ligaments. The articulation between the unciform and fourth and fifth metacarpals is not so firm as that between the other carpo-metacarpal bones, consequently there is greater freedom of motion forward, seen in deepening the palm and in shutting the hand. The movements at the metacarpo-

alangeal articulation are those of extension and flexion, of duction and abduction, the two latter being most marked in xtension of the finger. Between the thumb and trapezium all e movements of an enarthrodial joint exist, except that of otation; a little rotation probably takes place when the meta-arpal bone is flexed. In the interphalangeal and phalangeal oints, the only movements permitted are those of extension and lexion.

DISSECTION OF THE ABDOMEN.

urface Marking. The body should be sufficiently raised by placing blocks beneath the buttocks and the oulders, care being taken to have the chest higher than the lvis.

In the middle line, extending from the ensiform cartilage to the mphysis pubis, is a groove caused by the linea alba, the line of ion of the aponeurosis of the abdominal muscles. In this middle e, nearer the os pubis than the ensiform cartilage, is the umbilicus, ich corresponds, as a rule, with the body of the third lumbar ver-ra. The recti muscles can be distinguished on each side of the ddle line, and in well-developed subjects with little fat, the æ transversæ may be recognized, the lowest one being at the bilicus, the highest on a level with the ensiform cartilage, and third one midway between the two. On the outer border of rectus, about three inches from the middle line, is a concave e, the linea semilunaris, corresponding to the separation of the neurosis of the abdominal muscles to form the sheath of the tus. Above, and external to the spine of the os pubis, the ernal abdominal ring can be easily felt, the outer pillar being stronger; on it rests the spermatic cord passing to the testis. ssing from the spine of the os pubis to the anterior superior ne of the ilium, is a crescentic groove which indicates the line Poupart's ligament, and which can be felt as a firm and slightly rved cord; at about half an inch above the middle of the ament is situated the internal abdominal ring, which cannot, wever, be felt.

Arbitrary Division into Regions.

The abdomen is divided into arbitrary regions, that the situation of the viscera contained in it may be more easily described. For this purpose we draw the following lines: one horizontally across the abdomen, on a level with the cartilages of the ninth ribs; another on a level with the anterior superior spines of the ilia. These lines form the boundaries of three spaces, each of which is subdivided into three regions by a vertical line drawn on each side from the cartilage of the eighth rib to the middle of Poupart's ligament. Thus, there are a central and two lateral regions in each space. The central region of the upper space is termed the *epigastric;* the central one of the middle space is called the *umbilical* region; and the central of the inferior space the *hypogastric* region. The lateral regions of the spaces from above downward are termed the right and left *hypochondriac*, the right and left *lumbar*, and the right and left *inguinal* or *iliac* regions, respectively.

FIG. 96.



The viscera contained in these respective regions are as follows:—

In the *epigastric region* are, the left lobe of the liver, the round ligament of the liver, a small part of the right lobe, the middle and pyloric end of the stomach, the lobulus Spigelii, the pancreas, and the upper border of the transverse colon.

In the *umbilical region* are, the transverse colon, the great omentum and mesentery, the round ligament of the liver, the transverse portion of the duodenum, and part of the jejunum and ileum.

In the *hypogastric region* are, the small intestines, the urachus, the two obliterated hypogastric arteries, the bladder naturally in early life, and in the adult, if distended; and, lastly, the uterus in pregnancy.

n the *right hypochondrium* are, the right lobe of the liver, the e of the gall bladder, the descending duodenum, the hepatic xure of the colon, pancreas, supra-renal capsule, and the upper rt of the right kidney.

In the *left hypochondrium* are, the cardiac end of the stomach, e greater part of the spleen, the tail of the pancreas, the splenic xure of the colon, the supra-renal capsule, and the upper part the left kidney.

In the *right lumbar region* are, the ascending colon, the lower rt of the right kidney, and small intestines.

In the *left lumbar region* are, the descending colon, the lower t of the left kindey, omentum, and small intestines.

n the *right inguinal region* are, the cæcum and appendix miformis.

n the *left inguinal region* is the sigmoid flexure of the colon.

The abdomen should at this stage be distended with air by ans of a blowpipe inserted into the abdominal cavity through umbilicus, which, on the removal of the blowpipe, should be l with string to prevent escape of air.

Dissection. An incision is to be made from the ensiform cartilage to the os pubis, another from the ante- r superior spine of the ilium to a point midway between the bilicus and os pubis, and a third from the ensiform cartilage, nsversely outward toward the axilla, as far as the angles of the s. The skin should then be dissected from the subjacent pose and connective tissue, called the superficial fascia.

erficial Fascia. The subcutaneous tissue of the abdomen has the same general characters as that of other ts, and varies in thickness in different persons, according to amount of fat. At the lower part of the abdomen, it admits eparation into two layers, between which are found the sub- aneous blood vessels, the lymphatic glands, the ilio-inguinal ve, and the hypogastric branch of the ilio-hypogastric nerve.

Respecting the superficial layer, observe that it contains the and is continuous with the superficial fascia of the thigh, the otum and the perineum. The deeper layer is intimately con- ted with Poupart's ligament and the linea alba; but it is very

loosely continued over the spermatic cord and the scrotum, and becomes identified with the deep layer of the superficial fascia of the perineum. These points deserve attention, since they explain how urine, extravasated into the perineum and scrotum, readily makes its way over the spermatic cord on to the surface of the

FIG. 97.



SUPERFICIAL VESSELS AND GLANDS OF THE GROIN.

1. Saphenous opening of the fascia lata.
2. Saphena vein.
3. Superficial epigastric a.
4. Superficial circumflexa ilii a.
5. Superficial external pudic a.
6. External abdominal ring.
7. Fascia lata of the thigh.

abdomen; but from this it cannot travel down the thigh, on ac- count of the connection of the fascia with Poupart's ligament.

Superficial Blood Vessels and Lymphatic Glands.

Between the layers of the superficial fascia in the groin and upper part of the thigh, are seve al lymphatic glands and small blood vessels (Fig. 9). The glands are named, according to their sit a-

tion, inguinal or femoral. The *inguinal*, from three to four in number, are often small, and escape observation. They are of an oval form, with their long axis corresponding to the line of the crural arch (represented by the dark line in Fig. 97). They receive the superficial lymphatics from the lower part of the wall of the abdomen, from the integument of the scrotum, penis, perineum, anus and gluteal region, and are, therefore, generally affected in venereal disease. The lymphatics from the upper part of the abdominal parietes terminate in the lumbar glands.

The *superficial arteries* in the neighborhood arise from the femoral. One, the *superficial epigastric*, ascends over Poupart's ligament and ramifies over the lower part of the abdomen, as high as the umbilicus, inosculating with the deep epigastric and internal mammary arteries; another, the *superficial external pudic*, crosses the spermatic cord, and is distributed to the skin of the penis and scrotum, anastomosing with branches of the internal pudic; a third, the *superficial circumflexa ilii*, ramifies toward the spine of the ilium, and communicates with the deep circumflex iliac, the gluteal and external circumflex arteries. These subcutaneous arteries, the pudic especially, often occasion a free hemorrhage in the operation for strangulated inguinal and femoral hernia.

The corresponding *veins* join the internal saphena vein of the thigh. Under ordinary circumstances they do not appear in the living subject; but when any obstruction occurs in the inferior vena cava, they become enlarged and tortuous, and constitute the chief channels through which the blood would be returned from the lower limbs.[1]

Cutaneous Nerves. The skin of the abdomen is supplied with nerves after the same plan as the chest—namely, by lateral and anterior branches derived from the five or six lower intercostal nerves, as follows:—

a. The *lateral cutaneous nerves* come out between the digitations of the external oblique muscle, in company with small arteries, and divide, except the last, into anterior and posterior

[1] A cast in illustration of this is preserved in the Museum of St. Bartholomew's Hospital.

branches; the *anterior* pass forward as far as the rectus, and are distributed to the skin as far as its outer border, and to the digitations of the obliquus externus muscle; the *posterior*, small in size, run backward and supply the skin over the latissimus dorsi. The *lateral branch* of the *twelfth dorsal nerve* is larger than the others, and, piercing both the oblique muscles, passes over the crest of the ilium to the skin of the buttock, without dividing like the other nerves. The corresponding branch of the *first lumbar* has a similar distribution.

b. The *anterior cutaneous nerves* emerge with small arteries through the sheath of the rectus. They are not only smaller than the lateral nerves, but their number and place of exit is less regular.

c. The *ilio-hypogastric nerve* comes from the first lumbar nerve, pierces the transversalis at the iliac crest, and then divides into an iliac and hypogastric branch.

The *iliac branch* comes through both oblique muscles, and runs over the crest of the ilium, behind the last dorsal nerve, supplying the integument over the gluteal muscles.

The *hypogastric branch* lies at first between the transversalis and internal oblique; then, piercing the latter, it runs forward and comes through the aponeurosis of the external oblique, just above the external abdominal ring, and is distributed to the skin in the neighborhood.

d. The *ilio-inguinal nerve*, a branch also of the first lumbar nerve, is placed below the preceding nerve, with which it is connected near the crest of the ilium. It pierces the transversalis and internal oblique, runs down in the inguinal canal in front of the cord, and comes out through the external abdominal ring to be distributed to the skin of the inner part of the groin, to the scrotum and penis in the male, and to the labium pudendi in the female.

Dissection. The deep layer of the superficial fascia should now be removed from the external oblique, commencing at the fleshy portion of the muscle, and working in the course of its fibres. Care must be taken not to remove a part of its silvery aponeurosis, which is very thin, especially abo-

e digitations of this muscle with the serratus magnus and issimus dorsi must also be made out.

Muscles of the Abdominal Wall.

The abdominal muscles, three on each side, are arranged in strata, named, after the direction of their fibres, the external oblique, internal oblique, d transversalis. They terminate in front in strong aponeuroses, ranged so as to form a sheath for a broad muscle, called the ctus, which extends perpendicularly on each side the linea alba om the sternum to the os pubis.

ternal Oblique.

This muscle *arises* from the outer and lower surfaces of the eight or nine lower ribs, by as any pointed bundles, called *digitations*.[1] The upper five of these terdigitate with similar bundles of the serratus magnus; the ree lower correspond in like manner with the origin of the issimus dorsi; but they cannot be seen unless the body be ned on the side. The upper part of this muscle descends obuely forward, and terminates in the aponeurosis of the abdon; the lower proceeds almost perpendicularly from the last s, and is *inserted* into the anterior half of the st of the ilium.[2]

The *aponeurosis* of the external oblique increases in strength, adth and thickness, as it approaches the lower margin of the domen, this being the situation where the greater pressure of the cera requires the most effective support. Its tendinous fibres e the same direction as the muscle, and form by their decusion in the middle line the *linea alba*, which extends from the iform cartilage to the os pubis. Above, the aponeurosis benes much thinner, and is continued on to the pectoralis major l the ribs. The lowest fibres are strong, and form a thick der, called Poupart's ligament.

The posterior border of this muscle is fleshy and nearly vertical, l is overlapped in its upper half by the latissimus dorsi.

The upper digitations are attached to the ribs close to their cartilages; the er ones to the ribs some distance from the cartilages; the last to the apex of the lfth rib.

From its position and the direction of its fibres, it is manifested that the external ique represents, in the abdomen, the external intercostal muscles of the chest.

The aponeurosis is perforated by numerous cutaneous vessels and nerves for the supply of the skin and adjacent tissues; at its lower and inner aspect, close to the spine of the os pubis, there is a large oval opening called the external abdominal ring, transmitting in the male the spermatic cord, and in the female the round ligament.

Poupart's Ligament, or Crural Arch.

Along the line of junction of the abdomen with the thigh, the aponeurosis extends from the anterior superior spine of the ilium to the spine

Fig. 98.



1. External abdominal ring.
2. Gimbernat's ligament.
3. Poupart's ligament, or outer pillar of the ring.
4. Internal pillar of the ring.
5. Position of the internal ring, in dotted outline.

DIAGRAM OF POUPART'S LIGAMENT, OF THE APONEUROSIS OF THE EXTERNAL OBLIQUE, AND OF THE EXTERNAL ABDOMINAL RING.

of the os pubis, and forms an arch over the intermediate bony excavation (Fig. 98). This, which is termed the *crural arch*, or, more commonly, *Poupart's ligament*,[1] transmits the great vessels of the thigh, with muscles and nerves.

[1] This was first described by Fallopius, an Italian anatomist, in his *Observationes Anatomicæ*, published in 1561. It was subsequently described by Poupart in 1705 in the *Mém. de l' Acad. de Paris*, and is now commonly called "Poupart's ligament."

his ligament, when not separated from its fascial connections, s not run straight from the spine of the ilium to that of the os is, but is slightly curved, with its convexity toward the thigh. ove, and somewhat to the outer side of the spine of the os pubis, ituated an opening in the aponeurosis, called the *external ominal ring*. In the male it is a triangular opening about an h long, with its base at the os pubis, and will admit the sage of a finger; it transmits the spermatic cord. In the ale it is smaller, and transmits the round ligament of the rus. Its direction is downward and inward, and it is bounded ow by the crest of the os pubis, above by some arched fibres ich give strength to the apex of the opening, and on each e by the free margins of the aponeurosis, which are termed its umns or pillars. The *inner* or *upper pillar* (No. 4 in the gram) is thin, and is attached to the front of the os pubis, ussating with its fellow of the opposite side in front of the physis. The *outer* or *lower pillar* is thicker and stronger, and three attachments; one, into the spine of the os pubis—*part's ligament*—(No. 3); another, for three-quarters of an along the linea ilio-pectinea—*Gimbernat's ligament* (No. 2); third—or *triangular ligament*—consists of a few fibres which obliquely upward and inward beneath the spermatic cord the inner pillar as far as the linea alba, where they expand a triangular fascia in front of the conjoined tendon, and are inuous with the aponeurosis of the opposite side. At the r part of the aponeurosis of the external oblique, there are e arched fibres called *intercolumnar fibres*, which are strongest e the external ring. Their use is to strengthen the opening prevent the ring from enlarging.

ttached to the pillars of the external ring is a thin fascia, the *columnar* or *external spermatic fascia*, which is prolonged the spermatic cord and testis, and thus forms one of the rings of that organ.

he spermatic cord in its passage through the ring rests upon external pillar.

Dissection. The external oblique should now be detached from the ribs and the crest of the ilium, and

turned forward as far as this can be done without injuring its aponeurosis or the crural arch. In detaching this muscle from the ribs, care should be taken not to reflect with it the upper fibres of the rectus, and as the dissection is carried forward, the student should avoid injuring the thin aponeurosis of the internal oblique muscle. The second muscular stratum will thus be exposed and recognized by the difference in the direction of its fibres, which run upward and inward.

FIG. 99.



1. Conjoined tendon of internal oblique and transversalis.

2. Cremaster muscle passing down in loops over the cord.

DIAGRAM OF THE LOWER FIBRES OF THE INTERNAL OBLIQUE AND TRANSVERSALIS, WITH THE CREMASTER MUSCLE.

Internal Oblique. This is thinner than the last-named muscle, and *arises* by fleshy fibres from the outer half or more of Poupart's ligament, from the anterior two-thirds of the middle lip of the crest of the ilium, and from the posterior aponeurosis of the transversalis muscle (fascia lumborum). The fibres radiate from their origin, the anterior ones passing transversely forward, the posterior ones ascending nearly vertically. The fibres are *inserted* in the following manner: The anterior fibres (which arise from Poupart's ligament) pass inward, and arch over the spermatic cord, descending somewhat, to be

rted, in common with the tendon of the transversalis muscle, the crest of the os pubis, and for a short distance into linea ilio-pectinea immediately behind the external ring; middle fibres (which arise from the anterior iliac spine front of its crest) are directed transversely inward, to attached to an aponeurosis which passes to the linea alba; posterior fibres ascend nearly vertically to be attached into cartilages of the four lower ribs, and are continuous with internal intercostal muscles, which they represent in the lomen.

The aponeurosis of the internal oblique is the broad, expanded dinous tissue into which the muscle is anteriorly attached, and ontinued to the middle line, where its fibres join those of the posite side at the linea alba. It extends from the chest to the pubis, and its fibres run in the same direction as the muscle. the outer border of the rectus it splits into two layers—an erior, which passes in front of the rectus in conjunction with aponeurosis of the external oblique; and a posterior, which, common with the aponeurosis of the transversalis, passes ind the rectus. The point of division of the aponeurosis pre- ts a semilunar line extending from the os pubis to the cartilage he eighth rib. This is called the *linea semilunaris*, through ch a hernia occasionally protrudes. The two layers thus n a sheath for the rectus, which, except at the lower fourth nd, is complete. Midway between the umbilicus and the os s, the aponeuroses of all the three muscles pass in front of rectus, so that posteriorly in this situation it has no sheath. lower free border of the posterior part of the sheath—the *lunar fold of Douglas*—marks the situation where the deep astric artery enters the substance of the rectus.

Cremaster Muscle. The *cremaster* is a thin, pale muscle, or the reverse, according to the condition of the subject. best to regard it as a detachment of the lowest fibres of the rnal oblique, which arises from the middle of Poupart's liga- t. Passing along the outer side of the spermatic cord, the s descend with it through the external ring, and then arch up ront of the cord to the spine and crest of the os pubis, form-

ing loops of different lengths, some reaching only as low as the external ring, others lower still, whilst the lowest spread out over the tunica vaginalis of the testis. The muscular fibres are frayed out, being connected by loose cellular tissue, and form a covering for the testis, called the *cremasteric fascia*. This muscle is absent in the female. Its nerve comes from the genital branch of the genito-crural, and its artery (*cremasteric*) from the deep epigastric.

Dissection. The student should not now further dissect the structures on the left side, so that they may be left till a future period, for the complete demonstration of the parts concerned in inguinal hernia. On the right side, the internal oblique should be detached from the ribs and the crest of the ilium, and turned forward, without disturbing that portion of it connected with the crural arch. To avoid cutting away any part of the transversalis in reflecting the internal oblique, dissect near the crest of the ilium, and search for an artery which runs between these muscles, and may be followed as a guide. This artery, called the *deep circumflexa ilii*, is a branch of the external iliac, and supplies the abdominal muscles. Beneath the internal oblique the continuations of the intercostal nerves and vessels are brought into view, as are also the last dorsal, the ilio-hypogastric, and ilio-inguinal nerves near the crest of the ilium. These should be preserved.

The internal oblique is in relation, on its deeper surface, with the transversalis abdominis, the fascia transversalis, and with the spermatic cord near Poupart's ligament; on its inner side, at the division of its aponeurosis, with the outer border of the rectus; below, it forms the upper arched boundary of the inguinal canal.

Transversalis Abdominis. This muscle *arises* by fleshy fibres from the outer third of Poupart's ligament, from the anterior two-thirds of the inner lip of the crest of the ilium, from a strong fascia attached to the transverse processes of the lumbar vertebræ, and, lastly, from the inner surfaces of the six or seven lower costal cartilages, by digitations which correspond with those of the diaphragm. From this origin the fibres pass horizontally forward, and terminate anteriorly in a broad aponeurosis attached to its fellow at the linea alba. Some of its lower fibres arch down-

d are inserted with some fibres of the internal oblique, by
f a conjoined tendon, into the crest of the os pubis and
ilio-pectinea.
aponeurosis into which the fibres are inserted is broader
an above, and forms part of the posterior sheath of the
xcepting in the lower fourth, where it passes entirely in

dissection of the back we have fully described the lumbar
owing that it divides into three layers: the posterior
tached to the tips of the spinous processes, gives attach-
the internal oblique, and is continuous with the aponeu-

FIG. 100.



SE SECTION THROUGH THE ABDOMINAL MUSCLES, TO SHOW THE FORMATION OF THE ATH OF THE RECTUS, THE QUADRATUS LUMBORUM, AND THE ERECTOR SPINÆ.

the serratus posticus inferior and latissimus dorsi; the
ayer, attached to the tips of the transverse processes, and
rior layer—very thin—attached to the anterior aspect of
s of the transverse processes. Between the anterior and
layer is the quadratus lumborum; between the middle
erior, the erector spinæ.

tus minis. This long muscle is situated vertically in front of the abdomen, and is enclosed in a sheath formed aponeuroses of the lateral muscles of the abdomen, and d from its fellow by the linea alba. To expose it, there-

fore, slit up the middle of the sheath, and reflect the two halves. It *arises* by two tendons, the *inner* and *smaller* of which is attached to the front of the symphysis, the *outer* to the crest of the os pubis. As the fibres pass up, the muscle becomes broader and thinner, and is inserted into the fifth, sixth and seventh costal cartilages. Notice the tendinous intersections across the muscle, called *lineæ transversæ*, which are incomplete repetitions of the ribs in the wall of the abdomen.[1] Their number varies from three to five, but there are always more above than below the umbilicus. These tendinous intersections adhere closely to the sheath in front, but not behind; consequently, pus formed between the front of the rectus and its sheath would be confined by two intersections; not so on the back of the muscle, where pus might travel down the entire length of it. There is one intersection on a level with the umbilicus, one on a level with the ensiform cartilage, and an intermediate one between these two.

The *sheath of the rectus* consists, in front, of the aponeurosis of the external oblique, and half the thickness of that of the internal oblique; while the back of the sheath comprises the aponeurosis of the tranversalis, and half that of the internal oblique (Fig. 100). This, however, applies only to the upper three-fourths of the muscle; the lower fourth has no sheath behind, since all the aponeuroses pass in front of it; the only structure in contact with the muscle in this part is the fascia transversalis.

Pyramidalis. This small triangular muscle is situated near the os pubis, close to the linea alba, and has a sheath of its own. It *arises* by tendinous fibres from the front part of the os pubis and anterior pubic ligament in front of the rectus, and terminates in the linea alba about midway between the os pubis and the umbilicus. It is often absent on one or even both sides.

Linea alba.—The aponeuroses of the abdominal muscles decussate along the middle line and form a white fibrous band, extending from the ensiform cartilage to the os pubis. This is the *Linea alba:* it is the fibrous continuation of the sternum, and is broader above than below. A little lower than the middle is a large

[1] Some animals, *e. g.* the crocodile, have bony abdominal ribs.

erture in it—the *umbilicus*—through which a hernial protru-
n not infrequently takes place. It is in relation, behind, with
fascia transversalis, the urachus, and the bladder when
tended.

The linea alba, being the thinnest part of the abdomen, and
e from large blood vessels, is chosen as a safe line for tapping
dropsy, for puncturing the bladder in retention of urine, and
ovariotomy.

Lineæ semilunares.—These are the two slightly curved lines,
the front of the abdomen, corresponding with the outer mar-
s of the two recti muscles. They are formed by the junction
he aponeuroses of the lateral muscles.

The abdominal muscles serve many important purposes:—

Functions of the Abdominal Muscles.

1st. *In tranquil expiration* they push the diaphragm upward by gentle pressure on the abdominal viscera.

In forcible expiration the same process takes place, but with
ater energy. This is variously exemplified in coughing, sneezing
laughing.

2d. *In vomiting*, the diaphragm being fixed[1] by the closure
he glottis, the abdominal muscles contract, and assist the
nach to expel its contents.

3d. In conjunction with the contracted diaphragm, they assist
muscular walls of the bladder and rectum in the expulsion of
e and fæces, and the action of the uterus in parturition. They
rcise a gentle pressure and support on the abdominal viscera,
shield them from injury by strongly contracting when a blow
nticipated.

4th. They are movers of the trunk in various ways. For
mple, the right external oblique acting with the left internal
que will rotate the chest toward the left side, as in mowing,
vice versâ.

The rectus is chiefly concerned in raising the body from the
izontal position, as any one may ascertain by placing his hand
the abdomen while rising from the ground. The pyramidalis
kes the linea alba tense.

[1] By the term 'fixed,' it is meant that the diaphragm forms a resisting surface.

Dissection. By dividing the rectus transversely near the umbilicus, and raising it from its position, we have a complete view of the manner in which the sheath is formed; we observe, too, that this is absent behind the lower fourth of the muscle. Ramifying in the substance of the muscle is a large artery, called the *deep epigastric*, a branch of the external iliac; also the continuation of the internal mammary, which descends from the subclavian.

Nerves of the Abdominal Wall. These nerves are the anterior divisions of the six lower intercostal nerves, and of the first lumbar. They have the same general course and distribution, and are accompanied by small arteries derived from the intercostal and first lumbar arteries.

The *intercostal* or *abdominal* nerves come forward beneath the anterior extremities of the intercostal spaces, and then run between the internal oblique and transversalis, toward the edge of the rectus, which they enter, small twigs coming through it at the middle line to supply the skin. Each gives off a *lateral cutaneous branch*, which perforates the external intercostal and external oblique muscles, and divides, into an *anterior* branch—distributed to the skin and superficial fascia as far as the rectus, and into a *posterior* branch, smaller than the anterior, which supplies the skin over the latissimus dorsi.

The *last dorsal nerve*, larger than the other intercostals, is continued forward beneath the last rib, lying on the quadratus lumborum, and then piercing the transversalis aponeurosis, runs between this muscle and the internal oblique, and is finally distributed like the preceding nerves. Its *lateral cutaneous branch* is very large, and descends over the crest of the ilium.

The *ilio-hypogastric nerve*, a branch of the first lumbar, emerges from the outer border of the psoas, and then runs obliquely across the quadratus lumborum as far as the iliac crest, where it perforates the transversalis muscle and divides into an iliac and a hypogastric branch (p. 444).

The *ilio-inguinal nerve*, smaller than the former, and like it a branch of the first lumbar, runs along the iliac crest after piercing the psoas, and is here connected with the ilio-hypogastric. It

comes through the transversalis near the front of the ilium, and, after piercing the internal oblique, runs in front of the cord in the inguinal canal. Its cutaneous distribution has been described (p. 444).

Dissection. The transversalis muscle must now be reflected with the rectus by incisions similar to those for the reflection of the external oblique, when a thin, delicate fascia behind, the *fascia transversalis*, will be exposed. The dissection should take place from below upward, as the muscle is less intimately connected with the fascia below than it is higher up.

FIG. 101.

nternal abdominal ring.
osition of the external abdominal ring in dotted outline.
pigastric a. in dotted outline.

4. 4. Sheath of the femoral vessels, continued from the fascia transversalis.
5. Femoral a.
6. Profunda a.
7. Saphena v.
8, 8. Fascia transversalis.

DIAGRAM OF THE FASCIA TRANSVERSALIS, SEEN FROM THE FRONT.

Fascia Transversalis This fascia separates the transversalis muscle from the peritoneum, and is so called because it s in contact with the posterior surface of the muscle. It is mparatively thin, superiorly, where it is continuous with the scia on the under surface of the diaphragm. Inferiorly, it is ick and strong, and is attached to the crest of the ilium and to upart's ligament, where it is strengthened by fibres from the oneurosis of the transversalis; it becomes continuous with the

fascia covering the iliacus muscle (iliac fascia), and below with the pelvic fascia. About the middle of Poupart's ligament it sends a funnel-shaped prolongation downward into the thigh, forming the anterior part of the sheath of the femoral vessels. Internally, it is connected with the margin of the rectus, to the lower margin of the conjoined tendon, to the os pubis, and to the pectineal line. This fascia is strongest just behind the external abdominal ring, and, but for it and the conjoined tendon, there would be a direct opening into the abdominal cavity through the external ring. The outer half of the fascia is very firmly connected to Poupart's ligament and to the fascia iliaca; but the inner half is loosely connected with the crural arch, and passes down under it, as before stated, over the femoral vessels into the thigh, and forms the front of what is termed the crural sheath.

Internal Abdominal Ring. The opening in the fascia transversalis through which the spermatic cord passes is called the *internal abdominal ring* (or the inner aperture of the inguinal canal). It corresponds to a point midway between the anterior superior spine of the ilium and the spine of the os pubis, and about half an inch above Poupart's ligament. It is oval, with the long diameter nearly vertical; it is bounded above by the arched fibres of the transversalis muscle, and on the inner side by the deep epigastric vessels. Its margin is well defined on the inner, but not on the outer side, and from its border is continued forward a funnel-shaped prolongation over the spermatic cord, which passes through the ring. This covering, thin and delicate, is termed the *infundibuliform fascia*. (This is not seen in the diagram.) Close by the inner border of the internal ring, the deep epigastric artery ascends to enter the substance of the rectus.

Arteries of the Abdominal Walls. The abdominal walls are supplied with blood derived from the intercostal, lumbar, and deep circumflex iliac arteries, and, in front, also, from the internal mammary and deep epigastric arteries.

The *intercostal arteries* come from the descending thoracic aorta, and, like their accompanying nerves, enter the abdominal wall between the transversalis and the internal oblique muscles

y anastomose with the internal mammary, deep epigastric and
bar arteries.

he *lumbar artery* accompanies the last dorsal nerve.

he *internal mammary artery* divides, between the cartilages
e sixth and seventh ribs, into the musculo-phrenic and the
rior epigastric arteries. The superior epigastric artery
ends behind the cartilage of the seventh rib, and piercing the
th of the rectus, enters that muscle, supplying it and anasto-
ing with the deep epigastric.

he *musculo-phrenic artery* descends obliquely outward, behind
cartilages of the false ribs, and pierces the diaphragm about
ninth rib; it is then continued along the last intercostal space,
terminates in branches for the supply of the abdominal walls.

he *deep epigastric artery* arises from the external iliac, just
re this vessel passes under the crural arch to take the name of
oral. It ascends inward between the fascia transversalis and
peritoneum, forms a gentle curve on the *inner side* of the
rnal abdominal ring, and consequently on the inner side of
spermatic cord, and then enters the rectus muscle just below
fold of Douglas, which is the lower arched edge of the poste-
sheath of the rectus.

he artery runs in the substance of the rectus parallel with the
alba, and inosculates with the superior epigastric branch of
nternal mammary artery. It is accompanied by two veins, of
h the larger is on its inner side; these terminate in a single
k in the external iliac vein.

ıe deep epigastric gives off the following branches :—

ıe *pubic* is the most important branch. It runs inward, behind
rural arch, toward the os pubis, behind which it anastomoses
the pubic branch of the obturator. Sometimes the obturator
y is absent or small, in which case the pubic branch of the
astric enlarges and takes the place of the absent vessel. It
es its chief practical interest from the fact that it is liable to
ounded in dividing the stricture in femoral hernia.[1] But its

[1] here is a preparation in the Museum of St. Bartholomew's Hospital quite to
oint. The patient had profuse hemorrhage, which commenced five hours after
eration. He died from peritonitis.

size varies in different subjects, and is sometimes so small as to escape observation. The second branch is the *cremasteric*. It supplies the coverings of the cord, but chiefly the cremaster muscle. After giving off other unnamed *muscular branches*, some of which perforate the muscle to supply the skin, the main trunk terminates in the rectus by inosculations with the internal mammary.

Deep Circumflexa Iliі. The *deep circumflexa ilii* artery is a branch of the external iliac, just above the crural arch; it runs upward and outward, behind and parallel with Poupart's ligament, and at the middle of the crest of the ilium pierces the transversalis muscle, and, running in the same direction, lies between the transversalis and internal oblique. It anastomoses with the ilio-lumbar artery, and sends small muscular branches, which run upward, and communicate with the epigastric and the lumbar arteries.

The circumflex iliac veins join to form a single *vein*, which crosses the external iliac artery, and opens into the external iliac vein.

Dissection. To see that part of the peritoneum concerned in inguinal hernia, the fascia transversalis must be removed by incisions similar to those recommended before. The fascia is easily separable from the peritoneum which is situated immediately behind it, owing to the presence of more or less fat—*subperitoneal fat*. The peritoneum at the inner ring presents a well-marked depression, which varies, however, considerably; in some being scarcely visible; in others, being continued downward into the inguinal canal, in the form of a pouch. In some instances, a communication is found between the general cavity of the peritoneum and the tunica vaginalis testis.

Inguinal Canal. Having examined the several strata through which the spermatic cord passes, replace them in their natural position, and examine the inguinal canal as a whole. Its direction is obliquely downward and inward. Its length in a well-formed adult male is from one and a half to two inches. It commences at the inner ring, and terminates at the external abdominal ring. It is bounded *in front* by skin, superficial and deep fasciæ, by the aponeurosis of the external oblique, and

ternally by a small portion of the internal oblique; *behind*, by e fascia transversalis, by the conjoined tendon of the internal blique and transversalis, and by the triangular ligament; *above*, y the lower fleshy fibres of the internal oblique and transversalis; *low*, by the junction of the fascia transversalis with the crural rch.

Spermatic Cord. This round cord extends from the testis to the internal abdominal ring, and consists of numerous tructures, connected together by delicate areolar tissue, and is urrounded by the different strata from the abdominal muscles, hich are pushed down in the descent of the testis into the crotum in fœtal life. The cord lies in the inguinal canal; at the uter ring it rests on the outer pillar, and at the inner ring the ifferent constituents of the cord separate from each other.

The *arteries of the cord* are derived from the spermatic artery om the abdominal aorta, the deferential artery from the supe- or vesical, and the cremasteric branch from the deep epigastric.

The *veins of the cord* are chiefly the spermatic; they form on e cord a plexus of veins—the *pampiniform plexus*—and, ssing up in front of the cord, open on the right side into the erior vena cava; and on the left side, into the left renal vein.

The *lymphatics* pass into the lumbar glands.

The *nerves* are derived from the renal, aortic and hypogastric xuses. In front of the cord is the ilio-inguinal nerve, and ind it is the genital branch of the genito-crural nerve.

The cord, as will presently be described, receives coverings from external oblique, from the internal oblique and transversalis, l from the fascia transversalis.

The *vas deferens*, the excretory duct of the testis, passes ough the inguinal canal, being placed behind the other con- uents of the cord, and, after passing through the inner ring, ves round the epigastric artery in its descent into the pelvis.

and Ligament. In the female there is a round cord occupying the inguinal canal—the *round ligament*—which is , external to the outer ring, in the subcutaneous tissues of the ium majus. Its coverings are the same as those of the male, cepting the cremasteric fascia.

Deep Crural Arch. This structure, which is apparently a thickening of the fascia transversalis, has more to do with femoral hernia, and its description will, therefore, be deferred until this form of hernia is considered, in the dissection of the thigh.

DISSECTION OF THE PARTS CONCERNED IN INGUINAL HERNIA.

Dissection. The student has now completed the dissection on the right side, and, having mastered the general anatomy of this region, he may pass on to the special consideration of the anatomy of inguinal hernia. If the instructions before given have been duly observed, the left side is available for this purpose; and, although it may be well that the dissector (especially for the first time) should have the advantage of the parts being made clear by his demonstrator, there is no real difficulty in making out the different layers which constitute the coverings of a hernia, or the various parts through which a hernia travels.

It will be remembered that on the left side the dissection has been carried as far as the exposure of the internal oblique and cremaster. An incision must now be made through the *external oblique*, from a point midway between the umbilicus and the os pubis, transversely outward to the anterior superior iliac spine, and another from the same point downward in the middle line through the linea alba as far as the symphysis pubis. When this flap has been turned downward, the dissector will see that it is aponeurotic, and he can take the opportunity of making out the external abdominal ring and the external spermatic fascia which is prolonged downward from the pillars of the ring, as also the intercolumnar bands which strengthen the upper part of the ring.

The *internal oblique* now comes into view, and its origin from Poupart's ligament must be carefully made out, together with the cremaster muscle, which loops in front of the cord, and which can be seen coming under the arched fibres of this muscle. Next, the internal oblique should be reflected from Poupart's ligament and the iliac crest by a transverse incision to the extent of that through the external oblique. The flap should then be reflected inward, care being taken not to reflect with it the sub-

acent muscle (transversalis); this may be prevented by looking or a branch of the deep circumflex iliac artery, which runs along he crest of the ilium between these muscles. The turning back of this muscle exposes the transversalis, which in its turn will be reflected inward in the same manner. The inner parts of both these muscles are intimately connected by a common tendon, called the *conjoined tendon*, inserted in front of the rectus into the crest of the os pubis and the pectineal line. Observe that this tendon lies immediately behind the external abdominal ring, and that it varies in thickness in different subjects. The arching over of the lower fibres of the internal oblique and transversalis, so as to form the upper boundary of the inguinal canal, are now well seen.

Reflect the *transversalis* exactly in the same way as the internal oblique, when the fascia transversalis comes into view, presenting he funnel-shaped prolongation of fascia—*infundibuliform*—which s continued over the cord and testis.

The *transversalis fascia* should now be detached from the ubjacent peritoneum, in front of which is more or less fat (*subperitoneal*), and turned down, when the internal abdominal ring comes apparent, with its well defined inner margin.

During the reflection of these successive muscles, the student ill have been able to recognize the strata which are prolonged om them; viz.: from the external oblique is derived the *spermatic fascia;* from the internal oblique and transversalis is derived the *cremasteric fascia;* and from the fascia transversalis is olonged the transversalis fascia, which here takes the name of e *infundibuliform fascia.*

The extent and boundaries of the inguinal canal and the relan of the epigastric artery to the inner ring can now be clearly fined; and if the dissector passes his little finger into the ternal abdominal ring, down the canal and out through the ternal ring, he will easily see that it carries before it the three rata previously described, which constitute not only the coverngs of the cord, but likewise the coverings[1] of an oblique inguil hernia, when this exists.

[1] In the case of a hernia there is necessarily in addition a covering of subperitoal fat, and of peritoneum which forms the sac.

Practical Application. The testis, originally formed in the loins, passes, about the eighth month of fœtal life, from the abdomen into the scrotum, through an oblique canal in the wall of the abdomen, called the inguinal canal. A portion of peritoneum is pouched out before the descending testis, and constitutes the tunica vaginalis testis. The blood vessels, nerves, and vas deferens are drawn down with the testis, and constitute the spermatic cord. The inguinal canal runs obliquely through the abdominal wall, that it may the better resist the protrusion of intestine.

The wall of the abdomen, as previously stated, is composed of various strata, and the testis and cord in their passage through each stratum derive from each a covering similar in structure to the stratum itself. Of these strata there are three: the first, proceeding from within outward, is the *fascial stratum*, derived from the fascia transversalis; the second is the *muscular stratum* (cremasteric), from the internal oblique and transversalis muscles; the third is the *aponeurotic stratum*, from the external oblique.

The passage of the testis through the lower part of the abdominal parietes (inguinal canal) occasions, at this part of the belly, a natural weakness which, associated with other conditions, favors the protrusion of intestine in this situation.

A protrusion of intestine through any part of the inguinal canal is called an *inguinal hernia*, of which two chief varieties exist, the *indirect* or *oblique*, and the *direct;* the former protruding to the outer side of the deep epigastric artery; the latter coming out to the inner side of the artery.

Oblique Inguinal Hernia. The most common form of inguinal hernia is that in which a portion of intestine protrudes first through the internal ring, then, traversing the inguinal canal, emerges through the external ring, and thence may descend into the scrotum. This variety is called an *oblique inguinal hernia.*[1] If the intestine stops within the inguinal canal, it is called an *incomplete inguinal hernia;* if, however, the protrusion has emerged

[1] A hernia is sometimes called *external* or *internal*, according to the relation of the protrusion to the deep epigastric artery; thus, an oblique inguinal hernia which first protrudes through the inner ring is called an *external hernia*, and *vice versâ*.

through the external ring, it is called a *complete inguinal hernia;* and, lastly, if it descends into the scrotum, it is called a *scrotal hernia.*

Coverings of an Oblique Inguinal Hernia. A *complete* oblique inguinal hernia, passing as it does through the same structures as the testis did in fœtal life, receives the same coverings as that gland; they are:—

1. The *skin* and the *superficial fascia.*
2. The *intercolumnar fascia,* derived from the external oblique.
3. The *cremaster,* derived from the internal oblique and transversalis.[1]
4. The *infundibuliform fascia,* derived from the fascia transversalis.
5. The *subperitoneal fat,* and the *peritoneum* which constitutes the sac.

An *incomplete* oblique inguinal hernia is covered by—

1. The *skin* and *superficial fascia.*
2. The *aponeurosis* of the *external oblique.*
3. The *cremaster.*
4. The *infundibuliform fascia.*
5. The *subperitoneal fat* and the *peritoneum.*

irect Inguinal Hernia. The intestine, however, does not always escape through the internal ring. Sometimes it pro- des *internal* to the deep epigastric artery, through a triangular ak place, *Hesselbach's triangle,* bounded on the inner side by the tus, on the outer side by the deep epigastric artery, and below Poupart's ligament. This space is relatively weak, having in nt of it only the fascia transversalis and the conjoined tendon the internal oblique and transversalis; moreover, it is situated mediately behind the external abdominal ring. A portion of estine protruding through this triangle comes directly forward rough the external ring, and the hernia is then called a *direct* uinal hernia.

Coverings of a Direct Inguinal Hernia. A *direct* inguinal hernia protrudes immediately on the inner side of the epigastric artery through the external ring; and its course forward is

[1] The cremaster muscle is absent in the female.

mainly prevented by the resistance of the conjoined tendon.[1] This hernia is covered by—

1. The *skin* and *superficial fascia.*
2. The *intercolumnar fascia.*
3. The *conjoined tendon* of the internal oblique and transversalis.
4. The *fascia transversalis.*
5. The *subperitoneal fat* and the *peritoneum.*

A direct hernia, as has been said, emerges through Hesselbach's triangle, and if this triangle be viewed from its deeper aspect, it will be seen that, usually, the obliterated hypogastric artery cor-

FIG. 102.



VARIETIES OF CONGENITAL INGUINAL HERNIÆ, CONSEQUENT UPON SOME DEFECT IN THE DEVELOPMENT OF THE SPERMATIC PORTION OF THE TUNICA VAGINALIS. THE ARROWS MARK THE PROTRUSION.

1. Hernia in the tunica vaginalis testis.
2. Hernia in the funicular portion of the tunica vaginalis.
3. Infantile hernia.
4. Encysted hernia.

responds in its course to the deep epigastric artery. Occasionally, however, it is placed further inward, so that it divides the triangle into two smaller ones, in both of which there is a shallow pouching of peritoneum. Now, if a direct hernia protrudes through the outer of these two smaller triangles, it descends through the inguinal canal, and thus will have all the coverings of an ordinary oblique inguinal hernia.

In almost all cases, the immediate investment of the intesti[illegible]e is the parietal layer of the peritoneum. This constitutes the [illegible]ac

[1] In our experience the weakness of the conjoined tendon is, anatomically spe[illegible]k-ing, the determining cause of this form of hernia.

the hernia. The opening of the sac, communicating with the domen, is called its *mouth;* then comes the narrow, constricted rtion, or *neck;* and lastly, the *body*, or expanded part of the sac.

Congenital Herniæ.

Owing to the comparatively late descent of the testis in fœtal life, it frequently happens that ther no closure, or only a partial closure, takes place in the aginal portion of the tunica vaginalis. Under these conditions, hen a protrusion takes place, the intestine does not push forward sac derived from the parietal layer of the peritoneum, but it es in a sac formed by the tunica vaginalis, which still commu- icates with the peritoneal cavity. These herniæ are always blique, and are termed *congenital*.[1] There are four varieties, all f which are the result of, or associated with, some congenital efect. They are as follows:—

1. *Hernia* in the *tunica vaginalis testis*.—This occurs when a otrusion of intestine takes place through the narrow canal which rsists between the general cavity of the peritoneum and the ica vaginalis testis, in consequence of the non-obliteration of original communication between them. In this case, the stine surrounds the testis, and the sac is formed by the tunica inal testis (Fig. 102, 1).[2]

2. *Hernia* in the *funicular portion* of the *tunica vaginalis* urs when an incomplete closure of the tunica vaginalis takes e immediately above the testis; the canal above it being still losed and communicating with the peritoneal cavity. The is formed by the original pouch of the peritoneum in the cent of the testis, although shut off from the tunica vaginalis is by a thin septum (Fig. 102, 2).

3. *Infantile hernia* is rare, and occurs when the original peri- eal canal is occluded at the inner ring, so that the tunica inalis testis reaches up as high as the canal, or even as far as

[1] The term *congenital* applied to this form of hernia is apt to suggest the idea that curs at birth. But this is not of necessity so. Although the state of parts favor- to its occurrence exists at birth, the hernia itself may not take place till many s afterward—in fact, at any period of life.

[2] A good specimen of this variety, prepared by Percival Pott, is in the Museum of Bartholomew's Hospital.

the internal ring. The intestine in this variety protrudes a sac through the inner ring, but behind this abnormal extension of the tunica vaginalis; so that in front of the hernia there are three layers of peritoneum; two formed by the tunica vaginalis, the third by the sac (Fig. 102, 3).

4. *Encysted hernia* is still rarer than the preceding, and may occur in those cases in which the closing septum at the internal ring is so thin that an advancing hernia pushes before it this thin stratum (which forms its sac) as a diverticulum into an unclosed tunica vaginalis (Fig. 102, 4).

Position of Spermatic Cord. The spermatic cord is generally situated behind and to the outer side of a hernial sac. In some cases, however, the hernia separates the constituents of the cord, so that one or other of these comes to lie in front of the protrusion.

Seat of Stricture. The stricture may be seated either at the external ring, the internal ring, at any intermediate part between these, or at the neck of the sac. Sometimes there is a double stricture, one at the external ring, the other at the internal.

As stated, the stricture may be caused by the neck of the sac, independently of the parts outside it; for the peritoneum may become thickened and indurated, and sufficiently unyielding to strangulate the protruded intestine. The strangulation in a congenital hernia is nearly always caused by the neck of the sac itself.

In dividing the stricture, the surgeon should, in all cases, adhere to the golden rule laid down by Sir Astley Cooper—namely, to divide it directly upward. In this direction, there is the least likelihood of wounding the deep epigastric artery.

Changes Produced by an Old and Large Hernia. Whoever has the opportunity of dissecting an old hernia of some size, will observe that the obliquity of the inguinal canal is destroyed. The constant dragging of the protruded viscera upon the inner margin of the internal ring gradually approximates the internal ring to the external, so that at last the one gets quite behind the other, and there is a direct opening into the abdomen. But the position of the deep epigastric artery with regard to the

sac remains unaltered. It is still on the *inner side* of the neck of the sac.

In herniæ of long standing, all its coverings undergo a change. They become thickened and hypertrophied, and so altered from what they once were that they scarcely look like the same parts.

Umbilical Hernia. This is a hernia which protrudes through the umbilical opening in the middle line at the umbilicus, and is most commonly met with in infant life, and in the female sex in advanced life, especially in obese subjects. The hernia is frequently large, and, in its enlargement, it increases in an upward direction, so that the aperture through which it comes is not the centre of the hernia.

Its coverings are skin, subcutaneous fat and connective tissue, a thin fascia which covers in the umbilical aperture, fascia transversalis, subperitoneal fat and peritoneum.

The seat of strangulation in this variety of hernia is the fibrous margin of the umbilical aperture. This must be divided, and, as there are no vessels or other structures of importance near it, the margin may be divided at any part most convenient to the operator. Owing to the size which umbilical herniæ frequently attain, it is not always easy to divide the constricting margin, and great care is needed to prevent the intestines being injured during the operation.

Ventral Herniæ. This term is applied to those forms of hernia which protrude through the abdominal walls in situations not included in the inguinal and umbilical varieties. The most common variety is that which comes through the linea alba, usually above the umbilicus. They are small and mushroom-shaped, with narrow pedicles, and are commonly irreducible. In some cases, these protrusions are masses of subperitoneal fat emerging through foramina in the linea alba, growing after they protrude; so that these are not herniæ in the strict sense of the term. Ventral herniæ occasionally come through the linea semilunaris; others come through the triangular interval between the latissimus dorsi and posterior free border of the external oblique, and are termed *lumbar herniæ*. There is, in fact, no part of the abdominal walls through which a hernia may not protrude.

Dissection. Expose the contents of the abdomen, by an incision from the ensiform cartilage to the os pubis a little to the left side of the linea alba, so as to preserve a ligament, *ligamentum teres*, which passes from the umbilicus to the liver, and also a cord, the *urachus*, which ascends in the middle line from the bladder to the umbilicus; then make another incision transversely on a level with the umbilicus, and turn the flaps outward.

Urachus. Behind the linea alba, the peritoneum is raised into a fold by a fibrous cord, passing from the bladder to the umbilicus; this is the *urachus*, which in fœtal life is a tube connecting the bladder with the allantois. On either side of the urachus are two other folds, enclosing cords which ascend obliquely toward the umbilicus; these are the impervious remains of the *hypogastric arteries*.

On opening the cavity of the peritoneum, there are seen, in connection with the fibrous cords just mentioned, two fossæ in the neighborhood of Poupart's ligament—one on the inner side, the other on the outer side of the obliterated hypogastric artery. The depth of these fossæ depends upon the tension of this cord; so that, while sometimes they are scarcely visible, at others they are deep and well-marked. They are of importance, since they correspond with the internal and external abdominal rings; hence, the greater depth of one or other hollow may determine the locality of the protrusion of a hernia. Occasionally the deep epigastric artery lies nearer the middle line than it normally does, so that we may have three pouches instead of two, through either of which a hernia may emerge.

The abdominal cavity is seen to be composed of two divisions—an upper or abdomen proper, and a lower or the pelvis. It is the upper division that we are about to describe.

Take now a survey of the viscera before they are disturbed from their relative positions.

What is seen on opening the Abdomen. In the right hypochondrium, the right lobe of the liver is seen projecting more or less below the cartilages of the ribs, and the fundus of the gall-bladder below the edge of the liver, near the end of the ninth costa

cartilage. In the left hypochondrium is seen more or less of the stomach, according to its distention. Across the umbilical region extends a broad fold of the peritoneum containing fat, the great omentum, which descends from the lower curvature of the stomach, forming a curtain over the convolutions of the small intestine. The breadth of this fold varies; sometimes being so shrunk and crumpled as to be scarcely visible. The lower part of the abdomen and part of the pelvis are occupied by the small intestine. The urinary bladder is not apparent, unless distended sufficiently to rise out of the pelvis. In the right iliac fossa is the caput coli, the commencement of the large intestine; but the ascending part of the large intestine in the right lumbar region, and the descending part of it in the left, are not visible unless distended; they lie contracted at the back of the abdomen. Such are the viscera usually seen on opening the abdomen; but a certain latitude is to be allowed, as sometimes more of one organ is seen and less of another, according as this or that is distended or hypertrophied. Much also depends upon the amount of pressure which the ribs have undergone during life.

Particular Position of each Viscus. The position of each viscus should now be examined separately, and first that of the stomach.

The Stomach. The *stomach* is irregularly conical in shape. It is placed obliquely, the smaller or pyloric end being the lower; its great end is situated in the left hypochondrium; its narrow or pyloric end extends obliquely across the epigastrium into the right hypochondrium, where it is overlapped by the liver.

The left or cardiac end is situated behind the ribs, and in contact with the concavity of the spleen, to which it is connected by a fold of peritoneum. In front it has the abdominal wall, above it has the liver and the diaphragm, and below it is the transverse colon; the right or pyloric end extends to the gall bladder, and is in contact with the under aspect of the liver, where it is continuous with the duodenum; posteriorly, it rests on the pancreas, the aorta, and the two crura of the diaphragm. The stomach is connected with other viscera by the following peritoneal folds:

(1) the great omentum, attached to its lower convex border, forms a curtain of fat, more or less thick, in front of the transverse colon and small intestines; (2) the gastro-hepatic or lesser omentum, which connects the lesser curve of the stomach with the transverse fissure of the liver; and (3) the gastro-splenic omentum, which connects the cardiac end of the stomach with the hilum of the spleen. The relative position and size of the stomach vary according to the amount of distention; when much distended, the anterior surface, owing to the greater mobility of the great curve and the pyloric end of the stomach, is turned upward, and the lower border forward.[1]

Duodenum. The first part of the intestinal canal is termed *intestinum duodenum*, because it is about the breadth of twelve fingers. Commencing at the pyloric end of the stomach, the duodenum ascends as high as the neck of the gall bladder; then turning downward it passes in front of the right kidney; lastly, making another bend, it crosses the spine obliquely toward the left side of the second lumbar vertebra. Here the intestinum jejunum begins, and this part of the canal may be seen by raising the transverse colon. Thus the duodenum describes a horse-shoe curve, of which the concavity is toward the left, and embraces the large end or head of the pancreas. For convenience of description the duodenum is divided into an ascending, a descending and a transverse portion. The first is completely surrounded by a peritoneal covering; the second and third are only covered by peritoneum in front, and are fixed to the back of the abdomen. The relative anatomy of the duodenum will be more fully seen hereafter (p. 495).

Jejunum and Ileum. Pursuing its course from the left side of the second lumbar vertebra, the intestinal canal forms a number of convolutions, occupying the lower regions of the abdomen, and which are loosely connected to the spine by a broad peritoneal fold termed the *mesentery*. Of these convolu-

[1] Lesshaft states that the position of the stomach is *vertical*, and that when distended it does not alter its position, but that it is affected equally in all directions (*Lancet*, March 11, 1882). This opinion is not, however, generally entertained by anatomists.

he upper two-fifths constitute the *intestinum jejunum;* the hree-fifths, the *intestinum ileum.* This is an arbitrary divi- There is no definite limit; the character of the bowel grad- hanges—that is, it becomes less vascular, has fewer folds ining membrane, and its coats are therefore less substantial feel.

encement Large estine.

In the right iliac fossa, the small intestine opens into the left side of the cæcum, which is easily recognized by its sacculated appearance; here the

FIG. 103.

or caput coli.

3. Appendix vermiformis.

ON THROUGH THE JUNCTION OF THE LARGE AND SMALL INTESTINE, TO SHOW THE ILEO-CÆCAL VALVE AND APPENDIX VERMIFORMIS.

ntestine begins, and it is guarded by the ileo-cæcal valve 03). Immediately below the junction the large intestine is led into a blind pouch, about two and a half inches in length eadth, called the *cæcum* or *caput coli.* Into the back part of uch opens a little tube, closed at the other end, called the *lix vermiformis.* This tube varies from three to six inches gth, is about as thick as a large earthworm, and is either up behind the cæcum, or connected to it by a peritoneal as to hang loose in the abdomen. It is hollow, and its

opening into the cæcum is usually guarded by a valve of mucous membrane.

The large intestine is about five or six feet in length, and in its course it describes an arch which encircles the convolutions of the small intestines. It is largest at its commencement, and lessens in size until at the upper part of the rectum it becomes narrowest; below this it again forms a dilatation, the *ampulla*, just above its termination at the anus. It is successively divided into cæcum, ascending, transverse and descending colon, sigmoid flexure and rectum. The commencement of the large intestine is generally confined by the peritoneum to the iliac fossa, in which it lies, being

FIG. 104.

RELATIVE POSITION OF THE KIDNEYS AND THE LARGE INTESTINE, SEEN FROM BEHIND.

L. K. Left kidney.
R. K. Right kidney, crossed obliquely by the last dorsal artery and nerve.
A. C. Ascending colon.
D. C. Descending colon.

separated from the iliacus muscle by the iliac fascia and by more or less connective tissue and fat.[1] Tracing the large intestine from this point, it is continued as the *ascending colon*. We find it somewhat smaller than the cæcum, and it ascends through the right lumbar region in front of the right kidney, as high as the under surface of the liver, where it abruptly makes a bend to the left side—the *hepatic flexure* of the colon; it then crosses the umbilical region transversely from right to left, and is known as the *trans-*

[1] But this is not invariably so. The bowel is, in some subjects, connected to the fossa by a fold of peritoneum or a *meso-cæcum*. I have seen this fold sufficiently loose to allow the caput coli to travel over to the left iliac fossa.

e colon. Reaching the left hypochondriac region, it makes a rp bend downward beneath the lower border of the spleen, ming the *splenic flexure* of the colon;[1] thence it descends in nt of the left kidney,[2] through the left lumbar region into the iliac, as the *descending colon.* In the iliac fossa the intestine, the *sigmoid flexure*, becomes narrow and makes a curve like the er S. Lastly, the bowel enters the pelvis on the left side of the rum, and here takes the name of *rectum.* This term, so far as ncerns the human subject, is misapplied; the canal runs any- ng but a straight course through the pelvis, since it curves to pt itself to the sacrum.

Looking at the entire course of the colon, observe that it forms arch, of which the concavity embraces the convolutions of the ll intestines.

Let us now see to what extent the small and the large intestines invested with a peritoneal coat. The *small intestines*, with the ption of the duodenum, which cannot at present be examined, shall find are completely surrounded by peritoneum, except at mesenteric border, along which the vessels pass to the bowel; *cæcum* is covered, as a rule, only in front and on its sides, the erior surface being connected to the iliac fascia by connective ue; the *ascending colon* is also only covered on the front aspect sides, the posterior surface being loosely connected by areolar ue to the quadratus lumborum and right kidney; the *trans- e colon* is almost entirely surrounded by peritoneum, which is cted horizontally backward as a broad double layer—the sverse meso-colon—to the spine; the *descending colon*, smaller the transverse colon, and more deeply situated than the as- ding colon, is only invested with peritoneum on its anterior inner surfaces; the *sigmoid flexure* is completely invested peritoneum, which connects this part of the bowels with a

[1] This transverse part of the colon, in some instances, makes a coil behind the ach to the diaphragm; such a state of things, when the bowel happens to be nded, is apt to give rise to symptoms of diseased heart. See some observations int, by Dr. Copland, in *Lond. Med. Gaz.*, 1847, vol. v, p. 660.

[2] The contiguity of the ascending and descending colon to the right and left ey respectively, explains the occasional bursting of renal abscesses into the inal canal.

loose fold to the left iliac fossa; the *rectum* is completely surrounded by peritoneum in its upper half, and is connected to the front of the sacrum by a fold, the *meso-rectum;* the lower half of the rectum, for the first three inches, is only covered in front with peritoneum; the last inch and a half has no peritoneal investment at all.[1]

Relations of the Large Intestine.

At present we have only traced the course of the large intestine through the different abdominal regions, but now we are able to see the relations of the various portions of the bowel.

The *cæcum* rests in the right iliac fossa, separated from the iliacus by the iliac fascia and connective tissue; in front it has the anterior abdominal wall.

The *ascending colon* has behind it the quadratus lumborum and right kidney; in front, it has the abdominal wall and small intestines; above, it is in contact with the under aspect of the liver to the right of the gall bladder.

The *transverse colon* is concave posteriorly, and has above, the liver, gall bladder, the stomach and the lower border of the spleen; behind, it has the transverse meso-colon and the third part of the duodenum; in front, it is in contact with the abdominal wall and great omentum; below, with the small intestines.

The *descending colon*, deeper situated than the ascending colon, lies behind, in contact with the left crus of the diaphragm, the left kidney and quadratus lumborum; in front, with the small intestines, and on its left side with the abdominal wall.

The *sigmoid flexure* is in relation behind with the iliac fossa, the left spermatic artery and ureter, the left common iliac vessels; in front, with the small intestines and abdominal wall.

The relations of the *rectum* cannot at present be satisfactorily made out; the description of them has been deferred till the dissection of the side view of the pelvic viscera.

[1] It should be recollected, that the ascending and descending colon are not infrequently completely invested by peritoneum, and therefore, virtually speaking, have a mesentery. This occasional occurrence is important when the operation of right or left lumbar colotomy has to be performed.

ngth of the entary Canal. The small intestine, including the duodenum, varies from sixteen to twenty-four feet in length, the large intestine from five feet to five feet and a half; these surements are subject to some variation, according to the ht of the subject.

he average lengths of the different portions of the alimentary l are as follows:—

Duodenum,	8 to 10 inches.
Jejunum,	two-fifths of the remaining part
Ileum,	three-fifths of the small intestine.
Cæcum,	2½ inches.
Ascending colon,	8 inches.
Transverse colon,	12 inches.
Descending colon,	11 inches.
Sigmoid flexure,	22 inches.
Rectum,	6 to 8 inches.

ıation of the Liver. The liver occupies the whole of the right hypochondrium, and extends over the epigastric region, or less, into the left. Unless the individual be very corpuwe can ascertain during life the extent to which the liver cts below the costal cartilages, and the general dimensions ıe organ may be tolerably well told by percussion. Its *ior border* is sharp and thin, and presents in the epigastric n a deep notch for the round ligament, and generally projects le way below the ribs; its *posterior border* is broad and con-d to the diaphragm by the coronary ligament; it is in relabehind with the inferior vena cava, the aorta and the crura of liaphragm. Its *upper convex surface* ascends as high as the intercostal space, is accurately adapted to the arch of the ıragm, and is divided into two unequal parts by the falciform ıspensory ligament; its *under surface* overlies part of the ach, and of the duodenum, the right kidney and supra-renal ıle, and the hepatic flexure of the colon. Its *right border* is ; its *left* is thin and sharp. To the diaphragm the liver is ected by folds of peritoneum, called *ligaments*. One of these, y vertical in direction, and called the *suspensory*, or, from its e, the *falciform* ligament, is situated a little to the right of the

mesial line. The lower and free edge of it contains the impervious remains of the umbilical vein, called the *round* ligament. The suspensory ligament, traced backward, leads to another broad fold extending horizontally from the diaphragm to the posterior border of the liver; this constitutes the *lateral* ligament, right or left, according as we trace it on one or the other side of the falciform ligament.

The junction of the lateral and falciform ligaments is described by some authors as the *coronary* ligament.

Situation of the Gall bladder. The gall bladder is the reservoir for the bile, and is closely confined by the peritoneum in a slight depression on the under surface of the right lobe of the liver, to which it is connected by areolar tissue; occasionally the gall bladder is completely surrounded by peritoneum. It is pyriform in shape, and its broad end or fundus, covered with peritoneum, projects beneath the anterior border of the liver opposite the ninth costal cartilage. It measures three to four inches in length, is an inch and a half broad, and contains from eight to twelve drachms. The neck is inclined upward and toward the left, and is firmly connected to the liver by areolar tissue. The gall bladder is in relation *above* with the liver and small blood vessels; *below*, with the transverse colon and with the first portion of the duodenum; its *neck* is curved upon itself like the letter S, and bending downward terminates in the cystic duct. It sometimes happens that the gall bladder, in consequence of some obstruction to its duct, becomes unusually distended, and occasions a swelling below the margin of the ribs, which might be mistaken for an hepatic abscess.[1] The close proximity of the gall bladder to the duodenum and the transverse colon explains the occasional evacuation of gall stones by ulceration into the intestinal canal.[2]

Situation of the Spleen. The spleen is the dark, purple-gray, flattened organ deeply situated in the left hypochondrium, between the stomach and the ninth, tenth, and eleventh ribs. It

[1] See cases in point recorded by Andral, *Chir. Med.*, tom. iv; and Graves, *Dublin Hospital Reports*, vol. iv.

[2] See preparations in St. Bartholomew's Hospital Museum.

is placed nearly vertically; its outer surface is smooth and convex, to correspond with the diaphragm and ribs; its inner surface, where its great vessels enter, is concave, and connected to the great end of the stomach by a broad peritoneal fold, called the *gastro-splenic omentum*. Its *external surface* is in relation with the diaphragm, which separates the organ from the ninth, tenth, and eleventh ribs; its *internal surface* is concave, and presents a vertical fissure—the *hilum*—situated nearer the posterior than the anterior border; it is at this fissure that the two layers of peritoneum are reflected from the stomach to the spleen, and the splenic vessels enter and emerge; it is in relation with the cardiac end of the stomach, the tail of the pancreas, the left supra-renal capsule, and the left crus of the diaphragm; the *upper border* is rounded, and is connected to the diaphragm by a fold of peritoneum—the *suspensory ligament;* the *lower border* is in contact with the splenic flexure of the colon; the *posterior border* is thick, and is connected with the left kidney by areolar tissue; the *anterior margin* usually presents a more or less deep notch. Its hilum is connected with the cardiac end of the stomach by a fold of peritoneum—the *gastro-splenic omentum;* and with the under surface of the diaphragm by a small peritoneal fold—the *suspensory ligament*.[1]

Situation of the Pancreas. This is the large salivary gland of the abdomen. It is placed transversely across the back of the abdomen, in front of the spine, about the level of the first lumbar vertebra. It is about seven inches in length, and an inch and a half in breadth. Its *right end*, or *head*, is contained within the curve of the duodenum; its *left end*, or *tail*, extends as far as the spleen. The further connections and relations of the pancreas cannot at this stage of the dissection be satisfactorily seen.

Situation of the Kidneys. The kidneys are two large excretory glands, situated at the back of the abdomen in each lumbar region, nearly opposite the two last dorsal and the two upper lumbar vertebræ—the right, owing to the size of the liver, being a little lower than the left. They lie imbedded in fat, which maintains them in their proper position. *Behind*, they rest on the

[1] We find occasionally in the gastro-splenic omentum one or more small spleens, in addition to the large one.

crus of the diaphragm, on the quadratus lumborum and psoas, separated by the aponeurosis of the transversalis; *in front*, the right kidney is in relation with the peritoneum, the right lobe of the liver, the second part of the duodenum, and the ascending colon; the left kidney is in contact with the peritoneum, the cardiac end of the stomach, the spleen, the end of the pancreas, and the descending colon; *externally*, it is convex, and in contact with the abdominal parietes; *internally*, it is concave, and presents a deep hollow, the *hilum*, from which pass the ureter and large vessels; *above*, it is in relation with the supra-renal capsule; *below*, it extends nearly as low as the crest of the ilium.

Situation of the Supra-renal Capsules. These are two ductless glands, situated at the top of the kidneys and behind the peritoneum. The right one is triangular; the left oval and almond-shaped. The right supra-renal capsule is in relation, *in front*, with the under aspect of the liver; the left, with the pancreas and spleen; *behind*, it rests on the crus of the diaphragm; the *upper border* is convex and thin; the *lower border* is concave, and rests on the kidney; the *inner border* is in relation with the semi-lunar ganglion and splanchnic nerves, and with the vena cava on the right side, and with the aorta on the left side. The anterior surface is slightly indented, from which the supra-renal vein passes out to join on the right side the inferior vena cava, and on the left side the left renal vein.

Peritoneum. A certain range of motion being necessary to the abdominal viscera, they are provided with a serous membrane, called the *peritoneum*. This membrane, like other serous membranes, is a closed sac, one part of which lines the containing cavity, the other is reflected over the contained viscera. These are respectively termed the *parietal* and the *visceral* layers. In the female, however, it is not, strictly speaking, a closed sac, since it communicates with the cavity of the uterus through the Fallopian tubes. The internal surface of the peritoneum is smooth and polished, and lined by squamous endothelium; the external surface—the *sub-peritoneal tissue*—is composed of areolar tissue, which connects the internal layer to the invested viscus or abdominal parietes. There is nothing between the

rietal and the visceral layers—in other words, inside the sac—
t just sufficient moisture to lubricate its smooth and polished
rface. The viscera are all, more or less, outside the sac; some
altogether behind it, as the pancreas, kidneys, and supra-renal
psules; others, as the lower parts of the duodenum, cæcum,
cending and descending colon, are only partially covered by it;
hile others, as the stomach, liver, jejunum, ileum, and some
rts of the large intestine, are completely invested by it; these
tter push the visceral layer before them, and so give rise to
embranous folds; the larger the fold, the freer is the mobility
the viscus which occasions it.

Course of the Peritoneum. Now trace the peritoneum as a
ntinuous membrane. Since
e peritoneum is a perfect sac,
matters not where we begin;
must come back to the start-
point.

If a longitudinal section be
de through the viscera in the
dle of the body, one can trace
peritoneum thus—beginning
he diaphragm, and taking, for
vity's sake, two layers at a
e (Fig. 105).

rom the diaphragm two layers
peritoneum proceed to the
r, forming its *lateral ligaments;*
separate to enclose the liver,
t again on its under aspect,
pass on, under the name
he *gastro-hepatic omentum*, to
small curve of the stomach.
arating here, they embrace
stomach, and, meeting again
ts greater curve, pass down
a curtain over the small intestine to form the *great omentum*.

FIG. 105



DIAGRAM OF THE PERITONEUM.

At the lower margin of the great omentum, they are reflected upward (so that the great omentum consists of four layers) to the front of the transverse colon, which they enclose, and, after joining again at the back of the colon, proceed to the spine, forming the *transverse meso-colon.* At this situation the two layers diverge; the upper one ascends in front of the pancreas and the crura of the diaphragm to its under surface, at which point we started.[1]

The peritoneum passes from the under surface of the right lobe of the liver to the kidney, forming a slight fold—the *hepato-renal fold;* on the left side, where the peritoneum extends from the diaphragm to the cardiac end of the stomach, it passes as a slight duplicature, forming the *gastro-phrenic ligament;* an extension of this is seen, passing as a distinct fold—the *costo-colic ligament*—from the diaphragm to the splenic flexure of the colon.

The lower layer is reflected from the spine over the small intestine, back again to the spine, to form the *mesentery.* From the root of the mesentery it descends into the pelvis, and invests the upper two-thirds of the rectum. From the rectum, in the male, it is reflected to the posterior part of the bladder, forming the *recto-vesical pouch,* and thence to the wall of the abdomen, along which it can be traced up to the diaphragm. In the female, it is reflected from the rectum on to the posterior wall of the vagina, half an inch from the uterine extremity, constituting the recto-vaginal pouch (*Douglas' pouch*), and thence over all the back, but only about half way down the front of the uterus, to the posterior wall of the bladder; after which its reflections are the same as in the male.[2]

Such is the course of the peritoneum as seen in a longitudinal

[1] In fœtal life, the ascending layers of the great omentum may be traced back to the spine near the pancreas; and here the layers diverge from each other. The upper layer ascends in front of the pancreas to the diaphragm; the lower layer proceeds over the arch of the colon, and then back to the spine, thus forming the transverse meso-colon. Its reflections afterward are the same as in the adult. As the fœtus grows, the great omentum becomes adherent to the arch of the colon.

[2] For a detailed description of the development of the great omentum and the transverse meso-colon, see a paper by C. B. Lockwood, *Journal of Anatomy and Physiology*, vol. xviii.

but there are lateral reflections which cannot be seen a transverse section: thus, from the great end of the two layers proceed to the spleen, forming the *gastro-entum;* from the transverse meso-colon it is reflected side over the ascending and descending colon.

ructures *completely invested* with peritoneum are, the liver, first part of the duodenum, the jejunum and ileum, erse colon, sigmoid flexure, upper part of rectum, spleen, d ovaries.

llowing parts of the alimentary canal are only *partially* y peritoneum, namely, the descending and transverse of the duodenum, the cæcum, the ascending and

FIG. 106.

cavity of
toneum.
cavity of
toneum.

St. Stomach.
s. Spleen.
v. Inferior vena cava.
L. Liver.

M OF A TRANSVERSE SECTION THROUGH THE UPPER PART OF THE ABDOMINAL CAVITY, SEEN FROM ABOVE.

ng colon (with exceptional cases), the middle part of the he upper part of the vagina, and the hinder wall of the

scera *uncovered* by peritoneum are, the lower part of the he anterior and the lower part of the posterior wall of na, the anterior and part of the posterior wall of the

mists speak of the *lesser cavity of the peritoneum*, as shed from the greater. This lesser cavity, or *cavity of the ntum*, is situated behind the stomach and the descending the great omentum. If air be blown through the foramen ow (which is the constricted communication between the

I

greater and lesser cavities of the peritoneum), the lesser cavity becomes distended. It is bounded *in front* by the lesser omentum, the stomach, and the descending layers of the great omentum; *behind*, by the ascending layers of the great omentum, the colon, the upper layer of the transverse meso-colon and its ascending layer; *above*, by the liver; *below*, by the turn of the great omentum.

Foramen of Winslow. This foramen is the narrow circular opening between the greater and lesser cavities of the peritoneum, through which the two cavities communicate. It is situated behind the right edge of the gastro-hepatic or lesser omentum. By passing your finger into it, you will find the

FIG. 107.

DIAGRAM OF A TRANSVERSE SECTION THROUGH THE LOWER PART OF THE ABDOMINAL CAVITY.

foramen bounded *above* by the lobulus Spigelii or caudatus of the liver; *below*, by the commencement of the duodenum and by the curving forward of the hepatic artery; *in front*, by the lesser omentum, enclosing the hepatic artery and duct and the vena portæ; and *behind*, by the vena cava inferior.

The several folds, formed by the reflections of the peritoneum, which connect the viscera either to each other or to the back of the abdomen, are classified respectively as ligaments, mesenteries, and omenta; the *ligaments* are attached to the viscera, and help to maintain them in position; the *mesenteries* pass from the abdominal walls to the large and small intestines, and maintain them

position and allow of blood vessels to pass to and from them; d the *omenta* are broad, flat duplicatures which pass from the mach to the neighboring viscera.

Mesentery. This is the fold which suspends the small intestine from the back of the abdomen. To see it, ise the great omentum and the transverse arch of the colon. s attached part or root is about six inches in length, and extends m the left side of the second lumbar vertebra obliquely across spine to the right sacro-iliac symphysis. The loose part of mesentery is very broad, and curves like a ruffle, enclosing small intestine from the beginning of the jejunum to the end the ileum. Its shape resembles an open fan, and its length n the vertebral column to its attachment to the intestine is ut four inches. Above, it is connected with the under surface he transverse meso-colon; below, with that part of the perieum which lines the inner part of the cæcum and ascending on. We must trace between its two layers the mesenteric sels, nerves, glands, and lymphatics.

Transverse Meso-colon. This broad fold connects the transverse colon to the back of the abdomen, and between its ers the vessels pass to and from this portion of the larger . It forms an imperfect partition dividing the abdomen into upper compartment, containing the stomach, liver and spleen; a lower, containing the convolutions of the small intestines.

Ascending and Descending Meso-colon. As regards the cæcum, and the ascending and descending portions of the colon, they are, as a general rule, bound down by the peritoneum in ir respective situations (Fig. 107). The peritoneum covers y two-thirds or thereabouts of their anterior surface; their terior surface is connected by loose cellular tissue to the back he abdomen. The colon, ascending or descending, can there- be opened in the lumbar region, below the kidney, without in- to the peritoneum; a fact upon which is founded the operation olotomy for the relief of stricture of the rectum. In some cases, ascending and descending colon (more commonly the latter) completely surrounded by peritoneum and connected to the bar regions, respectively, by a right and a left *lumbar meso-colon*.

Sigmoid Meso-colon. The sigmoid flexure is, as a rule, completely invested by peritoneum, which passes as a thin fold to the iliac fossa, allowing a considerable amount of movement of this part of the intestine.

Meso-rectum. The upper third of the rectum is also surrounded by peritoneum, which passes to the sacrum and thus retains it in position. The hæmorrhoidal vessels pass between its layers.

Great Omentum. This broad peritoneal fold, known also as the *gastro-colic omentum*, is composed of four layers, and proceeds as a double layer from the lower border of the stomach, as far as the pelvis, where these two layers ascend to enclose the transverse colon. It lies like a curtain over the convolutions of the small intestines, and we find it in some bodies extending low into the pelvis; in others, small and crumpled up, usually, in the left hypochondrium. Its thickness varies considerably; in thin subjects it is often translucent; in corpulent persons, on the other hand, it is loaded with fat, and contributes in great measure to the size of the abdomen.

Gastro-hepatic or Lesser Omentum. This double fold passes from the transverse fissure on the under surface of the liver to the upper curve of the stomach. It is composed of two layers, and between them are the portal vein and hepatic artery with the nerves going to the liver, and the hepatic duct and lymphatics coming from it. The right border of this fold is free, and forms the anterior rounded margin of a constriction, called the *foramen of Winslow*, which leads into the lesser cavity of the peritoneum; its left border passes on to the œsophagus. In this fold the common bile duct lies to the right, the hepatic artery to the left, and the vena portæ behind and between them. If now the finger be introduced behind the right border, it passes through the foramen of Winslow into the lesser cavity of the peritoneum.

Gastro-splenic Omentum. This fold proceeds from the great end of the stomach to the spleen, and is continuous below with the great omentum. It contains between its layers the branches, *vasa brevia*, which proceed from the splenic artery to the great end of the stomach.

Ligaments. The reflections of the peritoneum from the abdominal walls to the liver, the spleen, the bladder and uterus, and constituting their *ligaments*, have been, or will be, described with the respective viscera.

Branches of the Abdominal Aorta. Our next object should be the examination of the arteries which supply the viscera. The *abdominal aorta* enters the abdomen between the pillars of the diaphragm in front of the last dorsal vertebra, and then, descend-

FIG. 108.



1. Phrenic.
2. Cœliac axis.
3. Superior mesenteric.
4. Supra-renal.
5. Renal.
6. Spermatic.
7. Inferior mesenteric.
8. Lumbar.
9. Sacra media.

BRANCHES OF THE ABDOMINAL AORTA.

ing a little to the left of the spine, divides on the body of the fourth lumbar vertebra, a little to the left of the middle line, into the two common iliac arteries. The relations of the aorta cannot at present be sufficiently made out, so that this will be described later on. In its course it gives off its branches in the following order (Fig. 108):—

1. The *phrenic*, for the supply of the diaphragm.
2. The *cœliac axis*, a short, thick trunk, which immediately

subdivides into three branches for the supply of the stomach, the liver and the spleen.

3. The *superior mesenteric*, for the supply of all the small intestines, and the upper half of the large.

4, 5. The *supra-renal* and the *renal* arteries.

6. The *spermatic*, for the testicles; the *ovarian*, for the ovaries.

7. The *inferior mesenteric*, for the supply of the lower half of the large intestine.

8. The *lumbar*, four branches analogous to the intercostals, for the supply of the back part of the abdomen.

9. The *arteria sacra media*, which is given off at the bifurcation of the aorta, supples the fifth lumbar artery, and running down in front of the sacrum, supplies the rectum and other structures.

By some anatomists the branches are arranged in two classes —those destined to supply the viscera, and those to supply the abdominal parietes; the former are the cœliac axis, supra-renal, renal, spermatic, superior and inferior mesenteric arteries; the latter are the phrenic, the lumbar and sacra media arteries.

Dissection. These branches are to be traced throughout in the following order. Take the cœliac axis first. To dissect this artery and its branches, the liver must be well raised and the stomach drawn down, as in Fig. 109, and the anterior layer of peritoneum removed from the gastro-hepatic omentum. A close network of very tough tissue surrounds the visceral branches of the aorta. This tissue consists almost entirely of plexuses of nerves, derived from the sympathetic system, each plexus taking the name of the artery which it surrounds. Of these plexuses, the largest surrounds the cœliac axis like a ring. This is the solar plexus, and is formed by the junction of the two semilunar ganglia (see Dissection of Thorax, p. 20[illegible]). From this, as from a root, other secondary plexuses branch off and surround the following arteries: the phrenic, coronary, hepatic, splenic, superior mesenteric, inferior mesenteric and renal; these plexuses receiving the names of the arteries around which they twine. It requires a lean subject and much patience to trace these.

Cœliac Axis and its Branches. The *cœliac axis* arises from the front of the aorta, between the pillars of the diaphragm, im-

ately above the upper border of the pancreas, to the left of obulus Spigelii, to the right of the cardiac end of the ach, and having the semilunar ganglia on each side. It is a , thick trunk, which runs between the two layers of the lesser itum, and after a course of about half an inch divides into branches, the hepatic running to the right, the splenic to ft, and the coronaria ventriculi upward and to the left side.

FIG. 109.



DIAGRAM OF THE BRANCHES OF THE CŒLIAC AXIS.

(Pancreas in dotted outline behind the stomach.)

1. Coronaria ventriculi.
2. Splenic a.
3. Hepatic a.
4. Pyloric a.
5. Gastro-duodenalis.
6. Gastro-epiploica sinistra.
7. Vasa brevia.
8. Superior mesenteric a.

e following is the plan of the cœliac axis and its branches:—

c AXIS . { Coronaria ventriculi { œsophageal. / gastric.
Hepatic . . . { pyloric. / gastro-duodenalis. { gastro-epiploica dextra. / pancreatico-duodenalis superior. / cystic.
Splenic . . . { pancreatic branches. / gastro-epiploica sinistra. / vasa brevia to stomach.

Coronaria Ventriculi. The *coronaria ventriculi*, the smallest of the three, ascends a little to the left, toward the œsophageal end of the stomach, where it gives off *œsophageal* branches, which anastomose with the œsophageal branches of the thoracic aorta; and others to the cardiac end of the stomach, which inosculate with the vasa brevia of the splenic artery. It then runs from left to right, along the lesser curvature of the stomach, toward the pylorus, supplying branches on both surfaces of the stomach, and finally anastomoses with the pyloric branch of the hepatic artery.

Hepatic Artery. The *hepatic artery* ascends to the right between the layers of the lesser omentum to the transverse fissure of the liver, where it divides into two branches, right and left, for the supply of the respective lobes of the liver.

In its course to the liver, it lies to the left of the common bile duct, and in front of the portal vein; all three are contained in the right half of the lesser omentum. The hepatic gives off—

a. The *pyloric*, which descends to the upper border of the pylorus, and runs along the lesser curve of the stomach from right to left, inosculating with the coronaria ventriculi.

b. The *gastro-duodenalis* descends behind the ascending portion of the duodenum, divides, after a short course, into (*α*) the *gastro-epiploica dextra*, which runs along the greater curve of the stomach, between the layers of the great omentum, from right to left, and anastomoses with the gastro-epiploica sinistra from the spleen, supplying both surfaces of the stomach and the great omentum, and (*β*) the *pancreatico-duodenalis superior*, which runs down tween the head of the pancreas and the descending portion of duodenum, and anastomoses with the pancreatico-duodenalis ferior, a branch of the superior mesenteric, and with the pancreatic branches of the splenic.

c. The *cystic*, commonly a branch of the right hepatic, ascends along the neck of the gall bladder, and divides into two branches, one of which ramifies on the under surface of the gall bladder, the other passes between the liver and the upper surface of the gall bladder.

Splenic Artery.

The *splenic*, the largest of the three, proceeds tortuously toward the left side, above its corresponding vein, along the upper border of the pancreas to the hilum of the spleen, which it enters by numerous branches.

It gives off: 1. Several small branches to the pancreas, *pancreaticæ parvæ;* one, rather larger than the rest, *pancreatica magna*, accompanies the pancreatic duct. These arteries anastomose with the pancreatico-duodenal branches of the hepatic and superior mesenteric arteries. 2. The *gastro-epiploica sinistra*, which runs to the right along the great curve of the stomach, between the layers of the great omentum, and inosculates with the gastro-epiploica dextra. 3. *Vasa brevia*, five to seven in number, which proceed between the layers of the *gastro-splenic* omentum, to the great end of the stomach, where they communicate with branches from the coronaria ventriculi, and the gastro-epiploica sinistra. The *splenic* branches are five or six in number, and enter the texture of the spleen.

Thus the stomach is supplied with blood by four channels, which by their inosculations form a main artery along its lesser curve, another along its greater; from these, numerous branches are furnished to both surfaces of the stomach. The artery of the greater curve also sends down numerous *omental* branches, which form a network between the layers of the great omentum.

The vein corresponding to the coronaria ventriculi artery, called *coronary*, commences close to the pylorus, runs along the lesser curve of the stomach as far as the œsophagus, and then ascending to the right, between the two layers of the gastro-hepatic omentum opens into the vena portæ.[1]

The *splenic vein* returns the blood from the spleen by five or six branches which unite to form a single trunk. This runs along the upper border of the pancreas below the artery, and after receiving the branches corresponding to the branches of the artery and the inferior mesenteric vein, joins the superior mesenteric vein to form the vena portæ.

The *hepatic veins* do not run with the hepatic artery, but return the blood from the liver and terminate in the vena portæ.

[1] See Walsham in the *Journal of Anatomy and Physiology*, vol. xiv, p. 399.

Vena Portæ; its Peculiarities. The veins which return the blood from the abdominal portion of the alimentary canal, the pancreas, and the spleen, do not empty themselves into the vena cava inferior, but all unite into one large vein, called the *vena portæ*, which ramifies throughout the liver, and secretes the bile. The trunk of the vena portæ itself is about three inches long. Tracing it downward, you find that it is formed behind the great end of the pancreas and in front of the inferior cava, by the confluence of the splenic and superior mesenteric veins (Fig. 110). In its passage to the liver, the vena portæ is accompanied by the hepatic artery and the common bile duct, lying behind and between them. At the transverse fissure of the liver it presents a slight enlargement, called the *sinus*, and then divides into two branches corresponding to the right and left lobes. The vein ramifies in the substance of the liver like an artery, and is surrounded, with the branches of the hepatic artery and duct, in a sheath of areolar hepatic tissue called *Glisson's capsule*. The vena portæ may, then, be compared to the stem of a tree, of which the roots arise in the digestive organs, and the branches spread out in the liver. After receiving the veins corresponding to the branches of the hepatic artery, the vena portæ returns its blood into the inferior vena cava through the venæ cavæ hepaticæ.

FIG. 110.



DIAGRAM OF THE VENA PORTÆ.

(The arrow is introduced behind the free border of the lesser omentum.)

The veins which empty themselves into the vena portæ have no valve. Therefore, if any obstruction arises in the venous circulation through the liver, the roots of the portal vein are apt to become congested; this is a common cause of hæmorrhoids, diarrhœa, hemorrhage from the bowels, and ascites.

Hepatic Duct. The hepatic duct is formed by the junction of the right and left hepatic ducts, which issue from the transverse fissure. The hepatic duct descends nearly vertically for about an inch and a half, when it is joined at an acute angle by the cystic duct. The *cystic duct*, about an inch in length, descends, from the neck of the gall bladder, toward the left in the gastro-hepatic omentum, lying to the right of the hepatic artery and in front of the vena portæ. The hepatic and cystic ducts unite to form the *ductus communis choledochus*, or the common bile duct; the duct thus formed passes downward and to the left, between the two layers of the lesser omentum, close to its right border. It is about three inches long, and if distended would be about the size of a crow quill. It descends behind the first portion of the duodenum; in front of the vena portæ; to the right of the hepatic artery; to the left of the descending portion of the duodenum; and behind the head of the pancreas. Then turning toward the right, it gets behind the descending duodenum, and opens obliquely into the back part of the second portion, near the junction with the third. The duct runs through the coats of the bowel for nearly three-quarters of an inch, and sometimes before doing so unites with the pancreatic duct (p. 490).

Cystic Duct.

Ductus Communis Choledochus.

Dissection. The great omentum, with the arch of the colon, must now be turned up over the chest, and the small intestines pushed toward the left side. Then, by removing the anterior layer of the peritoneum from the mesentery, we expose the mode in which the superior mesenteric artery ramifies so as to supply the small intestines. In making this dissection, the mesenteric glands immediately attract notice. They lie in great numbers between the layers of the mesentery, and vary considerably in size. The fine tubes, called lacteal vessels, which

traverse the glands, are too thin and transparent to be seen under ordinary circumstances. But in cases where sudden death has taken place during digestion, they are found distended with chyle, and can be traced into the glands from all parts of the small intestine.[1] After traversing the glands, they all eventually empty their contents into the receptaculum chyli (p. 203).

Superior Mesenteric Artery and Branches.

This large artery arises from the front of the aorta just below the cœliac axis, descends beneath the pancreas, in front of the transverse part of

FIG. 111.



1. Superior mesenteric a.
2. Colica media.
3. Colica dextra.
4. Ileo-colica.

5. Inferior mesenteric a.
6. Colica sinistra.
7. Arteria sigmoides.
8. Superior hæmorrhoidal a.

PLAN OF THE MESENTERIC ARTERIES AND THEIR COMMUNICATIONS.[2]

the duodenum (p. 487), and then runs between the layers of the mesentery toward the right iliac fossa, where it terminates in branches for the supply of the cæcum. Thus it describes a gentle curve from left to right. It is crossed by the pancreas and splenic vein, and will be seen to supply the descending and transverse duodenum, the jejunum, ileum, and the ascending and trans-

[1] The arrangement of the chyliferous vessels is well displayed in the plates of Mascagni.

[2] The inferior pancreatico-duodenal artery is not represented.

verse colon. It is accompanied by its corresponding vein, and is surrounded by the superior mesenteric sympathetic plexus. It gives off the following branches :—

1. The *inferior pancreatico-duodenal* branch, which runs up behind the pancreas, within the concavity of the duodenum, to inosculate with the *superior pancreatico-duodenal* branch of the hepatic.

2. *Vasa intestini tenuis* of the small intestine, from ten to sixteen in number, are given off from the left or convex side of the curve, and are distributed to the jejunum and ileum; while from the concave side come—

3. The *ileo-colic;*

4. The *right colic*, for the supply of the ileum, cæcum, and ascending colon; and

5. The *middle colic*, for the supply of the transverse colon.

The student should now trace the branches to the small intestine, in order to see the series of arches which they form by their mutual inosculations. There are three or four tiers of them, each tier composed of smaller and more numerous branches than the preceding. The ultimate branches ramify in circles round the intestine. This circular arrangement of the vessels in the coats of the bowel is practically interesting, because it enables one in almost all cases to distinguish the intestine from the hernial sac.

The *colic* branches of the superior mesenteric are the *ileo-colic*, which is the continuation of the main trunk, and divides into two branches; one supplies the lower part of the ileum, and the other thecæcum;

The *right colic*, which proceeds toward the ascending colon; and

The *middle colic*, which ascends between the layers of the meso-colon to the arch. They are arranged after the same plan as those of the small intestine; that is, they inosculate and form a series of arches, which successively decrease in size, and finally terminate in circles round the bowel.

The *superior mesenteric vein* joins the splenic behind the pancreas, and forms the vena portæ (p. 40).

Dissection of the Inferior Mesenteric Artery and Branches.

To trace this artery, the small intestine must be drawn over toward the right side, and the peritoneum covering the artery removed, since the artery lies behind the peritoneum. It is given off from the front of the aorta, about two inches above its bifurcation, and is surrounded by the inferior mesenteric plexus of sympathetic nerves. Descending toward the left iliac fossa, it crosses obliquely over the left common iliac artery, passes between the layers of the meso-rectum, and, taking the name of *superior hæmorrhoidal*, is finally distributed to the upper part of the rectum. Its branches are :—

1. The *colica sinistra*, which crosses behind the peritoneum, over the left kidney, and supplies the descending colon.

2. The *sigmoidea*, which runs over the psoas, is distributed to the sigmoid flexure.

3. The *superior hæmorrhoidal*, which supplies the upper part of the rectum, and will be dissected with the side view of the pelvis.

These branches of the inferior mesenteric inosculate in the form of arches, like the colic branches of the superior mesenteric. The colica sinistra, too, forms a large arterial arch with the colica media, so that there is a chain of arterial communications from one end to the other of the intestinal canal (Fig. 111).

The *inferior mesenteric vein* ascends nearly vertically behind the peritoneum, passes in front of the left psoas, behind the third portion of the duodenum and the pancreas, and joins the splenic behind the pancreas.

Dissection.

To see the relations of the duodenum and the pancreas, two ligatures, about an inch apart, should be placed on the upper end of the jejunum, and two others, at a similar distance apart, on the lower end of the sigmoid flexure of the colon. After the jejunum and the sigmoid flexure have been divided between the ligatures respectively, the small and large intestines are to be removed by cutting through the peritoneal folds which connect them to the abdominal walls. By turning up the stomach, we expose the duodenum curving round the great end of the pancreas.

Duodenum; Relations of.

The duodenum (p. 487) commences at the pyloric end of the stomach, and terminates on the left side of the second lumbar vertebra, where the intestinum jejunum begins. It is about eight to ten inches in length, and is divided into three parts, an ascending, descending and transverse.

The first portion ascends obliquely as high as the neck of the gall bladder; then, making a sudden bend, it descends in front of the right kidney as low as the third lumbar vertebra. Lastly, making another bend, it ascends obliquely across the spine to the left side of the second lumbar vertebra; here the intestine takes the name of jejunum. Thus the duodenum describes a horse-shoe curve, the concavity of which is directed toward the left side, and embraces the head of the pancreas.

The first or *ascending* portion is about two inches long, and is completely invested by peritoneum. It is comparatively free, so that the movements of the stomach may not be restricted. In front of it are the liver and the neck of the gall bladder. Behind it are the bile duct, the hepatic artery, and the vena portæ. The second or *descending* portion is about three inches long, and is covered by peritoneum only on its anterior surface. It is firmly connected to the deeper structures behind, and to the pancreas on its left side, so that no movement is permitted in this portion. It descends from the neck of the gall bladder to the right side of the body of the third lumbar vertebra. It lies behind the transverse colon, in front of the right kidney and the ductus communis choledochus; on the left side it is in relation with the head of the pancreas, its duct, and the superior and inferior pancreatico-duodenal arteries. The third or *transverse* portion, about four inches long, is situated behind the transverse meso-colon, just above the mesentery and the superior mesenteric vessels. Above are the pancreas, and the superior mesenteric artery and vein which pass between the pancreas and the duodenum; behind, it rests upon the crura of the diaphragm, the inferior vena cava and the aorta. This portion, like the second, is only covered in front by peritoneum. Notice how firmly the duodenum is braced up on the left side of the second lumbar vertebra; and how the jejunum begins here by an abrupt downward bend.

Pancreas; Relations of. The pancreas is a large, compound racemose gland, situated immediately behind the stomach (p. 490). It is of an elongated form, and of pinkish-white color. It is placed transversely across the spine; its larger end, or *head*, is embraced by the duodenum; its lesser end, or *tail*, is in contact with the spleen. It is about six to eight inches in length, its average breadth is one inch and a half, and its thickness from half an inch to an inch. Its weight is from 2¼ oz. to 3½ oz., although it frequently exceeds the latter weight.

In *front*, the gland has the ascending layer of the transverse meso-colon and the stomach; its *right extremity*, or *head*, is embraced by the duodenum, separated from it by the pancreatico-duodenal arteries; *behind* the head is the ductus communis choledochus, while the body is in relation posteriorly with the inferior vena cava, the superior mesenteric vein and artery, the aorta, the beginning of the vena portæ, the crura of the diaphragm, the left kidney, the supra-renal capsule, and the inferior mesenteric vein; its *left extremity*, or *tail*, touches the concavity of the lower surface of the spleen, and is in front of the left supra-renal capsule; the *upper border* is in relation with the cœliac axis, the splenic artery and vein lying in a groove in the gland, and on the right side with the ascending portion of the duodenum and the hepatic artery; the *lower border* is in relation with the transverse portion of the duodenum, from which it is separated by the superior mesenteric vessels, and to the left side with the inferior mesenteric vein.

Its *duct* (*canal of Wirsung*) runs from left to right, near the lower border and anterior surface of the gland, and empties itself into the back part of the descending portion of the duodenum, conjointly with, or close to, the opening of the common bile duct. It receives numerous branches from the splenic artery, which runs along its upper border; some from the superior mesenteric, which lies immediately beneath it, and others from the gastro-duodenalis.

Dissection. The liver, stomach, duodenum, pancreas and spleen should now be collectively removed. For this purpose it is necessary to cut through the ligaments of the liver, the venæ cavæ hepaticæ, and the branches of the cœliac

axis. These viscera, with the remainder of the intestinal canal, should be macerated in water, while you examine all that is to be seen at the back of the abdomen; namely, the deep-seated muscles, the aorta, the inferior vena cava, the kidneys, the lumbar plexus of nerves and the sympathetic nerve.

Kidneys and Ureter; Relations of. The kidneys, two large glands which excrete the urine, are situated in the lumbar region, behind the peritoneum, one on each side of the spine. They extend from the eleventh rib nearly as far as the crest of the ilium, and lie embedded in more or less fat, on the quadratus lumborum, the psoas and the crura of the diaphragm. The weight of the kidney is from 4½ oz. to 8 oz., the left weighing a little more than the right, and the right is somewhat the lower of the two. Surmounting each is a small body, called the supra-renal capsule.

The *anterior surface* is convex, and the right kidney has *in front* he liver, the ascending colon, the descending portion of the uodenum, and the colica dextra artery; the left kidney has *in front* he lower part of the spleen, the cardiac end of the stomach, the escending colon, the tail of the pancreas and the colica sinistra rtery. This explains how it is that a renal abscess or calculus sometimes evacuated by the rectum. *Above*, the right kidney is contact with the under surface of the liver, and its upper end eaches as high as the lower border of the eleventh rib; the left idney is in contact above with the spleen, and reaches to the level f the upper border of the eleventh rib. The *posterior surface* is at and lies on the corresponding crus of the diaphragm, the uadratus lumborum, and the psoas, separated, however, by the nterior layer of the aponeurosis of the transversalis; the *outer order* is convex, and looks toward the parietes; the *inner border* resents a deep notch, the *hilum*, continuous with a cavity, the *inus*, through which pass the renal artery and vein, the ureter, he renal plexus of nerves and lymphatics, surrounded by conective tissue and fat. The vessels and duct have the following elations: anteriorly is the renal vein, posteriorly is the ureter, the enal artery being between them. The ureter descends almost ertically on the psoas muscle, enters the pelvis over the division

of the common iliac artery, and empties itself into the lower part of the bladder after running obliquely through its coats. The kidney, below, descends nearly as low as the crest of the ilium, and is not so broad as the upper extremity. The kidneys and supra-renal capsules must be removed and reserved for further examination.

Dissection.

Semilunar Ganglia. The *semilunar ganglia*, two in number, are contained in the solar plexus, and are situated one on each side of the cœliac axis, in the neighborhood of the supra-renal bodies; that on the right side will be found lying under the vena cava inferior. They consist of irregular ganglionic masses. Above, each ganglion receives the great splanchnic nerve (p. 208), and the two ganglia are connected on their inner sides. Filaments are distributed to the supra-renal and renal plexuses and to the plexuses which surround the branches of the abdominal aorta. The branches of the solar plexus will be described later on.

Diaphragm.[1] This is a partly muscular and partly tendinous arch, so constructed as to form a complete movable partition between the chest and the abdomen; a floor for the one, and a roof for the other. Its upper or thoracic surface is convex; its lower or abdominal, concave. On removing its peritoneal coat, and a thin fascial covering from the transversalis fascia, we observe a broad tendon in the centre, and that muscular fibres converge to it from all sides (Fig. 112). The diaphragm *arises*, 1. From the ensiform cartilage by fleshy fibres; 2. From the inner surfaces of the cartilages of the six lower ribs by as many digitations, which correspond with those of the transversalis; 3. From two thin tendinous arches, called, respectively, the *ligamenta arcuata*, *externum* and *internum* (the external arch is the thickened upper border of the anterior layer of the transversalis fascia, and extends from the last rib to the transverse process of the first lumbar vertebra, and arches over the quadratus lumborum; the internal passes from the transverse process of the first lumbar vertebra to the body of the same vertebra, and arches over the psoas); and 4. From the front of the bodies of the lumbar vertebræ by two elongated bundles, called the *crura* of the diaphragm.

[1] διάφραγμα, a partition wall.

ra have tendinous origins; the right crus is, however, a ger than the left; the former arises from the first, second l lumbar vertebræ and their intervening cartilages; the not descend so low, by one vertebra. The inner fibres crus decussate; those of the right being the more ante- their decussation the fibres separate the aortic from the eal openings. Between the two crura the aorta enters men.

these various origins the fibres ascend, at first nearly ver- nd then all arch inward, and converge to be *inserted* central tendon.

FIG. 112.

ing be- crura. of œso- or vena

4. Quadratus lumborum.
5. Psoas magnus.

The dark arches above the quadratus and psoas are the 'arcuate ligaments.'

AM OF THE DIAPHRAGM, THE OPENINGS IN IT, AND THE PHRENIC ARTERIES.

ntral tendon is nearly the highest part of the diaphragm. ts a white, glistening surface, owing to the crossing of its s fibres; its shape may be compared to that of a trefoil it is composed of right and left leaflet and a middle parated from each other by indentations. Of the three he right is the largest and the left the smallest. The nt of interest about the tendon is that, in consequence nections with the pericardium, below which it lies (p. is always maintained nearly on the same level; so elps to support the heart, and serves as a fixed point sertion of the muscular fibres of the diaphragm.

FIG. 113.

DIAPHRAGM FROM ITS UPPER SURFACE.
(The dotted lines show the amount of descent on contraction.)

Openings in the Diaphragm. There are three large openings in the diaphragm for the transmission of the aorta, the œsophagus, and the inferior vena cava, respectively, and several smaller apertures for the transmission of nerves and vessels. The *aortic opening* is osso-aponeurotic, and lies in the middle line between the two crura in front of the spine; it transmits, also, the vena azygos major and the thoracic duct, both of which lay rather to the right of the aorta. Trace the crura upward, and observe that the inner fibres of each cross each other in front of the aorta, somewhat like the letter X.[1] Above the decussation, and a little to the left of it, is the *œsophageal opening;* this is oval and entirely muscular, and transmits the œsophagus and the pneumogastric nerves. The *opening* for the *vena cava* (*foramen quadratum*) is situated in the highest part of the central tendon, rather to the right of the middle line, and is quadrate in shape. Through this opening pass the inferior vena cava, some lymphatics from the convex surface of the liver, and usually branch from the right phrenic nerve. Observe that the vein intimately connected to its margin, and kept permanently open. Lastly, there pass *through the crus*, on each side, the sympathetic and the greater and lesser splanchnic nerves; and in addition, on the left side, the vena azygos minor. The arch of the diaphragm, in expiration, extends about as high as the fifth rib on the right side, and the sixth rib on the left.

The nerves of the diaphragm are the phrenic (p. 199), and five or six lower intercostal nerves. The diaphragm also rec

[1] This decussation is not always complete. But the right crus always more or less over the left, so that the crura are never strictly parallel.

nute filaments from the diaphragmatic plexuses, which come
n the semilunar ganglia. On its under surface, on the right
e, close to the supra-renal capsule, the plexus joins some
nches of the right phrenic nerve, at which spot there is a small
glion (*ganglion diaphragmaticum*), from which filaments are
en off to the liver, vena cava, and supra-renal capsule. It is
ent on the left side. Its blood vessels are the two phrenic,
ived from the aorta, the internal mammary (p. 178), and the
er intercostal.

unction of the Diaphragm. The diaphragm is the great muscle concerned in inspiration. It may be said, with Haller, that s "musculus post cor nobilissimus." During inspiration the scular sides of the diaphragm contract, and become less arched shown by the dotted line in Fig. 113); the floor of the chest sinks consequence, and more room is made for the expansion of the gs. During expiration the diaphragm relaxes, and the air is elled, partly by the elasticity of the lungs and the thoracic walls, tly by muscular action. This alternate sinking and rising of diaphragm constitutes a chief part of the mechanism of breath-. But the diaphragm conduces to the performance of many er functions. Acting in concert with the abdominal muscles, ssists in the expulsion of the fæces and the urine, also in partu-on and in vomiting; for in all these operations we first take in eep breath, that the diaphragm may be in a state of contraction, l so form a resisting surface, against which the viscera may be npressed by the abdominal muscles. Moreover, by its rapid or smodic contractions it is one of the chief agents concerned in ghing, sneezing, coughing, hiccough.

Dissection. The student should now dissect the large vessels and the muscles of the back part of the lomen. To do so, the mesentery which lies in front of the ta and vena cava is to be removed, as well as the fat and con-tive tissue. The dissection should include the parietal branches the abdominal aorta; afterward, its great primary divisions—common and external iliac arteries—should be cleaned as far Poupart's ligament. The quadratus lumborum, the psoas l iliacus muscles should be carefully cleaned, care being taken

not to injure the nerves and arteries lying in front of them; thus, in front of the quadratus lumborum are the last dorsal, the ilio-hypogastric, and ilio-inguinal nerves, which cross the muscle obliquely; in front of the iliacus are the external cutaneous and anterior crural nerves; and coming through, and then lying in front of the psoas, is the genito-crural nerve, while to the inner side of the muscle is the obturator nerve. The gangliated cord of the sympathetic nerves, situated on each side of the bodies of the vertebræ, must also be made out; and, lastly, the sheath which invests the psoas should be examined, and the branches of the lumbar plexus preserved as they emerge from beneath the outer border of the muscle.

Before examining the course of the aorta, notice that a chain of lymphatic glands extend along the brim of the pelvis and the bodies of the lumbar vertebræ, following the course of the great blood vessels. Generally speaking, they are small; only one here and there attracts observation. They transmit the lymphatics from the lower limbs, the abdominal wall and the testicle; and all eventually lead to the *receptaculum chyli*, or beginning of the thoracic duct (p. 203). This is usually found on the right of the aorta, close to the second lumbar vertebra.

Course and Relations of the Abdominal Aorta. The *abdominal aorta* enters the abdomen between the crura of the diaphragm in front of the body of the last dorsal vertebra, and descends a little to the left side of the front of the spine, as low as the middle of the fourth lumbar vertebræ, where it divides into the two common iliac arteries. It follows the curve of the lumbar convexity, attaining its greatest curve on a level with the third lumbar vertebra. It rapidly lessens in size, owing to the large branches it gives off in its course. The division of the aorta into the two common iliac arteries is about the level of the highest point of the crest of the ilium, and just below the left side of the umbilicus. The artery has *in front* of it the stomach and lesser omentum, the solar plexus surrounding the cœliac axis, splenic vein, the pancreas, the transverse portion of the duodenum, the left renal vein, the mesentery, the aortic sympathetic plexus and a chain of lymphatic glands. To the *right side* of

ie the right crus of the diaphragm, the inferior vena cava, the thoracic duct, the vena azygos and the right semilunar ganglion. To the *left side* of it are the left crus, the left semilunar ganglion and the sympathetic nerves. *Behind*, it rests on the receptaculum chyli, the thoracic duct, the left lumbar veins, and the anterior common ligament.

The branches of the aorta still to be examined arise from it in airs—namely, the phrenic, capsular, renal, spermatic and lumbar ee diagram, p. 504).

hrenic Arteries. These arteries supply the under surface of the diaphragm, and occasionally arise separately, usuy by a common trunk, from the aorta, after its passage under e crura of the diaphragm (Fig. 114). The *right phrenic* passes tward, behind the liver and the inferior vena cava, and then gets to e right side of the caval opening in the diaphragm; the *left phrenic* ends behind the œsophagus, and gets to the left side of the sophageal opening. Each ascends, lying on its corresponding us, as far as the central tendon, where it divides into two branches: ie, the external, passes transversely across the tendon to the side the diaphragm; the other, the internal branch, which seems to be e continuation of the artery, runs forward to the anterior part of e muscle. Their first branches are to the supra-renal capsules; en, the internal branch of the right gives off a small branch to e vena cava, the corresponding branch of the left sends one to e œsophagus. Moreover small branches are distributed respecely to the liver and to the spleen. They inosculate with each ier, with the musculo-phrenic branches of the internal mamary, and the intercostal arteries. The *right phrenic vein* termites in the inferior vena cava; the *left vein* in the renal vein, if t in the vena cava.

Supra-Renal Arteries. The *supra-renal* or *capsular arteries* are two very small branches, given off from the aorta, one each side, opposite to the superior mesenteric artery; each ns upon the crus of the diaphragm, the right behind the inferior va, and is distributed to the supra-renal body, inosculating with anches from the phrenic and renal arteries. The *right capsular in* terminates in the inferior cava, the *left* in the left renal vein.

FIG. 114.



DIAGRAM OF THE COURSE AND RELATIONS OF THE ABDOMINAL AORTA AND VENA CAVA INFE[illegible]IOR.

1. Ilio-hypogastric nerve.
2. Ilio-inguinal n.
3. External cutaneous n.
4. Anterior crural n.

tenal Arteries and Veins. The *renal* arteries arise from the aorta immediately below the superior mesenteric artery, and n transversely to the hila of the kidneys. Both are covered by ir corresponding veins. The right is longer and rather lower n the left, and passes behind the vena cava. Each, after sending a small branch to the supra-renal body and ureter, enters kidney, not as a single trunk, but by several branches, corresding to the original lobes of the organ. The *renal veins* lie ront of the arteries, and join the vena cava at right angles. The is longer than the right, and crosses over the aorta; it also re-es the spermatic, capsular, and the phrenic veins of its own side.

Spermatic Arteries and Veins. The *spermatic arteries*, two in number, arise from the front of the aorta, a little below the renal, and pass to the testes in the male and to ovaries in the female. Each runs down behind the peritoneum, iquely over the psoas, crossing over the ureter, and the front of external iliac artery immediately above the crural arch; the ht artery in addition lying over the vena cava. Each then passes ough the internal abdominal ring and inguinal canal, with the er constituents of the spermatic cord, to the testicle, where it omes tortuous, and divides into several branches, some of which mpany the vas deferens and supply the epididymis; others ply the testis by piercing the tunica albuginea. Each artery is mpanied, below the external abdominal ring, by a very convod plexus of veins—*pampiniform plexus*. At the inner ring they inate in two tortuous veins, which unite before they empty nselves, on the *right* side, obliquely, into the vena cava; on *left* side, at right angles, into the left renal vein, after passing nd the sigmoid flexure of the colon. In the female, the *ian arteries* descend toward the pelvis, and lie between the layers of the broad ligament to be distributed to the ovaries, e branches also going to the Fallopian tubes, and one to the of the uterus to anastomose with the uterine artery of the rnal iliac. They likewise send small offsets to the round ligat, and thence to the skin of the pubes and groin.

mbar Arteries d Branches. There are usually five of these arteries on each side; four arise from the back of the aorta, the

fifth comes from the arteria sacra media. They are analogous to the intercostal arteries on a small scale. They proceed outward over the bodies of the vertebræ beneath the sympathetic nerve and the arches formed by the psoas muscle; the two upper pass beneath the crura of the diaphragm; those on the right side being also behind the vena cava inferior. Passing toward the intervertebral foramina, they, like the intercostals, divide into dorsal and abdominal branches.

The *dorsal branches* pass between the transverse processes of the vertebræ, accompanied by the posterior branches of the corresponding nerves, and are of a size proportionate to the large development of the muscles of the back which they supply. They also send *spinal branches*, which enter the spinal canal through the intervertebral foramina; some of these are distributed to the anterior part of the cauda equina, and others to the bodies of the lumbar vertebræ, forming a series of arches behind them.

The *abdominal branches* all run outward behind the quadratus lumborum, except the last, which usually runs in front. After supplying the quadratus and psoas, they pass forward between the abdominal muscles and supply the walls of the abdomen.[1] They anastomose, *laterally*, with the ilio-lumbar and circumflex iliac arteries; *in front*, with the internal mammary and epigastric arteries; and *above*, with the intercostals.

The *lumbar veins* empty themselves into the vena cava inferior, the left passing behind the aorta.

The *arteria sacra media*, a diminutive continuation of the aorta, proceeds from its bifurcation, and runs down, behind the left common iliac vein and in front of the sacrum, to the coccyx. It sends off the fifth lumbar artery and lateral branches, which anastomose with the lateral sacral arteries; it also supplies small vessels to the posterior part of the rectum, which run forward between the layers of the meso-rectum. In close connection with the terminal branch of this artery at the tip of the coc-

[1] Just as the thoracic intercostals, by communicating with the internal mammary, form an arterial ring round the chest, so do the lumbar, by communicating with the epigastric, form a similar, though less perfect, ring round the walls of the abdomen.

x is a small, roundish body called the *coccygeal* or *Luschka's ind*.[1] It is about the size of a pea, and is placed between the ·atores ani and the sphincter ani. It is probably composed of plexus of small arteries, which are surrounded by one or more yers of granular polygonal cells. The body itself is invested · connective tissue, in which also some branches from the gan- ion impar have been traced. This gland should be considered an arterial gland, of which the intercarotic ganglion is another xample.

The *vena sacra media* empties itself into the left common iliac ein.

Vena Cava Inferior.

The *vena cava inferior* is formed by the junction of the two common iliac veins, a little to the right de of the intervertebral cartilage, between the fourth and fifth mbar vertebræ. It ascends in front of the spine, in the greater rt of its course lying to the right of the aorta. As it approaches diaphragm, the vena cava inclines a little to the right, sepa- ed from the aorta by the right crus of the diaphragm. It is n received into a deep groove on the posterior border of the er, and afterward passes through the tendinous opening in the phragm to reach the right auricle of the heart. Its relations, ginning from below, are—*in front*, the mesentery, the third part the duodenum, the pancreas, the right spermatic artery, the rtal vein, and the liver; *behind* it are the right renal artery, the ht lumbar arteries, the sympathetic of the right side, and the ht phrenic artery; on its *left* is the aorta, and higher up the ht crus. It receives the lumbar veins, the right spermatic (the t joins the renal), the renal, the right supra-renal, the right renic and the hepatic veins, which are usually three in number, e each from the right and left lobes and one from the lobulus igelii.

The student should now direct his attention to the three large uscles which are seen at the back of the abdomen; the quad- us lumborum, situated between the last rib and the crest of the um; the iliacus, occupying the iliac fossa; and the psoas

[1] Luschka, *Anatomie des Menschen*, 1864, vol. ii, pt. 2, p. 187; Arnold, *Virchow chiv*, 1864, 1865, and 1866; Callender, *British Medical Journal*, June 13, 1874.

magnus, passing from the sides of the lumbar vertebræ, along the brim of the pelvis, and beneath Poupart's ligament, to be inserted into the lesser trochanter of the femur. The nerves should at the same time be cleaned, and, if more convenient, the student might dissect the muscles on one side and the nerves and arteries on the other.

Psoas Fascia. The fascia covering the iliacus and psoas muscles is seen to be thin above and thicker below. It consists of two portions, the psoas and the iliac fascia. The *psoas fascia* is attached to the sides of the lumbar vertebræ and their intervertebral cartilages, internally to the sacrum, and above to the ligamentum arcuatum internum; externally it is thinner, and is continuous with the fascia lumborum. It is this sheath which determines the ordinary course of a psoas abscess —namely, beneath the crural arch into the upper part of the thigh; for it is a rare exception when the matter travels into the pelvis.

Iliac Fascia. The *iliac fascia* covers the iliacus muscle, and is attached to the inner lip of the crest of the ilium, and to the brim of the pelvis, through its connection with the sheath of the psoas and the tendinous insertion of the psoas parvus. Its most important attachment is to the outer half of the crural arch, and it is here directly continuous with the fascia transversalis (p. 455), so that together they present an effectual barrier to the escape of intestine beneath this part of the arch.[1] On the inner half of Poupart's ligament, these two fasciæ are separated by the femoral vessels, so that the fascia transversalis lies in front of the fascia iliaca, behind the artery and vein, thus forming their sheath as they pass down the thigh. This portion of the iliac fascia also forms a sheath for the psoas and iliacus as far as their insertion, and becomes continuous with the iliac portion of the fascia lata. Internally the iliac fascia is attached

[1] The iliac fossæ are liable to be the seat of suppuration, and the course which the pus takes depends upon its position with regard to the iliac fascia. If the pus be seated in the loose cellular tissue between the peritoneum and the fascia, it usually advances just above the crest of the ilium, or toward the groin through the inguinal canal; but, if seated beneath the fascia, the matter usually makes its way under the crural arch toward the upper and outer part of the thigh.

to the linea ilio-pectinea, where it becomes continuous with the pubic portion of the fascia lata. The external iliac artery and vein lie in front of the fascia, while the anterior crural nerve is behind it.

These fasciæ are now to be dissected off, when the psoas and iliacus will be completely exposed.

Psoas Magnus. This long muscle *arises* by five muscular fasciculi from the transverse processes of all the lumbar vertebræ, also from the bodies of the last dorsal and all the lumbar vertebræ and their intervening fibro-cartilages, but only from the projecting borders of their bodies, not from the central concave part; here the fibres arise from tendinous arches thrown over the lumbar vessels. The muscle descends vertically along the brim of the pelvis, beneath the crural arch into the thigh, and is *inserted* by a strong tendon into the back part of the lesser trochanter of the femur. In *front*, the psoas has in relation with it the psoas fascia, the ligamentum arcuatum internum, the common and external iliac artery and vein, the kidney and ureter, the spermatic vessels, the genito-crural nerve, the colon, the renal vessels, the vas deferens, the psoas parvus when present; *behind* it are the ilium, the transverse processes of the lumbar vertebræ, the quadratus lumborum, the anterior layer of the transversalis aponeurosis, the lumbar plexus of nerves, and the obturator nerve, which lower down runs along the inner border of the muscle; on its *inner side* are the crus of the diaphragm, the bodies of the lumbar vertebræ, the lumbar arteries, the lumbar glands, and the sympathetic nerves; to the inner side of the left psoas is the aorta, and to that of the right psoas is the inferior vena cava. Toward its insertion the tendon of the psoas lies between the iliacus and pectineus.

As it passes under the crural arch, the tendon of the psoas lies immediately over the capsule of the hip joint, and there is a large bursa between them to facilitate the play of the tendon. It should be borne in mind that occasionally, even in young subjects, but more frequently in old ones, in consequence of wear and tear, this bursa communicates with the hip joint. The fact is important,

for it explains how a psoas abscess sometimes makes its way into the hip joint, a result frequently fatal.

Psoas Parvus. Once in about eight or ten subjects there is a small muscle called the *psoas parvus* placed superficially to the preceding muscle. It *arises* from the sides of the bodies of the last dorsal and the first lumbar vertebræ, and the intervening fibro-cartilage; thence, descending in front, and to the inner side of the psoas magnus, it ends in a long, flat tendon, which spreads out, and is *inserted* into the linea ilio-pectinea.

Iliacus. The iliacus *arises* from the iliac fossa, the inner lip of the crest of the ilium, the ilio-lumbar ligament,[1] the base of the sacrum, and the anterior superior spine of the ilium; below, also, from the anterior inferior spine and from the capsule of the hip joint. The fibres converge beneath the crural arch, and are *inserted* mainly into the outer side of the tendon of the psoas, and partly into the triangular surface of the femur, below and anterior to the lesser trochanter. Thus the two muscles, so far as their action goes, may be considered as one, and are sometimes called the ilio-psoas.

The iliacus in the abdomen is in relation, *in front*, with the iliac fascia, the external cutaneous and anterior crural nerves, with the cæcum on the right, and with the sigmoid flexure on the left side; *behind*, it is in contact with bone; on its *inner side* is the psoas. In the thigh it is in relation, *in front*, with the rectus, the sartorius, the fascia lata, the anterior crural nerve, the profunda femoris and the external circumflex arteries; *behind*, with the capsular ligament of the hip joint; on the *inner side*, with the psoas; and on the *outer*, with the crureus and vastus internus muscles.

The combined *action* of the psoas and iliacus is to assist in raising the body from the recumbent position, and to fix the pelvis steadily on the thigh; this supposes the fixed point to be at the trochanter minor. If the fixed point be at the spine, then the muscle flexes and rotates the femur outward. It is this action which often occasions so much displacement in fractures of the upper third of the femur.

[1] This ligament extends from the transverse process of the last lumbar vertebra to the ilium.

uadratus nborum and s Sheath.

This quadrilateral muscle extends from the crest of the ilium to the last rib, and is contained in a sheath formed for it by the aponeurotic in of the transversalis (p. 451). The anterior layer of its ath is attached to the roots of the *transverse* processes of the nbar vertebræ, and the posterior layer to their summits. The uscle, broader below than above, *arises* by two portions—one m the ilio-lumbar ligament and from the crest of the ilium for o inches external to it, and is *inserted* into the last rib, and by ndinous slips into the apices of the transverse processes of the per four lumbar vertebræ; the other portion of the muscle ises from the transverse processes of the third, fourth and fifth mbar vertebræ, and is inserted into the lower margin of the last , in front of the preceding portion. The principal use of the uscle is to steady the spine; it also steadies the last rib, and nables it to serve as a fixed point for the action of the intercostal uscles and the diaphragm.[1]

By raising the quadratus we observe the aponeurotic origin of he transversalis from the summits of the transverse processes; his constitutes the posterior part of its sheath, and separates the uscle from the erector spinæ.

Common Iliac Arteries and Veins.

The abdominal aorta divides, in front of the left side of the fourth lumbar vertebra, into two great branches, termed the *common iliac arteries*. They diverge at an acute angle, and after a course of about two inches downward and outward, each divides, over the sacro-iliac symphysis, into the external and internal iliac arteries. They lie upon the bodies of the fourth and fifth lumbar vertebræ. The right common iliac is rather larger than the left. They are covered *in front* by peritoneum; they are crossed by branches of the sympathetic to form the hypogastric plexus, and they are

[1] The respective attachments of the quadratus lumborum, the crossing of its fibres, and its mode of action, lead to the inference that it is a large intercostal muscle. It is worth remembering that the outer edge of the quadratus lumborum, in a well-grown adult, is about three inches from the spines of the lumbar vertebræ, and midway between the last rib and the crest of the ilium. It is just outside the edge of this muscle that we can cut down to open the large bowel without wounding the peritoneum.

crossed at or near their division by the ureters; on the *outer side* they are in relation with the psoas. So far, then, the relations of both common iliac arteries are similar. But each has its special relations, as follows :—

The special relations of the right common iliac are, that it has *behind* it the two common iliac veins, which separate it from the fifth lumbar vertebræ; on its *outer side*, it has, above, the inferior vena cava; below, the right common iliac vein.

The special relations of the left common iliac are, that it has *in front* of it the end of the sigmoid flexure of the colon, and the superior hæmorrhoidal artery; and to its *inner side*, the left common iliac vein, which gradually gets more behind it toward the sacro-iliac symphysis.

The relations of these arteries with regard to their corresponding veins are, practically, important, in reference to the operation of tying them. This operation is, obviously, easier on the left side than the right.[1]

If the common iliac artery were ligatured, the collateral circulation would be maintained through the following vessels: by the internal mammary anastomosing with the deep epigastric a.; by the lumbar arteries with the circumflex iliac and the ilio-lumbar a.; by the lumbar with the gluteal; by the middle sacral with the lateral sacral a.; by the spermatic with the deferential, cremasteric, external pudic and superficial perineal arteries; by the superior hæmorrhoidal with the middle and inferior hæmorrhoidal a.; by the lower intercostals with the epigastric a. (superficial and deep); and by the middle and the inferior hæmorrhoidal, the pudic and its branches, and the vesical arteries communicating in the middle line with the corresponding branches of the opposite side.

[1] The length of the common iliac artery is apt to vary in different persons. We have seen it from three-fourths of an inch to three and a half inches long. These varieties may arise either from a high division of the aorta, or a low division of the common iliac, or both. It is impossible to ascertain, beforehand, its length in any given instance, as there is no necessary relation between its length and the height of the adult individual. It is often very short in men of tall stature, and *vice versâ*. The left is usually described as rather longer than the right; but, from the examination of 100 bodies, our conclusion is that their average length is the same.

External Iliac Artery. The *external iliac artery*, the continuation of the preceding artery, passes along the brim of the lvis, first on the inner side, and then in front of the psoas. ower down it passes under the crural arch, midway between the terior superior spine of the ilium and the symphysis pubis, here it takes the name of femoral. The artery has *in front* of the peritoneum, the intestines, and a sheath derived from the ac fascia investing the artery and the vein; it has also the ermatic vessels in front, and it is crossed by a branch of the enito-crural nerve, the deep circumflexa ilii vein and the vas eferens; *behind*, it is in relation with the psoas magnus and corsponding vein, which lies also on the inner side of the artery; e iliac fascia also lies behind the vessels, but a thin layer of cia derived from it is continued over them; *internally*, it has corresponding vein, and low down toward Poupart's ligament, vas deferens; *externally*, it has the psoas magnus and the iliac cia. In front of, and on the inner side of the artery are a chain lymphatic glands.

The branches given off by this artery are:—

a. *Small branches* to the psoas and lymphatic glands.

b. The *deep epigastric*, already described (p. 457).

c. The *deep circumflexa ilii*, which arises from the outer side the artery, just above the crural arch, and, ascending upward ward the anterior superior spine of the ilium, in a sheath formed the fascia iliaca, runs along the inner aspect of the iliac crest, d subsequently perforates the transversalis muscle.[1] In the section of abdominal muscles (p. 450), the continuation of it is seen skirting the crest of the ilium between the internal lique and the transversalis, and sending a branch upward tween these muscles for their supply. The main trunk, much duced in size, inosculates with the ilio-lumbar, derived from the ternal iliac.

Ligature of the External Iliac. The best way of tying the external iliac is to make a curved incision at the lower part of the domen, beginning a little above the middle of the crural arch, d ending about an inch to the inner side of the spine of the ium. The strata of the abdominal muscles, with the fascia

[1] The course of this artery should be borne in mind in opening iliac abscesses.

transversalis, should then be divided to the same exte which, the peritoneum with the spermatic vessels must rated by the fingers from the iliac fossa. It is necessary a small incision through the sheath of the vessels, to the passage of the needle. Remember that the vein connected to its inner side,[1] that the genito-crural ne far off, and that the circumflex iliac vein crosses the a above Poupart's ligament.

After ligature of the artery the collateral circulation maintained by anastomoses between the internal mam the deep epigastric; between the lumbar arteries and th flex iliac; between the pubic branch of the obturato branch of the epigastric; between the gluteal and th and external circumflex; between the sciatic and the f rating and the internal circumflex; between the obtu the internal circumflex; between the spermatic, the d and the cremasteric and the external pudic; between intercostals and the lumbar arteries and the epigastric a

Sympathetic Nerve.

The general plan upon which the sy nerve is arranged has been noticed in t tion of the neck (p. 170). The lumbar portion of it mu examined.

The lumbar portion of the sympathetic descends on in front of the bodies of the lumbar vertebræ, along

[1] This relative position of the vessels does not always exist. In old frequently in adults, it is sometimes found that the external iliac artery r tuously, instead of nearly straight, along the brim of the pelvis. But t not follow the artery in its windings, and may possibly lie outside th where we propose to place the ligature.

The mode of performing the operation described in the text is recon Sir A. Cooper. Mr. Abernethy, however, who first set the example artery in 1796, adopted a somewhat different proceeding. He says: " an incision, about three inches in length, through the integuments of t in the direction of the artery, and thus laid bare the aponeurosis of oblique muscle, which I next divided from its connection with Poupar in the direction of the external wound, for the extent of about two i margins of the internal oblique and transversalis muscles being thus introduced my finger beneath them for the protection of the peritoneu divided them. Next, with my hand, I pushed the peritoneum and upward and inward, and took hold of the artery."

border of the psoas. The nerve has an oval ganglion of grayish color, opposite each lumbar vertebra, so that there are either four or five of them on each side. These ganglia are connected together by small filaments of a white color, and each ganglia receives, on its outer side, two branches from the corresponding spinal nerve, as in the chest; other branches pass inward, and form, in front of the aorta, the *aortic plexus;* and some pass downward over the common iliac arteries to form the hypogastric plexus.

Solar Plexus and the Semilunar Ganglia.

The *solar* or *epigastric plexus* is situated in front of the aorta, and surrounds the cœliac axis in a dense network of nerve filaments, in which are several ganglia. It receives the great splanchnic nerves, part of the lesser splanchnic, and some branches from the pneumogastric nerves. The solar plexus gives off filaments which form plexuses surrounding the various branches of the abdominal aorta, and are as follows:—

Diaphragmatic.	Hepatic.	Renal.	Inferior mesenteric.
Cœliac.	Splenic.	Spermatic.	Hypogastric.
Gastric.	Supra-renal.	Superior mesenteric.	

The *diaphragmatic plexus* is derived from the upper part of the semilunar ganglion, and is larger on the right than on the left side. It joins with some filaments of the phrenic nerve, and, while in relation with the supra-renal body, it gives off some branches to it. The right plexus has a small ganglion in it—*diaphragmatic ganglion*—and sends off filaments to the vena cava inferior and to the hepatic plexus.

The *cœliac plexus* receives branches from the lesser splanchnic nerve, and on the left side it receives a filament from the right pneumogastric nerve. It divides into the gastric, hepatic and splenic plexuses, which ramify on the corresponding arteries and their branches; the *gastric plexus* receives in addition filaments from the pneumogastric nerves; the *hepatic plexus*, the largest, is joined by branches from the left pneumogastric and right phrenic nerves, and it distributes filaments to the right supra-renal plexus; the *splenic plexus* is formed by branches from the left semilunar ganglion and right pneumogastric nerve.

The *supra-renal plexus* is formed by branches from the so[illegible] plexus, the semilunar ganglion, and the diaphragmatic plexus, a[illegible] is larger on the right than on the left side.

The *renal plexus* consists of numerous filaments from the se[illegible] lunar ganglion, and from the solar and aortic plexuses; it is a[illegible] reinforced by branches from the splanchnic nerves. From t[illegible] plexus is given off—

The *spermatic plexus*, which likewise receives filaments low[illegible] down from the aortic plexus; in the female, it takes the name the *ovarian plexus*.

The *superior mesenteric plexus* receives, in addition to i[illegible] branches from the solar plexus, some filaments from the rig[illegible] pneumogastric nerve and the cœliac plexus. It is the densest [illegible] all the plexuses derived from the solar plexus, and it breaks u[illegible] into secondary plexuses corresponding to the branches of t[illegible] artery of the same name.

The *inferior mesenteric plexus* comes mainly from the left si[illegible] of the aortic plexus.

Hypogastric Plexus. The *hypogastric plexus* is situated between th[illegible] common iliac arteries, on the last lumbar ver[illegible] tebra and the sacrum. It consists of an intricate interlacement o[illegible] sympathetic filaments, which pass down into the pelvis, for the supply of the pelvic viscera. Although this plexus is so intricate it presents no distinct ganglia. As it passes down it receives branches from some of the spinal nerves, but mainly from the third and fourth sacral nerves. From this large plexus are derived secondary plexuses, which ramify around branches of the internal iliac artery: thus there are the inferior hæmorrhoidal plexus, the vesical, the uterine, the ovarian, the prostatic, and the vaginal, all of which send filaments which accompany the smallest branches of the arteries.

Lumbar Plexus of Nerves. The *lumbar plexus* is formed by the union of the anterior branches of the four upper lumbar nerves, and is frequently connected with the last dorsal nerve by a small loop—the dorsi-lumbar nerve. The fifth does not enter into the formation of this plexus, but joins the sacral plexus under the name of the lumbo-sacral cord. The plexus lies over the transverse processes of the corresponding vertebræ, embedded

in the substance of the psoas, so that this muscle must be dissected away before the plexus can be seen. Like the brachial plexus, the nerves composing it successively increase in size from above. Its branches are five in number, and arise in the following order (Fig. 115).

a. The *first lumbar nerve* generally divides into two branches; the upper being the ilio-hypogastric, the lower the ilio-inguinal. They pass downward and outward nearly parallel with each other,

FIG. 115.



a. Ilio-hypogastric n.
b. Ilio-inguinal n.
c. External cutaneous n.
d. Anterior crural n.
e. Crural branch of genito-crural n.
f. Obturator n.
g. Genital branch of genito-crural n.
h. Lumbo-sacral n.

1. First lumbar
2. Second " "
3. Third " "
4. Fourth " "
5. Fifth " "

PLAN OF THE LUMBAR PLEXUS AND BRANCHES.

cross obliquely over the quadratus lumborum to the crest of the ilium, and then separate.

The *ilio-hypogastric nerve* emerges from the outer border of the psoas, and, crossing the quadratus lumborum, passes forward to the crest of the ilium, where it pierces the transversalis, and divides into its two terminal branches—the *iliac* branch, which pierces the internal and external oblique muscles, and supplies the skin over the gluteal region, behind the last dorsal nerve; and the *hypogastric* branch, which runs forward between the

transversalis and internal oblique, and then perforates the aponeurosis of the external oblique to supply the skin of the hypogastric region.

The *ilio-inguinal nerve*—the smaller—comes through the psoas, and perforates the transversalis close to the front of the iliac crest, where it communicates with the preceding nerve. It then pierces the internal oblique, and, lying in front of the spermatic cord, comes out through the external abdominal ring, and supplies the skin of the inner and upper part of the thigh, of the penis and scrotum in the male, and of the labium in the female.

The *genito-crural nerve* is small, and comes from the second lumbar, and by a few filaments from the communicating branch of the first. After perforating the psoas, it lies for a short distance upon its anterior surface, and then runs down along the outer side of the external iliac artery. Near the crural arch it divides into the *genital* branch (*g*), which runs down on the external iliac artery, and, piercing the fascia transversalis, descends through the internal abdominal ring, along the inguinal canal, on the posterior aspect of the spermatic cord; it supplies the cremaster in the male, and the round ligament in the female; and the *crural* branch (*e*), which proceeds under the crural arch, enters the sheath of the femoral vessels, and, piercing the anterior layer of the sheath, just external to the artery, is lost in the skin of the upper part of the front of the thigh; here it communicates with the middle cutaneous nerve and supplies also a few filaments to the femoral artery, where it perforates the sheath of that vessel.

The *external cutaneous nerve of the thigh* (*c*) is generally derived from the loop between the second and third lumbar nerves. It runs through the psoas, then, crossing obliquely over the iliacus toward the anterior superior spine of the ilium, passes beneath the crural arch, and is finally distributed to the skin on the outside of the thigh. If the external cutaneous be not found in its usual situation, look for it as a distinct branch of the anterior crural, nearer the psoas muscle.

The *anterior crural* (*d*), the largest and most important branch, is formed by the union of the third and fourth lumbar nerves, receiving a small branch from the second. It descends in a groove between the psoas and the iliacus, behind the fascia iliaca, supplie

these muscles and a branch to the femoral artery, and then,
ng under the crural arch to the outer side of the femoral
y, is finally distributed to the extensor muscles of the knee,
e sartorius and pectineus, and the skin of the thigh and leg.
he *obturator nerve* (*f*), next in size to the anterior crural,
eeds from the third and fourth lumbar nerves, and some-
s from the second. It descends through the psoas muscle,
then, getting to its inner border, runs along the brim of the
s, above its corresponding vessels, to the obturator foramen,
gh the upper part of which it passes to the adductor muscles
e thigh.

e *accessory obturator nerve*, by no means a constant branch,
rived from the third and fourth lumbar nerves, and sometimes
the obturator nerve. It runs down along the inner border
e psoas, passes in front of the horizontal ramus of the os
, supplies the pectineus, and gives off a small branch to the
oint, and another to communicate with the anterior branch of
bturator nerve.

stponing the minute anatomy of the abdominal viscera, begin
xamination of the contents of the pelvis.

DISSECTION OF THE PELVIC VISCERA.

e functions of the pelvis are to protect its own viscera; to
ort those of the abdomen; to give attachment to the muscles
steady the trunk; to transmit the weight of the trunk to
wer limbs, and to give origin to the muscles which move
. In adaptation to these functions, the form of the pelvis
t of an arch, with broadly expanded wings at the sides,
rojections in appropriate situations to increase the leverage
e muscles. The sacrum, impacted between the ilia, repre-
the keystone of the arch, and is capable of supporting not
the trunk, but great burdens besides. The sides or pillars
epresented by the ilia; these transmit the weight to the
s of the thigh bones, and are thickest and strongest just in
ine, *i. e.*, the brim of the pelvis, along which the weight is
mitted. Moreover, to effect the direct transmission of the
ht, the plane of the arch is oblique. This obliquity of the
s, its hollow expanded sides, its great width, the position and

strength of the tuberosities of the ischia, are so many proofs that man is adapted to the erect posture.

The general conformation of the pelvis in the female is modified, so as to be adapted to utero-gestation and parturition. Its breadth and capacity are greater than in the male. Its depth is less. The alæ of the iliac bones are more expanded. The projection of the sacrum is less perceptible, and consequently the brim is more circular. The depth of the symphysis pubis is less, the span of the pubic arch is wider. The bones, too, are thinner, and the muscular impressions less strongly marked.

The cavity of the pelvis being curved, the axis, or a central line drawn through it, must be curved in proportion. For all practical purposes, it is sufficient to remember that the axis of the pelvis corresponds with a line drawn from the anus to the umbilicus.[1]

Contents of the Male Pelvis. The male pelvis contains the last part of the intestinal canal (named the rectum), the bladder, with the prostate gland at its neck, and the vesiculæ seminales. If the bladder be empty, some of the small intestine will be in the pelvis; not so if the bladder be distended.

The relative positions of these pelvic viscera are as follows: most posteriorly is the rectum, which follows the curve of the sacrum and coccyx, and ends at the anus; immediately in front is the oval hollow viscus, the bladder, which alters in size and position according to the amount of urine it contains; in front of the bladder, surrounding its neck, and behind the os pubis, is the pros-

[1] In a well-formed female the base of the sacrum is 3¾′ inches higher than the upper part of the symphysis pubis, and the point of the coccyx is rather more than half an inch higher than the lower part of the symphysis. The obliquity of the pelvis is greatest in early life. In the fœtus, and in young children, its capacity is small, and the viscera, which subsequently belong to it, are situated in the abdomen.

The relative diameters of the male and the female true pelvis are as follows:—

	Male.			Female.		
	Brim.	Cavity.	Outlet.	Brim.	Cavity.	Outlet.
Transverse . . .	4½	4½	3½	5¼	5	4¾
Oblique . . .	4¼	4½	4	5	5¼	4¾
Antero-posterior . .	4	4½	3¼	4½	5¼	5

ate gland. Placed beneath the bladder are the vesiculæ seminales, nd curving round the sides of the bladder are the vasa deferentia, hich subsequently lie beneath the base of the bladder. Passing lownward and inward over the brim of the pelvis are the ureters, hich likewise get beneath the fundus vesicæ. The bladder and ie rectum are partially invested with peritoneum. Besides the lvic viscera just enumerated, there are found, to the outer part the pelvic cavity, the internal iliac artery and its branches, and e sacral plexus of nerves, with the obturator nerve running ward to the obturator foramen. All the pelvic viscera are re or less invested by prolongations from the pelvic fascia, ich constitute some of the true ligaments of the bladder; also apsule for the prostate, and coverings for the pelvic muscles. e superior hæmorrhoidal artery, the continuation of the in- or mesenteric, passes down into the pelvis, and supplies the per half of the rectum. Some of these structures will now be scribed, while others can be better dissected in the side view the pelvis.

Course of the Rectum. The rectum enters the pelvis on the left side of the sacrum, and, after describing a curve rresponding with the concavity of the sacrum, terminates at the us. In the first part of its course it is loosely connected to the ck of the pelvis by a peritoneal fold, called the *meso-rectum*; tween the layers of this fold, the superior hæmorrhoidal ssels, the continuation of the inferior mesenteric, with nerves d lymphatics, runs to the bowel.

The rectum does not take this course in all cases; sometimes makes one, or even two lateral curves. In some rare cases it ers the pelvis on the right side instead of the left. Since se variations from the usual arrangement cannot be ascer- ned during life, they should make us cautious in the intro- ction of bougies.[1]

Recto-Vesical Pouch. While the parts are still undisturbed, intro- duce the finger into the *recto-vesical peritoneal uch* (Fig. 116). This is a cul-de-sac formed by the peritoneum

[1] In old age the rectum has sometimes a zigzag appearance immediately above anus. These lateral inclinations are probably produced by the enormous disten- as to which the bowel has been occasionally subjected.

in passing from the front of the rectum to the lower and back part of the bladder. In the adult male, the bottom of this pouch is about one inch distant from the base of the prostate gland;[1] therefore part of the under surface of the bladder is not covered by peritoneum; and since this part is in immediate contact with the rectum, it is practicable to tap the distended bladder through the front of the bowel, without injuring the peritoneum. The operation has, of late years, been revived, and with great success.[2] It is easily done, and not attended with risk, provided all the parts be in their regular position. But this is not always the

FIG. 116.



DIAGRAM OF THE RELATIVE POSITION OF THE PELVIC VISCERA.

case. It sometimes happens that the peritoneal pouch com nes down nearer to the prostate than usual—we have seen it in actual contact with the gland; so that, in such a case, it wou uld be impossible to tap the bladder from the rectum without goi ing through the peritoneum. In children the peritoneum com nes down lower than it does in the adult, because the bladder in t the child is not a pelvic viscus.

The recto-vesical pouch is permanent. But there is anoth her

[1] The bottom of the pouch is from three to four inches distant from the anus.

[2] See a paper in the *Med.-Chir. Trans.*, vol. xxxv, by Mr. Cock.

ritoneal pouch on the front part of the bladder, which is only oduced when the bladder is distended. To produce it, the adder should be blown up through one of the ureters. The adder soon fills the pelvis, and then, rising into the abdomen, casions the pouch between it and the abdominal wall. At first e pouch is shallow, but it gradually deepens as the bladder rises. f the bladder be distended half way up to the umbilicus, which is ommonly the case when it has to be tapped, we find that the ottom of the pouch would be about two inches from the symphysis ubis (Fig. 116). Within this distance from the symphysis the ladder may be tapped in the linea alba, without risk of wounding e peritoneum. Thus, the surgeon has the choice of two situ-tions in which he may tap the bladder—above the os pubis, or om the rectum. Which of the two is more appropriate, must decided by the circumstances of the case.

alse Ligaments f the Bladder. The reflections of the peritoneum from the pelvic walls to the bladder constitute the *false aments of the bladder*, and they can be best examined before viscera are disturbed, although they will be described when bladder itself is dissected.

he *two posterior* pass forward from the sides of the rectum to back of the bladder, forming the lateral boundaries of the p recto-vesical pouch. Each contains within its duplicature obliterated hypogastric artery, the ureter, together with some sels and nerves.

he *two lateral* pass inward from the sides of the pelvis to the s of the bladder.

he *superior* passes upward from the summit of the bladder to back of the anterior abdominal wall, covering the urachus l the obliterated hypogastric arteries.

ontents of the emale Pelvis. The relative positions of the pelvic viscera in the female should now be examined, leaving the special description till a later stage.

neral Position f the Uterus and its Appendages. The uterus is interposed between the bladder in front, and the rectum behind. From each side of it a broad fold of peritoneum extends transversely to the side of the pelvis, dividing that rity into an anterior and a posterior part. These folds are called

the *broad ligaments* of the uterus (Fig. 135). On the posterior surface of the ligament are the ovaries, one on each side. They are completely covered by peritoneum, and suspended to the ligament by a small peritoneal fold. Each ovary is attached to the uterus by a cord termed the *ligament of the ovary*. Along the upper part of the broad ligament we find between its layers a tube about four inches long, called the *Fallopian tube*, which conveys the ovum from the ovary into the uterus. For this pur-

FIG. 117.



VERTICAL SECTION THROUGH THE FEMALE PELVIC VISCERA.

pose one end of it terminates in the uterus, while that near[er] to the ovary expands into a wide mouth, furnished with prehen[sile] fringes—*fimbriæ*—which, like so many tentacles, grasp the ov[um] as soon as it escapes from the ovary. One of these fimbri[æ is] attached to the ovary. Lastly, there run up to the ovary, betw[een] the layers of the broad ligament, the ovarian vessels, which ar[ise] from the aorta in the lumbar region, like the spermatic arteries in the male, because the ovaries are originally formed in the lo[ins.]

On the anterior surface of the broad ligament, on either s[ide,]

:n its layers, is the *round ligament* of the uterus. This cord
ds from the fundus of the uterus, anterior to the Fallopian
hrough the inguinal canal, like the spermatic cord in the
and terminates in the mons Veneris. Besides one or two
blood vessels, it contains muscular fibres analogous to those
uterus; these increase very much in pregnancy, so that,
the full term, the cord becomes nearly as thick as the end
little finger. In early life the round ligament receives a
ng from the peritoneum, which advances in a tubular form
ıe inguinal canal. It corresponds to the processus vaginalis
peritoneum in the male. It is called the *canal of Nück*, and
erally obliterated in the adult. It is sometimes the seat of
uinal hernia.

ections of eritoneum. From the front of the rectum the peritoneum is reflected on to a small part of the posterior wall vagina, thus forming what is called the *recto-vaginal pouch.* the vagina the peritoneum is continued over the posterior :, but only about half-way down the *front* of the uterus; it is reflected over the posterior surface of the bladder, on wall of the abdomen. Laterally, it is reflected from the to the sides of the pelvis, forming the *broad ligaments* ig. 135).

:ases of ascites the fluid might distend the recto-vaginal , and bulge into the vagina, so that it would be practicable w it off through this channel.

DISSECTION OF THE MALE PERINEUM.

ɔre dissecting the perineum, it is expedient first to examine seous and ligamentous boundaries of the lower aperture of lvis. Looking at the male pelvis (with the ligaments pre-), we observe that this aperture is of a lozenge shape; that ɔunded *in front* by the pubic arch and the sub-pubic liga- *laterally*, anteriorly, by the rami of the os pubis and ischium, ıe tuberosity of the ischium, posteriorly, by the great sciatic :nt; and *behind*, by the tip of the coccyx.

s space, for convenience of description, is subdivided into y a line drawn from one tuber ischii to the other. The ɔr forms a nearly equilateral triangle, of which the sides

are from three to three and a half inches long; and, since it transmits the urethra, it is called the *urethral region* of the perineum. The posterior, containing the anus, is called the *ischio-rectal* or *anal region* (Fig. 118).[1]

FIG. 118.

DIAGRAM OF THE FRAMEWORK OF THE PERINEUM.

The subject should be placed in the usual position for lithotomy, with a block placed beneath the pelvis. A full-sized staff should now be passed into the bladder, the rectum moderately distended with tow, and the scrotum raised by means of hooks. A central ridge, named the *raphé*, extends from the anus, along the perineum, scrotum, and under surface of the penis. Between the tuberosities of the ischia and the anus are two depressions, one on each side, marking the *ischio-rectal fossæ*, which are found immediately beneath the skin, filled with more or less fat. In the lateral operation of lithotomy, the incision should commence at a point midway between the anus and the posterior fold of the scrotum, close to the left side of the raphé; it should be carried downward and outward to a point midway between the tuber ischii and the anus. In the bilateral operation, the incision is semilunar, the horns being made on either side, between the tuber ischii and the anus, equidistant from these points respectively; while the centre of the incision runs about three-quarters of an inch above the anus.

Anal Glands. At the anus the skin becomes finer and more delicate, forming a gradual transition toward the mucous membrane; during life it is drawn into wrinkles by the permanent contraction of the cutaneous sphincter. Moreover, the skin at the margin of the anus is provided with numerous minute glands,[2] which secrete an unctuous substance to facilitate

[1] The dimensions of the lower outlet of the pelvis are apt to vary in different subjects, and the lithotomist must modify his incision accordingly.

[2] These glands are the analogues of the anal glands in some animals, *e.g.* the dog and the beaver. They are found not only about the anus, but also in the subcutaneous tissue of the perineum, a fact for the demonstration of which we are indebted to the late Professor Quekett. They are large enough to be seen with the naked eye.

e of the fæces. When this secretion becomes defective or
l, the anal cutaneous folds are apt to become excoriated,
d, or fissured; and then defecation becomes very painful.
margin of the anus a thin white line can be distinguished,
ing, not only the junction of the skin with the mucous mem-
but also the linear interval between the external and
l sphincters.[1]

section. The skin should be reflected, by making an incision along the raphé, round the margin of the
o the coccyx. Two others must be made on each side, at
ngles to the first, the one at the upper, and the other at
wer end of it. The skin of the perineum must then be
utaneous issue. reflected outward, with much care, otherwise the superficial sphincter ani may be reflected with
in. In reflecting the skin, notice the characters of the sub-
ous structure.[2] Its characters alter in adaptation to the
cies of each part. On the scrotum the fat constituent of
sue is entirely absent; while the connective tissue element
st abundant, and during life elastic and contractile. But,
n Ischio-l Fossæ. toward the deeper part of the anus, the fat accumulates more and more, and on either side of the
it is found in the shape of large masses, filling up what
otherwise be two deep hollows in this situation— namely,
chio-rectal fossæ. These fossæ are pyramidal, with their
toward the skin, and their apices at the divergence of the
tor internus and levator ani. They are about two inches
pth, and much deeper posteriorly than in front. This
ulation of fat on each side of the anus permits the
listention and contraction of the lower end of the bowel
and after the passage of the fæces. Over the tuberosities
ischia are large masses of fat, separated by tough,
s septa, passing from the skin to the bone, so as to make
astic padding to sit upon. Occasionally, too, there are

[1] ton, *Lectures on Rest and Pain*, p. 280.

[2] probable thickness of this subcutaneous tissue is a point which ought to be
ned by the lithotomist in making his first incision. Its great thickness in
ses explains the depth to which the surgeon has to cut in letting out pus from
io-rectal fossa.

one or more large *bursæ* interposed between this padding and the bone.

So much respecting the general characters of the subcutaneous tissue of the perineum. Some anatomists describe it as consisting of three, four, or even more layers, but in nature we do not find it so. It may, indeed, be divided into as many layers as we please, according to our skill in dissection; but this only complicates what is in itself simple.

Dissection. The external sphincter ani must now be cleaned, care being taken not to remove any of its fibres, which are intimately connected with the skin. Posteriorly, the lower border of the gluteus maximus must be displayed, and the vessels and nerves crossing the perineum, toward the anus, carefully dissected.[1]

External Sphincter Ani. The *external sphincter* of the anus is elliptical, and is composed of a thin layer of striped muscular tissue about an inch in breadth. It *arises* from the tip of the coccyx and the ano-coccygeal ligament. The muscular fibres surround the anus, and are *inserted* in a pointed manner in the tendinous centre of the perineum, in conjunction with the transversus perinei, the accelerator urinæ, and the levator ani (p. 530). It is called the external sphincter, to distinguish it from a deeper and more powerful band of muscular fibres which surrounds the last inch or more of the rectum, and is situated next to the mucous membrane.

Cutaneous Vessels and Nerves. The cutaneous vessels and nerves of the perineum come from the internal pudic artery and nerve, and chiefly from that branch of it called the *superficial perinei*. This will be traced presently.

The *external* or *inferior hæmorrhoidal arteries* cross transversely through the ischio-rectal fossa, from the ramus of the ischium toward the anus. They come from the pudic (which can be felt on the inner side of the ischium), and, running inward, divide into numerous branches, which supply the rectum, levator ani and sphincter ani. The *nerves* which accompany the arteries com-

[1] Radiating outward from the margin of the anus is a thin stratum of involuntary muscular fibres, called the *corrugator cutis ani*, which by its action produces the radiating ridges of skin from the anus.

m the pudic nerve, and supply the sphincter ani and the skin the perineum.

The *fourth sacral nerve* emerges through the coccygeus close to tip of the coccyx, and, through its *hæmorrhoidal* or *perineal nch*, supplies the external sphincter and the skin of the peeum between the coccyx and the anus.

The *inferior pudendal nerve* comes through the muscular fascia the thigh, a little above the tuber ischii, and ascends, dividing o filaments, which supply the front and outer part of the scrom and perineum. It is a branch of the lesser sciatic nerve, and mmunicates in front with the posterior branch of the superficial ineal nerve.

erficial Fascia the Perineum. The subcutaneous fascia of the perineum is composed of a *superficial* and a *deep* layer. The *erficial* layer contains more or less fat, and is continuous with t of the scrotum, the thighs, and the posterior part of the ineum. The *deeper* layer is a stratum of considerable strength, d is best demonstrated by blowing air beneath it with a blowe; its connections are as follows: It is attached on each side the anterior lip of the ramus of the os pubis and ischium, erficial to the crus penis; traced forward, it is directly conuous with the *tunica dartos* of the scrotum; traced backward, at base of the urethral triangle it is reflected beneath the transse perinei muscle, and joins the *deep perineal fascia* or *triangu- ligament.* These connections explain why urine effused into perineum does not make its way into the ischio-rectal fossæ, down the thighs, but passes readily forward into the connective ue of the scrotum, penis and groins.

Dissection. Remove the fascia, to see the muscles which cover the bulb of the urethra and the crura of penis. The bulb of the urethra lies in the middle of the ineum, and is covered by a strong muscle, called accelerator næ. The crura penis are attached, one to each side of the bic arch, and are covered each by a muscle, called erector is. A narrow slip of muscle, called transversus perinei, exds on either side from the tuber ischii to the *tendinous centre* the perineum. This point is about one inch and a quarter in

front of the anus; and serves for the attachment of muscular fibres from all quarters of the perineum.

Thus the muscles of the perineum describe on each side a triangle, of which the sides are formed by the accelerator urinæ and the crus penis respectively, and the base by the transversus perinei. Across this triangle run up, from base to apex, the super-

FIG. 119.



MUSCLES, WITH SUPERFICIAL VESSELS AND NERVES, OF THE PERINEUM.

ficial perineal vessels and nerves. External to the ramus of ischium is seen the *inferior pudendal* nerve, a branch of the sciatic.

Superficial Perineal Vessels and Nerves.

The *superficial perineal* artery lies beneath deep layer of the superficial perineal fascia, comes from the internal pudic as it runs up

r side of the tuber ischii. Though the main trunk cannot be
, it can be easily felt by pressing the finger against the bone.
artery comes into view a little above the level of the anus,
es up usually in front of the transversus perinei muscle, and
to the perineal triangle lying to the inner side of the erector
s. It distributes branches to all the muscles, and is finally
on the scrotum. The only named branch is called *transver-
perinei* (Fig. 119). This is given off near the base of the
gle, and runs transversely inward with the transversus perinei
:le toward the central tendon of the perineum, where it anas-
ses with its fellow. It is necessarily divided in the first
ion in lithotomy, and deserves attention, because it is some-
s of considerable size.

ıe artery is accompanied by two *veins*, which are frequently
ed and tortuous, especially in diseased conditions of the
um.

ıe *nerves*, two in number, are derived from the internal pudic,
w the course of their corresponding arteries, and give off
ar branches. They not only supply the skin of the perineum
scrotum, but each of the perineal muscles.

ccelerator Urinæ. This muscle embraces the bulb of the urethra, and is composed of two lateral symmetrical halves. *ises* from a fibrous median raphé beneath the bulb, and from endinous centre of the perineum. Starting from this origin, ibres diverge, and are *inserted* as follows:—The *upper* ones ed on either side round the corpus cavernosum penis, like the :hes of the letter V, and are fixed on its dorsal surface, in of the erector penis, and expanding also into a broad apo-osis, which covers the dorsal vessels of the penis; the *middle* letely embraces the bulb and adjacent part of the corpus giosum like a ring, and meet in an aponeurosis on the upper ce of the urethra; the *lower* are fixed to the anterior surface ıe deep perineal fascia, often called the triangular ligament . 120).[1]

ıus, the entire muscle acts as a powerful compressor of the , and expels the last drops of urine from this part of the

[1] This muscle is called also the *ejaculator urinæ* or the *bulbo-cavernosus*.

urethra.[1] By dividing the muscle along the middle line and turning back each half, its insertion, as above described, can be clearly made out.

Erector Penis. This muscle is moulded upon the crus of the penis. It *arises* by musculo-tendinous fibres from the inner surface of the tuber ischii, from the crus itself, and from the ramus of the os pubis; the fibres ascend, completely covering the crus, and terminate on a strong aponeurosis, which is *inserted* into the external and inferior aspect of the crus penis. The *action* of this muscle is to compress the root of the penis, and so, by pre-

FIG. 120.



DIAGRAM TO SHOW THE ACCELERATOR URINÆ IN PROFILE.

venting the return of the venous blood, contributes to the erection of the organ.[2]

Transversus Perinei. This muscle is of insignificant size, and sometimes absent. It *arises* from the inner aspect of the tuber ischii, and proceeds forward and inward toward the central point of the perineum, where it is blended with the muscle of the opposite side, with the fibres of the accelerator urinæ in front, and with the external sphincter behind. This muscle with its artery is divided in lithotomy.

[1] The middle fibres assist in the erection of the corpus spongiosum, and the upper fibres in that of the penis; the former by compressing the bulb, the latter by compressing the dorsal vein.

[2] This muscle is sometimes called the *ischio-cavernosus*.

The *deep transversus perinei* is a small muscle occasionally present; it arises more deeply from the pubic arch than the superficial muscle, and passes inward behind the bulb to the central tendon.

The next stage of the dissection consists in reflecting and

FIG. 121.



DIAGRAM TO SHOW THE TRIANGULAR LIGAMENT OF THE URETHRA OR DEEP PERINEAL FASCIA.

removing the accelerator urinæ from the bulb of the urethra, the erectores penis with the crura penis from the rami of the os pubis and ischium, and the transversi perinei muscles. This done, the *triangular ligament* or *deep perineal fascia* is fairly exposed.

Triangular Ligament of the Urethra.

Understand that the triangular ligament of the urethra and the deep perineal fascia are synonymous terms.

The *triangular ligament*, shown in Fig. 122, is a strong, fibrous

FIG. 122.



DIAGRAM OF THE PARTS BEHIND THE ANTERIOR LAYER OF THE TRIANGULAR LIGAMENT OF THE URETHRA.

(The anterior fibres of the levator ani are hooked down, to show part of the prostate; the rest tracked by a dotted line.)

membrane stretched across the pubic arch. It is about an inch and a half in depth, with the base directed backward. It consists of two layers—an anterior and a posterior. The *anterior layer* is firmly attached on each side to the *posterior* lip of the rami of the os pubis and ischium, beneath the crus penis; superiorly—

e., toward the symphysis of the os pubis—it is connected with ıe sub-pubic ligament; inferiorly, it does not present a free ɔrder, but is connected to the tendinous centre of the perineum, ıd is continuous with the deep layer of the superficial perineal scia which curves backward under the transversus perinei muscle, d with the ischio-rectal or anal fascia (p. 529).

The anterior layer of the triangular ligament is perforated ɔut one inch below the symphysis pubis, for the membranous rt of the urethra. The aperture through which the urethra sses does not present a distinct edge, because the ligament is ɔlonged forward over the bulb, and serves to keep it in position. also presents apertures for the transmission of the dorsal vein, d outside this for the pudic arteries and nerves.

The *posterior layer* cannot at present be seen. It belongs, ictly speaking, to the obturator prolongation of the pelvic fascia, d slopes somewhat backward from the anterior layer so as to ıve an interval between them, in which are found structures ıich will be presently described.

Points of ırgical Interest. The triangular ligament is very important surgically, for these reasons:—

1. Here we meet with difficulty in introducing a catheter, unıs we can hit off the right track through the ligament. The ft and spongy tissue of the bulbous part of the urethra in front the ligament readily gives way if force be used, and a false ssage results.

2. By elongating the penis we are much more likely to hit off e proper opening through the ligament.

3. When, in retention of urine, the urethra gives way *anterior* this ligament, it is this which prevents the urine from traveling to the pelvis. Its connection with the superficial perineal fascia events the urine from getting into the ischio-rectal fossæ; nor n the urine make its way into the thighs. The only outlet for is into the connective tissue of the scrotum and penis.

4. When suppuration or extravasation of urine takes place *hind* the ligament, the pus is pent up and should be speedily let ıt; if not, it may find its way into the connective tissue of the lvis, and may burst into the urethra or the rectum.

5. The ligament is partially cut through in lithotomy.

Parts Divided in Lateral Lithotomy. The parts divided in the lateral operation of lithotomy are: the skin, the superficial fascia, the transverse perineal muscle, vessels and nerve, the inferior hæmorrhoidal vessels and nerves, the inferior fibres of the accelerator urinæ, the anterior fibres of the levator ani, the triangular ligament (anterior layer), the compressor urethræ, the membranous and prostatic parts of the urethra, and a small portion of the prostate.

Parts to be Avoided. The incision in lateral lithotomy should not be made too far forward, for fear of wounding the artery of the bulb; nor too far inward, for fear of injuring the rectum; nor too far outward, for fear of cutting the pudic artery.

Structures Between the Layers of the Triangular Ligament. The anterior layer of the triangular ligament must now be cut away, to see what lies between its two layers. These parts are shown in Fig. 122, namely: (1) the membranous part of the urethra, surrounded by (2) the compressor urethræ muscle; (3) Cowper's glands and their ducts; (4) the pudic artery and its branch, the artery of the bulb; the artery of the crus and the dorsal artery of the penis being given off in front of the anterior layer; (5) the pudic nerve and its branches; (6) the dorsal vein of the penis; (7) the sub-pubic ligament.

Dissection. To obtain the best perineal view of the compressor urethræ muscle, cut through the spo[illegible] part of the urethra about three inches above the end of the b[illegible] and dissect it from the corpus cavernosum. Thus the upper fib[illegible] of the constrictor will be exposed; to see the lower, it is o[illegible] necessary to raise the bulb. The most perfect view, however, the muscle is obtained by making a transverse section through t[illegible] rami of the ossa pubis, so as to get at the muscle from above, [illegible] shown in Fig. 123.

Compressor or Constrictor Urethræ. This muscle consists of transverse fibres whic[illegible] surround and support the whole length of th[illegible] membranous portion of the urethra in its passag[illegible] between the two layers of the triangular ligament. It *arises* fro[illegible] the ramus of the os pubis on either side for about half an inch [illegible] from thence its fibres pass, some above, some below the urethra[illegible]

ng the whole length of its membranous part. It forms a com-
te muscular covering for the urethra between the prostate and
bulb. It is chiefly through its agency that we retain the
ne. This muscle is the chief cause of spasmodic stricture of the
thra.[1] Besides this muscle, the membranous portion of the
thra is surrounded by involuntary circular muscular fibres,
ced beneath the compressor urethræ, and continuous with the
scular fibres of the bladder.

wper's Glands. These small glands are situated, one on either side, immediately behind the bulb between the layers of the triangular ligament, in the substance of the pressor urethræ. Their size is about that of a pea, but it es in different individuals. They are compound racemose nds, consisting of several lobules firmly connected together by ular and some muscular tissue. From each a slender duct s forward, and, after a course of about one inch, opens obliquely the floor of the bulbous part of the urethra (Fig. 120). They ish a secretion accessory to generation.

ic Artery and s Branches. The *internal pudic artery* is a branch of the anterior division of the internal iliac. It leaves pelvis through the great sciatic foramen between the pyri-is and coccygeus muscles, above the sciatica artery, winds nd the spine of the ischium, re-enters the pelvis through the er sciatic foramen, and then runs along the inner side of the r ischii, between the layers of the obturator fascia, up toward pubic arch. About an inch and a half above the tuber ii, the trunk of the pudic artery can be felt; but we cannot it, nor draw it out, for it is securely lodged in a fibrous canal ned by the obturator fascia. It subsequently pierces the terior layer of the triangular ligament, runs along the inner gin of the ramus of the os pubis, and lastly, piercing the erior layer of the triangular ligament, it divides into the artery he corpus cavernosum and the dorsal artery of the penis. In present dissection we find the artery between the two layers he triangular ligament, where it gives off the artery of the

[1] The compressor urethræ was first accurately described and delineated by orini (septemdec. tabulæ), and afterwards by Müller, in his monograph (Ueber rgan. Nerv. der männlich. Geschlechtsorgane).

bulb of the urethra, and then pierces the triangular ligament (Fig. 121).

Taken in order, the branches of the pudic artery as seen in this dissection are :—

a. The *external hæmorrhoidal*, the *superficial perineal*, and the

FIG. 123.



DIAGRAM OF THE RELATIONS OF THE COMPRESSOR URETHRÆ, SEEN FROM ABOVE.

transverse perineal branches have already been described (pp. 528–531).

b. The *artery of the bulb* is of considerable size, and passes transversely inward between the two layers of the triangular ligament; it runs inward through the substance of the compressor

ethræ, and before it enters the bulb divides into two or three anches. It also sends downward a small branch to Cowper's and. From the direction of this artery it will at once strike e attention that there is great risk of dividing it in lithotomy. the artery runs along its usual level, and the incision be not ade too high in the perineum, then indeed it is out of the way harm. But, supposing the reverse, the vessel must be divided. is deviation from the normal distribution is met with about ce in twenty subjects, and there is no possibility of ascertaining s anomaly beforehand.

. The *artery of the corpus cavernosum*, one of the terminal nches, ascends for a short distance near the pubic arch, and n enters the crus, running forward in its cavernous structure the side of the septum pectiniforme.

. To see the *dorsal artery of the penis*, the crus should be dis-ted from its attachment to the symphysis pubis. The artery rces the suspensory ligament, and can be traced upon the dor-n of the penis down to the glans. It forms a complete arterial le, with its fellow, round the corona glandis, and gives numer-ramifications to the papillæ on the surface.

The *veins* corresponding with the branches of the pudic artery minate in the pudic vein, with the exception of the *dorsal vein the penis*. This vein is of large size and results from the union two small veins in front of the dorsum of the penis which re-ve the blood from the glans, the corpus spongiosum, and the puce. The vein runs along the middle of the dorsum, pierces, t, the suspensory ligament, and then the triangular ligament der the symphysis, and divides into two branches which open o the prostatic plexus.

Pudic Nerve. The *pudic nerve* comes from the lower part of the sacral plexus, and corresponds, both in its rse and branches, with the artery. It gives off, close to its gin, (*a*) the *external* or *inferior hæmorrhoidal*, which communi-e in front with the superficial perineal and inferior pudendal ves; (*b*) the *perineal*, which accompanies the superficial perineal ery, and divides into a posterior and an anterior branch; the mer runs to the front of the ischio-rectal fossa, distributing nches to the sphincter and the skin in front of the anus; the

latter lies in front of the preceding, and supplies the scrotum and under aspect of the penis; both communicate with each other and with the inferior pudendal nerve; (*c*) *muscular branches* to the transversus perinei, the accelerator urinæ, the erector penis and the compressor urethræ; (*d*) the *dorsal nerve*, which is the main trunk of the nerve, runs with the pudic artery, and with it pierces the posterior, and then the anterior layer of the triangular ligament; then perforating the suspensory ligament of the penis, it accompanies the dorsal artery on its outer side, along the dorsum of the penis to the glans. In its passage it supplies the integuments of the penis, and sends off one or two branches into the corpus cavernosum. This part of the penis also receives filaments from the sympathetic system.

Ischio-rectal Fossa. This is the deep hollow, on each side, between the anus and the tuber ischii. When all the fat is removed from it, observe that it is lined on all sides by fascia. Introduce the finger into it to form a correct idea of its extent and boundaries. *Externally*, it is bounded by the tuber ischii and the fascia covering the obturator internus muscle; *internally*, by the rectum, levator ani and coccygeus; *posteriorly*, by the gluteus maximus; *anteriorly*, by the transversus perinei. The fossa is crossed by the external hæmorrhoidal vessels and nerves.

These deep spaces on each side of the rectum explain the great size which abscesses in this situation may attain. The matter can be felt only through the rectum. Nothing can be seen outside. Perhaps nothing more than a little hardness can be felt by the side of the anus. These abscesses should be opened early; else they form a large cavity, and may burst into the rectum, and result in a fistula.

DISSECTION OF THE FEMALE PERINEUM.

The *pudenda* in the female consist of folds of the integument, called the labia. Between these is a longitudinal fissure which leads to the orifices of the urinary and genital canals.

Labia Majora. The pubic region is generally surmounted by an accumulation of fat, called *mons Veneris*, which is covered with hair. From this, two thick folds of skin descend, one on either side, constituting the *labia majora*, and gradually

diminish in thickness toward the perineum. Their junction, about an inch above the anus, is called the *posterior commissure*, or *frænulum labiorum*, within which is a transverse crescentic fold, the *fourchette*; it is generally torn in the first labor. Between the fourchette and the posterior commissure is an oval depression, called the *fossa navicularis*. The inner layer of the skin of the labium is thinner, softer, and more like mucous membrane than the outer; for this reason, whenever pus forms in the labium, the abscess bursts on the inner side. Where the labia are in contact, they are provided with small sebaceous glands, of which the minute ducts are observable on the surface. They are the analogues of the scrotum in the male, and occasionally contain extruded ovaries, forming a hernia of the ovary.[1]

Clitoris. In form and structure the clitoris resembles the penis on a diminutive scale, being about an inch and a half long. It has, however, no corpus spongiosum, or urethra. Like the penis, it is attached to the sides of the pubic arch by two crura (Fig. 124, p. 544), each of which is grasped by its special *erector clitoridis*. The crura are continued forward, like the corpora cavernosa of the male, and unite to form the body of the organ, which is surmounted by a small *glans*. It has also, like the penis, a suspensory ligament. The glans is provided with extremely sensitive papillæ, and covered by a little prepuce. Its dorsal arteries and nerves are large in proportion to its size, and have precisely the same course and distribution as in the penis. Its internal structure consists of a plexus of blood vessels, which freely communicate with those of the labia minora; for one cannot be injected without the other.

Labia Minora or Nymphæ. By separating the external labia, two small and thin folds of mucous membrane, about an inch and a half in length, are exposed, one on either side, termed *labia minora*. These folds converge anteriorly, and form a covering for the clitoris, called *preputium clitoridis;* posteriorly, they are gradually lost on the inside of the labia majora. They, unlike the labia majora, do not contain fat, but are composed of

[1] See paper on "Hernia of the Ovary," *St. Bartholomew's Hospital Reports*, vol. xviii.

minute veins. Between the nymphæ and about the clitoris are a number of sebaceous glands.

Between the labia minora, and below the clitoris, is an angular depression called the *vestibule*, at the back of which is the *meatus urinarius*. Immediately below this is the vagina, of which the orifice is partially closed in the virgin by a thin fold of mucous membrane called the *hymen*.

Hymen. The *hymen* is a thin fold of mucous membrane which, in the virgin, extends across the lower part of the entrance of the vagina, about half an inch behind the fourchette. In most instances its form is crescent-shaped, with the concavity upward. There are several varieties of hymen; sometimes there are two folds, one on either side, so as to make the entrance of the vagina a mere vertical fissure;[1] or there may be a septum perforated by several openings, *hymen cribriformis*, or by one only, *hymen circularis*. Again, there may be no opening at all in it, and then it is called *hymen imperforatus*. Under this last condition no inconvenience arises till puberty. The menstrual discharge must then necessarily accumulate in the vagina; indeed the uterus itself may become distended by it to such an extent as even to simulate pregnancy.[2]

When the hymen is ruptured, it shrivels into a few irregular eminences, called *carunculæ myrtiformes*.

The presence of the hymen is not necessarily a proof of virginity, nor does its absence imply the loss of it. Cases are related by writers on midwifery in which a division of the hymen was requisite to facilitate parturition. In Meckel's Museum, at Halle, are preserved the external organs of a female in whom the hymen is perfect even after the birth of a seven months' child.

Bartholin's or Duverney's Glands. Between the orifice of the vagina and the erector clitoridis is imbedded in the loose tissue on either side a small gland,[3] which corresponds to Cowper's gland in the male. Each is about half an inch in length. Its long slender duct runs forward and opens on the inner side of the nympha external to the hymen. In cases of virulent gonorrhœa

[1] Such a one may be seen in the Museum of the College of Phys. Series No. 2843.

[2] See Burn's *Midwifery*.

[3] See Tiedemann, *Von der Duverneyschen Drüsen des Weibs*. Heidelberg, 1840.

these glands are apt to become diseased, and give rise to the formation of an abscess in the labium, very difficult to heal.

Urethra.

A smooth channel, called the *vestibule*, three quarters of an inch in length, leads from the clitoris down to the orifice of the urethra. This orifice, *meatus urinarius*, is not a perpendicular fissure like that of the penis, but rounded and puckered, and during life has a peculiar dimple-like feel, which assists us in finding it when we pass a catheter. You should practice the introduction of the catheter in the dead subject, for the operation is not so easy as might at first be imagined, provided the parts are not exposed. The point of the forefinger of the left hand should be placed at the entrance of the vagina and the meatus felt for, when the catheter, guided by the finger, slips, after a little manœuvring, into the urethra. The canal is about one inch and a half in length, and runs along the upper wall of the vagina. The two canals are in such close apposition that you can feel the urethra imbedded in the vagina like a thick cord. The urethra is slightly curved, with the concavity upward; but for all practical purposes it may be considered straight. Its direction, however, is not horizontal. In the unimpregnated state it runs nearly in the direction of the axis of the outlet of the pelvis; so that a probe pushed on in the course of the urethra would strike against the promontory of the sacrum. But after impregnation, when the uterus begins to rise out of the pelvis, the bladder is more or less raised also, in consequence of their mutual connection; therefore the urethra, in the latter months of utero-gestation, acquires a much more perpendicular course.

The female urethra is provided with a *compressor* muscle, similar in origin and arrangement to that which surrounds the membranous part of the urethra in the male. It also passes through the triangular ligament. The prostate gland is wanting, but there are minute glands scattered around the neck of the bladder. In consequence of the wider span of the pubic arch and the more yielding nature of the surrounding structures, the female urethra is much more dilatable than the male. By means of a sponge-tent, it may be safely dilated to admit the easy passage of the forefinger into the bladder. Advantage is taken of this great dilatability in the extraction of calculi from the bladder.

The mucous coat of the urethra is pale and arranged in longitudinal folds, and is lined by squamous epithelium, which changes to the spheroidal variety near the bladder. Next to the mucous coat is a layer of elastic and non-striped muscular fibres intermixed. The muscular tissue is arranged in two layers—an outer, consisting of circular fibres, and an inner, of longitudinal fibres. Externally there is a plexus of veins bearing a strong resemblance to erectile tissue.

Vagina. The vagina is the canal which leads to the uterus; at present, only the orifice of it can be seen. It is surrounded by a sphincter muscle, easily displayed by removing the integument. The muscle is about three-fourths of an inch broad, and connected with the cutaneous sphincter of

FIG. 124.

1. Meatus urinarius
2. Vagina.

3. Bulb of vagina.
4. Clitoris with its two crura.

BULB OF THE VAGINA.[1]

the anus in such a manner that they together form something like the figure 8.

On each side of the orifice of the vagina, between the mucous membrane and the sphincter, is a plexus of tortuous veins, termed the *bulb of the vagina*, from its analogy to the bulb of the urethra in the male. This vaginal bulb is about an inch long and extends across the middle line between the meatus urinarius and the clitoris, as shown in Fig. 124.

The description of the perineal branches of the pudic vessels and nerves, given in the dissection of the male perineum, applies, *mutatis mutandis*, to the female, excepting that they are proportionably small, and that the artery which supplies the bulb of the urethra in the male is distributed to the bulb of the vagina in the female.

[1] Taken from an injected preparation in the Musée Orfila, at Paris.

ANATOMY OF THE SIDE VIEW OF THE PELVIC VISCERA.

Dissection. To obtain a side view of the pelvic viscera, the left innominate bone should be removed, thus: Detach the peritoneum and the levator ani from the left side of the pelvis, cut through the external iliac vessels, the obturator vessels and nerve, and the nerves of the lumbar plexus; then saw through

FIG. 125.



VERTICAL SECTION THROUGH THE PERINEUM AND PELVIC VISCERA.

(The arrows point out where the bladder can be tapped.)

the os pubis about two inches external to the symphysis, and cut through the sacro-iliac symphysis; now draw the legs apart, and saw through the base of the spine of the ischium; after cutting through the pyriformis, the great sacro-sciatic ligament, the great and small sciatic nerves and the gluteus maximus muscle, the innominate bone can be easily detached. This done the rectum should be distended with tow, and the bladder blown up through the ureter. A staff should be passed through the urethra into the bladder, and a block placed under the sacrum.

The reflection of the peritoneum as it passes from the front of the rectum to the lower part of the bladder (forming the recto-vesical pouch), and thence over the back of the bladder to the wall of the abdomen, has been already described. You see where the distended bladder is bare of peritoneum, and that it can be tapped either through the rectum or above the pubes without injury to the serous membrane, as shown by the arrows in Fig. 125.

FIG. 126.



TRANSVERSE SECTION OF THE PELVIS, TO SHOW THE REFLECTIONS OF THE PELVIC FASCIA. (AFTER GRAY.)

False Ligaments of the Bladder.

The peritoneal connections of the bladder are called its false ligaments; *false* in contradistinction to the *true*, which are formed by the fascia of the pelvis, and really *do* sustain the neck of the bladder in its proper position. The *false ligaments* are five in number, two posterior, two lateral, and one superior. The *posterior* are produced by two peritoneal folds, one on either side the recto-vesical pouch; the *two lateral*, by reflections of the peritoneum from the sides of the

pelvis to the sides of the bladder; the *superior* is produced by the passage of the peritoneum from the front of the bladder to the abdominal wall. These have been already described (p. 523).

Pelvic Fascia. To expose the *pelvic fascia*, the peritoneum must be removed from that side of the pelvis which has not been disturbed; in doing so, notice the abundance of loose connective tissue interposed between the peritoneum and the fascia, to allow the bladder to distend with facility. Whenever urine is extravasated into this loose tissue, it is sure to produce the most serious consequences; therefore, in all operations on the perineum, it is of the utmost importance not to injure this fascia.

The pelvic fascia is a thin but strong membrane, and constitutes the true ligaments of the bladder and the other pelvic viscera, supporting and maintaining them in their proper position.

Examine, first, to what parts of the pelvis the fascia is attached; secondly, the manner in which it is reflected on the viscera.

Beginning, then (Fig. 126), we see that, in front, the fascia is continuous with the transversalis fascia, and laterally with the iliac fascia, and that superiorly it is attached to the body of the os pubis, to the brim of the pelvis, and to the side of the bone just above the attachment of the obturator internus, close to the obturator foramen and the great sciatic notch. Here it becomes gradually thinner, covers the pyriformis and the sacral plexus, and is gradully lost on the front of the sacrum.

Traced forward, we find that it is attached to the bone along the upper border of the obturator internus, and, as it passes forward over the obturator foramen, completes the canal through which the obturator vessels pass to the foramen; anteriorly, it is attached to the posterior surface of the lower part of the symphysis pubis. From this attachment the fascia descends as far as a line drawn from the spine of the ischium to the pubic symphysis, where it forms a dense white line which marks the division of the fascia into two layers, an outer, the obturator, and an inner, the recto-vesical fascia. It also serves for the attachment of a considerable part of the middle portion of the levator ani.

The *obturator fascia*, the outer layer, is the continuation of the pelvic fascia, and descends on the inner surface of the obturator

internus, forming at the same time a sheath for the pudic vessels and nerve, the nerve being the lowest. It is attached to the pubic arch, to the tuberosity of the ischium, and to the margin of the great sacro-sciatic ligament. It is continuous in front, below the symphysis pubis, with the corresponding layer of the opposite side, and here forms the posterior layer of the triangular ligament. From this fascia is derived the *ischio-rectal* or *anal fascia*, which lines the under or perineal surface of the levator ani, and is subsequently lost upon the side of the rectum.

The *recto-vesical fascia* descends on the upper or internal surface of the levator ani, and invests the bladder, prostate and rectum. From the symphysis pubis it is reflected over the prostate and the neck of the bladder to form, on either side of the symphysis, two well-marked bands—the *anterior true ligaments* of the bladder. From the side of the pelvis it is reflected on to the side of the bladder, constituting the *lateral true ligaments* of that viscus, and encloses the prostate and the vesical plexus of veins. A prolongation from this ligament encloses the vesicula seminalis, the lower layer of which passes between the bladder and the rectum, to join its fellow from the opposite side. The continuation of the recto-vesical fascia covers the remainder of the upper surface of the levator ani as far as its attachment to the rectum, where it is reflected round this tube.

General Position of the Pelvic Viscera in the Male.

The pelvic viscera are so surrounded by veins and loose areolar tissue, that he who dissects them for the first time will have a difficulty in discovering their definite boundaries. The rectum runs at the back of the pelvis, and follows the anterior curve of the sacrum and coccyx. The bladder lies in front of the rectum, immediately behind the symphysis pubis. At the neck of the bladder is the prostate gland through which the urethra passes. In the cellular tissue, between the bladder and the rectum, there is, on each side, a convoluted tube, called the vesicula seminalis, and on the inner side of each vesicula is the seminal duct or vas deferens. Before describing these parts in detail, it is necessary to say a few words about the large, tortuous veins which surround them.

Vesico-prostatic Plexus of Veins.

Beneath the pelvic fascia surrounding the prostate and the neck of the bladder are large and

tuous veins, which form the prostatic and the vesical plexuses. ey empty themselves into the internal iliac. In early life they not much developed, but as puberty approaches they gradually rease in size, and one not familiar with the anatomy of these ts would hardly credit the size which they sometimes attain in persons. They communicate freely behind with the inferior morrhoidal plexus, or veins about the anus, and they receive the

FIG. 127.



SIDE VIEW OF THE PELVIC VISCERA.

(Taken from a Photograph.)

1. External sphincter.
2. Internal sphincter.
3. Levator ani cut through.
4. Accelerator urinæ.
5. Membranous part of the urethra, surrounded by compressor muscle.
6. Prostate gland.
7. Vesicula seminalis.
8. Ureter.
9. Vas deferens.
10. Crus penis divided.
11. Triangular ligament.
12. Superficial perineal fascia.
13. Rectum.

od returning from the penis through the large veins which pass er the pubic arch.

, in lithotomy, the incision be carried beyond the limits of

the prostate, the great veins around it must necessarily be divided; these, independently of any artery, are quite sufficient to occasion serious hemorrhage.

Rectum and its Relations. The rectum is about eight inches long. It is a continuation of the sigmoid flexure of the colon, enters the pelvis at the left sacro-iliac articulation, describes a curve corresponding to the sacrum and coccyx, and terminates at the anus. The rectum also inclines from the left side to the middle line, and before its termination the bowel turns downward so that the anal aperture is dependent. Although it loses the sacculated appearance, it is not throughout of equal calibre, and its capacity becomes greater as it descends into the pelvis; immediately above the sphincter it presents a considerable dilatation, the *ampulla* (Fig. 125). This dilatation is not material in early life, but it increases as age advances. Under such circumstances the rectum loses altogether its cylindrical form, and bulges up on either side of the prostate and the base of the bladder. For this reason the rectum should always be emptied before the operation of lithotomy.

The rectum is conveniently divided into three portions, the upper, the middle, and the lower.

The *upper portion* is about three inches and a half in length, and extends as low as the third bone of the sacrum, to which bone it is connected by a fold of peritoneum, termed the *meso-rectum*. In this fold, the terminal branch of the inferior mesenteric artery with its vein runs down to supply the bowel. This portion of the rectum has behind it the sacral plexus of nerves, the pyriformis, and some branches of the left internal iliac artery; in front, it has the bladder and the recto-vesical pouch.

The *middle portion* comprises three inches in length, and is continuous with the lower portion at the tip of the coccyx. It is connected posteriorly to the sacrum and coccyx by loose connective tissue, and is covered by peritoneum only in front in the upper part, which forms the recto-vesical pouch. It has in front the fundus of the bladder, the vesiculæ seminales, the vasa deferentia and the prostate; while in the female it is closely connected to the posterior wall of the vagina.

The *lower portion* comprises the lowest inch and a half of the

. It is entirely destitute of peritoneum, and is supported levatores ani, the larger portions of which are inserted into e; it has also surrounding it the internal, and lastly the al sphincters. There is a considerable interval between it e membranous portion of the urethra in the male, and the in the female.

al Examination of the ectum.

The relations of the front part of the rectum—that, namely, included between the recto-vesical pouch and the anus—are most important. If the ger be introduced into the anus, and a catheter into the a, the first thing felt through the front wall of the bowel is embranous part of the urethra (Fig. 125). It lies just the sphincter, and is about ten lines in front of the gut. one and a half or two inches from the anus the finger upon the prostate gland; this is in close contact with the d is readily felt on account of its hardness; by moving the from side to side we recognize its lateral lobes. Still up the finger goes beyond the prostate, and reaches the e of the bladder; the facility with which this can be exam- lepends, not only upon the length of the finger and the t of fat in the perineum, but upon the degree of distention bladder; the more distended the bladder, the better can the te be felt. These several relations are practically important. explain why, with the finger in the rectum, we can ascertain er the catheter is taking the right direction—whether the te be enlarged or not. We might even raise a stone from the n of the bladder so as to bring it in contact with the forceps. rectum is supplied with blood by the superior, middle and r hæmorrhoidal arteries. The superior comes from the r mesenteric (p. 494); the middle from the anterior division internal iliac artery, and the inferior from the pudic artery. perior hæmorrhoidal veins join the inferior mesenteric, and quently the portal system; the middle and the inferior rrhoidal veins join the internal pudic, and thence the internal ein. They are very large, and form loop-like plexuses about wer part of the rectum. Having no valves, they are liable come dilated and congested from various internal causes; the frequency of hæmorrhoidal affections.

Bladder.

This viscus, being a receptacle for the urine, must necessarily vary in size, and according to the nature of its connections and coats is such as to permit this variation. When contracted, the bladder sinks into the pelvis behind the pubic arch, and is completely protected from injury. But, as it gradually distends, it rises out of the pelvis into the abdomen, and, in cases of extreme distention, may reach up to the umbilicus.[1] Its outline can then be easily felt through the walls of the abdomen. The form[2] of the distended bladder is oval, and its long axis, if prolonged, would pass superiorly through the umbilicus, and inferiorly through the end of the coccyx. The axis of a child's bladder is more vertical than that of the adult; for in children the bladder is not a pelvic viscus. This makes lithotomy in them so much more difficult.

FIG. 128.

POSTERIOR VIEW OF THE BLADDER.

1. Ureter.
2. Vas deferens.
3. Vesicula seminalis.
4. Trigone.
5. Prostate.

The quantity of urine which the bladder will hold without much inconvenience varies. As a general rule, it may be stated at about a pint. Much depends upon the habits of the individual; but some persons have, naturally, a very small bladder, and are obliged to empty it more frequently.

In young persons the lowest part of the bladder is the neck, or that part which joins the prostate. But as age advances, the bottom of the bladder gradually deepens so as to form a pouch behind the prostate. In old subjects, particularly if the prostate

[1] When the bladder is completely paralyzed it becomes like an inorganic sac, and there seems to be no limit to its distention. Hall found, in a drunkard, the bladder so dilated that it would hold twenty pints of water. (*Elem. Phys.*, art. Vesica.) Frank saw a bladder so distended as to resemble ascites, and evacuated from it twelve pounds of urine. (*Oratio de Signis Morborum, etc., etc.* Ticini, 1788.)

W. Hunter, in his *Anatomy of the Gravid Uterus*, has given the representation of a bladder distended nearly as high as the ensiform cartilage.

[2] In all animals with a bladder, the younger the animal the more elongated is the bladder. This is indicative of its original derivation from a tube, *i.e.* the *urachus*. In the infant, the bladder is of a pyriform shape, as it is, permanently, in the quadruped; but as we assume more and more the perpendicular attitude, the weight of the urine gradually makes the lower part more capacious.

be enlarged, this pouch becomes deep, micturition becomes tedious, and the bladder cannot completely empty its contents. It sometimes happens that a stone in the bladder is not felt; the reason of which may be that the stone, lodged in such a pouch below the level of the neck of the bladder, escapes the detection of the sound. Under these circumstances, if the patient be placed on an inclined plane, with the pelvis higher than the shoulders, the stone falls out of the pouch, and is easily struck.

The bladder is divided into a summit, a body, a base, and a neck.

The *summit* is its highest part, and to it is attached a thin, fibrous cord, the urachus, which passes up to the umbilicus, and is the obliterated remains of a canal connecting the fœtal bladder with a sac external to the fœtus, called the allantois.

The *body* on its anterior aspect is not covered with peritoneum, and is in relation with the symphysis pubis, the triangular ligament, and the obturator internus; posteriorly, it is covered with peritoneum, and is in relation in the male with the rectum, and in the female with the uterus; laterally, it is only covered with peritoneum behind, and is in relation with the obliterated hypogastric arteries, the vasa deferentia and ureters.

The *base* is the lowest part of the bladder resting upon the middle portion of the rectum, and is only slightly covered behind with peritoneum; below, it is in contact with the vesiculæ seminales and vasa deferentia, which latter passes forward as far as the prostate; the reflection of the peritoneum posteriorly with the vasa deferentia converging toward the front, forms a triangular space through which the bladder is tapped in cases of retention of urine.

The *neck* is the narrow portion where the urethra begins, and its direction is downward and forward. It is embraced by the prostate gland.

Ureter.

The *ureter*[1] is about seventeen inches long, and conveys the urine from the kidney to the bladder. In the dissection of the abdomen (p. 497), it was seen descending along the psoas muscle, behind the spermatic vessels, and crossing the common iliac artery into the pelvis. Tracing it downward, in the posterior false ligament of the bladder, below the obliterated

[1] From οὐρέω, I pass urine.

hypogastric artery, we find that it runs along the side of the bladder, external to the vas deferens, and enters it about an inch and a half behind the prostate, and about two inches from its fellow of the opposite side (Fig. 128). It perforates the bladder very obliquely, so that the aperture, being valvular, allows the urine to flow into, but not out of it. The narrowest part of the ureter is at the vesical orifice; here, therefore, a calculus is more likely to be arrested in its progress than at any other part of the canal.

Vas Deferens. This tube, about twenty-four inches in length, conveys the seminal fluid from the testicle into the prostatic part of the urethra. Taking its origin at the lower part of the globus minor, behind the testis, it ascends at the back part of the testis and epididymis, along the back of the spermatic cord, through the inguinal canal into the abdomen; then, leaving the cord at the inner ring, it curves round the deep epigastric artery, then crosses over the external iliac vessels, and descends into the pelvis on the side of the bladder, gradually approaching nearer the middle line. Before it reaches the prostate, it passes between the bladder and the ureter; then, becoming very sacculated, it runs forward internal to the vesicula seminalis, and is joined by the duct of this vesicle. The common duct thus formed, *ductus communis ejaculatorius*, terminates in the lower part of the prostatic portion of the urethra (Fig. 128, p. 552). In point of size and hardness, the vas deferens has very much the feel of whipcord,[1] its canal not being more than $\frac{1}{10}$ of an inch in diameter.

Vesiculæ Seminales. These are situated, one on either side, between the base of the bladder and the rectum, and serve as reservoirs for the fluid secreted by the testes, and also secrete themselves a fluid accessory to that of the testicles (Fig. 127). Each is a tube, but so convoluted that it is like a little sacculated bladder. When rolled up, the tube is about two and a half inches long, and a quarter of an inch in breadth; unrolled, it would be

[1] The description in the text assumes the bladder to be distended. But when the bladder is empty, the vas deferens runs down upon the side of the pelvis. In this course it may be seen, through the peritoneum, crossing—1, the external iliac vessels; 2, the remains of the umbilical artery; 3, the obturator artery and nerve; 4, the ureter.

more than twice that length, and about the size of a small writing quill. Several cæcal prolongations proceed from the main tube, after the manner of a stag's horn. The vesiculæ seminales do not run parallel, but diverge from each other, posteriorly, as far as the reflection of the recto-vesical peritoneal pouch, like the branches of the letter V; and each lies immediately on the outer side of the vas deferens, into which it opens.

The vesiculæ seminales contain a brownish-colored fluid, presumed to be in some way accessory to the function of generation.[1]

Prostate Gland. The prostate gland[2] is situated at the neck of the bladder, and surrounds the first part of the urethra (Fig. 127). In the healthy adult it is about the size and shape of a chestnut. Its *apex* is directed forward as far as the deeper layer of the triangular ligament. It is surrounded by a plexus of veins, and is maintained in its position by the pelvic fascia (p. 548). Its upper surface is about three-quarters of an inch below the symphysis pubis; its apex is about one inch and a half from the anus; the base is about two and a half.

Above the prostate are the pubo-prostatic or anterior ligaments of the bladder, with the dorsal vein of the penis between them; *below*, and in contact with it, is the rectum; on *each side* of it is the levator ani; in *front* of it are the membranous part of the urethra (surrounded by its compressor muscle), and the triangular ligament; *behind* are the neck of the bladder and the vesiculæ seminales with the ejaculatory ducts.

The transverse diameter is about one inch and a half; the vertical is about half an inch less. But the gland varies in size at different periods of life. In the child it is imperfectly developed; it gradually grows toward puberty, and generally increases in size with advancing age.

To ascertain the size and condition of the prostate during life, the bladder should be at least half full; the prostate is then pressed down toward the rectum, and readily within reach of the finger.

[1] The vesiculæ seminales are imperfectly developed till the age of puberty. In a child three years of age they can hardly be inflated with the blowpipe.

[2] From προΐστημι, to stand before.

Anatomy of the Urethra in its Passage under the Pubic Arch.

The urethra is a canal about eight inches in length, and leads from the bladder to the end of the penis. It is divided into three portions—the *prostatic*, the *membranous* and the *spongy*. At present only the relations of the *membranous part*, which comprises that part of the canal between the prostate and the bulb, can be examined. The urethra in this part is the narrowest part of the canal, and measures three-quarters of an inch on its upper surface and one-half on its lower surface, in consequence of the encroachment of the bulb. In its passage under the pubic arch it is surrounded by the compressor urethræ, and below it are Cowper's glands. It traverses the two layers of the triangular ligament, and is about an inch below the symphysis pubis, from which it is separated by the dorsal vessels and nerves of the penis, and by some connective and muscular tissue; it is nearly the same distance above the rectum; it is not, however, equidistant from this portion of the intestine at all points, because of the downward bend which the rectum makes toward the anus.[1]

The membranous part of the urethra in children is very long, owing to the smallness of the prostate at that period of life; it is also composed of thin and delicate walls, and lies close to the rectum. In sounding a child, therefore, it is very necessary not to use violence, else the instrument is likely to pass through the coats of the urethra and make a false passage.

Levator Ani.

This muscle supports the anus and lower part of the rectum like a sling; and, with the coccygeus and compressor urethræ, forms a muscular floor for the cavity of the pelvis. To see the muscle, the pelvic fascia must be reflected from its upper surface. It *arises* in front, from the posterior aspect of the os pubis near the symphysis; behind, from the inner surface of the spine of the ischium; and between these bones, from the tendinous line which marks the division of the pelvic fascia into the obturator and recto-vesical layers (p. 546). From this long origin the fibres descend inward toward the middle line, and are *inserted* thus: the anterior, the longest, passing under

[1] If a clean vertical section were made, we should see that the two canals form the sides of a triangular space, of which the apex is toward the prostate. This is sometimes called the recto-urethral triangle.

the prostate, meet their fellow in the middle line of the perineum in *front* of the anus (forming the *levator prostatæ*), joining the fibres of the transversus perinei and the external sphincter at the central tendon of the perineum; the middle, the most numerous, are inserted into the side of the rectum; the posterior are inserted partly into the coccyx and partly into the median raphé between the coccyx and the anus, and meet their fellow beneath the rectum.

The levator ani is supplied by the inferior hæmorrhoidal, the two lower sacral and the coccygeal nerves.

The action of the levatores ani is to retract the anus and the rectum after it has been protruded in defecation by the combined action of the abdominal muscles and the diaphragm.

Coccygeus. This muscle is placed behind the levator ani, and should be regarded as a continuation of that muscle. It is triangular in shape, and *arises* by its apex from the spine of the ischium and the lesser sacro-sciatic ligament, gradually spreads out, and is *inserted* into the side of the lower part of the sacrum and the coccyx. Its posterior fibres are in relation with the pyriformis, its anterior fibres are continuous with the levator ani. This muscle is supplied by the two lower sacral and the coccygeal nerves.

Dissection. At this stage of the dissection, the bladder should be drawn downward, and the branches of the internal iliac artery and the sacral plexus clearly displayed on the right side, by carefully clearing away the prolongations of the pelvic fascia which surround them.

Internal Iliac Artery and Branches. From the division of the common iliac artery, the *internal iliac* descends into the pelvis, and, after a course of about an inch and a half, divides, opposite the great sacro-sciatic notch, into two large branches, an anterior and a posterior (Fig. 129). The artery lies upon the lumbo-sacral cord, the pyriformis muscle, the external and internal iliac veins; the ureter, enclosed in the posterior false ligament of the bladder, passing in front; the psoas lies to its outer side at the commencement of its course.

The *posterior* division gives off the ilio-lumbar, lateral sacral, and gluteal arteries; the *anterior* gives off the superior vesical,

obturator, inferior vesical, middle hæmorrhoidal, sciatic and pudic; also the uterine and vaginal in the female. Such is their usual order; but these branches, though constant as to their general distribution, vary as to their origin.

The branches of the posterior division are—

a. The *ilio-lumbar* is analogous to the lumbar branches of the aorta. It ascends beneath the psoas and the external iliac vessels to get to the superficial surface of the iliacus. Here it divides into an iliac and a lumbar branch; the *iliac branch* supplies branches to the iliacus, a branch to the diploë of the ilium, and a large branch along the iliac crest, which finally inosculates with the

FIG. 129.



PLAN OF THE BRANCHES OF THE INTERNAL ILIAC ARTERY.

deep circumflexa ilii, the epigastric, the gluteal, and the external circumflex arteries; the *lumbar branch* supplies the psoas and the quadratus lumborum, and anastomoses with the last lumbar artery; it distributes a small branch to the cauda equina, through the foramen between the last lumbar and first sacral vertebræ.

b. The *lateral sacral*, usually two in number, an upper and a lower, descend in front of the sacral foramina, and inosculate on the coccyx with the middle sacral artery; the upper enters one of the upper sacral foramina, and, after supplying the structures in the sacral canal, emerges on the back through one of the posterior foramina and supplies the muscles in the neighborhood, anastomosing with the gluteal artery; the lower descends in front

of the pyriformis, supplying branches to this muscle, the bladder and rectum, and others which enter the anterior sacral foramina for the supply of the cauda equina, and finally emerge through the posterior sacral foramina to end in the muscles and skin of the back; this branch inosculates with the middle and lateral sacral arteries and the gluteal.

c. The *gluteal* is the largest branch. It passes immediately out of the pelvis through the great sciatic notch, above the pyriformis muscle, and then divides into a superficial and deep branch; the former passes beneath the gluteus maximus; the latter passes between the gluteus medius and minimus, and then divides into two branches, one running along the upper attached border of the gluteus minimus, the other passing obliquely across the same muscle as far as the great trochanter, to anastomose with the external circumflex artery. These will be dissected with the gluteal region.

The anterior division gives off—

a. The *superior vesical artery*, which comes off from the unobliterated portion of the hypogastric, and supplies the upper part of the bladder. It gives off the *middle vesical artery*, which supplies the base of the bladder and the vesicula seminalis; a small branch, the *deferential*, which accompanies the vas deferens to the testis and inosculates with the spermatic artery; and smaller branches to the ureter.

b. The *inferior vesical artery*, which ramifies on the under surface of the bladder, the vesiculæ seminales and the prostate, anastomosing with branches of the corresponding artery of the other side.

c. The *middle hæmorrhoidal artery*, which usually arises in conjunction with the preceding, and supplies the rectum, inosculating with the other hæmorrhoidal arteries.

d. The *obturator artery*, which runs along the side of the pelvis, below the corresponding nerve, to the upper part of the obturator foramen, through which it passes to be distributed to the muscles of the thigh. *In the pelvis* it lies between the peritoneum and the pelvic fascia, and gives off a small branch to the iliacus, which anastomoses with the ilio-lumbar; a *vesical branch*, which passes backward to supply the bladder; and another, the *pubic branch*,

which ramifies on the back of the os pubis, and inosculates with the corresponding branch of the deep epigastric artery and with its fellow of the opposite side. *External to the pelvis* it divides into an external and internal branch, which respectively skirt the outer and inner margins of the obturator foramen.

The obturator artery does not, in all subjects, take the course above stated, since, in one case in three and a half, it arises from the deep epigastric, and in one out of seventy-two cases it has its origin by a branch from the obturator joining a branch from the epigastric. It may arise from the external iliac near the crural arch, or by a short trunk in common with the epigastric,[1] Under

FIG. 130.



A B

VIEW OF THE DIFFERENT DIRECTIONS WHICH AN ABNORMAL OBTURATOR ARTERY MAY TAKE. (SEEN FROM ABOVE.)

A. 1. Gimbernat's ligament.
2. Femoral ring.
3. Abnormal obturator artery.
4. External iliac vein.
5. External iliac artery.
6. Diminutive obturator artery arising from its normal source.

B. 1. Gimbernat's ligament.
2. Abnormal obturator artery.
3. Femoral ring.
4. External iliac vein.
5. External iliac artery.
6. Diminutive obturator artery.

these circumstances, in order to reach the obturator foramen, it generally descends on the *outer* side of the femoral ring. Instances, however, occasionally occur, where it makes a sweep

[1] In most subjects a small branch of the obturator ascends behind the ramus of the os pubis to inosculate with the epigastric. The variety in which the obturator arises in common with the epigastric is but an unusual development of this branch. The branch derives additional interest from the fact, that after ligature of the external iliac it becomes greatly enlarged, and carries blood directly into the epigastric. See a case in *Med.-Chir. Trans.*, vol. xx, 1836.

round the *inner* side of the ring; so that three-fourths of the ring, or, what comes to the same thing, of the neck of a femoral hernia, would in such a case be surrounded by a large artery.[1]

e. The *sciatic artery* is the larger of the two branches into which the anterior trunk divides. It proceeds over the pyriformis and the sacral plexus, to the lower border of the great sciatic notch, through which it passes out of the pelvis between the pyriformis and coccygeus to the buttock, where it runs with the great sciatic nerve between the great trochanter and the ischial tuberosity. It gives off small *muscular branches* in the pelvis to the pyriformis, coccygeus and levator ani; *vesical branches* to the bladder, prostate and vesiculæ seminales; and *hæmorrhoidal branches* to the rectum.

f. The *internal pudic artery* supplies the perineum, scrotum and penis. In the pelvis it usually lies above the sciatic, and rests upon the pyriformis and sacral plexus, having the rectum to its inner side. It passes out of the pelvis through the great sciatic foramen, below the pyriformis and above the coccygeus, crosses over the spine of the ischium, and re-enters the pelvis through the lesser foramen. It then ascends on the inner side of the obturator internus toward the pubic arch, where it gives branches to the several parts of the penis. In its passage on the inner side of the obturator muscle it is enclosed in a strong tube of fascia, formed by the obturator fascia, and is situated about an inch and a quarter above the tuberosity of the ischium. It now ascends under cover of the ascending ramus of the ischium, where it pierces that part of the pelvic fascia which forms the posterior layer of the triangular ligament, and continues its course close to the ramus of the os pubis, between the two layers of the ligament, the anterior layer of which it pierces, and then divides into the artery of the corpus cavernosum and the dorsal artery of the penis. Throughout its course it is accompanied by the pudic nerve and veins. The branches of the pudic artery were described in the dissection of the perineum (p. 537).

[1] The Museum of St. Bartholomew's Hospital contains two examples of double femoral herniæ in the male, with the obturator arising on each side from the epigastric. In three out of the four ruptures the obturator runs on the inner side of the mouth of the sac.

The pudic artery, however, sometimes takes a very different course. Instead of passing out of the pelvis, it may run by the side of the prostate gland to its destination; or, one of the large branches of the pudic may take this unusual course, while the pudic itself is regular, but proportionably small. Anatomists are familiar with these varieties, and a winter session rarely passes without meeting with several examples of them. It need hardly be said that lithotomy, under such conditions, might be followed by a large hemorrhage.

The *middle sacral artery* is a small branch of the abdominal aorta at its point of bifurcation. It descends in front of the body of the fifth lumbar vertebra, the sacrum and the coccyx. In its course it gives off small branches to the rectum, to the anterior sacral foramina, and it finally inosculates on the sacrum and the coccyx with the lateral sacral arteries. It gradually becomes smaller as it passes down and terminates near the tip of the coccyx in a small body about the size of a pea, called the *coccygeal* or *Luschka's gland*, which has been previously described (p. 507).

Respecting the *veins* of the pelvis, they correspond with the arteries and empty themselves into the internal iliac vein. The remarkable plexus of veins about the prostate, neck of the bladder and rectum, has been described (p. 548).

Nerves of the Pelvis. Those which proceed from the spinal cord should be examined first, afterward those derived from the sympathetic system

Sacral Nerves. Five sacral nerves proceed from the spinal cord through the anterior sacral foramina. The upper four, from their large size, at once attract observation; but the fifth is small; it perforates the coccygeus muscle, supplying it and the skin over the coccyx.

Sacral Plexus. The anterior divisions of three upper sacral nerves, and part of the fourth, with the lumbo-sacral cord, form the sacral plexus. The great nerves of this plexus lie on the anterior surface of the pyriformis, covered by the pelvic fascia, which separates it from the branches of the internal iliac vessels and the pelvic viscera. The large cords, diminishing in size from above downward, converge from the sacral foramina to the great sacro-sciatic foramen, where they

coalesce to form a broad, flat cord, which passes out of the pelvis beneath the pyriformis muscle, for the supply of the flexor muscles of the inferior extremity.

Before describing the branches of the sacral plexus, it will be best to trace those sacral and coccygeal nerves which do not enter into the formation of the sacral plexus.

The lower part of the *fourth sacral nerve* lies on the coccygeus muscle, and divides into muscular and visceral branches, sending a filament downward to join the fifth sacral nerve. It distributes branches to the pelvic viscera, and muscular twigs to the levator ani, the coccygeus and sphincter, the latter of which also furnishes a small cutaneous filament to the skin between the bone and the anus.

The *fifth sacral nerve* emerges between the sacrum and the coccyx, pierces the coccygeus, and lies on its anterior surface. It is joined by a twig from the fourth sacral, and, after running a short distance, pierces the coccygeus again, and is distributed to the skin over the back of the coccyx. It communicates with the coccygeal nerve, and supplies the coccygeus muscle.

The *coccygeal nerve*, not easily found, emerges through the end of the sacral canal, and comes forward through the coccygeus, between the first and second pieces of the coccyx. It pierces the great sacro-sciatic ligament, and, after receiving the communicating twig from the fifth sacral nerve, it passes backward to supply the integument over the back and side of the coccyx. The communications between these three last nerves are sometimes described as the *coccygeal plexus*.

The *muscular branches* of the sacral plexus are as follows:—

a. Muscular branches, distributed to the pyriformis, the gemelli, the quadratus femoris, and the obturator internus. The *nerve to the obturator internus* is given off from the anterior aspect of the plexus (sometimes from the pudic), leaves the pelvis through the great sciatic foramen with the pudic artery, winds with it round the ischial spine, and re-enters the pelvis with the artery to reach the inner aspect of the obturator internus; it distributes a small twig to the gemellus superior. The branch to the quadratus femoris is derived from the plexus near the preceding nerve; it passes down, beneath the gemelli and obturator internus, to enter

the anterior or deep aspect of the quadratus femoris, lying between this muscle and the capsule of the hip-joint; it sends off a small twig to the inferior gemellus, and another to the hip joint.

b. The *superior gluteal nerve* proceeds from the lumbo-sacral cord and the first sacral nerve, leaves the pelvis through the great sacro-sciatic foramen with the gluteal artery, above the pyriformis.

FIG. 131.



12. N. of pyriformis.
13. N. of gemellus superior.
14. N. of gemellus inferior.
15. N. of quadratus femoris.
16. N. of gluteus maximus.
17. Long pudendal n.
18. Cutaneous n. of the buttock.
19. N. of the long head of the biceps.
20. N. of semi-tendinosus.
21. N. of semi-membranosus.
22. N. of short head of the biceps.

1, 2, 3, 4, 5. Sacral nn.
6. Superior gluteal n.
7. Great sciatic n.
8. Lesser sciatic n.
9. Pudic n.
10. N. of obturator inter
11. N. of levator ani.

PLAN OF THE SACRAL PLEXUS AND BRANCHES.

and there divides into two branches: the *upper* passes along the iliac attachment of the gluteus minimus, supplying it and the gluteus medius; the *lower* accompanies the lower branch of the gluteal artery, and supplies the glutei medius and minimus and the tensor fascia femoris.

c. The *pudic nerve* runs with the pudic artery, and is contained in the same sheath of the obturator fascia; it divides into two branches—the perineal nerve, and the dorsal nerve of the penis; the *former* accompanies the superficial perineal artery, and sup-

plies cutaneous branches and muscular branches to the external sphincter, the accelerator urinæ, the transversus perinei, the erector penis, and the compressor urethræ; the *dorsal nerve* accompanies the last part of the pudic artery, and after piercing the anterior layer of the triangular ligament and the suspensory ligament, runs along the dorsum of the penis external to the dorsal tery, and is distributed to the glans and the prepuce.

d. The *small sciatic nerve* is formed by the junction of two anches from the sacral plexus, and passes through the great ro-sciatic foramen below the pyriformis, and then divides into o branches: one, a motor—the *inferior gluteal*—supplies the teus maximus; the other, a sensory, supplies cutaneous nches to the back of the thigh and leg, to the skin over the teus maximus, and to the perineum and scrotum. These will dissected later on, with the lower extremity.

c. The *great sciatic nerve* is the large nerve cord which passes ng the back of the thigh beneath the gluteus maximus and e hamstring muscles, and will be dissected at a later stage.

Pelvic Sympathetic Plexus. From the lumbar region the sympathetic nerve descends into the pelvis along the inner side of he anterior sacral foramina. In this part of its course its ganglia ary in number, from four to five. The nerves of opposite sides nite in front of the coccyx, where they form the *ganglion impar*.

The arrangement of the sympathetic nerves in the pelvis is similar to that in the abdomen, each ganglion receiving a branch from the ganglion above and another from the ganglion below. The external branches communicate with the sacral nerves, one probably going to, and the other coming from, the spinal nerves; the internal branches pass partly to join the pelvic plexus, and partly to the plexus around the arteria sacra media.

The *pelvic plexuses* are two in number, and are situated one one each side of the rectum, being derived from the hypogastric plexus, which passes downward between the common iliac arteries into the pelvis, reinforced by filaments from the second, third and fourth sacral nerves and ganglia. The visceral branches are exceedingly delicate, and cannot be traced unless the parts have been previously hardened in spirit. They accompany the arteries supplying the respective organs, and are—the *inferior hæmorrhoidal*

plexus to the rectum; the *vesical plexus* to the sides and base of the bladder, and secondary plexuses to the vas deferens and vesicula seminalis; the *prostatic plexus* to the prostate, the vesicula seminalis, and the cavernous structure of the penis; and, in the female, the *vaginal plexus* to the vagina and its erectile tissue, and the *uterine plexus* to the neck and lower part of the body of the uterus, running between the layers of the broad ligament. It also distributes numerous filaments to the fundus of the uterus and the Fallopian tubes.

STRUCTURE OF THE BLADDER, PROSTATE, URETHRA AND PENIS.

It is assumed that the parts have been collectively taken out of the pelvis, and that the partial peritoneal covering of the bladder has been removed.

The bladder, in a fairly dilated condition, measures about five inches in length and three in breadth, and when moderately full will contain about a pint of urine.

Structure of the Bladder. The bladder is composed of a partial peritoneal coat, a muscular and a mucous; between the last two there is a layer of connective tissue, which is called the cellular coat.

The *serous* or *peritoneal coat* invests the posterior, lateral and superior surfaces of the bladder; it is absent on the anterior and inferior aspect.

The *muscular coat* is situated beneath the serous, and consists of unstriped muscular fibres, which interlace with each other in all directions. Their general arrangement is as follows: An *outer* or *longitudinal* layer arises from the pubo-prostatic ligaments, the upper half of the circumference of the prostate and the neck of the bladder, and thence its fibres spread out longitudinally over the summit of the bladder, pass round its posterior aspect and base, to be inserted into the prostate in the male, and the vagina in the female. This layer is especially marked on the anterior and posterior surfaces of the bladder. There are also some lateral longitudinal fibres which pass backward from the sides of the prostate and interlace in all directions. Between these is a thin layer of *circular fibres*, especially developed near the neck and

he commencement of the urethra, where they form a sphincter—*phincter vesicæ*. Toward the sides of the bladder the two sets of bres have a less definite arrangement and form a kind of network; ese, therefore, are the weakest parts of the bladder, and more able to the formation of pouches.[1] The development and color the muscular fibres depend upon how far the subject has suffered om irritation of the bladder, or any obstruction to the expulsion the urine.

The *cellular coat* loosely connects the muscular with the mucous oat, and is firmly adherent to the latter.

The bladder must be laid open by an incision along its front, examine its interior. In a recently contracted bladder, the ucous membrane is disposed in irregular folds, which disappear hen the bladder is distended. In a healthy state, it is pale; hen inflamed, it becomes of a bright red. Under the microscope, s surface is seen to be studded with mucous follicles. These llicles secrete the thick, ropy mucus in inflammation of the ladder.

The *mucous coat* is loosely connected with the subjacent muscular layer, except at the *trigone* of the bladder, where it is firmly dherent. The epithelium is composed of flattened polyhedral ells of the transitional variety, and beneath these there have been escribed by Klein a layer of large club-shaped cells arranged at ght angles to the surface.

When the interior of the bladder is examined, there is seen nmediately behind the urethra a triangular smooth surface, the pex being at the urethra. This surface is called the *trigonum esicæ*, and is paler and smoother than the vesical mucous membrane elsewhere; laterally, it is bounded by ridges extending om the urethra to the orifices of the ureters, the base being between the two ureters. This space corresponds with another one

[1] These pouches arise in the following manner: A portion of mucous membrane protruded through one of the muscular interstices, so as to form a little sac. This small at first, but gradually increases in size, because, having no muscular coat, it as no power of emptying itself; generally speaking, several such sacs are met with the same bladder; and they sometimes contain calculi. If a calculus, originally ose in the bladder, happen to become lodged in a pouch by the side of it, a sudden mission of the symptoms may ensue. This explains our occasional inability to etect its presence at each examination with the sound.

FIG. 132.



BLADDER AND URETHRA, LAID OPEN BY AN INCISION ALONG THE UPPER SURFACE.

eady described, external to the bladder, and which is bounded erally by the vesiculæ seminales, and behind by the reflection the peritoneum. It is more richly provided with blood vessels d nerves than the rest of the bladder, and is endowed with more ate sensibility. This is why a stone is more painful when the adder is empty; and in the erect than in the recumbent position.

The *vesical orifice of the urethra* is situated at the lower and terior part of the bladder, not at the most dependent part, which rms the pouch behind the orifice, in which urine is apt to accu- ulate in old persons. It appears small and contracted in the fresh adder, but if the little finger be introduced into it, it will dilate nsiderably. Immediately behind the orifice there is, in some ladders, a slight elevation called the *uvula*. It is composed of a rtion of the mucous membrane raised up by an accumulation of e prostatic and submucous tissue, but is rarely of sufficient size interfere with the passage of the urine. This elevation must distinguished from enlargement of the third or middle lobe of e prostate.

The *orifices of the ureters* are situated about an inch and a half hind the urethra, and about two inches apart. These tubes rforate the coats of the bladder obliquely, and slant toward ch other, standing out in relief under the mucous membrane.[1] slight ridge proceeds from the orifice of each ureter to the ck of the bladder, looking like a continuation of the ureter elf. If the mucous membrane be removed from these ridges, we d that they are produced by muscular fibres. Sir Charles ll,[2] who first drew attention to them, believed them to be of use regulating the orifices of the ureters, and named them *the uscles of the ureters.*

[1] This slanting of the ureters serves all the uses of a valve. The urine enters bladder, drop by drop, but cannot return, because the internal coat is pressed ainst the other side of the orifice, so as to stop it. When the bladder becomes ckened, in consequence of difficulty in passing urine, it sometimes happens that ureters lose their valvular direction, so that the urine, when the bladder con- cts, is partly forced back up the ureters; the result is that they become dilated, d the pelvis of the kidney also.

[2] *Med-Chir. Trans.*, vol. iii. He says: "These muscles guard the orifices of the eters by preserving the obliquity of the passage, and pulling down the extremities the ureters according to the degree of the contraction of the bladder generally."

FIG. 132.

superior, middle and
nes from the unob-
dle, from the superior
from the anterior divi-
Small branches are also
r and sciatic arteries.
plexuses around its neck,
nto the internal iliac veins.
ne veins.
gastric and sacral plexuses;
e top, the latter to the neck

amined the form, size and
e (p. 555), we have now to
wo lateral lobes presenting on
median longitudinal furrow, the
a deep cleft; and a third or
s furriform in shape, unites the
between them and the urethra. In
a separate lobe; but when abnor-
ward the cavity of the bladder, and
the urethra.

through the upper surface of the
This canal runs rather nearer to
and is not of the same calibre
urethra is about an inch and a
in diameter. It forms a sinus
described by anatomists as the *sinus*
the ducts of the prostate open. Along
ngitudinal ridge, about three-quarters

the middle lobe, in England, by Sir Everard
the *Phil. Trans.* for 1806. The prepara-
preserved in the Museum of the Royal
Series. No. 2583 A. But the anatomy and
the prostate gland is not a discovery of
described by Santorini in 1739, and subsequently
by Morgagni in the third book of his Epistles. Dr.
Trans., has shown that in subjects over
ent.

BLA

broad and elevated behind, but gradually is called the crest of the urethra, and the of it is named the *veru montanum* or *caput* supposed resemblance to the head of a wood- de of this prominence the common ejaculatory

in front of the caput gallinaginis, in the middle opening which will admit a probe. It leads back- a little cul-de-sac or pouch in the substance of the This pouch is described as the analogue of the uterus, the *utriculus* or *sinus pocularis*. It is also called the *culinus*. It is of a pyriform shape, running backward ward with the narrowest part at the orifice, and its length three lines. It ascends between the lateral lobes of the and beneath the middle; its coats are comparatively with some muscular tissue enclosed in them, and it is lined squamous epithelium. The minute orifices of the ducts are opening into the floor of the prostatic sinus. The substance the gland is permeated by the divisions and subdivisions of the ducts. They are not visible to the naked eye, but if traced out with the microscope they are seen to terminate in blind sac- culated extremities, upon which the capillaries ramify in rich profusion.[1]

Structure of the Prostate. The prostate is surrounded by a firm capsule of fibrous tissue, and is composed of muscular as well as glandular tissue. Nearly two-thirds of it is made up of unstriped *muscular fibres*, which constitute the stroma of the gland, and have the following arrangement: externally, beneath the capsule, they form a thick layer, continuous behind with the external muscular layer of the bladder; and in front they are ar- ranged in a circular manner round the urethra at its vesical orifice, so as to form in conjunction with the vesical muscular tissue, a sphincter; the next layer forms a dense interlacing stratum, in the

[1] This was first demonstrated by the late Mr. Quekett. The same anatomist has discovered that the secreting cells of the gland contain calculi of microscopic ness. He finds them, almost without exception, in the prostate at every period For further detail concerning them, consult the article "Prostate," in Todd's

With ganglion cells; the *lymphatics* pass to the internal nds.

ıcular scales. The external appearance of these bodies, each of which consists of a tube coiled upon itself, has eady described (p. 554). Respecting their structure, we they have an *external* or *connective-tissue coat* derived recto-vesical fascia; a *middle* or *muscular*, consisting of al fibres arranged transversely, and of deep fibres arranged inally, and continuous with those of the urethra; and an or *mucous*, which is lined by a scaly epithelium, and pre- oneycombed structure, not unlike that of the gall bladder. t emerges from the anterior part of the vesicula, and joins ute angle the vas deferens behind the prostate, to form mon ejaculatory duct (p. 554). Its *arteries* come from the vesical and middle hæmorrhoidal; its *veins* pass to the iliac vein, and its *nerves* are derived from the hypogastric The function of these bodies is twofold—they act voirs for the semen, and secrete a fluid accessory to on.

Glands. The glands of Cowper have been examined *in situ* in the discussion of the perineum (p. 537). e placed close to the urethra, one on either side, imme- behind the bulb and between the two layers of the tri- ligament. They consist of a number of lobules united by nective tissue, and their collective size is somewhat larger ea. Each pours its secretion by a minute duct, about an g, into the bulbous part of the urethra. The use of these s analogous to that of the vesiculæ seminales and the —namely, to pour into the urethra a fluid accessory in ıy to generation. They are found in all mammalia, and , *e.g.* the mole, they increase in size periodically with the

thra. The urethra is the canal which extends from the bladder to the end of the penis, and serves as the outlet for the urine, but to transmit the secretion sticles and the several glands accessory to generation. It length from eight to nine inches, and is divided into rtions, according to the different structures by which it is

surrounded in different parts of its course. The first inch and a quarter is surrounded by the prostate gland, and is called the *prostatic portion* (p. 549); the next three-quarters of an inch, which passes under the pubic arch, is surrounded by the compressor urethræ, and is termed the *membranous portion* (p. 549); the remainder of its course, about six inches in length, is contained in the corpus spongiosum, and is called the *spongy portion.* The length of the urethra will vary much in different subjects, and according to the condition of the penis.

The direction of the urethra, when the penis hangs flaccid, is like the letter S reversed; but if the penis be held straight, the canal forms only one curve through the pubic arch, with the concavity upward. The degree of this curvature varies at different periods of life. In the child, the bladder being more an abdominal than a pelvic viscus, the curve forms part of a much smaller circle than in the adult; but it gradually widens as age increases, and catheters are shaped accordingly.[1] However, the parts, when in a sound state, will yield sufficiently to admit the introduction of a straight instrument into the bladder. A straight staff is sometimes used in lithotomy.

In its contracted state the sides of the urethra are in close apposition; the appearance it presents on a transverse section differs in the different parts of its course. Through the glans it is flattened vertically; through the prostate it is crescentic, with its convexity upward, owing to the veru montanum. But throughout the rest of its course the canal exhibits on section the appearance of a transverse slit (Fig. 133).

The urethra must be laid open from end to end along its roof, to see that the canal is not of uniform calibre throughout. The external orifice is the narrowest and the least dilatable part; so that the urine may be expelled in a jet. Therefore, any instrument which will enter the meatus ought to pass into the bladder, if there be no stricture. The junction of the membranous with the bulbous part is almost as narrow.

[1] The sharper curve of the urethra in the child was well known to Camper. "In recenter natis, vesica basi sua elatius sita, pedetentim descendit, unde necessario sequitur curvaturam urethræ majorem esse in junioribus quam in adultis."—*Demon. Anat. Pathol.*, lib. ii, p. 13.

The *prostatic portion* of the urethra has been described with the prostate (p. 570); the *membranous portion* with the anatomy of the side view of the pelvic viscera (p. 556).

The *spongy portion*, so termed because it is surrounded by the erectile tissue of the corpus spongiosum, is about six inches long. That part of it running through the bulb is called the *bulbous portion*, and is the most dilatable part of the urethra except the prostatic. In the centre of the glans penis the canal widens into a sinus, termed *fossa navicularis;* its termination, at the *meatus urinarius*, is the most contracted part of the urethra.

The most *dilatable* part of the urethra is the prostatic. Even the narrowest parts of the canal must admit of considerable dilatation, since calculi of from three to four lines in diameter can pass through it.

The common ejaculatory ducts open into the prostatic part of

FIG. 133.

TRANSVERSE SECTIONS OF THE URETHRA.

A. Through the prostate. B. Through the corpus spongiosum. C. Through the glans penis.

the urethra, by the side of the veru montanum. The ducts of Cowper's glands open into the bulbous part. Besides these glands, a number of ducts open into the urethra, proceeding from small glands situated in the submucous tissue. These ducts, called the *glands of Littré*, or *lacunæ*, are large enough to admit a bristle, and run in the same direction as the stream of the urine. Most of them are on the lower surface of the urethra; but one, called *lacuna magna*, is on the upper surface, about one inch and a half down the canal.

The urethra is composed of three coats—a mucous, muscular, and erectile.

The *mucous coat* is continuous posteriorly with that of the bladder, and it sends down prolongations into the various ducts which open into it. It is arranged in longitudinal folds in the

surrounded in … is lined by columnar epi-
quarter is su… there are papillæ, covered
prostatic portio… ure, is the most sensitive
passes under …
urethræ, and … a double layer of *unstriped*
mainder of it… arranged longitudinally,
the corpus … superficial fibres are continu-
length of the … external fibres of which
according to … thra, being placed between
The direction … fibres of the bladder pass
like the letter … thra, and subsequently the
canal forms … the mucous membrane.
cavity upward … coats is a layer of areolar
periods of life …
than a pelvic … erectile tissue derived from
than in the … this body round the mem-
catheters are … the canal.
a sound state … with a closely-set network of
a straight in… demonstrated by quicksilver in-
times used … and join the lymphatics
In its con… their contents are transmitted
apposition; … penis to the inguinal glands.
differs in the …
is flattened … fibulous organ through which
with its conv… for three-fourths of its course;
throughout t… the glans penis. The *root* is the
appearance … by two crura to the rami of the
The urethr… supported by a strong elastic
to see that t… attached to the symphysis pubis.
external orif… of the two corpora cavernosa
that the urin… The glans is the expanded extremity
ment which … orifice of the urethra, and at its
if there be … body there is a deep circular
the bulbous … margin in front being called the

[1] The sharp… &c. Pavia, 1830. This anatomist has
recenter natis… the network of lymphatics which covers
sequitur curvat… network are smaller than the diameter of
Anat. Pathol.…

rona glandis. In these situations are a number of minute baceous glands, *glandulæ Tysonii odoriferæ*, which secrete a baceous substance, called smegma preputii. The surface of the lans has no sebaceous glands, but is covered with minute vascular apillæ, endowed with keen sensibility by the dorsal nerves of the enis. The skin of the penis is remarkably thin and extensible, nd connected to the body of the organ by loose areolar tissue, estitute of fat. At the extremity the skin forms the prepuce, or reskin, for the protection of the glans;[1] and the thin fold which asses from the under surface of the glans to the prepuce is called *enum preputii.* The skin, altered in character, is reflected over e glans, to which it is intimately adherent, and at the orifice of e urethra is continuous with the mucous membrane.

The bulk of the penis consists of two parallel cylindrical bodies, erectile structure, named, from the appearance of their interior, *rpora cavernosa.* In a groove along their under surface is dged a third cylindrical body, the *corpus spongiosum*, composed vascular spongy tissue, through which runs the urethra; an pansion of this at the end of the organ forms the *glans.* These uctures, then—the corpora cavernosa and the corpus spon-osum—together form the penis; though the corpus spongi-um appears closely united to the corpora cavernosa, yet it quite distinct from them, as shown in the transverse section ig. 134).

Corpora Cavernosa.

The *corpora cavernosa*, placed side by side, constitute more than two-thirds of the bulk of the nis. Each commences posteriorly by a gradually tapering rtion, called the *crus penis*, which is attached along a groove in : **rami** of the ischium and os pubis, where it is embraced by the **:ctor penis** (p. 532). The two crura converge, come into appo-**ion** at the root of the penis, prior to which each presents an

[1] **When the foreskin is, from birth, so tight that the glans cannot be uncovered, h a state is called a congenital phimosis. This condition occasions no inconveni-:e in childhood, but is apt, after puberty, to become troublesome and painful, so t it may become necessary to slit up the prepuce and set the glans at liberty. In sons who have a tight foreskin, it sometimes happens that, when the glans has en uncovered, the prepuce cannot be again drawn over it: this is called a para-imosis. The neck of the glans becomes tightly girt; great distention and inflam-tion are the consequences, unless the foreskin be reduced.**

ference than in the centre of the corpora cavernosa, at the root than toward the glans. Each corpus cavernosum thus consists of innumerable spaces mainly occupied by dilated venous sinuses, from which the blood is conveyed by the dorsal vein, the prostatic plexus and the pudendal veins. When the penis is flaccid, these spaces are empty; when it is erect, they are distended with blood.

The arteries of the corpora cavernosa come from the branches of the pudic (p. 539), which enter the inner side of each crus, at its bulbous enlargement, and proceed forward near the septum, distributing numerous ramifications. These are supported in the middle of the fibrous trabeculæ, and end, some in capillaries which convey their blood at once into the intertrabecular spaces; others in tendril-like prolongations with dilated extremities which project into the spaces, called *helicine arteries* by Müller, and which open directly into the cavities of the veins by funnel-shaped orifices. The helicine arteries are absent near the glans, and are best marked at the root of the penis. The peculiar appearance which they present when distended with injection is due to the fact that they are bound down to the trabecular tissue by fibrous bands.

The blood from the intertrabecular spaces of the penis returns, partly through veins which pass out on the upper surface of the penis into the dorsal vein (which joins the prostatic plexus), partly through the deep veins which leave the inner side of each crus, and the bulb, to join the internal iliac.

Corpus Spongiosum. The *corpus spongiosum* is the erectile tissue which surrounds the urethra as it runs along the penis. It commences in the middle of the perineum, anterior to the triangular ligament, in a bulb-like form—*the bulb*—and at the end of the penis it expands to form the *glans penis*. It receives posteriorly an expansion from the triangular ligament, and presents a median groove, marking its development from two lateral halves. The urethra does not pass through the middle of the spongy body, but runs nearer to its upper surface. The bulb hangs more or less pendulous from the urethra, and is surrounded by the accelerator urinæ muscle (p. 531). In old

tends lower down than in children, and is, conse-

exposed to injury in lithotomy.

The corpus spongiosum has a fibrous coat resembling very much the external fibrous investment of the corpus cavernosum, but it is thinner, whiter, and composed of more elastic tissue. The reticular structure is also finer, and the cavernous meshes smaller, and arranged in a longitudinal direction. Plain muscular fibres surround the urethra, and they are also found in considerable amount in the external fibrous coat.

Its interior consists of erectile tissue, composed of a plexus of minute tortuous veins, lined by a single layer of flattened endothelial cells, and which communicate very freely with each other. This is easily demonstrated by injecting the dorsal vein of the penis with wax. In this way we not only fill the spongy body, but also the glans, and the large veins which form the plexus round the corona glandis.

The *veins* return the blood; some by small veins, which emerge from the glans and collect on the dorsal surface to form the dorsal vein; others pass into the dorsal vein, either through the corpora cavernosa, or by curving round the sides of the corpora cavernosa; but by far the larger number join the prostatic and the pudic veins, communicating also with the subcutaneous veins of the penis and the scrotum.

The *nerves* of the penis are the *pudic* and its *superficial perineal branch.* The largest branches run along the dorsum to the surface of the glans; a few only enter the erectile tissue of the organ. The pudic nerve and its branch supply the skin and the mucous membrane. Some of the filaments distributed to the glans have connected with them Pacinian bodies, and some end in simple and compound end-bulbs. The erectile tissue is supplied by numerous filaments proceeding from the *hypogastric plexus.*

The *lymphatics* consist of a superficial and a deep set; *superficial*, proceeding from the glans and the integument of penis, join the inguinal glands. The lymphatics of the g communicate freely all round it; this explains why a vene sore on one side sometimes affects the inguinal glands on other. The *deep* lymphatics from the corpora cavernosa and corpus spongiosum pass beneath the pubic arch and join lymphatics of the pelvis.

DISSECTION OF THE FEMALE PELVIC VISCERA.

de View of the Female Pelvic Organs. After the removal of the left innominate bone, as described in the dissection for the side view of the male pelvic viscera, the vagina, rectum and ıdder should be moderately distended, the two former with tow, e latter with air. This done, the reflections of the peritoneum ust be traced, the description of which will be found in the ısection of pelvic viscera from above (Fig. 117). After this, ɜan off the peritoneum and make out the pelvic fascia and its olongations.

Pelvic Fascia. To the description of the fascia already given in the dissection of the male pelvis (p. 547) nothing ed be added, except that from the side of the pelvis it is reflected 'er the side of the vagina and the uterus, as well as the bladder. It is this fascia which in great measure supports the uterus in proper level in the pelvis. When, from any cause, the fascia comes relaxed, there is a liability to prolapsus uteri.

Levator Ani. For the description of this muscle, see p. 556.

Bladder. The female bladder is broader transversely, and upon the whole, more capacious than the ıle. The vesical plexus of veins is not so large, and there are vasa deferentia or prostate gland. The short urethra has a nstrictor muscle, as in the male, and is supported in a similar ınner by the pelvic fascia.

Venous Plexus out the Vagina. Though the veins round the neck of the bladder are comparatively small in the female, attention ould be directed to the plexus of large veins which surround e vagina. They communicate freely with the veins about the ɔtum, and empty themselves into the internal iliac. Their con-stion in pregnancy sufficiently accounts for the dark color of e vagina and the external organs, and the frequent occurrence hæmorrhoidal tumors.[1] These veins must be removed, with e connective tissue in which they are imbedded, before a clear ɔw of the parts can be obtained.

[1] During pregnancy varicose tumors may form even in the vagina. In the Berlin *d. Zeitung*, 1840, No. 11, a case is related of a woman who, at the sixth month, d to death from the bursting of a large vein in the vagina. Other cases of the ıd are related by Siebold.

The *mucous membrane* is of a pale rose-color, continuous above with that of the uterus, and below with the integument of the labia majora. It is rough and furrowed, especially near the orifice, and presents two longitudinal ridges—*columnæ rugarum*—which run, one along the anterior, the other along the posterior wall. From each side of these proceed a series of transverse ridges—*rugæ*—with rough margins directed forward. They are well marked in virgins, but repeated parturition and increasing age gradually smooth them down. The use of the vaginal rugæ is to excite the sensibility of the glans in coition. They themselves also possess keen sensibility, being richly endowed with papillæ. The mucous membrane is provided with numerous papillæ, conical in shape, and covered with a thick lining of squamous epithelium. In the submucous tissue, which is very loose, there is a good deal of muscular tissue, with a considerable venous plexus, forming a kind of erectile tissue; in it, also, are found an abundant supply of muciparous glands, which increase in size and number toward the uterus.

The *muscular coat* is arranged in two layers, a longitudinal and a circular, between which there may be demonstrated a number of interlacing fibres passing from one to the other layer. The longitudinal fibres are continuous with the superficial muscular fibres of the uterus, while the latter are chiefly aggregated at the orifice of the vagina, forming a kind of sphincter muscle, which is continuous with the external sphincter ani. Superiorly, the vagina is intimately attached to the neck of the uterus, while to the rectum it is but loosely connected.

The *erectile tissue* found in the connective tissue forms the chief strength of the vagina, being about one-twelfth of an inch in thickness. If this coat be minutely injected, we find that it is composed mainly of a plexus of veins surrounded with numerous fasciculi of unstriped muscular fibres.

Uterus. The uterus is the hollow muscular organ which receives the ovum, retains it for nine months to bring it to maturity, and then expels it, by virtue of its muscular walls. Its situation and peritoneal connections have been described (p. 523). Its axis slants forward, so that, upon the whole, the axis of the vagina and uterus describes a curve nearly parallel

Urethra

then, is so situated that it

opportunity

connected to

tate is pyriform, or rather

how, in cas

nd is somewhat flattened

the contigu

position by the broad and

that a fistul

three inches in length, two

which urine

ick in its upper part, and

Vagina

and a half; but there is a

manner in

e, the effect of pregnancy,

the extent

The leng

uterus is divided into the

average, a

est part, which lies above

and six al

is completely invested by

may be m

the depth

radually narrows down to

The vagi

nearly straight, and give

feel the ne

wnward, to the Fallopian

or lower

t of the ovary, and the

little low

s flat, and for full three-

again, in

peritoneum; its posterior

a little into

sted with peritoneum.

reverse h

rt which projects into the

pelvis.

The

ched round the neck of the

upward

her up behind than in front.

The

a transverse slit, the *os uteri*,

narrow

ehind by the posterior lip.

the neck

mination of the interior of

here a fr

n to the vessels and nerves

broad,

bivalve

ied by the uterine arteries

speculu

rnal iliac, and also by the

afterwa

inal arteries from the same

Structu

arteries (which correspond

Va

ven off from the abdominal

arteries.

vessels from the anterior division of the

nal iliac, toward the neck of the uterus, between the layers he broad ligament, and then ascends tortuously by the side he uterus, giving off numerous branches to it, which anasto- e freely with each other, and with a small branch from the rian artery. The fundus of the uterus is mainly supplied with nches from the ovarian arteries.

The *vaginal artery* ramifies along the side of the vagina, and tributes branches to the lower part of the bladder and the ctum.

The *veins*, of large size, corresponding with the arteries, form he uterine sinuses and the vaginal plexuses, which empty them- elves into the internal iliac vein.

Nerves of the Uterus. The *nerves* of the uterus are derived from the third and fourth sacral nerves, from the hypogastric and ovarian plexuses (p. 516). They accompany the blood vessels in the broad ligament to the neck of the uterus, and ascend with them along its sides.

Some small filaments continue with the vessels, and form around them plexuses, upon which minute ganglia are found.[1] But most of the nerves soon leave the vessels, and, subdividing, sink into the substance of the uterus, chiefly about its neck and the lower part of its body. A branch may be traced passing up to the fundus of the uterus, and another to the Fallopian tube.

The nerves of the uterus enlarge during pregnancy, like the arteries. Surgically speaking, the os uteri may be said to have no nerves; for it is insensible to the cautery and to the knife.

The *lymphatics* of the uterus are small in its unimpregnated state, but greatly increase in size when it is gravid. Those from the fundus and the ovaries proceed with the ovarian vessels to the lumbar glands; thus explaining the affection of these glands in ovarian diseases. Those from the body and the lower part of the uterus accompany the uterine arteries, and join the glands in the pelvis; some, however, run with the round ligament to the ...in; hence, in certain conditions of the uterus the inguinal ...s may be affected.

... uterus, vagina, Fallopian tubes and the ovaries should

[1] Beck, *Philosophical Transactions* for 1846.

now be collectively removed from the pelvis for the purpose of examining their internal structure.

The structure of the vagina has already been described (p. 582).

Structure of the Uterus.

Before the uterus is laid open, examine the shape of that portion of the neck which projects into the vagina. The back part of the cervix appears to project into the vagina more than the front; but this arises from the vagina being attached higher up posteriorly. If the vagina were cut away from the cervix, the anterior lip of the uterus would appear to project a trifle more than the posterior. For this reason, as well as on account of the natural slope forward of the uterus, the front lip is felt first in an examination per vaginam.[1] The length, however, and the general appearance of the vaginal part of the cervix vary according to the age; it is also considerably altered by parturition In the adult virgin it is smooth and round, and projects about half an inch; its mouth is a small, transverse fissure. But after parturition it loses its plumpness, the lips become flaccid and fissured, and the mouth larger than it was before.[2]

The uterus must now be laid open by a longitudinal incision, to examine its interior. In doing so, observe the thickness of its walls, which is greatest toward the fundus. Before coming into the proper cavity in the body of the uterus, slit up a long narrow canal which leads up into it through the neck. This canal, which is about an inch in length, is not of the same dimensions throughout; it is dilated in the middle, and gradually narrows toward the end. The upper end, which leads into the body of the uterus, is called *os internum;* the lower end, which leads into the vagina,

[1] This is the only way to reconcile the discrepancies one meets with in anatomical works, respecting the comparative length of the lips of the uterus. Krause, Weber, Busch and others, say the anterior is the longer; Mayer, Meckel, Quain and others, the posterior.

[2] Instances are recorded in which the neck of the uterus is preternaturally long. It has been known to project even as much as an inch and a half into the vagina. In such cases it gradually tapers and terminates in a very narrow mouth. This is said to be one cause of sterility, and it is recommended either to dilate the mouth, or to cut off a portion of the neck. In support of this opinion, it is stated that Dupuytren was once consulted by a lady on account of barrenness; finding the neck of the uterus unusually elongated, he removed a portion of it, and shortly the lady became pregnant. (Hyrtl, *Handbuch der top. Anatom.*)

os externum. The passage is called the *canal of the cervix*. It remains unchanged in pregnancy for some time after the cavity in the body has expanded, but gradually disappears with the increasing size of the embryo.

The shape of the *cavity* in the body of the uterus is triangular, with the apex toward the cervix. In a virgin uterus the cavity is very small, and its sides are convex; but in a uterus which has borne many children, the cavity has lost the convexity of its sides, and has increased in capacity. Each angle at the base is somewhat prolonged, and leads to the minute opening of the Fallopian tube. This prolongation of the angles is noticed more or less in different females, and is the last indication of the two horns of the uterus in some orders of mammalia.

The interior of the uterus is smooth at the fundus; but the reverse at the cervix. Here there is a central longitudinal ridge, both in front and behind (as in the vagina); from these, other closely set oblique ridges curve off laterally, like the branches of a palm-tree, called *arbor vitæ uterina*. The roughness produced by these ridges occasions an impression as though we were touching cartilage when a sound is introduced into the uterus.

The *neck* of the uterus is provided with small muciparous glands, of which the minute ducts open in the furrows between the ridges referred to. The secretion of these glands is glairy, albuminous, and slightly alkaline. Soon after impregnation, the secretion becomes so firm as to plug the mouth of the uterus, but shortly before and during parturition it is poured out in great quantity, to facilitate the passage of the child. It happens occasionally that one or more of the ducts of the glands become obstructed, and then dilate into small transparent vesicles, which gradually rise to the surface and burst.[1]

The walls of the uterus consist of an outer serous coat derived from the peritoneum, an inner mucous lining, and an intermediate layer of unstriped muscular tissue.

The *serous coat* has been already described.

The *muscular coat* forms the greater part of the thickness of the walls of the uterus, and consists of non-striped or involuntary

[1] These were first described by Naboth, and supposed to be true ova; hence their name, *ovula Nabothi*. (*De Sterilitate Mulierum*. Lips., 1707.)

muscular fibres, chiefly aggregated at the fundus, less so at the junction of the Fallopian tubes. The texture of these fibres is very close, and interwoven together with blood vessels, nerves, lymphatics, and connective tissue; so that in the unimpregnated uterus it is almost impossible to trace them. In the impregnated condition it is less difficult to trace them, and we can make out that the fibres are arranged in three layers—an external, a middle, and an internal.[1]

The *external layer*, placed immediately beneath the peritoneum, is thin, and its fibres, beginning as longitudinal at the cervix, run transversely round the uterus, some of them being continued in an oblique direction over the body into the broad ligaments; these are continued on to the Fallopian tubes, the round ligaments and the ligaments of the ovaries. A band of longitudinal fibres passes from the anterior surface of the uterus round the fundus to its posterior aspect, beneath the recto-uterine folds of the peritoneum.

The *middle layer* runs in all directions, having no definite arrangement of its fibres.

The *internal layer* is composed mainly of concentric circles which surround the orifices of the Fallopian tubes; at the cervix its fibres are arranged transversely, forming a sphincter. It is this layer which forms the thickest stratum and is closely connected with the mucous membrane; it is called the muscularis mucosa.

Upon the whole, the collective disposition of the muscular layers is such as to exert equal pressure on all sides, when called into action. At the same time that they expel the fœtus, the muscular fibres perform another very important function; they close the large venous sinuses consequent upon the great increase in the amount of blood during prégnancy. Therefore, little hemorrhage accompanies the expulsion of the placenta, provided it have been attached to the fundus or the side of the uterus. every one knows the danger of what is called *placenta pre* Here, the placenta, placed entirely or partly over the orifice o uterus, is attached to a part of the organ which must of nece expand during labor; and every uterine contraction increa instead of checking, the bleeding. For the same reason, para

[1] In the unimpregnated uterus the muscular fibres are about $\frac{1}{110}$ of an in length; in the gravid uterus they increase to $\frac{1}{40}$ of an inch.

the muscular fibres in immediate connection with the placenta, it where it may, is likely to be a source of serious hemorrhage parturition.

The *mucous membrane* of the uterus is more delicate and softer in that of the vagina, with which it is continuous, and is closely ited to the subjacent tissue. The greater part of it is lined by olumnar ciliated epithelium, but that which lines the lower part the cervix is squamous, like that of the vagina. Examined h a lens, the mucous membrane lining the body of the uterus seen to be covered with minute follicles or tubes (*uterine nds*) arranged at right angles to its surface. These tubes pass ward in a more or less spiral manner, some of them appearing nched and dilated at their extremities. They become greatly

FIG. 135.

IAGRAM OF THE UTERUS, ITS BROAD LIGAMENTS, THE OVARIES AND FALLOPIAN TUBES. (SEEN FROM BEHIND.)

1. Uterus.
2. Ovary, with its ligaments.
3. Fallopian tube.
4. Fimbriated extremity of Fallopian tube.
5, 5. Broad ligament.
6. Vagina.

eloped shortly after impregnation, and take an important part he formation of the *membrana decidua*.

'he arrangement of the mucous membrane in the cervix has n already described (p. 587), when the uterus was laid open to ose its cavity.

llopian Tubes. The *Fallopian tubes* or *oviducts* are situated, one on each side, along the upper free border of, and losed by, the broad ligament of the uterus, and convey the m from the ovary to the uterus (Fig. 135). They are about inches in length; one end, the *ostium internum*, opens by a

minute orifice into the upper angle of the cavity of the uterus; the other terminates in a wide, funnel-shaped mouth, the *ostium abdominale*, surrounded by fringe-like processes called the *fimbriæ*. This termination of the Fallopian tube is called the *fimbriated extremity*,[1] and its canal opens into the peritoneal cavity, so that thus the cavity of the peritoneum in the female communicates, through the uterus and the vagina, indirectly with the exterior. The fimbriated extremity extends about an inch beyond the ovary, and, by floating it in water, one or two of the fimbriæ may be seen connected with the outer end of the ovary. If the Fallopian tube be opened from the dilated end, and a probe introduced into it, you will find that the tube runs very tortuously at first, then straight into the uterus, gradually contracting in size, so that the uterine orifice scarcely admits a bristle. The free end of the tube communicates with the cavity of the peritoneum. This is the only instance where a mucous membrane is directly continuous with a serous one. It explains how the embryo may escape into the peritoneal cavity; though this is an extremely rare occurrence. It also explains what is said to have occurred, namely, the escape of the fluid in dropsy through the Fallopian tubes. In a well injected subject, the Fallopian tubes are seen to be well supplied with blood from the ovarian arteries. The coats of the Fallopian tube are three—an *external serous coat*, derived from the peritoneum; a *middle muscular coat*, consisting of plain muscular fibres; an external layer, longitudinal, and an internal layer, circular, both being continuous with those of the uterus; and an *internal mucous coat*, arranged in longitudinal folds, especially at the ovarian end, and covered with a columnar ciliated epithelium.

Ovaries. The *ovaries* (called by Galen, *testes muliebres*, being the analogues of the testes in the male) are two oval bodies situated between the two layers of the broad liga- ment of the uterus, in its posterior part. They are connected o their inner side to the uterus by a thin cord, called the *ligament the ovary*, and at their outer end they are usually attached to o of the fimbriæ of the Fallopian tube by fibrous tissue. The ova are of whitish color, with the long axis transverse, flattened fr

[1] The fimbriated extremity is sometimes called the *morsus diaboli*, since it embr in a peculiar way the ovary during sexual excitement.

above downward, and in females who have not often menstruated their surface is smooth and even; in after-life they become puckered and scarred by the repeated escape of the ova.

The position of the ovaries is described by Professor His as being nearly vertical, and he states that the Fallopian tube curves round the outer to the lower border of the ovary, so that the fimbriated extremity lies beneath the ovary, with its fimbriæ directed upward; thus the ova on their escape from the ovary fall into the Fallopian tube.[1]

The ovary is about an inch and a half long, three-quarters of an inch wide, and about half an inch thick; its weight being from one to two drachms. It consists of a dense, soft stroma, imbedded in which are numerous small vesicles (*Graafian vesicles*), muscular tissue, blood-vessels and nerves, the organ being invested by a serous covering.

The *serous layer* covers the ovary, but does not present the ordinary features of a peritoneal investment, for the covering is dull and not shining, and the epithelium consists of a single layer of columnar cells which are the remains of the germ epithelium, from which the ova and the other cells in the Graafian vesicles have been originally developed.[2]

The *stroma* composes the substance of the ovary, and consists of some connective tissue associated with a large amount of spindle-shaped cells, resembling in their appearance unstriped muscle cells. It contains also elastic tissue, and is abundantly supplied with blood vessels, which are larger at the hilum of the ovary, diminishing in size toward its surface. The outer part of the stroma is much condensed, so as to give a white appearance to the organ; this has been described as a proper fibrous coat, the *tunica albuginea ovarii*, but which does not actually exist as a separate layer.

If a section be made through the ovary, you will find that imbedded in the stroma are a large number of small, transparent vesicles, which are more abundant at the circumference of the ovary, while in the central part there are comparatively few, it being composed almost entirely of the stroma.

[1] His, "Lage der Eierstöcke," *Archiv. f. Anat.*, 1881.

[2] Waldeyer, *Eierstock u. Ei*, Leipzig, 1870; and in Stricker's *Handbuch*, 1871.

The transparent vesicles just alluded to are the *Graafian follicles*, or the ovisacs, which contain the ova.[1] In the outer part or cortical layer of the stroma of the ovary may be observed a large number of closely set minute vesicles, about $\frac{1}{100}$ of an inch in diameter, more numerous in the ovaries of young children and in some animals. In the central part or medullary portion of the stroma are seen larger and less numerous vesicles, the largest being placed most deeply; but these, as they become mature, gradually make their way toward the surface, probably by absorption, and when fully developed measure from $\frac{1}{10}$ to $\frac{1}{6}$ of an inch in diameter. One, or perhaps more than one, Graafian vesicle ruptures at each menstrual period, and the little ovum it contains escapes from the vesicle, and is either grasped by, or falls into the fimbriated end of the Fallopian tube, and is thus conveyed into the uterus. The ruptured vesicle from which the ovum has escaped becomes filled with blood, and subsequently also with an exudation from its walls, so as to constitute a reddish-yellow substance, called the *corpus luteum*, which persists for awhile, and then degenerates into a small, stellate fibrous cicatrix.

The Graafian vesicles are very numerous, especially in the young subject, the smaller ones being most numerous, their average diameter being about $\frac{1}{100}$ of an inch.[2] External to the larger vesicles there can be distinguished a *membrana propria* or basement membrane; internal to this, the stroma becomes altered so as to constitute a distinct wall to the follicle. Within this, and lining the wall of the vesicle, there is a layer of nucleated cells, called the *membrana granulosa*, which surrounds a transparent albuminous fluid in which the *ovum* or *germ* is contained. The vesicle, as it approaches the surface of the ovary, developes an additional layer of granular cells, called the *discus proligerus*, within which the ovum is imbedded, lying usually toward the free surface of the ovary.

The ramifications of the *ovarian artery* through the ovary are remarkable for their convolutions; they run in parallel lines, as in

[1] So called after De Graaf, a Dutch anatomist, who discovered them in 1672, and believed they were the true ova.

[2] It has been computed that in the ovaries of a child at birth there are no less than 70,000 Graafian follicles.

e testicle. Its *nerves* are derived from the ovarian plexus, hich comes from the renal. The *ovarian veins* form, like the ermatic veins, near the ovary, the pampiniform plexus, and then rminate, the right in the inferior vena cava, the left in the renal in.

Parovarium. The *parovarium*, or the *organ of Rosenmüller*, is the remains of a fœtal structure situated in the oad ligament, between the Fallopian tube and the ovary. It nsists of a series of convoluted closed tubules, lined with epi- elium, converging from beneath the Fallopian tube to the ovary. t their ovarian end the tubules are separate, but at their broader d they are joined by a longitudinal tube running parallel to e lower border of the Fallopian tube. It is the vestige of a tal structure, and is the analogue of the epididymis in the ale, and is connected at its uterine end with the remains of the Wolffian duct.

DISSECTION OF THE ABDOMINAL VISCERA.

The Liver. The liver is the largest glandular organ in the body, and in the adult weighs from fifty to sixty nces. It serves for the secretion of the bile, and moreover ers some of the constituents of the blood in its passage through e organ. Its diameter in the transverse direction is from ten to elve inches; from before backward it measures from six to en inches, and its greatest thickness, which is at its back, is out three inches.

Its surface is entirely covered with peritoneum, except a small rt behind, which is connected to the diaphragm and the upper rt of the right kidney by cellular tissue, and again in the llow which lodges the gall bladder.

The *upper surface* is smooth and convex, in adaptation to the aphragm, and is marked by a fold of peritoneum running from hind forward, dividing this surface into two unequal lobes, a ght and a left, the right being the larger. The fold of perito- um is the *suspensory* or *broad ligament.*

The *under surface* is concave and irregular, and is divided into right and left lobe by the longitudinal fissure.

The *posterior border* is thick and round, having attached to it the *coronary ligament.*

The *anterior border* is thin and sharp, and presents a notch indicating the division into a right and left lobe; the notch lodges the round ligament, which is the remains of a fœtal structure, the umbilical vein. There is also to the right side of the notch a slight groove corresponding to the base of the gall bladder.

The *right border* is thick and round, the *left* is flat and thin.

Fissures. The under surface is irregular, and is marked by five fissures which map out the five lobes (Fig. 136). They are, the longitudinal fissure, the fissure for the ductus venosus, the fissure for the gall bladder, the fissure for the inferior vena cava, and the transverse fissure. The relative position of these fissures (the liver being *in situ*) may be best impressed on the memory by comparing them collectively to the letter H. The transverse fissure represents the cross-bar of the letter; the longitudinal fissure and the fissure for the ductus venosus represent the left bar; the fissures for the gall bladder and the vena cava make the right bar.

The *longitudinal fissure* divides the right from the left lobe, and contains the round ligament, which is the remains of the umbilical vein in the fœtus. It is deeper in front than behind, and is not infrequently bridged over by liver tissue, constituting the *pons hepatis.*

The *fissure for the ductus venosus* is the continuation backward of the longitudinal fissure to the posterior border of the liver, and contains a fibrous cord, which is the obliterated remains of what was, in the fœtus, the ductus venosus.

The *fissure for the gall bladder* is a shallow depression to the right of the longitudinal fissure, and lodges the gall bladder.

The *fissure for the inferior vena cava* runs obliquely backward and joins at an acute angle the fissure for the ductus venosus at the posterior border of the liver. It begins close to the right extremity of the transverse fissure, and is not uncommonly surrounded by hepatic tissue.

The *transverse* or *portal fissure*, about two inches in length, unites the other fissures, and transmits the large vessels which enter the

liver in the following order: in front is the hepatic duct, behind is the vena portæ, and between them the hepatic artery.

The lobes of the liver, five in number, are also seen on its under surface.

Lobes.

The *right lobe*, much larger than the left, is separated from it by the longitudinal fissure on its under aspect, and by the falciform ligament on its upper surface. Its superior surface is smooth and convex, somewhat quadrilateral in shape; on the under surface it is marked by three fissures—the transverse, and those for the gall bladder and vena

FIG. 136.

DIAGRAM OF THE UNDER SURFACE OF THE LIVER.

cava. It has also on its under surface two shallow fossæ; the anterior (*impressio colica*) is for the hepatic flexure of the colon; the posterior (*impressio renalis*) for the right kidney and the suprarenal capsule.

The *left lobe* is the smaller; its upper surface is smooth and convex, its lower is concave and rests on the stomach.

The remaining lobes may be considered as forming parts of the right lobe, and are the lobulus Spigelii, the lobulus caudatus and the lobulus quadratus.

The *lobulus Spigelii* is a prominent quadrilateral lobe, placed between the transverse fissure and the fissures for the ductus venosus and vena cava; behind the transverse fissure it is connected to the right by a ridge, the lobulus caudatus.

The *lobulus caudatus*, which passes obliquely forward and to the right, separating the fissure for the vena cava from the transverse fissure.

The *lobulus quadratus* is a square lobe situated between the gall bladder, the longitudinal and the transverse fissures. This lobe is occasionally connected to the left lobe by a bridge of hepatic substance arching over the longitudinal fissure, and alluded to before as the pons hepatis.

Ligaments. The liver has five ligaments, of which the *coronary*, the *right* and *left lateral* and the *falciform* are reflections of the peritoneum; the fifth is the *round ligament*, placed in the anterior free border of the falciform ligament in the longitudinal fissure; it consists of the remains of the umbilical vein of the fœtus. The ligaments have been already described (p. 475).

Vessels. The vessels which pass to and from the liver are five also in number: the hepatic artery, the vena portæ, the hepatic veins, the hepatic duct and the lymphatics. The consideration of these is deferred till we have examined the capsule of the liver.

The *fibrous coat* surrounds the liver, and is best seen on those parts of it not covered with peritoneum. This coat is connected on the surface of the gland to the areolar tissue which surrounds the lobules, but does not send down partitions to form a framework for the interior. It is continuous, at the transverse fissure, with the sheath of loose areolar tissue, called *Glisson's capsule*, which surrounds the vessels as they enter that fissure, and encloses them in a common sheath in their ramifications through the liver.

The interlobular connective tissue is exceedingly delicate; hence the great liability of the liver to be lacerated by external violence, and consequent hemorrhage into the peritoneal cavity.

Lobules. The liver consists of an aggregation of small polyhedral masses, called *lobules*, about the size of a millet seed, which range from $\frac{1}{24}$ to $\frac{1}{12}$ of an inch (1 to 2 millimetres) in diameter. These lobules are more or less distinctly marked out by septa of areolar tissue, and in a transverse section have the appearance of mosaic pavement (Fig. 137); but in a per-

pendicular section they somewhat resemble an oak leaf (Fig. 138). Each lobule is mapped out by and separated from the adjacent lobules by delicate connective tissue, in which runs a plexus of vessels—*interlobular plexus*—consisting of branches of the portal vein. From this plexus passes inward into the lobule a fine capillary network, whose branches converge to the centre of the lobule and end in a single vein, the *interlobular vein*, which, in its turn, opens into the *sublobular vein*, and thence into the *hepatic vein*. With the interlobular plexus run the biliary ducts. The lobules themselves consist of a minute plexus of blood vessels, ducts, and cells—*hepatic cells*—which latter fill up the spaces

FIG. 137.



a. Interlobular vein.

b. Intralobular or central vein.

TRANSVERSE SECTIONS OF THREE LOBULES OF THE LIVER, MAGNIFIED TO SHOW THE PORTAL VENOUS PLEXUS. (After Kiernan.)

between the ramifications of the vessels. It will facilitate the understanding of the different hepatic vessels, if it be borne in mind, (1) that the portal vein, hepatic artery and hepatic duct ramify together from first to last, enclosed in a common sheath of connective tissue, called *Glisson's capsule;* (2) that the hepatic veins run alone from first to last, and terminate in the inferior vena cava as it passes under the liver.

The *portal vein* enters the transverse fissure of the liver, accompanied by the entering hepatic artery and the emerging right and left hepatic ducts, which, as before stated, are surrounded in the liver by a common sheath called Glisson's capsule. In the

liver, the portal vein ramifies between the lobules and gives off numerous branches, called *interlobular* or peripheral veins, which pass between the lobules and communicate freely with each other. These receive the *vaginal* and *capsular veins* which convey the blood from the corresponding branches of the hepatic artery. The interlobular plexus of veins give off a minute capillary network, which penetrates into the interior of the lobule, and converges toward the centre of the lobule to open directly into a single central vein called the *intralobular vein*. The capillaries which compose this network run in a nearly parallel direction from the circumference to the centre of the lobule, and communicate freely with each other by small transverse branches. The intervals between the branches of the capillary network are very small, varying from $\frac{1}{1000}$ to $\frac{1}{250}$ of an inch, while the vessels themselves are comparatively large, being about $\frac{1}{2500}$ of an inch in diameter. The intralobular vein returns the blood from the centre of the lobule, and opens immediately into a *sublobular vein*, larger or smaller, as the case may be, upon which the lobule is sessile (Fig. 138).

FIG. 138.



LONGITUDINAL SECTIONS OF THE LOBULES OF THE LIVER. INTRALOBULAR VEINS SEEN JOINING THE SUBLOBULAR.

The sublobular veins, increasing in size, empty themselves into the smaller *hepatic veins;* these unite to form the main hepatic trunks, usually three in number, one each from the right and left lobes and the lobulus Spigelii. These hepatic veins open into the inferior vena cava, as this vessel passes to the posterior border of the liver.

The *hepatic artery*, entering the transverse fissure of the liver, divides and subdivides with the portal vein and biliary ducts, and ramifies with them between the lobules. The artery distributes branches—*vaginal branches*—while within the portal canals which supply the coats of the hepatic vessels and Glisson's capsule; also *capsular branches* to the fibrous coat of the liver which appear on the surface of the liver in stellate plexuses; and *interlobular branches* which accompany the corresponding veins and pass into

the lobules to join the capillary network which leads to the radicles of the intralobular vein.

The interior of each lobule—that is, the space left between the several vessels—is filled by the *hepatic cells*. They are of spheroidal or polyhedral shape, with a diameter varying from $\frac{1}{1100}$ to $\frac{1}{850}$ of an inch, and have no cell wall. They consist of a granular substance, of a light yellow tinge, containing one or more distinct nuclei having a highly refracting nucleolus. In some cases the cells contain fat globules; when these accumulate in large quantities, they constitute what is called a fatty liver. The office of the hepatic cells is to separate the bile from the blood, and, when filled with bile, to discharge their contents into the biliary ducts.

The *biliary ducts* are the channels through which the bile is conveyed into the hepatic ducts. They commence as minute passages—*bile canaliculi*—which are formed between and around the hepatic cells, and then pass outward to the circumference of the lobule, where they open into the bile ducts between the lobules.[1] Here they form a plexus—*interlobular*—which opens into ducts which run with the hepatic artery and portal vein in the portal canals; eventually these join with other ducts to form the right and left hepatic ducts, which leave the transverse fissure of the liver, and, after a course of about an inch, unite to form a single duct, the *common hepatic duct.*

The *lymphatics* of the liver are arranged in two sets, superficial and deep. The *superficial* may be divided into those on the upper convex surface and those on the under aspect; the lymphatics on the convex surface are: (1) those which run backward between the layers of the falciform ligament, and then pass through the diaphragm behind the ensiform cartilage to enter the anterior mediastinal glands, and thence to the right lymphatic duct; (2) those which pass forward over its anterior margin to the under aspect, to enter the glands in the gastro-hepatic omentum; (3) those which run to the right lateral ligament, and then either pierce the diaphragm to join the anterior mediastinal glands,

[1] The biliary ducts between the lobules are lined with a short columnar epithelium, resting on a basement membrane; but it is doubtful whether the bile canaliculi are very minute ducts, or whether they are not simple intercellular passages between the hepatic cells.

or pass inward to open into the thoracic duct at its commencement; and (4) those which run to the left lateral ligament, which, after piercing the diaphragm, open into the anterior mediastinal glands. The lymphatics on the under aspect are arranged as follows: (1) those on the right side of the gall bladder open into the lumbar glands; (2) those surrounding the gall bladder accompany the hepatic artery to join the glands in the gastro-hepatic omentum; (3) those on the left side enter the glands on the lesser curve of the stomach and the œsophagus. The *deep lymphatics* originate partly in the connective tissue between the lobules, and there accompany the hepatic artery and portal vein and duct to the transverse fissure, to join the lymphatic glands on the lesser curve of the stomach and behind the pancreas; some lymphatics also accompany the branches of the hepatic veins.

The *nerves* of the liver are derived from the pneumogastric, chiefly the left, and from the hepatic plexus which comes from the cœliac plexus. These plexuses enter the liver at the transverse fissure, surround the hepatic artery and the portal vein, and accompany these vessels in their ramifications through it. The ultimate termination of these nerves is not known.

The *functions* of the liver may be thus briefly expressed: 1. It renders the albuminous matter (albuminose), brought to it by the portal vein, capable of being assimilated. 2. It forms a substance, *glycogen*, easily converted into sugar, which passes into the hepatic veins, and, being consumed, helps to maintain animal heat. 3. It secretes the bile, which assists in converting the chyme into chyle and reducing it into a state fit to be absorbed by the lacteals. 4. The bile acts as a natural aperient. 5. The bile is an antiseptic, and probably prevents the decomposition of the food during its passage through the intestine.

Gall Bladder. The gall bladder, or reservoir for the bile, is pyriform in shape, and is confined in a slight depression on the under surface of the right lobe of the liver (p. 476). It is about four inches long, an inch in its broadest part, and is capable of holding about 1½ oz. of fluid. Its broadest part projects beyond the anterior margin of the liver, and it is maintained in its position by the peritoneum, which usually is continued over it from the liver, but which may occasionally com-

pletely invest it, so as to form a kind of mesentery. It is divided into a fundus, a body and a neck. The *fundus* is its broadest part, and usually projects a little beyond the front border of the liver; from this it gradually narrows, forming the *body;* and this again still further contracts to form the *neck*, which makes a bend downward, curving upon itself like the letter S. The neck terminates in a duct called the *cystic duct*, which, after a course of about an inch, joins the common hepatic duct at an acute angle (Fig. 109, p. 487). The common duct thus formed, called the *ductus communis choledochus*, is about three inches long, and of the size of a crow quill; it opens into the inner side of the back of the descending part of the duodenum, after running very obliquely through the coats of the bowel.

The gall bladder consists of two coats, and of a partial peritoneal covering which only completely surrounds the fundus.

The *middle coat* consists of strong connective tissue, whose fibres interlace in all directions, and in which involuntary muscular fibres can be traced, running mainly in the long axis of the gall bladder.

The *mucous coat* can only be seen when the gall bladder has been opened, which should now be done, by laying it open from the fundus to the neck. It is loosely connected to the middle coat, and it is gathered into ridges, which give it a honeycombed appearance, more or less tinged yellowish-brown by the bile. This appearance is most marked in the middle of the gall bladder, where the surface is covered with polygonal ridges enclosing depressions, in which may be seen with a lens numerous openings leading down to mucous follicles. It is covered with columnar epithelium, which secretes an abundance of viscid mucus. At the bend of the neck of the gall bladder both its coats project very much into the interior, making the opening considerably narrower that it appears to be outside. In the cystic duct the mucous membrane presents a series of folds, so arranged one after the other as to form a complete spiral valve. The probable use of this is to prevent the too rapid flow of the bile. The gall bladder appears to serve mainly as a reservoir for the bile while digestion is not going on. The bile becomes during its sojourn in the gall bladder very viscid and intensely bitter.

The gall bladder is supplied with blood from the cystic branch of the right hepatic artery; its blood is returned by the cystic vein, which opens into the vena portæ; its nerves are derived from the cœliac plexus, which accompany the hepatic artery.

Pancreas. The *pancreas* is a large gland belonging to the acino-tubular class. It is placed transversely across the posterior wall of the abdomen, and measures from six to eight inches in length, about an inch and a half in breadth, and from half an inch to an inch in thickness; its weight being usually from two to three and a half ounces.

It presents for examination a head, a body and a tail.

The *head*, situated to the right side, is turned down, and is embraced by the descending and transverse portions of the duodenum, the superior and inferior pancreatico-duodenal arteries running between them. A considerable prolongation usually extends upward from the posterior part of the gland, and reaches the lesser curve of the stomach; this constitutes sometimes a separate mass, and is then termed the *lesser pancreas*.

The *tail* is the narrow end of the pancreas, which extends to the left as far as the hilum of the spleen.

The *body* is convex in front, and is covered by the ascending layer of the transverse meso-colon.

The pancreas has a *posterior surface*, which is concave, and lies on the vena cava, the aorta, the crura of the diaphragm, the superior mesenteric artery and vein, the commencement of the vena portæ, and the inferior mesenteric vein; an *upper border*, which is thick, and is in relation with the splenic artery and vein, the cœliac axis, the hepatic artery, and the first portion of the duodenum; and a *lower border*, which is thin, and is in relation with the superior mesenteric artery and vein, and on its left with the inferior mesenteric vein.

The *duct of the pancreas*, called also the *duct of Wirsung*, passes from left to right in the pancreas, nearer its lower than its upper border, and nearer its anterior than its posterior surface. Commencing at the tail by the junction of the smaller ducts, it receives, in its course to the right side, repeatedly, other ducts, and thus gradually increases in size until it reaches the head, where it usually receives a large branch from the lesser pancreas; the large

duct then curves downward, and comes into relation with the common bile duct; it then passes very obliquely through the coats of the descending duodenum on its posterior aspect, and then either opens separately, or in conjunction with the common bile duct, into this portion of the intestine.

The *structure* of the pancreas resembles, in most of its characters, that of the salivary glands. The alveoli are tubular, frequently convoluted, and are lined by columnar epithelium, which leaves only a narrow lumen, which is often occupied by spindle-shaped cells called the *centro-acinar cells*.[1] The ducts are very constricted at their commencement from the alveolus, and are lined by short columnar epithelial cells, which become larger toward the termination of the duct.

The *arteries* of the pancreas are derived from the splenic, the superior and inferior pancreatico-duodenal branches, respectively, of the hepatic and the superior mesenteric arteries. The veins open into the splenic and superior mesenteric veins. The *lymphatics* end in the lumbar glands; its *nerves* are derived from the solar plexus.

The *uses* of the pancreatic fluid are: to convert the starchy matters into dextrine and grape sugar; to emulsify (in conjunction with the bile) the fatty particles, and, by its alkaline salts, partly saponify the fatty acids, thus enabling them to be taken up by the lacteals; and, lastly, to convert into peptones the albuminous and gelatinous substances of food. It is an alkaline fluid, very similar to that secreted in the salivary glands, but contains no sulpho-cyanogen.

Spleen.

The *spleen* is a very vascular, sponge-like organ, and belongs to the class of ductless glands.[2] It varies in size according to the amount of blood in it, fluctuating in weight, consistently with health, between five and ten ounces.[3] It is of a reddish-blue color, is more or less elliptical in shape, and in its natural position is placed with its long axis nearly vertical.

[1] Langerhans, *Beiträge*, etc., Berlin, 1869.

[2] The other ductless glands are, the thyroid, thymus, and supra-renal capsules.

[3] In ague and other forms of fever, it sometimes attains a weight of from 18 to 20 pounds; it diminishes in weight after forty years of age, and is enlarged during and after digestion.

[illegible] in length, three to four inches in breadth, [illegible] and a half inches in thickness. Its *outer* [illegible] and convex, and corresponds to the ninth, tenth [illegible] on the left side, being in relation with the under [illegible] diaphragm; its *inner surface* is concave, and is [illegible] cardiac end of the stomach; this surface is divided [illegible] and a smaller posterior portion by a vertical [illegible]—at the bottom of which are large openings, [illegible] the vessels enter and emerge from the spleen. [illegible] an *upper*, thick and rounded; a *lower*, pointed; [illegible] rounded; and an *anterior*, also rounded, and often [illegible].

[illegible] mentioned, the spleen is connected to the stomach [illegible] *gastro-splenic omentum*, and to the under aspect of the [illegible] by the *suspensory ligament*.

The spleen is invested with two coats—a serous or peritoneal, [illegible] fibro-elastic. The *outer* or *serous coat* entirely covers the [illegible], except at the hilum, from which it is reflected to the stom[illegible]. It is thin and smooth, and is intimately connected to the [illegible] fibrous coat. Its *fibro-elastic coat* (tunica propria), thick [illegible] strong, not only covers the spleen, but sends inward through- [illegible] substance fibrous bands (*trabeculæ*), which interlace in all [illegible], and thus form an intricate network of what are termed [illegible] *spaces* or *areolæ*; this coat consists of a strong, white [illegible] tissue, and is consequently exceedingly elastic, to ad- [illegible] the varying size of the spleen; it moreover contains more [illegible] unstriped muscular fibres, so that it contracts faintly on the [illegible] of the galvanic current. Besides this, the trabeculæ [illegible] sheaths and supports for the splenic vessels throughout their [illegible].

The areolæ, above described, are filled with what is termed [illegible] *pulp*. This pulp is a soft, reddish-brown substance, [illegible] under the microscope is seen to consist of connective-tissue [illegible], which, with their branched communicating processes, [illegible] the *sustentacular cells of the pulp*, make up a fine reticular [illegible], the interstices of which are filled with red and white blood [illegible]. Thus the areolæ are divided into a large number of [illegible] spaces by these sustentacular cells, and the white blood

cells contained within them are more numerous than in normal blood, especially in the neighborhood of the Malpighian corpuscles. The cells have either one or more nuclei, according to their size, and present distinct amœboid movements. In these cells, frequently, small, yellowish granules may be distinguished, which are probably derived from blood cells, for they present all the characters of hæmatin. Blood corpuscles in all stages, from an unchanged disc to one of complete disintegration, may be seen; and it has been shown by Klein that some of them present knob-like projections, as if from budding of small nuclei by a process of gemmation.

The *splenic artery* enters the hilum of the spleen by several branches which ramify in its substance, ensheathed and supported by its fibrous framework.[1] The artery is remarkable for its large size, as compared with the organ to which it is distributed, and also for its serpentine course. The smaller branches leave the trabeculæ, still invested by a sheath derived from the fibro-elastic coat of the spleen; but, before they terminate in penicillate tufts, the sheath becomes changed into a thick investment of lymphoid tissue, which surrounds the smallest arterioles. The lymphoid tissue which forms the sheath of the arterioles is here and there dilated into oval enlargements, called the *Malpighian corpuscles*, varying in size from $\frac{1}{70}$ to $\frac{1}{25}$ of an inch in diameter. These bodies are sometimes thickenings on the side of the arterioles, but more commonly they completely surround the vessels. They are visible in a fresh spleen, and look like white spots scattered through the dark pulp. There do not appear to be any definite boundaries between them and the reticular tissue; their interior consists of a fine reticulum, denser at the circumference than at the centre, and is filled with lymphoid cells possessing amœboid movements. The smaller arteries, after branching in all directions, enter the spleen pulp, and their lymphoid walls alter in character, presenting numerous branched processes which communicate with the branched cells of the sustentacular tissue. Through this connection they pour their blood directly into the pulp tissue, and thus into relation with the constituents of the

[1] The ramifications of the splenic artery may be seen by washing away the pulp, and floating the flocculent-looking spleen in water.

pulp tissue, by which means it is subjected to important changes. The *veins* commence in the pulp tissue in the same way as the arteries, and are at first formed by the arrangement into rows of the connective-tissue corpuscles, which subsequently become spindle shaped and overlap each other, so constituting a variety of endothelial lining to the venous passages. Assuming more the ordinary character of veins, they travel along the trabeculæ like the arteries, but do not accompany them, and freely communicate with each other, and so far are unlike the arteries. The small veins present transverse lines or markings, caused by the encircling elastic fibres around the vessels of the sustentacular tissue of the spleen. After entering the trabecular tissue which forms sheaths for the veins, they gradually join and form four to six large veins, which leave the hilum to constitute the splenic vein.

The *lymphatics* of the spleen are arranged in two sets—a *trabecular* and a *perivascular;* the former originate in the trabeculæ, and are connected with the lymphatics beneath the capsule; the latter arise in the lymphoid tissue around the arteries, and subsequently run on each side of the arteries, anastomosing frequently by transverse branches. The trabecular and perivascular lymphatics join at the hilum, and run between the layers of the gastro-splenic omentum to the lymphatic glands.

The *nerves* come from the solar plexus and the right pneumogastric nerve.

The function of the spleen appears to be that of a great blood gland, and thus concerned in the development of white corpuscles; for the blood which is conveyed from the spleen contains a large excess of white corpuscles. The large number of red blood corpuscles, in various stages of disintegration, also points to another use of the spleen as the gland for the degeneration of red blood cells into pigment, which is conveyed through the spleen to the liver to be used in the secretion of the bile. It is also presumed that the gland elaborates the albuminous materials of food, and stores them up for a time before they pass into the circulation.

Kidneys.

The kidneys, two in number, are situated in the lumbar region, behind the peritoneum, imbedded in fat. The left, usually situated higher than the right, is

generally longer and somewhat heavier. Their color is reddish-brown. Each is about four inches in length, two inches and a half in breadth, and one inch and a quarter in thickness. Each weighs about 4½ ounces in the male and rather less in the female.

The kidney presents for examination two surfaces, two borders and an upper and a lower end.

The *anterior surface* is convex and is covered with peritoneum, and looks somewhat outward; the *posterior surface* is rather flattened and rests on the anterior layer of the lumbar aponeurosis and psoas magnus; the *outer border* is convex and rounded; the *inner border* presents, about its middle, a deep notch about an inch in length, the *hilum*, leading to a hollow in the kidney, the *sinus*, for the entrance and exit of the renal vessels and ureter, the nerves and lymphatics; these have the following relations to one another: in front lies the renal vein; behind is the ureter; between them is the renal artery; the *upper end* is large and thick, and looks upward and inward, and upon it rests its corresponding supra-renal capsule; the *lower end*, smaller and flatter than the upper end, looks downward and outward.

The kidney is surrounded by a thin, *fibrous capsule* of thick connective tissue, to which it is loosely connected by areolar tissue and minute vessels, except at the hilum; here it is reflected inward and becomes continuous with the walls of the renal vessels and ureter. The capsule can be readily stripped off when healthy, leaving the surface perfectly smooth.[1]

A longitudinal section should be made through the kidney, from the outer to the inner border, to demonstrate the interior. This section displays two distinct parts—an *outer* or *cortical portion*, and an *inner* or *medullary portion*.

The *cortical structure* is deeper in color than the medullary portion, and is soft and easily lacerated. It forms the external portion of the kidney to the extent of four to five lines, arching over the bases of the pyramids. It moreover sends down prolongations between the pyramids as far as the sinus, forming the *septula renum* or the *columns of Bertini*. The cortical sub-

[1] Under the capsule there have been traced unstriped muscular fibres forming an incomplete layer.

FIG. 139.



SECTION OF THE KIDNEY.

1. Ureter.
2. Pelvis of the kidney.
3, 3, 3. Papillæ.

stance consists of convoluted and straight tubes, *tubuli uriniferi*, of little reddish granules called *Malpighian bodies*, and of blood vessels, nerves and lymphatics (Fig. 139).

The *medullary structure* is composed of numerous conical masses, the *pyramids of Malpighi*, having their bases directed to the surface, their sides in relation with the columns of Bertini, and their apices, termed *papillæ* or *mammillæ*, projecting into one of the calices of the ureter. The pyramids, of which there are from eight to sixteen, are surrounded by the cortical substance; they are composed of minute straight tubules (which proceed from the cortical portion to end in the papillæ), of looped tubes described by Henle, and of arteries and veins.[1]

At the hilum is the dilated commencement of the ureter, called the *pelvis of the kidney*. It is funnel-shaped, and its broad part divides into three principal channels, *infundibula*, an upper, middle and lower, which again branch and form from eight to twelve cup-like excavations, called *calices*. Into each of these calices one, sometimes two or more, papillæ project. Between the calices the branches of the renal artery ascend to ramify in the kidney, lying imbedded in fat. The pelvis and the calices are composed of three layers—an *external fibrous layer* continuous with the reflected part of the capsule into the sinus; a *middle* or *muscular*,

[1] Each pyramid represents what was, in the early stage of the kidney's growth, a distinct and independent lobe. In the human subject the lobes gradually coalesce, and no trace of their primordial state remains, except the pyramidal arrangement of the tubes. But in the kidneys of the lower mammalia, of birds and reptiles, the lobes are permanently separate.

consisting of longitudinal and circular fibres, the former extending nearly as far as the calices, the latter encircling the calyx round the papillæ; and an *internal* or *mucous coat*, reflected over the papillæ.

Structure of the Kidney. With a lens, each papilla may be seen to be studded with minute apertures, which are the terminations of the *tubuli uriniferi*. These apertures open into the bottom of about twenty shallow depressions on a papilla, called *foveolæ*. The orifices are from $\frac{1}{150}$ to $\frac{1}{100}$ of an inch in diameter. These tubes as they pass out into the pyramidal structure run straight, bifurcate repeatedly at very acute angles, their subdivisions running parallel, and reach the bases and sides of the pyramids, from which they pass into the cortical substance, greatly increased in number. These, termed the *straight* or *collecting tubes*, pass into the cortical substance still as straight tubes, the central ones passing nearly to the surface, the outer ones being very short, and only run a short distance into the cortex, so that they are arranged as a series of cones, with their apices to the surface of the organ. These bundles are called the *pyramids of Ferrein*, or the *medullary rays*, and receive on each side the curved extremities of the convoluted tubes. We find the cortical substance is arranged between and around these medullary rays, which, from the intricate arrangement of its tubes, receives the name of the *labyrinth of the cortex*.

Each uriniferous tubule commences in a dilated extremity, termed the *Malpighian capsule*, in which is enclosed an arterial vascular tuft, the *Malpighian tuft*, of about $\frac{1}{130}$ of an inch in diameter, and is visible to the naked eye as a minute red point. At the point of union of the tubule with the capsule, it presents a narrow portion, called the *neck*, beyond which the tubule becomes convoluted for a considerable distance, forming the *first* or *proximal convoluted tube* (Fig. 140, 2). As it descends toward the medullary ray, the tubule becomes nearly straight, but having a slight spiral tendency; this portion of the tube is termed the *spiral tubule* (Schachowa). The tubule now enters the medullary portion, narrowing very suddenly in its calibre, and descends toward the apex of the pyramid, constituting the *descending limb of Henle's loop* (3). The tubule here bends upon itself, forming a loop, the *loop of Henle*, and

ascends to re-enter the cortical substance as the *ascending limb of Henle's loop* (4), which is larger than the descending limb. On passing out of the medullary ray of the cortical portion, the tubule becomes irregularly dilated, and takes the name of the *irregular tubule* (5); this is continued on into another convoluted portion, called the *second* or *distal convoluted tubule* (2^b), which, before entering the

FIG. 140.



1. Malpighian capsule.
2. First or proximal convoluted tubule.
2b. Second or distal convoluted tubule.
3. Descending limb of Henle's loop.
(Between 2 and 3 the tube is called the spiral tubule of Schachowa)
4. Ascending limb of Henle's loop.
5. Irregular tubule.
6. Collecting tube.
(Between 2 and 6 the tube is called the curved or junctional tubule.)
6, 7, 8. Different portions of the collecting or straight tube.
a. Apex of pyramid.
b. Base of pyramid.
c. Cortical portion.

DIAGRAM OF THE COURSE AND ARRANGEMENT OF THE URINIFEROUS TUBES.

straight tube, becomes much narrowed and curved, called the *curved* or *junctional tubule*. We have thus traced the straight tubes from their termination at the papillæ to their commencement at the pyramids of Ferrein, and have also traced the convoluted tubules from their origin in the Malpighian capsules to their junction with the commencement of the straight tubes.

The tubuli uriniferi consist of a basement membrane lined with

epithelium, which varies in the different parts of the tubuli. The *capsule* is lined with flattened cells, having oval nuclei; the *neck* has cubical epithelium; the *first convoluted tubule* is lined with polyhedral epithelium, presenting numerous rod-like processes, resting at one end on the basement membrane, while the other extends toward the lumen of the tubule,[1] and thus presents the appearance of striation; the *spiral tubule of Schachowa* has similar epithelium; the *descending limb* is lined with flattened epithelium, like that in the capsule; the *ascending limb* presents epithelium similar to that found in the first convoluted and spiral portion of the tubule, although smaller and with shorter rod-like processes; the *irregular tubule* is furnished with the rod-like cells of unequal length, which, however, render the lumen more uniform; the *second convoluted tubule* has epithelium somewhat like that of the first convoluted tubule, but having long cells with large nuclei, and possessing high refractive properties; the *curved* or *junctional tubule* has a large lumen, and is lined by angular or fusiform cells with short processes; the *collecting* or *straight tubes* are lined with cubical epithelium, which in the larger tubes becomes distinctly columnar.

The *renal artery* enters the hilum between the pelvis and the renal vein. It shortly divides into four or five branches, which pass outward between the papillæ, and then enter the cortical portion between the pyramids. From these there pass to each Malpighian pyramid two branches, which ascend along its sides as far as its base, distributing in their course small vessels which pass to the Malpighian capsules. At the base of the pyramid they form arches, and make a bend from which two sets of branches are given off, the interlobular arteries and the arteriolæ rectæ.

The *interlobular arteries* pass off at right angles between the medullary rays, and then run among the convoluted tubes, some to enter the Malpighian capsules, and others to reach the surface and supply the capsule, ending in the stellate veins beneath the capsule. The arteriole which passes to the Malpighian capsule is termed the *afferent vessel*, and, entering the dilated extremity of the uriniferous tube, breaks up into a number of convoluted capillary vessels, constituting the *glomerulus* of Malpighi. The

[1] Heidenhain, *Archiv. f. mikr. Anatom.*, x, 1873.

FIG. 141.



a. Artery.
v. Vein, or efferent vessel.
c. Capsule.
d. Urinary tube.

blood is returned from the glomerulus by a small *efferent vein*, which emerges from the capsule close to the entrance of the artery. This vein, after a short course, breaks up, like an artery, into a dense network of capillaries, which ramify over the convoluted tubules. Some of the veins from the lower glomeruli break up into straight vessels, and then pass from the medullary rays into the pyramid (Fig. 141).

The *arteriolæ rectæ* are destined for the supply of the Malpighian pyramids, entering them at their bases, and then pass downward to their apices, where they join the venous plexuses.

The *Malpighian bodies* are small, red, granular masses, about $\frac{1}{170}$ of an inch in diameter, and are only found in the cortical substance. Each is composed of the dilated commencement of a uriniferous tube forming the *Malpighian capsule*, containing within it a coil of small blood vessels called the *Malpighian tuft* or *glomerulus*. The capsule is composed of a homogeneous membrane, and is pierced by a small artery, *afferent vessel*, which enters it opposite the commencement of the urinary tubule. In the capsule the artery breaks into a coil of minute blood vessels, *glomerulus*, and returns its blood by a vein, *efferent vessel*, which emerges from the capsule close to where the artery entered (Fig. 141). The capillary plexus within the capsule is surrounded by the epithelium lining the interior of the capsule.[1]

The *renal veins* return the blood from three sources; from the veins situated beneath the capsule and those corresponding to the interlobular arteries which pass between the medullary rays, and at the bases of the Malpighian pyramids join the venæ rectæ; the venæ rectæ return the blood from the arteriolæ rectæ, and be-g

[1] Histologists differ with respect to the disposition of the epithelium over glomerulus; some assert that it has no epithelial covering, but that it hangs within the capsule; some, that the tuft is completely invested with epithelium except where the afferent and efferent vessels pass in; others, again, that only portion of the glomerulus which looks toward the neck of the tubule is covered epithelium.

ı plexuses at the apices of the pyramids; they then pass outward etween the tubuli recti, and join the interlobular veins to form ıe proper renal veins; these pass down along the sides of the falpighian pyramids, accompanied by their corresponding arte-ies, and in their course to the sinus receive the efferent veins om the adjacent cortical substance. At the sinus they com-ıunicate freely with each other and join to form the renal vein.

The *nerves*, about fifteen in number, forming the renal plexus, re derived from the lesser and smallest splanchnic nerves, the olar plexus and the semilunar ganglion.

The *lymphatics*, consisting of a deep and a superficial set, pass o the lumbar glands.

Supra-renal Capsules. These bodies, situated at the top of the kidneys, belong to the class of ductless glands. They are f yellow-ochre color; the right is triangular, and resembles a ocked hat; the left is more almond shaped and rather the larger f the two. They measure about an inch and a half in their long iameter, about three-quarters in breadth, and two or three lines ı thickness; they weigh from one to two drachms. The gland ı surrounded by connective tissue and fat, and is invested by a hin fibrous covering, which sends down partitions into the inte-ior through furrows on the surface.

A perpendicular section shows that it consists of a firm exterior ır cortical part, and of an anterior or medullary substance, soft ınd pulpy.

The *cortical portion* is of a yellow color and forms the principal ıart of the organ. It consists of parallel columns arranged per-ɔendicularly to the surface, due to the capsule sending processes nto the interior of the gland, which communicate at frequent ntervals by transverse bands. There are thus formed numerous ıpaces which communicate with each other; the spaces at the ıurface are smaller, while those toward the centre are longer; the ıection through the cortex gives the appearance of a fine network, he external portion taking the name of the *zona glomerulosa*, the nternal portion that of the *zona reticularis*, the intermediate ɔortion that of the *zona fasciculata*. The reticular tissue is made ıp of fibrous tissue with longitudinal bundles of unstriped mus-ːular tissue. The interspaces are occupied by granular, polyhe-

dral cells, from $\frac{1}{2000}$ to $\frac{1}{1350}$ of an inch, which do not fill the spaces, so that between them and the walls of the spaces there are intervals, believed to be spaces which communicate with the lymphatics in the trabecular tissue.

The *medullary portion* varies in color, according to the amount of blood contained in it, being sometimes of a dark brown color, sometimes nearly white. Not infrequently the medullary part is converted into a cavity, but this is probably a post-mortem change. It consists of a plexus of minute veins, supported by the delicate areolar tissue containing muscular fibres, and presents a reticular aspect. Among these are numerous granular and branched cells.[1]

The *arteries* to the gland are conveyed along the trabecular tissue, and, after supplying the gland tissue, converge to the centre, where the blood is returned into the venous plexuses in the medullary portion. They are derived from the abdominal aorta, the phrenic and the renal arteries. The *vein* begins in the centre as a single vessel, and joins, on the right side, the inferior vena cava, and on the left side the left renal vein.

The *lymphatics* terminate in the lumbar glands.

The *nerves* are derived from the solar and renal plexuses, and in them are found numerous ganglia. They are distributed chiefly to the medullary portion.[2] Of late years the minute structure and functions of the supra-renal capsules have been much investigated, in consequence of the discovery, made by Dr. Addison, of the close relation which exists between certain diseases in these bodies and a brown discoloration of the skin. Their precise function is still unknown.

Stomach and Intestines.

The alimentary canal is composed of four coats; a serous, a muscular, a submucous, and a mucous. First, is the *serous* or *peritoneal* coat, described at p. 479. Secondly, under the serous is a *muscular* coat, upon which the chief strength of the canal depends. It consists of two distinct strata of plain muscular fibres; the outer stratum is longitudinal, the

[1] The medullary cells are stained a deep brown color on the addition of bichromate of potash, the cortical cells being scarcely affected by it.

[2] Luschka states that the branched granular cells of the medullary portion are connected with the nerve fibres.

inner circular. This arrangement not only makes the bowel stronger, but regulates its peristaltic action, for the longitudinal fibres, by their contraction, tend to shorten and straighten the tube, while the circular fibres contract upon and propel its contents to greater advantage. Connecting this coat and the mucous, is a layer of areolar tissue called the *submucous coat*, in which the arteries break up before entering the mucous membrane. The *mucous* is the most complicated of all the coats, for it presents different characters in different parts, according to the functions which it has to perform.

Stomach. The stomach should be moderately distended, to see its size, which varies in different subjects, according to the habits of the individual. When distended, an average stomach would be about ten or twelve inches in length, and four in depth and width; its weight is stated to be about four and a half ounces. It is conical in shape; the left part forms a large bulge, called the *bulge* or *splenic end;* and on the right side, where the food passes out, it becomes small and contracted, and is called the *pyloric end.* The stomach presents for examination two surfaces, two borders, two ends, and two orifices.

The *anterior surface* is convex, and looks upward and forward; the *posterior surface* looks downward and backward.

The *upper border* or the *lesser curvature* is concave and short, and extends from the œsophagus to the pylorus; it is connected to the liver by the gastro-hepatic omentum. The *lower border* or the *greater curvature* is convex, and affords attachment to the great omentum.

The *left end* is the larger, and is called the *cardiac* or *splenic end;* it bulges out to the extent of two or three inches to the left of the entrance of the œsophagus, and is called the *great cul-de-sac* or *fundus.* The *right end* is narrow, and makes a double bend; near the first it bulges into a pouch, called the *antrum pylori*, or the *small cul-de-sac.*

The *œsophageal* or *cardiac orifice*, situated at the highest part of the stomach, is on the left, and receives the œsophagus; the *pyloric orifice* is continued on into the duodenum, and is narrow, being guarded by a musculo-mucous ring, the *pylorus.*[1]

[1] The position of the stomach within the abdomen and its relations with surround-

The stomach is connected at its borders by peritoneal folds extending to neighboring structures; thus, its lesser curve is connected with the transverse fissure of the liver by the gastro-hepatic or lesser omentum; its cardiac end is connected with the hilum of the spleen by the gastro-splenic omentum; to the left of the œsophagus it is connected with the diaphragm by the gastro-phrenic ligament; to its greater curve is attached the great omentum, which is continuous on the left side with the gastro-splenic omentum.

The *pylorus* is the narrow, circular ring, composed of circular muscular fibres and mucous membrane, through which the food passes into the duodenum. The serous and longitudinal muscular fibres take no part in its formation, being continued over it on to the duodenum.

The stomach consists of four coats: serous, muscular, submucous and mucous.

The *serous* or *peritoneal coat* covers the surface of the stomach, except at the borders where the peritoneum is continued as omenta to other organs; it is along these borders that the vessels run.

The *muscular coat* can be seen when the serous coat is removed. The fibres are of the unstriped variety, and arranged in three layers: an external or longitudinal, a middle or circular, and an internal or oblique.

The *longitudinal fibres* are continuous with the longitudinal fibres of the œsophagus, and spread out over the stomach, being most numerous along the curvatures of the stomach; they are at the pyloric end continuous with the longitudinal fibres of the duodenum.

The *circular fibres* are well marked about the middle of the stomach, but are most abundant at the pylorus, where they form a powerful sphincter.

The *oblique fibres* are scattered over the sides of the stomach and are most distinct at the entrance of the œsophagus, crossing

ing structures are matters of much dispute. Dr. Lesshaft has come to the conclusion that the stomach is nearly vertical, so that its fundus touches the diaphragm. (*Lancet*, March 11th, 1882, p. 406.) His, and most anatomists, are of opinion that the long axis is placed obliquely from left to right within the abdomen.

bliquely from left to right and from right to left. They are ontinuous with the well marked circular fibres of the œsophagus.

The *submucous coat* serves to connect the muscular with the ıucous coats. It consists of areolar tissue, and permits the ıuscular and mucous coats to move freely on each other, and llows the blood vessels to ramify minutely before they enter the ıucous membrane.

When the stomach is laid open from the œsophageal to the yloric orifice, the *mucous membrane* is seen to be thick, of pale ink or straw color, and is gathered into longitudinal folds—*rugæ* —which disappear when the stomach is distended.

If a portion of the mucous membrane be examined under the ıicroscope, its surface will be seen to be mapped out into small exagonal pits or *alveoli*, surrounded by ridges, giving it a honey-ombed appearance. The pits vary from $\frac{1}{100}$ to $\frac{1}{150}$ of an inch in iameter. At the bottom of them are a number of minute aper-ıres, the orifices of the *gastric follicles*. In a perpendicular ection the follicles are arranged in parallel lines at right angles ɔ the surface, and terminate in blind sacculated ends set in the ubmucous tissue. The entire thickness of the mucous mem-ɪrane is made up of these tubular glands. The follicles consist ɪf two kinds, *mucous* and *peptic glands*. Tubular in shape, they ıave a basement membrane lined with epithelium, and average bout $\frac{1}{10}$ of an inch long, and $\frac{1}{150}$ of an inch in diameter. The *ıucous glands* are found over the whole surface, but are most ıumerous at the pyloric end of the stomach. They are composed ɪf tubes, each consisting of two or three short tubules, opening ıto a common duct, which itself opens into the bottom of an lveolus. They are lined with columnar epithelium continuous ɪith that lining the mucous membrane. The *peptic glands* are lso found over the entire surface of the mucous membrane, and onsist of tubules with branched cæcal extremities opening into . common duct, which is, however, shorter than that of a mucous ;land. They are lined with columnar epithelial cells, called the *ɛntral cells*, and are supposed to be concerned in the secretion of he gastric juice; these cells become, at the neck of the gland, nuch shorter and more granular. The lumen of the gland is ɪery small, but is somewhat larger at the free and cæcal ends

than in the middle. Toward the lower part or fundus of the gland there are found spheroidal and granular cells between the epithelium and the basement membrane, called *parietal cells.*

In the stomachs of young children there is a large amount of lymphoid tissue found between the gastric glands. It is aggregated into small masses in the mucous membrane, and resembles in many respects the solitary glands of the intestine, although not so well defined.

The mucous membrane of the stomach is lined by columnar epithelium, which also extends into the glands. A thin layer of unstriped muscular tissue (muscularis mucosæ) is found between the mucous membrane and its submucous tissue, varying in amount and in the number of its layers.

The glands of the stomach are richly supplied with blood, which is derived from the gastric, the vasa brevia, the right and left gastro-epiploica, and the pyloric arteries. The arteries form a stratum of minute inosculations in the submucous tissue, in which the closed ends of the tubes are set; from this stratum the capillary plexuses run up between the tubes to the surface of the stomach, where they again form a larger capillary plexus and form the hexagonal spaces before alluded to. The *veins*, corresponding to the pyloric and gastric arteries, end in the vena portæ; those corresponding with the vasa brevia and the epiploic arteries open into the splenic vein.

The *lymphatics* enter the glands along the lesser and great curvatures of the stomach, and may be divided into a superficial and a deep set.

The *nerves* are derived from the pneumogastric nerves and from the solar plexus.

Small Intestine. The small intestine, consisting of the duodenum, jejunum and ileum, forms a tube averaging twenty feet in length, which gradually lessens in calibre until it opens into the cæcum. The duodenum is about twelve fingers' breadth in length (whence its name); the jejunum[1] comprises two-fifths, the ileum[2] three-fifths, of the remaining part of the small intestine. As regards their external characters, the duodenum and jejunum are more vascular than the ileum, and feel thicker, in consequence

[1] From *jejunus*, empty. [2] From εἱλειν, to roll or twist.

of the peculiar arrangement of their mucous membrane; but there are no defined limits between the different portions of the intestinal canal.

Its coats are four in number: serous, muscular, submucous and mucous.

The *serous coat* consists of peritoneum, and forms a complete investment, except in the case of the descending and transverse portions of the duodenum, which are only covered in front.

The *muscular coat* consists of an outer longitudinal layer and an inner, circular, thicker layer, which, however, becomes thinner toward the end of the ileum.

The *submucous coat* connects the muscular and mucous coats; immediately beneath the mucous membrane there is a very thin layer of non-striped muscular fibres, termed *muscularis mucosæ*.

The *mucous coat* can only be seen when the intestine is cut open from the upper end, and is composed of the following strata: the muscularis mucosæ, a layer of retiform tissue with lymph corpuscles, with blood vessels and nerves; and lastly, a layer of columnar epithelium.

When the intestine is laid open we see that the mucous membrane is arranged in close transverse folds, called *valvulæ conniventes* or *valves of Kerkring*. These differ from other folds in the alimentary canal—*e. g.* in the œsophagus and stomach—in that they are not obliterated when the tube is distended. Each fold extends about one-half or two-thirds round the intestine, but they are not all of equal size, and are about one-third of an inch in depth. They commence immediately below the opening of the biliary and pancreatic ducts, and are most developed in the duodenum and the upper part of the jejunum. Below this part of the tube they gradually decrease in size, and become wider apart, till they finally disappear near the middle of the ileum. The use of the valvulæ conniventes is to increase the extent of surface for the absorption of chyle; to prevent the food passing too rapidly through the intestine, and for secretion.

If a portion of small intestine be washed and placed in water, the surface of the mucous membrane appears like the soft fur or pile upon velvet. This appearance is produced by small processes called *villi*. These are extremely vascular projections of the



g,
the
cting
umber
he sub-
hoid tis-
ontain an
traced from
neir contents
les are found
ches is covered
rated in typhoid
old age.
rts of the small and
ymphoid structure as
them in being solitary

ose of the muscular and
ceive those from the villi,
te plexus, and after pierc-
former, which are chiefly
the circular layers of mus-

ne superior mesenteric plexus,
senteric artery and its branches,
tery; after reaching the intestinal
ate from the arteries. They then
l muscular fibres, and form a very
uerbach's plexus or *plexus mesenteri-*
ents to the muscular layer of the

m, 1682. These glands were first described by

mucous membrane, about a fourth of a line in length, and are so numerous that a square line contains from forty to ninety of them.[1] Their size, however, and their number, bear a direct ratio to those of the valvulæ conniventes. Under the microscope a villus is seen to consist of an outstanding process of the mucous membrane, covered by a layer of columnar epithelium, which rests upon a basement membrane. Each villus is furnished with an artery, which forms a network of inosculations in it, and then returns its blood by a single vein. Down its middle runs a *lacteal* or absorbing vessel, which commences in a closed end near the summit of the villus, where it is surrounded by a layer of pale, non-striped muscular fibres proceeding from the *muscularis mucosæ*. This is surrounded by a plexus of capillaries, external to which is the basement membrane supporting columnar epithelium. Forming the matrix of the villus is a fine network enclosing large, flattened cells, with oval nuclei and lymph cells.

Intestinal Glands. There are four kinds of *glands*[2] in the small intestine, called the glands of Lieberkühn, Brunner, Peyer and the solitary glands. The first and last are distributed over the whole tract of the intestinal mucous membrane; the other two over particular parts.

The *simple follicles* or *crypts of Lieberkühn*,[3] the most numerous of all, are minute tubes with blind ends, very thickly distributed over the small and large intestines. Under the microscope their orifices are seen between the villi, like so many minute dots. They vary in depth from $\frac{1}{10}$ to $\frac{1}{10}$ of a line, and about $\frac{1}{50}$ of a line in diameter; their walls consist of a delicate basement or endothelial membrane, and are lined with columnar epithelium.

The *duodenal* or *glands of Brunner*[4] are found only in the duodenum and a small part of the beginning of the jejunum. They are just visible to the naked eye, and may be seen by removing

[1] Krause estimates the total number of villi at four millions.

[2] A satisfactory examination of the intestinal glands can be made only in specimens quite recent, taken from young persons who have died suddenly, or from rapidly fatal disease.

[3] J. N. Lieberkühn, *Diss. de Fabric. et Actione Villorum Intestin. ten.*, 1782.

[4] J. C. Brunner, *Gland. Duoden. seu Pancreas Secundarium*, 1715.

the muscular coat. Their structure exactly resembles the round compound glands of the mucous membrane of the mouth.

The *glands of Peyer*[1] (*glandulæ agminatæ*) abound most in the ileum, and are seen most distinctly in children. They are arranged in groups, from twenty to forty in number, on that part of the intestine most distant from the attachment of the mesentery. These groups are from half an inch to three inches long, of an oval form, and increase in size and number toward the lower part of the ileum. If a group be examined by dissecting away the muscular coat, you find that it is composed of a number of small oval vesicles, like Florence flasks, imbedded in the submucous tissue. They are composed of masses of lymphoid tissue, of about three-fourths of a line in diameter, and contain an opaque, grayish fluid. No excretory ducts have been traced from these vesicles, but they are supposed to discharge their contents by rupture of their capsules. Between the vesicles are found Lieberkühn's follicles; and the surface of the patches is covered with villi. These glands are liable to be ulcerated in typhoid fever. They diminish in number and size with old age.

The *solitary glands* are scattered over all parts of the small and large intestines. They consist of the same lymphoid structure as the glands of Peyer, and only differ from them in being solitary instead of being aggregated into groups.

The *lymphatics* consist of two sets—those of the muscular and those of the mucous coats; the latter receive those from the villi, at the base of which they form a minute plexus, and after piercing the muscular coat, join with the former, which are chiefly found between the longitudinal and the circular layers of muscular fibres.

The *nerves* are derived from the superior mesenteric plexus, and accompany the superior mesenteric artery and its branches, between the layers of the mesentery; after reaching the intestinal walls the nerve filaments separate from the arteries. They then pierce the external longitudinal muscular fibres, and form a very minute gangliated plexus—*Auerbach's plexus* or *plexus mesentericus*—which distributes filaments to the muscular layer of the

[1] Peyer, *De Glandulis Intestinorum*, 1682. These glands were first described by Nehemiah Grew, in 1681.

entire intestinal canal. From this plexus numerous branches perforate the internal circular muscular layer, and unite to form a largely gangliated plexus—*Meissner's plexus*—in the submucous tissue. The *intermuscular plexus* probably supplies the muscular coat and regulates the peristaltic action of the bowel; the *submucous plexus* determines the calibre of the blood vessels.

Large Intestine. The principal external characters of the large intestine are that it is pouched or sacculated, and that it has attached to it little pendulous portions of fat covered by peritoneum, called *appendices epiploicæ*. The pouches (sacculi) are produced by a shortening of the longitudinal muscular fibres, and by their being collected into three bands, about half an inch wide, nearly equidistant from each other. One of these bands corresponds with the attached part of the circumference of the bowel; another with the front part; a third with its concavity. If at any given part the three bands be divided, the pouches immediately disappear.

In a colon moderately distended and dried, we observe that the mucous membrane forms numerous ridges or incomplete septa (see Fig. 142); they correspond to the grooves on the external surface of the bowel, and disappear, like the sacculi, when the longitudinal bands are divided.

The *rectum* differs from the rest of the large intestine in that its longitudinal muscular fibres are not collected into bands, but distributed equally over its whole circumference. Moreover, both the longitudinal and circular fibres are of considerable strength, like those of the œsophagus, as one might expect from the particular functions which these parts of the alimentary canal have to perform. For an inch and a half, or thereabouts, above the anus, the circular fibres are remarkably developed, and constitute the *internal sphincter ani*.

The mucous membrane of the large intestine differs considerably from that of the small. There are neither valvulæ conniventes, nor villi, but the *glands of Lieberkühn* and the *lymphoid follicles* may be seen studding the mucous membrane. The follicles are more abundant in the cæcum and in the appendix vermiformis than in any other part of the alimentary canal. The blood vessels present the same hexagonal arrangement on the surface

as that of the stomach. That the mucous membrane of the large intestine may be temporarily used as a substitute for the stomach is proved by the fact of persons having been nourished for many weeks solely by injections. The mucous membrane is lined throughout with columnar epithelium.

Ileo-cæcal Valve. At the junction of the small with the large intestine the mucous membrane is folded so as to form a valve; but it is not a perfect one, as is proved by pouring water into the large intestine, or by the occasional vomiting of

FIG. 142.

1. Ileum.
2. Cæcum or caput coli.

3. Appendix vermiformis.

SECTION THROUGH THE JUNCTION OF THE LARGE AND SMALL INTESTINE, TO SHOW THE ILEO-CÆCAL VALVE.

injections. The arrangement of the valve is best examined in a dried preparation. The opening is a transverse fissure like a button hole; and the two flaps are arranged like an upper and a lower eyelid. The upper lid of the valve projects more than the lower, so that the contents of the ileum drop naturally down into the caput coli, where they are apt to collect and form hard lumps. The flaps of the valve consist of mucous membrane and the circular fibres of the ileum. The longitudinal fibres of the ileum are continued directly on to the cæcum;

if these be divided, the ileum can be drawn out, and the valve disappears.[1]

Folds in the Rectum. In many subjects we observe that transverse or oblique folds of the mucous membrane project into the rectum. These cannot be seen to advantage unless the bowel be hardened by alcohol in its natural position. Three, more prominent than the rest, and half an inch, or thereabouts, in width, were first pointed out by Mr. Houston.[2] One projects from the upper part of the rectum, opposite the prostate gland; another is situated higher up, on the side of the bowel; while the third is still higher. When thickened or ulcerated, these folds are apt to occasion great pain and obstruction in defecation.

Arterial Supply of the Alimentary Canal. The present opportunity is the best for reviewing the arterial supply of, and the anastomoses round, the alimentary canal, from *the mouth to the anus*. Part of the blood supply has been examined in the dissection of the head and neck; part in the dissection of the œsophagus as it passes through the thorax; and the remainder in that of the abdomen. The following table represents the arteries in their order, beginning at the mouth:—

LOWER LIP	Submental (deep branch).
	Mental.
	Inferior labial.
	Inferior coronary.
UPPER LIP	Superior coronary.
CHEEK	Buccal.
	Superior coronary (slightly).
	Facial.
	Transverse facial.
	Infra-orbital.
	Superior alveolar.
MOUTH, ROOF OF	Descending palatine.
	Ascending palatine.
	Pharyngeal br. of ascending pharyngeal.
MOUTH, FLOOR OF, AND TONGUE	Artery of the frænum.
	Ranine.

[1] It is interesting to note that the surface of the valve, toward the ileum, is covered with villi; not so the surface toward the large intestine.

[2] *Dublin Hospital Reports*, vol. v, p. 163.

I, FLOOR OF, AND TONGUE	Sublingual. Dorsales linguæ. Tonsillar. Ascending palatine. Ascending pharyngeal.
OTTIS	Superior laryngeal artery.
'NX	Pterygo-palatine. Branches of ascending pharyngeal. Branches of ascending palatine. Superior thyroid.
HAGUS, CERVICAL . . .	Superior thyroid. Inferior thyroid.
HAGUS, THORACIC . . .	Inferior thyroid. Thoracic aorta. Gastric. Left phrenic.
HAGUS, ABDOMINAL . .	Gastric. Left phrenic.
ACH	Gastric. Pyloric. Gastro-epiploica dextra. Gastro-epiploica sinistra. Vasa brevia. Gastro-duodenalis.
ENUM	Pancreatico-duodenalis superior. Pancreatico-duodenalis inferior.
M	Superior mesenteric.
. . . . .	Superior mesenteric.
M	Colic br. of ileo-colic.
DING COLON	Colica dextra.
SVERSE COLON	Colica media.
NDING COLON	Colica sinistra.
ID FLEXURE	Sigmoid arteries.
M	Superior hæmorrhoidal (inferior mesenteric). Middle hæmorrhoidal (internal iliac). Inferior hæmorrhoidal (internal pudic). Arteria sacra media.

DISSECTION OF THE LOWER EXTREMITY.

The body must be placed on its back, with a block placed beneath the buttocks, and the thigh should then be slightly flexed and abducted.

Surface Marking. The student, before commencing to reflect the skin, should notice the irregularities of the surface which are produced by subjacent structures. The upper part of the thigh is marked off from the abdomen by a more or less well-marked curved furrow, having its convexity downward. This furrow corresponds with Poupart's ligament, which is attached externally to the anterior superior iliac spine, and internally to the spine of the os pubis. The spine of the os pubis can, even in the fattest subject, be distinctly felt, and is a very valuable landmark in the diagnosis between an inguinal and a femoral hernia; for the aperture through which an inguinal hernia emerges is the external abdominal ring, situated *above* the spine; the aperture through which a femoral hernia comes out is the saphenous opening, situated *outside* the spine. In front of the thigh is a large triangular depression corresponding with Scarpa's triangle, which has its base at Poupart's ligament. This depression, which is best seen in thin subjects, contains the large vessels and nerves passing down to the leg, the femoral artery being nearly in the centre of the space; a furrow indicating the course of these vessels may be observed extending obliquely down the inner side of the thigh. About three or four inches below the anterior superior iliac spine, there is seen on the outer side of the thigh the well-marked prominence of the great trochanter, which is nearly on the same level as the spine of the os pubis. The sartorius can be seen passing obliquely inward from the iliac spine, and crossing over the femoral vessels about four inches below Poupart's ligament; in the latter two-thirds of its course it descends nearly vertically. The well-defined ridge, extending from the os pubis to the middle of the femur, when the thigh is abducted, is caused by the adductor longus muscle.

The prominence in front of the knee is produced by the patella, to which is attached, above, the tendon of the quadriceps muscle, and below, the ligamentum patellæ, both of which can be distinctly felt. On each side of the patella is a deep depression, which leads

on the outer side to a rounded prominence, the external condyle; and on the inner to the internal condyle, the latter being the larger. The synovial membrane which lines the knee joint usually extends about two fingers' breadth above the patella, and is a little higher on the inner than on the outer side of the joint.

An incision should be made along the groin, extending from the anterior superior spine of the ilium to the spine of the os pubis; another, from the middle of the first down the front of the thigh for about six inches. The skin only should be reflected, outward and inward, when the superficial fascia will be exposed.

Superficial Fascia. The *superficial fascia* varies in thickness, according to the condition of the body. Like other superficial fasciæ, it is divisible into two or more layers, between which are situated the inguinal glands and the cutaneous vessels and nerves. The superficial layer is continuous with that of the abdomen, and becomes firmer below Poupart's ligament, to which, however, it is not connected; the deeper layer is best marked in the upper part of the thigh, especially where it stretches across the saphenous opening, to the margins of which it is closely attached; this portion is called the *cribriform fascia*, and is protruded forward by a femoral hernia, forming one of its coverings; this layer is also attached to Poupart's ligament.

The superficial layer of this fascia should now be reflected, by searching for one of the subcutaneous veins (the internal saphena will do) which run between the upper and the deeper layers of the fascia. The cutaneous vessels can thus be traced, and come from the common femoral artery; they are three in number, the *superficial epigastric*, the *superficial external pudic* and the *superficial circumflexa ilii* arteries. The first ascends over Poupart's ligament to the abdomen (p. 628); the second crosses inward toward the os pubis; the third passes outward to the ilium. Each artery is accompanied by one, sometimes by two veins, which empty themselves, either directly into the femoral, or into the great cutaneous vein of the thigh, called the saphena.

Superficial Inguinal Glands. These glands are easily recognized, by their oval form and reddish-brown color. There are two sets: one set runs parallel to Poupart's ligament, and receives the lymphatics from the skin of the penis, the scrotum, the perin-

eum, the anus, the buttock, the lower part of the abdominal wall, and the upper and outer aspect of the thigh; the outer and lower set lies along the saphena vein, chiefly around the saphenous opening, and receives the lymphatics from the foot, the leg, and the lower part of the thigh. This explains why in cancer of the scrotum and syphilitic disease of the penis the first set becomes enlarged; and the second, in diseases of the lower extremity. The lymphatic vessels which pass to and from the glands are small, and may escape observation, unless specially looked for. They all pass through the femoral ring into the abdomen, and eventually empty themselves into the receptaculum chyli.

The glands mentioned in the preceding paragraph are all superficial. There are others, more deeply seated, close to the great vessels of the thigh; these are much smaller, and sometimes cannot be found.

Superficial Arteries of the Groin.

The *superficial epigastric* artery comes through the fascia lata, sometimes through the saphenous opening, half an inch below Poupart's ligament. It ascends over Poupart's ligament to pass to the subcutaneous tissue of the abdomen, as high as the umbilicus, and supplies the inguinal glands, and anastomoses with the deep epigastric and internal mammary arteries. Its further course is described at p. 443.

The *superficial circumflexa ilii* emerges through the fascia lata, runs parallel to Poupart's ligament toward the crest of the ilium, and ends in the subcutaneous tissue and inguinal glands. It anastomoses with the deep circumflex iliac, the gluteal, and the ascending branches of the external circumflex arteries.

The *superficial external pudic* comes through the saphenous opening, crosses over the spermatic cord, and supplies the penis and scrotum in the male, and the labium in the female. This artery is usually divided in the operation for femoral hernia; also in that for phimosis, since it runs along the penis to supply the prepuce. Arising directly from so large an artery as the femoral, it sometimes bleeds profusely; for it is an admitted fact that when even a small branch, coming directly from a principal artery, is divided near its origin, it will sometimes pour out as much blood as if an opening were punched out of the trunk as large as the

:a of the divided branch.[1] There is another pudic artery, called : *deep* or *inferior external pudic;* this runs between the fascia a and the pectineus, supplying that muscle, the scrotum in the ıle, and the labium in the female. They both anastomose with anches of the internal pudic artery.

The incision should be prolonged down the thigh, over the ıee, to the tubercle of the tibia. The skin must then be

FIG. 143.



SUPERFICIAL VESSELS AND GLANDS OF THE GROIN. SAPHENOUS OPENING WITH THE CRIBRIFORM FASCIA.

1. Saphenous opening of the fascia lata.
2. Saphena vein.
3. Superficial epigastric artery.
4. Superficial circumflexa ilii artery.
5. Superficial pudic artery.
6. External abdominal ring.
7. Fascia lata of the thigh.

flected, to expose the subcutaneous tissue over the whole of ıe front of the thigh. The cutaneous vessels and nerves should

[1] Mr. Liston had occasion to tie the external iliac artery for a supposed injury (by pistol ball) to the femoral. It was discovered, after the death of the patient, that e ball had injured only one of the superficial branches of the femoral, about an ch from its origin. See his paper in the *Med.-Chir. Trans.*, vol. xxix, 1846.

be looked for in the subcutaneous fat in the following situations: on the *inner side* are the inguinal branch of the ilio-inguinal nerve passing down through the external abdominal ring, internal to the saphenous opening; *lower down* are the two branches of the internal cutaneous nerve supplying the skin on the inner aspect of the thigh as far as the knee, the lower branch accompanying the internal saphena vein, which ascends to pierce the saphenous opening; there are also low down some filaments from the long saphena nerve; on the *front* of the thigh there is found the crural branch of the genito-crural nerve, and lower down, as far as the knee, are the middle cutaneous nerves; on the *outer side* are seen filaments of the external cutaneous nerve.

Internal Saphena Vein. This is the chief subcutaneous vein of the lower limb. Its roots, arising from the inner side of a venous arch on the dorsum of the foot, unite into a single trunk, which ascends in front of the inner ankle, along the inner side of the leg, behind the knee, along the inner and front part of the thigh, where it passes through an opening—the *saphenous opening*—in the fascia lata, to join the femoral vein, immediately below the crural arch (Fig. 143). In this long course it receives many tributary veins, some of which are often large, especially one which, coursing round the inner part of the thigh, is frequently as large as the main trunk. Just before its termination it is joined by the superficial veins, which accompany the arteries of the groin, already alluded to, p. 627. Like all subcutaneous veins, it is provided with valves, chiefly where joined by other veins, to support the column of the blood.

Cutaneous Nerves. The distribution of the cutaneous nerves of the thigh varies considerably, but they are always found more abundantly on the inner than on the outer aspect of the thigh. The nerves are divided into *external*, *middle* and *internal*. All directly or indirectly proceed from the lumbar plexus, and, perforating the fascia lata, divide in the subcutaneous tissue.

a. The *external cutaneous nerve* is a branch of the second and third lumbar nerves. It enters the thigh beneath Poupart's ligament close to the anterior superior spine of the ilium. Here it divides into two branches, an anterior and a posterior. The

nterior branch comes through the fascia lata about four inches elow Poupart's ligament, and can be traced down the outer side f the thigh as far as the knee, giving off numerous branches. he *posterior* branch, after coming through the fascia lata, ivides into filaments, which are distributed to the skin over the ates and the posterior part of the thigh.

b. The *middle cutaneous nerves*, one or two in number, are iven off by the anterior crural in the thigh. They pass through ne sartorius about four inches below Poupart's ligament, erforate the fascia lata, and descend along the front and inner art of the thigh as far as the knee, distributing branches on ither side; some of which communicate with the long saphenous erve. In its course along the front of the thigh it joins with he crural branch of the genito-crural and the internal cutaneous erves.

c. The *internal cutaneous nerve*, also a branch of the anterior rural, crosses obliquely over the sheath of the femoral artery. It hen divides into two branches, an anterior and an internal; the *nterior branch* comes through the fascia lata in the lower third of he thigh, where it terminates in two branches, one being distri- uted to the inner side of the knee, the other crossing over the atella to the outer side of the joint; the *internal branch* perforates he fascia lata just above the knee joint, after running down along he posterior border of the sartorius, and supplies the integument n the inner side of the leg. While still beneath the fascia lata, he internal cutaneous nerve unites below the adductor longus, in plexiform manner, with the long saphenous and obturator nerves.[1]

d. The *crural branch* of the *genito-crural nerve* perforates the nterior layer of the sheath of the femoral vessels, comes through he fascia lata immediately below Poupart's ligament, and supplies he skin in front of the thigh. About two or three inches below he crural arch it usually communicates with the middle cutane- ous nerve. It also distributes a few filaments to the femoral rtery in its passage under the crural arch.

[1] It is important to note that one, sometimes two, of these branches of the inter- al cutaneous crosses the sheath of the femoral artery, just where the sartorius be- ins to overlap it, and therefore at the spot where it is usually tied. See diagram . 644.

c. The *inguinal branch* of the *ilio-inguinal nerve*, after emerging from the external abdominal ring, supplies the skin on the inner aspect of the upper third of the thigh.

Fascia Lata.

Remove the subcutaneous fat and the deeper layer of the superficial fascia, to examine the dense white fascia—the *fascia lata*—of the thigh. The use of this fascia is to cover the muscles of the thigh collectively, and to form separate sheaths for each; so that it not only keeps them together, but maintains each in its proper position. A knowledge of these sheaths is important, because they interfere with the progress of deep-seated matter toward the surface, and cause it to burrow in this or that direction, according to the part in which it forms.

The fascia is not of equal strength all round the thigh. It is comparatively thin on the inner side; exceedingly thick and strong down the outer side; here, indeed, it has the appearance of a dense expanded aponeurosis, strapping down the vastus externus muscle, and is sometimes called the *ilio-tibial band;* and it certainly performs the office of a tendon, for it gives insertion between its two layers to two powerful muscles—namely, the tensor fasciæ femoris, and the gluteus maximus (Fig. 144).

The fascia lata is attached to the margin of the bones which constitute the framework of the lower extremity. Beginning from above, its attachment can be traced from the posterior surface of the sacrum and coccyx, along the crest of the ilium, thence along Poupart's ligament to the body of the os pubis and the linea ilio-pectinea, and along the rami of the os pubis and ischium. Proceeding down the thigh, it penetrates on each side of the limb to the linea aspera, forming what are called the *external* and *internal intermuscular septa;* the external one, stronger, separates the vastus externus anteriorly from the short head of the biceps, both of which have origin from the fascia; the internal one separates the vastus internus in front from the adductor muscles behind. Below, it can be traced round the knee joint, and is particularly strong, especially on the outer side, where it is attached to the head of the tibia and fibula, and forms the insertion of the tensor fasciæ femoris. The fascia lata is very strong over the gluteus medius—the *gluteal aponeurosis*, and at

Fig. 144.

Fascia on the outside of the thigh.

1. Tensor fasciæ femoris.
2. Gluteus maximus.
3. Lower fibres of ditto.
4. Fascia lata.

pper border of the gluteus maximus divides ito two layers, one superficial to the muscle, ie other deep, which separates this muscle from he deeper muscles, and becomes connected with ie great sacro-sciatic ligament. The fascia lata lso furnishes thinner sheaths for the separate uscles.

There are numerous small apertures in the ascia, through which the cutaneous nerves and essels are transmitted; but the most important ne is the large opening—the *saphenous opening* —through which the saphena vein passes to join he femoral. The part of the fascia situated exernal to the saphenous opening is termed the liac portion of the fascia lata; that internal to t, the pubic portion.

The *iliac portion* is attached to the crest of he ilium, to the whole length of Poupart's ligament, and, in conjunction with Gimbernat's ligament, to the linea ilio-pectinea; from this ttachment it arches downward and outward, s inner margin forming the outer falciform dge of the saphenous opening; this border asses over the anterior sheath (formed by he transversalis fascia) of the femoral artery, nd is seen to be continuous below with the *ubic portion*, which can be traced upward over the pectineus and dductor longus muscles, behind the posterior sheath (formed by he iliac fascia) of the femoral vessels, where it is connected with he sheath of the iliacus and psoas muscles and the fibrous tructures of the hip joint. Above, it is attached to the linea lio-pectinea, to the body and the ramus of the os pubis.

Saphenous Opening in the Fascia Lata.

The *saphenous opening* is an oval aperture in the fascia lata, immediately below the crural arch, on the inner side of the front of the thigh, through which the saphena vein passes to join the femoral. There is no definite border to the saphenous opening until the fascia—*cribriform*, which covers the opening and blends with its margin—has been

removed. It is situated just below the crural arch and external to the spine of the os pubis; it is oval, with the long axis vertical, and is about one inch and a half long and an inch broad. Its border on the inner side is not defined; for here the fascia lata ascends under the femoral vessels, and is continuous with the iliac fascia of the pelvis.[1] But the outer or iliac border is clearly defined. This lies in front of the femoral vessels, is crescent-shaped, with the concave upper end toward the os pubis, and is called the *falciform process*, while its deeper fibres are known as *Burns' ligament*. The lower horn of the crescent curves under the saphena vein with a well-defined border, and on being traced

FIG. 145.

1. Crural arch.
2. Saphenous opening of the fascia lata.
3. Saphena vein.
4. Femoral vein.



5. Gimbernat's ligament.
6. External abdominal ring.
7. Position of the internal ring in dotted outline.

DIAGRAM OF THE FEMORAL RING AND THE SAPHENOUS OPENING.
(The arrow is introduced into the femoral ring.)

upward becomes less well marked until it is gradually lost in the fascia on the inner side of the opening. The upper horn, *Hey's ligament*,[2] arches over the femoral vein, and then descending slightly is continued uninterruptedly into Gimbernat's ligament—*i. e.* into that part of the crural arch which is inserted into the linea ilio-pectinea. The upper horn deserves especial attention, because it forms the upper boundary of the aperture through which a femoral hernia takes place; and, being chiefly concerned

[1] On the inner side of the femoral vessels the pubic portion of the fascia is attached to the linea ilio-pectinea.

[2] This upper horn is sometimes called *Hey's ligament*, after the surgeon who first drew attention to it. (*Observations in Surgery*, by W. Hey, F.R.S., London, 1810.)

the constriction of the rupture, must be divided for its relief. is may be easily ascertained by introducing the little finger der the crural arch, on the inner side of the femoral vein—in ıer words, into the femoral ring (see the arrow in the diagram). el how the upper horn of the crescent would gird the neck of ernia, and that its tension is greatly influenced by the position he limb; for if the thigh be bent and brought over to the other e, the tension of all the parts is materially lessened.[1]

briform Fascia. The *cribriform fascia* is so called because it is perforated with numerous apertures for the passage the superficial vessels and lymphatics. It is a thin membranous vering over the saphenous opening, and is prolonged from the ter edge of the opening over the sheath of the femoral vessels, l adheres on the inner side to the fascia lata, over the pectineus ıscle. Some anatomists describe this fascia as a portion of the per layer of the superficial fascia; others consider it as a thin longation of the fascia lata itself across the opening. Its chief gical importance is derived from the fact that it forms one of coverings of a femoral hernia.

The cribriform fascia must now be removed on one side, so as to play the saphenous opening, which will appear as represented Fig. 145.

NATOMY OF THE PARTS CONCERNED IN FEMORAL HERNIA.

The anatomy of the parts concerned in femoral hernia cannot thoroughly understood without the assistance of special dissec- ns. The following demonstration, therefore, takes for granted t the student has the opportunity of seeing the parts, not only their femoral, but also on their abdominal side.

The different parts of the subject should be examined in the lowing order:—

1. The formation of the crural arch.
2. The arrangement of the parts as they pass under the arch.

[1] We must always bear in mind that, though the crural arch and the fascia attached have received particular names, they are not, on that account, distinct and sepa- ; but all are intimately connected, and portions merely of one continuous ınsion. Thus all the parts are kept in a condition of mutual tension, which de- ds very much on the position of the thigh.

c. The sheath of the femoral vessels.

d. The crural canal and ring.

e. The practical application of the subject.

Poupart's Ligament or Crural Arch.

The lower border of the aponeurosis of the external oblique muscle extends from the anterior superior spine of the ilium to the spine of the os pubis, and forms over the bony excavation beneath the *crural arch* or *Poupart's ligament.* (It is marked by the dark line in Fig. 145.) The direction of the arch is at first somewhat oblique, but toward its inner half becomes nearly horizontal. In consequence of its intimate connection with the fascia lata of the thigh, the line of the arch describes a gentle curve with the convexity downward. The arch is attached to the spine of the os pubis, and also for some distance along the linea ilio-pectinea (Fig. 145). This additional attachment, called *Gimbernat's ligament,*[1] is of importance, for it is frequently the seat of stricture in femoral hernia.

Gimbernat's Ligament.

The best view of Gimbernat's ligament is obtained from within the abdomen; it being only necessary to remove the peritoneum. It is that portion of the aponeurosis of the external oblique muscle which is inserted into the linea ilio-pectinea for about an inch in length. It is placed nearly horizontally in the erect posture, and is triangular with its apex at the os pubis and its base directed outward. In front, it is continuous with the crural arch; behind, it is inserted into the linea ilio-pectinea; externally, it is continuous with the fascia lata through Hey's ligament (Fig. 145). Its length is from three-quarters of an inch to one inch; but it is usually longer in the male than in the female.

On putting your finger into the femoral ring, you feel the sharp and wiry edge of this ligament; observe, too, that as the body lies on the table, the plane of the ligament is perpendicular, and therefore that it *recedes from the surface.*

An incision should now be made through the fascia lata along the entire length of Poupart's ligament; another also through the fascia vertically, from the anterior superior iliac spine down the

[1] Don Antonio de Gimbernat was a Spanish surgeon, who published, in 1793, *A New Method of Operating for the Femoral Hernia.* Madrid.

thigh for about four inches; and the fascia lata carefully dissected downward and inward from the subjacent structures. This will expose the structures as they pass under Poupart's ligament in their course down the thigh.

Arrangement of the Parts which pass under the Arch.

The crural arch transmits from the abdomen into the thigh (proceeding in order from the outer side) the following objects shown in Fig. 146: 1. The external cutaneous nerve. 2. The iliacus with the anterior crural nerve lying on it near its inner border. 3. The femoral artery resting on the psoas muscle. 4. The crural branch

FIG. 146.

POSITION OF PARTS UNDER THE CRURAL ARCH (VERTICAL SECTION).

of the genito-crural nerve. 5. The femoral vein. 6. The crural sheath surrounding the femoral vessels, formed in front by the fascia transversalis, behind by the fascia iliaca. 7. The lymphatics passing upward through the femoral canal. 8. The pectineus. These muscles and vessels fill up the space beneath the crural arch, except on the inner side of the femoral vein, where a space is left for the passage of the lymphatics; this is called the *crural* or *femoral ring*. The muscles are separated from the vessels by a strong, vertical, fibrous partition passing from the arch to the bone, which is nothing more than a continuation of the sheath of the psoas. The artery, too,

is separated from the vein by a similar, although a much weaker partition, and there is a third close to the inner side of the vein. These three partitions not only keep all the parts in their right place, but confine the arch down to the bone, and prevent its being uplifted by any protrusion between it and the muscles and vessels. This, coupled with the close attachment of the fascia iliaca to the crural arch, explains why a femoral hernia rarely takes place in any other situation than on the inner side of the femoral vein.[1]

FIG. 147.



1, 1. Fascia transversalis.
2. Internal ring.
3. Crural arch reflected.
4. Sheath of the femoral vessels.
5. Saphena vein.

DIAGRAM OF THE SHEATH OF THE FEMORAL VESSELS.

Sheath of the Femoral Vessels. The femoral vessels descend beneath the crural arch, enclosed in a funnel-shaped membranous sheath. This sheath appears to be derived immediately from the arch itself, but it is really formed *in front* by a prolongation from the fascia transversalis of the abdomen. This prolongation, uniting with the continuation from the fascia iliaca (to join the fascia lata) *behind* the femoral vessels, forms a funnel, with the wide part uppermost, into which the femoral vessels enter. This is the funnel-shaped sheath of the femoral vessels.

[1] If the partitions from any cause yield, or become slack, then a rupture may descend in front of the vessels, or even (though this is rare) on the outer side of the artery.

The fascia transversalis, descending over the femoral vessels, forms the *front part* of their sheath; the *hind part* of the sheath is formed by the fascia iliaca, which runs down behind the vessels to join the pubic portion of the fascia lata. The sheath descends as low as the lower horn of the saphenous opening, where it is gradually lost upon the external cellular coat (tunica adventitia) of the femoral vessels. The outer part of the sheath, in front, is perforated by the crural branch of the genito-crural nerve, and the superficial arteries of the groin; the inner part, by the saphena vein and some lymphatic vessels.

The sheath of the femoral vessels is divided into three compartments separated from each other by partitions; the outer is oocupied by the femoral artery; the middle, by the femoral vein; the inner is the crural canal, into which a femoral hernia descends.

The *deep crural arch* is the thickened band of fibres connected with the front of the crural sheath; the fibres run in the same direction as the crural arch, but quite independently of it, as shown in Fig. 147; these bands lie over the neck of the sac of a femoral hernia, and are often the seat of the stricture.

Practically, the sheath is important, for many reasons:—

1. A femoral hernia descends within it. 2. It constitutes, therefore, one of the coverings, *fascia propria*, of the hernia. 3. It contains within its substance the *deep crural arch*, which not infrequently forms the stricture of a femoral hernia, and has, therefore, to be divided before the intestine can be returned.

Crural Canal and Femoral Ring.

The hollow under the crural arch is completely occupied by the structures before mentioned, except for a small triangular space, forming the inner compartment of the femoral sheath, called the *crural canal*. The canal is on the inner side of the femoral vein, and is from a quarter to half an inch in length. Its base commences above in the femoral ring, and its apex ends below at the saphenous opening. *In front*, it has Poupart's ligament and the falciform process of the iliac portion of the fascia lata, and is formed by the fascia transversalis; *behind*, it is formed by the fascia iliaca; *internally*, it is formed by the junction of the fascia transversalis and the fascia iliaca, and is in relation with the base of Gimbernat's liga-

ment; *externally*, it is separated from the femoral vein by the septum of fascia which divides the middle from the inner compartment of the crural sheath.

The *femoral ring* is the upper opening of the crural canal, and is bounded, *in front*, by the superficial and deep crural arches; *behind*, by the horizontal ramus of the os pubis, the pectineus, and the pubic portion of the fascia lata; on the *outer side*, by the fascial septum separating it from the vein; on the *inner side*, by the thin, wiry edge of Gimbernat's ligament, the conjoined tendon of the internal oblique and transversalis, the fascia transversalis, and the fibres of the deep crural arch. In the undisturbed condition of the parts there is no gap; it is only a weak place, which, when a hernia escapes through it, feels like a ring; hence the name of *femoral ring*.[1]

The femoral ring is surrounded on all sides by unyielding structures. This accounts for the little benefit afforded by the warm bath in cases of strangulation. Sir W. Lawrence was in the habit of saying that he never saw a strangulated femoral hernia where the warm bath was of any avail.

Practical Application of the Subject. From what has been said, the student ought now to understand—1, at what aperture a femoral hernia escapes from the abdomen; 2, the course which it takes, and its relations to the surrounding parts; 3, the proper mode of attempting the reduction; 4, the structure and arrangement of its coverings; and, 5, the probable seat of stricture.

The hernia escapes from the abdomen through the femoral ring—that is, under the weak part of the crural arch, between the femoral ring and Gimbernat's ligament. Here is the mouth of the hernial sac, or that part of it which communicates with the abdomen. It descends a short distance nearly perpendicularly, and projects as a small tumor in front of the pectineus muscle. Its progress downward, however, is soon arrested, partly by the very close adhesion of the subcutaneous structures to the lower

[1] The femoral ring is naturally occupied by a little fat and cellular membrane, by lymphatic vessels, and often by a small lymphatic gland. But we have never met with anything deserving the name of a diaphragm or membranous septum, such as is described by Cloquet as the *septum crurale*, and is, surgically, of no importance.

margin of the saphenous opening; partly by the flexion of the thigh. Consequently, if the hernia increases in size, it usually rises over the crural arch, where the subcutaneous tissue offers less resistance; and the bulk of the hernia extends outward toward the ilium, assuming more or less of an oblong form, with the long axis parallel to the crural arch. Since, then, the body of the hernia forms a very acute angle with the neck, the right mode of attempting its reduction is, to draw it, first, down from the groin, and then to make pressure on it, backward, in the direction of the femoral ring.

Coverings of a Femoral Hernia. The *coverings* of a femoral hernia are as follows:—It first protrudes before it the *peritoneum*, technically called the hernial sac.[1] The sac is covered by more or less fat, according to the condition of the patient, called the *subperitoneal fat.* It next pushes before it the *sheath of the femoral vessels*, which forms an investment more or less thick. In front of this is the *cribriform fascia.* Lastly, there is the *subcutaneous tissue* and *skin.*

Seat of Stricture. The *seat of stricture* is usually at the femoral ring, and the position of the neighboring blood vessels indicates that the proper direction in which to divide the structure is, either directly inward, through Gimbernat's ligament, as recommended by Sir W. Lawrence, or upward, through Hey's ligament, as recommended by Sir A. Cooper.[2] There is no risk of wounding an artery, supposing the vessels to take their ordinary course. But it occasionally happens (Fig. 148), that the obturator artery runs *above* (in the recumbent position) the femoral ring; in such a case, the neck of the sac would be encircled by a large blood vessel.[3] From the examination of two hundred bodies, the

[1] In some cases the fascia propria so much resembles the hernial sac, that it is not easy to distinguish between them. Generally speaking, they are separated by a small quantity of fat.

[2] The operation recommended by Sir A. Cooper is that usually performed now; because, if Gimbernat's ligament be divided, its cut edges often retract to such an extent that no truss can possibly retain the hernia when the patient assumes the erect posture.

[3] The museum of St. Bartholomew's Hospital contains two examples of double femoral herniæ in the male, with the obturator arising on each side from the epigastric. In three out of four herniæ the obturator runs on the inner side of the mouth of the sac.

chances are about seventy to one against this unfavorable distribution. But the possibility of it has given rise to *this rule* in practice—not to cut deeply in any one place through the stricture, but rather to notch it in several. By this proceeding we are much less likely to wound the abnormal artery, because it does not run at the base of Gimbernat's ligament, but about a line and a half from the margin of it.[1]

Such is an outline of the anatomy of the parts concerned in a femoral hernia. The normal anatomy in each case being similar, it might be supposed that all operations for the relief of this kind of hernia would be straightforward and pretty much alike; but this is very far from being the case; indeed, surgeons agree that they never operate without the expectation of meeting some peculiarity.

FIG. 148.



A B

VIEW OF THE DIFFERENT DIRECTIONS WHICH AN ABNORMAL OBTURATOR ARTERY MAY TAKE (SEEN FROM ABOVE.)

A. 1. Gimbernat's ligament.
2. Femoral ring.
3. Abnormal obturator artery.
4. External iliac vein.
5. External iliac artery.
6. Diminutive obturator artery arising from its normal source.

B. 1. Gimbernat's ligament.
2. Abnormal obturator artery.
3. Femoral ring.
4. External iliac vein.
5. External iliac artery.
6. Diminutive obturator artery.

Dissection. The fascia must now be removed from the front of the thigh, without disturbing the subjacent muscles from their relative positions. The mass of muscles on

[1] During the session of 1867–68 more than half a dozen instances occurred where the obturator artery was given off by a common origin with the epigastric artery. In all these cases, however, the artery passed close by the bone, that is, *behind* the sac, so that it would not have been injured in the operation for the relief of strangulation.

the inner side of the thigh consists of the adductors; that in the middle, of the extensors; the long, thin muscle crossing obliquely in front from the outer to the inner side is the sartorius. In the middle are seen the femoral vessels, and the anterior crural nerve emerging beneath the crural arch.

Sartorius. This, a narrow, flat muscle (Fig. 149, 1), *arises* from the anterior superior spine of the ilium, and from the ridge below to the extent of an inch. It passes obliquely like a strap over the front of the thigh toward the inner side; and then descends almost perpendicularly on the inner side of the thigh as far as the knee, where it terminates in a flat tendon which expands and is *inserted* into the inner and front part of the tibia just below its tubercle. The tendon appears all the wider on account of its broad connection with the fascia of the leg, which extends as low as the internal malleolus. The broad insertion of this muscle lies anterior to and covers the tendinous insertions of the gracilis and semi-tendinosus, and between them is a bursa. A large bursa[1] is interposed between the tendon and the internal lateral ligament. The chief *action* of the muscle is to fix the pelvis steadily on the thigh.[2] It first bends the leg upon the thigh, and then bends the thigh upon the abdomen. It crosses one leg over the other, as tailors sit when at work. If the leg be the fixed point, it will bend the trunk upon the thigh and rotate the pelvis inward. Its nerve comes from the middle cutaneous branch of the anterior crural.

Scarpa's Triangle. In consequence of the oblique direction of the upper third of the sartorius, a triangle is formed, which has this muscle and the adductor longus for its two sides, and the crural arch for its base; the triangle is called *Scarpa's*.[3]

Its floor is formed by the iliacus, the psoas, the pectineus and the adductor longus, with sometimes the adductor brevis between

[1] In persons, females especially, who are in the habit of riding, this bursa sometimes becomes enlarged.

[2] Hence the name, given to it by Spigelius (*De Corporis Hum. Fabric.*), "Quem ego sartorium musculum vocare soleo, quod sartores eo maximè utuntur, dum crus cruri inter consuendum imponunt."

[3] So called in compliment to the Italian anatomist who first tied the femoral in it for popliteal aneurism.

the borders of the two latter muscles. The contents of this important space should be carefully displayed, and their relative positions well studied. This triangle contains all the parts which pass under the crural arch: namely, from without inward, the external cutaneous nerve, close to the anterior spine of the ilium; the iliacus and psoas; the anterior crural nerve and its divisions, especially the long saphenous nerve; the crural branch of the genito-crural nerve, the common femoral artery with its two large

FIG. 149.

DIAGRAM OF SCARPA'S TRIANGLE.

divisions, the superficial femoral and the profunda, which run down, nearly parallel to each other, the latter being the more external and giving off the internal and external circumflex arteries; the femoral vein, joined by the profunda vein and the internal saphena, and the pectineus muscle with the deep external pudic artery.

The triangle is important in a surgical point of view, since it is in this space that the femoral artery is usually ligatured for popliteal aneurism. The guide to the artery is the inner border

of the sartorius. The situation at which this muscle crosses over the femoral artery varies from one and a half to four and a half inches below Poupart's ligament; so that no rule can be laid down as to the exact situation where the artery disappears beneath the sartorius. The best way to find the inner border of the muscle during life is to make the patient put it in action.

Adductor Muscles. A strong group of muscles, called the *adductors*, extends along the inner side of the thigh, from the pelvis to the femur. Their two most important actions are to co-operate in balancing the pelvis steadily on the thigh, as in standing on one leg; and (if the fixed point be reversed) to draw together or adduct the thighs, at the same time rotating the thigh externally. They are five in number, and are supplied, with one exception—the pectineus—by the same nerve, namely, the obturator. They are termed, respectively, the gracilis, adductor longus, pectineus, adductor brevis, and adductor magnus. The innermost is the gracilis; to clean it properly, it should be stretched by separating one thigh from the other.

Gracilis. This long, flat muscle *arises* by a broad, ribbon-like tendon, two to three inches in breadth, from the os pubis close to the symphysis, and from the inner margin of the rami of the os pubis and ischium. It descends almost perpendicularly on the inner side of the thigh, and terminates in a thin, round tendon, which subsequently spreads out and is *inserted* into the inner side of the upper part of the tibia below the tubercle, immediately behind the sartorious and above the semitendinosus. The tendon plays over the internal lateral ligament of the knee joint, and there is a bursa common to it and the semitendinosus, to diminish friction. This muscle assists in fixing the pelvis, and in adducting the thigh; it further helps to bend the knee. Its nerve comes from the anterior division of the obturator.

Adductor Longus. This triangular muscle lies between the gracilis and the pectineus, and *arises* by a round tendon from the front of the body of the os pubis, below the crest. As it descends, the muscle becomes broader, and passing downward, outward, and backward, is *inserted* by a broad aponeurosis into the middle third of the inner margin of the linea aspera of the

femur. It forms with the sartorius the triangular space called Scarpa's triangle, above described. It rests upon the adductor brevis and magnus, the profunda vessels, and the anterior branches of the obturator vessels and nerve. It is supplied by the anterior division of the obturator nerve.

Pectineus. This muscle lies on the same plane, but external to the adductor longus, from which it is separated by a slight interval, in which may be seen the adductor brevis and the anterior division of the obturator nerve. It *arises* from the linea ilio-pectinea, from the triangular surface of the os pubis in front of the line, and from the fascial prolongation of Gimbernat's ligament covering the muscle; it passes downward, outward and backward, and is *inserted* into the upper part of the ridge leading from the lesser trochanter to the linea aspera. It lies upon the capsular ligament of the hip joint, the adductor brevis, the obturator vessels and nerve, and the obturator externus. Its nerve comes from the anterior crural, which runs under the femoral vessels to enter it close to its outer border; sometimes also from the obturator, and the accessory obturator if present (p. 519).

By separating the contiguous borders of the pectineus and the adductor longus, the adductor brevis is exposed with the anterior division of the obturator artery and nerve lying upon it. To obtain a complete view of it, the pectineus and adductor longus must be reflected from their origins and turned downward. The obturator nerve supplies all the adductors. It leaves the pelvis through the upper part of the obturator foramen, and soon divides into an anterior and posterior branch; the *anterior* runs in front of the adductor brevis, and supplies the hip joint, the adductor longus, the gracilis, and sometimes the adductor brevis and the pectineus; the *posterior* runs behind the adductor brevis, and supplies it as well as the obturator externus, the adductor magnus, and the knee joint.

The student should now, before the parts are distributed, examine the femoral artery as it passes down along the centre of Scarpa's triangle; its further course will be described later on, as well as the branches which come off from it.

Course and Relations of the Femoral Artery. The *femoral artery* is a continuation of the external iliac. Passing beneath the crural arch at a point midway between the spine of the ilium

and the symphysis pubis, it descends along the front and inner side of the thigh. At the junction of the upper two-thirds with the lower third of the thigh, it passes through an opening in the tendon of the adductor magnus, and, entering the ham, takes the name of popliteal. A line drawn from the point indicated of the crural arch to the adductor tubercle on the internal condyle corresponds with the course of the artery. Its distance from the surface increases as it descends. Immediately under, and for a short distance below the crural arch, it is supported by the inner border of the psoas; lower down it runs in front of the pectineus, but separated from it by the profunda vessels; still lower down it lies upon the adductor longus, and then upon the adductor magnus.

That part of the artery which extends from the crural arch to the giving off of the profunda is called the common femoral artery; its continuation beyond the profunda is termed the superficial femoral; and it is the latter vessel which is ligatured for aneurism of the popliteal artery.

In the *upper third of the thigh* the artery is situated in Scarpa's triangle, and is comparatively superficial, having *in front* the skin, superficial fascia and fat, inguinal glands, deep fascia, the fascia lata, the crural branch of the genito-crural nerve, and the sheath of the femoral vessels. About the *middle third* it is more deeply seated, and is covered in addition by the sartorius; and lower down by a tendinous aponeurosis, which stretches from the adductor longus and magnus over to the vastus internus. This, which forms part of Hunter's canal, will be examined presently.

The femoral artery in Scarpa's triangle *lies upon* the psoas, the two branches of the anterior crural nerve to the pectineus, the profunda vein, and the pectineus; to its *outer side* it has the anterior crural nerve (separated from it by a few fibres of the ilio-psoas), the profunda artery, and the long saphena nerve; to its *inner* side it has the femoral vein.

Adductor Brevis. This muscle *arises* from the front surface of the body of the os pubis below the spine, and from its descending ramus for about an inch, between the gracilis and the obturator externus; it widens as it descends outward and backward, and is *inserted* behind the pectineus into the whole

length of the ridge leading from the lesser trochanter to the linea aspera. Behind, it rests upon the posterior division of the obturator vessels and nerve, and the adductor magnus. Its nerve is derived from the obturator. By reflecting it from its origin, the following muscle is exposed.[1]

Adductor Magnus. This muscle arises from the lower part of the descending ramus of the os pubis between the adductor brevis and obturator externus, from the margin of the ascending ramus of the ischium, and from the lower and anterior part of the tuberosity of the ischium. Its fibres spread out, and are *inserted*, behind the other adductors, into the lower part of the linea quadrati, into the ridge leading from the great trochanter to the linea aspera, also into the whole length of the linea aspera, and the ridge leading from it to the inner condyle; while those fibres which arise from the tuberosity of the ischium pass vertically downward, and are inserted by a rounded tendon into the adductor tubercle on the inner condyle of the femur. Between the muscular fibres of the middle and lower thirds of the insertion of this muscle the femoral artery passes to the back of the thigh. The upper fibres pass transversely outward to their insertion, while the lower fibres descend nearly vertically. In front of the muscle are, the adductor longus and brevis, the vastus internus, the obturator nerve and artery, and the profunda artery; above it are the internal circumflex artery, the obturator externus, and the quadratus femoris; behind it the biceps, semi-tendinosus and semi-membranosus, the great sciatic nerve, and the gluteus maximus. Its nerve comes from the posterior division of the obturator and the great sciatic. Observe that all the adductor muscles are inserted into the femur by flat tendons more or less connected.

About the junction of the upper two-thirds with the lower third of the thigh, the femoral artery passes through an oval opening in the tendon of the adductor magnus.

Psoas Magnus and Iliacus These muscles have been fully described in the dissection of the abdomen (p. 509).

[1] Beneath the adductor brevis, and running parallel with the upper border of the adductor magnus, is seen the obturator externus. But the description of this muscle is deferred till the dissection of the external rotators of the thigh.

Tensor Fasciæ Femoris. This muscle is situated at the upper and outer part of the thigh. It *arises* from the anterior part of the external lip of the crest of the ilium, and from the surface below the anterior superior spine. It descends with a slight inclination backward, and is *inserted*, at the junction of the upper with the middle third of the thigh, between two layers of the strong aponeurosis, generally described as part of the fascia lata, which is continued downward to the head of the tibia, and is called the *ilio-tibial band* (p. 632).[1] Its chief use is to fix the pelvis steadily on the thigh, and to rotate the thigh inward; in this last action it co-operates with the anterior fibres of the gluteus medius, with which it is almost inseparably connected. Any one may convince himself of this by placing his hand on the hip, and rotating the thigh inward. Both these muscles are supplied by the same nerve—the superior gluteal.

To form an adequate idea of the strength, extent and connections of the aponeurosis on the outer side of the thigh, it should be separated from the vastus externus muscle upon which it lies. There is no difficulty in doing so, for it is united to the muscle by an abundance of loose connective tissue.[2] With a little perseverance the aponeurosis can be traced to the linea aspera, the head of the tibia and the fibula, completely protecting the outer side of the knee joint.

Extensor Muscles or Quadriceps Extensor. The powerful fleshy muscles occupying the front of the thigh, and situated between the tensor fasciæ on the outer side, and the adductors on the inner, are the extensors of the leg. One of them, the *rectus*, arises from the pelvis; the other, the triceps, arises from the shaft of the thigh bone by three portions, called, respectively, the *crureus*, the *vastus internus* and *externus*. All are supplied by the anterior crural nerve.

To see the origins of the rectus femoris, dissect between the

[1] The deeper of these two layers runs up to be strongly connected with the tendon of the rectus and the front of the capsule of the hip joint.

[2] When this tissue becomes the seat of suppuration, the pus is apt to extend all down the outside of the thigh, not being able to make its way to the surface, by reason of the dense fascia.

origin of the sartorius and the tensor fasciæ; in doing so, avoid injuring the branches of the external circumflex artery.

Rectus Femoris. This bipenniform muscle *arises* from the pelvis by two strong tendons, which soon unite at an acute angle; one, the straight tendon, is round, and arises from the anterior inferior spine of the ilium; the other, the reflected tendon, is flat, and comes from the rough surface of the ilium, just above the acetabulum. The muscle descends along the front of the thigh, and is *inserted* into the common extensor tendon, which will be presently examined. The structure of this muscle is remarkable. A tendon runs down the centre, and the muscular fibres are inserted on either side of it, like the vane on the shaft of a feather. Notice, also, that the surface of the upper part of the rectus is aponeurotic in front and muscular behind, while the reverse is seen at the lower part, not far from its insertion. Its nerve comes from the anterior crural.[1]

Triceps Extensor. This mass of muscle invests, like a cloak, the greater part of the front and sides of the shaft of the femur; therefore, the whole of it cannot be seen without completely dissecting the thigh. It consists of an outer, middle and inner portion, called, respectively, the vastus externus, the crureus and the vastus internus.

The *vastus externus arises* by a strong, glistening aponeurosis from the outer side of the base of the great trochanter, from the upper third of the anterior intertrochanteric ridge, from the rough line leading from the greater trochanter to the linea aspera, external to the gluteus maximus, from the outer lip of the linea aspera nearly down to the external condyle, and also slightly from the external intermuscular septum. From this origin the fibres pass downward, forward and inward, and end in a flattened tendon, which is inserted into the outer border of the patella to form part of the common extensor tendon, to be presently described.

The *vastus internus* and *crureus* should be described as one muscle, for they are inseparably connected. They arise conjointly by an aponeurosis commencing a short distance below the inner

[1] An accurate description of this muscle is given by W. R. Williams, *Journ. of Anat. and Phys.*, vol. xiii, p. 204, in which he states that the reflected head of the rectus is its real origin, as it alone exists in early fœtal life.

two-thirds of the anterior intertrochanteric ridge, from the upper three-fourths of the front and inner surfaces of the shaft of the femur, from the entire length of the inner lip of the linea aspera, and from the internal intermuscular septum. The outer bundle of muscular fibres, the *crureus*, passes vertically downward; the inner, the *vastus internus*, descends forward and outward, and both are inserted by a common aponeurosis into the upper and the inner borders of the patella. The muscular fibres of the vastus internus extend lower than those of the vastus externus.

A few of the deeper fibres of the crureus are inserted into the fold of the synovial membrane of the knee joint which rises above the patella. These are described as a distinct muscle, under the name of the *sub-crureus*. Their use is to raise the synovial membrane, so that it may not be injured by the play of the patella. Since the triceps is connected to the lower part of the shaft of the femur only by loose connective tissue, there is nothing to prevent the distention of the synovial membrane, in cases of inflammation, to the extent of several inches above the patella.

Common Extensor Tendon. The tendon of the rectus, gradually expanding, becomes connected on its under surface with the tendon of the crureus, and on either side with that of the vasti, and is firmly fixed into the upper part and sides of the patella. From this bone the common extensor tendon, the *ligamentum patellæ*, descends over the front of the knee joint, and is *inserted* into the rough part of the tubercle of the tibia. Besides this, the lower fibres of the vasti terminate on a sheet-like tendon, which runs wide of the patella on either side, and is directly inserted into the sides of the head of the tibia and fibula, so that the knee is completely protected all round. The patella is a large sesamoid bone, interposed to facilitate the play of the tendon over the condyles of the femur; it not only materially protects the joint, but adds to the power of the extensor muscles, by increasing the angle at which the tendon is inserted into the tibia.

To facilitate the play of the extensor tendon there are two bursæ. One is placed between the ligamentum patellæ and the smooth part of the tubercle of the tibia, the other between the crureus and the lower part of the femur. This last is of consid-

erable size. In early life it is, as a rule, distinct from the synovial membrane of the knee joint; but after a few years a wide communication frequently exists between them.

Action of the Extensor Muscles. The extensor muscles of the thigh are among the most powerful in the body. Great power of extending the knee is one of the essential conditions of the erect attitude. Without it how could we rise from the sitting position? When erect, how could we walk, run or spring? The rectus, by taking origin from the pelvis, gains a double advantage; it acts upon two joints simultaneously, bending the thigh while it extends the knee, as when we advance the leg in walking; it also contributes to balance the pelvis on the head of the thigh bone, and thus prevents the body from falling backward. We cannot have a better proof of the power of the extensor muscles than when the patella is broken by their sudden contraction—an injury which sometimes happens when a man, slipping backward, makes a violent effort to recover his balance.

Bursa over the Patella. The skin over the patella is exceedingly loose, and in the subcutaneous tissue is a bursa of considerable size. Since this bursa is apt to enlarge and inflame in females who are in the habit of kneeling at their work, it is generally called the housemaid's bursa. The bursa is not seated precisely over the patella, but extends some way down the ligamentum patellæ; indeed, in some cases it is entirely confined to this ligament. This corresponds with the position of the tumor which the bursa occasions when enlarged. Generally speaking, in subjects brought for dissection, the wall of the bursa is more or less thickened, and its interior intersected by numerous fibrous cords, remnants of the original cellular structure altered by long continued friction. Again, the wall of the bursa does not always form a complete sac; sometimes there is a wide opening in it; this explains the rapidity with which inflammation, in some cases, extends from the bursa into the surrounding areolar tissue.

Below the bursa is a layer of fascia lata, and under this is a network of arteries. The immediate covering of the bone, or what may be called its periosteum, is a strong expansion derived from the extensor tendon. This is interesting, for the following

reason; in ordinary fractures of the patella from muscular action the tendinous expansion over it is torn also; the ends of the bone gape widely, and never unite except by ligament. But in fractures from direct mechanical violence, the tendinous expansion, being entire, maintains the fragments in apposition, so that there is commonly a bony union.

The remaining part of the femoral artery can now be examined after the sartorius has been cut through near its middle, and both ends reflected. This part of its course corresponds to the middle third of the thigh, and is contained in Hunter's canal.

Femoral Artery in Hunter's Canal. *In front* of the artery are the skin, superficial and deep fasciæ, the long saphena vein, the sartorius, the long saphenous nerve, and the aponeurotic layer forming the anterior boundary of Hunter's canal; to its *outer side* are the femoral vein and the vastus internus; to its *inner side* are the adductor longus, the adductor magnus and the sartorius; *behind* it are the adductor longus, the femoral vein, and the adductor magnus. The artery and vein lie close together, and are enclosed in a common sheath.

Hunter's Canal. In the middle third of the thigh the femoral artery is contained in a tendinous canal[1] beneath the sartorius, called *Hunter's canal.* This canal at its upper part is rather indistinct; but it gradually becomes stronger toward the opening in the tendon of the adductor magnus. Its boundaries are formed by the tendons of the muscles between which the artery runs. On the inner side are the tendons of the adductor longus and magnus; on the outer side is the tendon of the vastus internus; in front the canal is completed by an aponeurotic expansion thrown obliquely across from the adductors to the vastus internus, as shown in Fig. 150. In a horizontal section the canal appears triangular. The adapt-

FIG. 150.

SECTION THROUGH HUNTER'S CANAL.

1. Vastus internus.
2. Adductor longus.
3. Aponeurosis thrown across.

[1] Called Hunter's canal, because it was in this part of its course that John Hunter first tied the femoral artery for aneurism of the popliteal, in St. George's Hospital, A. D., 1785. The particulars of this interesting case are published in the *Trans. for the Improvement of Med. and Chir. Knowledge.*

ation of this shape to the exigencies of the case is manifest when we reflect that the muscles keep the sides of the triangle always tight, and thereby prevent any compression of the vessels.

Hunter's canal contains not only the femoral artery and vein, but the internal saphenous nerve. The vein lies behind and to the outer side; the nerve crosses over the artery from the outer to the inner side.

A ligature can be placed around the artery, in the upper third of the thigh, with comparative facility; not so easily in the middle third. The artery is tied for an aneurism of the popliteal, just when the sartorius begins to overlap it, for three reasons: (1) it is more accessible; (2) the coats of the artery at this distance are less likely to be diseased; (3) the origin of the profunda is sufficiently far off to admit of the formation of a clot. An incision, beginning about three inches below the crural arch, should be made, about three inches long, over the line of the artery. The muscular fascia should be divided on a director to the same extent. Then, by gently drawing aside the inner border of the sartorius, the artery is seen enclosed in its sheath with the vein. An opening should be made into the sheath, which must be carefully separated from the artery to an extent sufficient to allow the passage of the aneurismal needle. The needle should be turned round the artery from within outward, great care being taken not to injure the vein. The nerves to be avoided are, the long saphenous, which runs along the outer side of the artery, and the internal cutaneous, which crosses obliquely over it.

Having already traced the superficial branches of the femoral artery in the groin, namely, the superficial epigastric, the external pudic, and the superficial circumflexa ilii (p. 628), we pass now to the profunda.

Profunda Artery and Branches. The *profunda femoris*, the chief branch of the femoral, is the proper nutrient artery of the muscles of the thigh, and is considered as a division rather than a branch, of the common femoral artery. It is given off from the outer and back part of the femoral, from one and a half to three inches below the crural arch, lying to the outer side of the artery for about two inches, and then runs down behind the femoral till it reaches the tendon of the adductor longus; here the

profunda passes behind the adductor, and piercing the adductor magnus as a small branch, is finally lost in the hamstring muscles.[1] In most subjects the profunda, for a short distance after its origin, lies rather on the outer side of the femoral and on a deeper plane, on the iliacus; in this situation it might be mistaken for the superficial femoral itself—indeed, such an error has occurred in practice. It soon, however, gets behind the femoral, and lies upon the pectineus, the adductor brevis and magnus; it is separated from the femoral artery, at first, by their corresponding veins; lower down, by the adductor longus.

The branches of the profunda generally arise in the following order: (1) the internal circumflex; (2) the external circumflex; (3) the perforating.

The *internal circumflex* is given off from the inner and back part of the profunda, and then sinks deeply into the thigh between the psoas and pectineus. At the lower border of the obturator externus it divides into two branches: one, the *ascending*, supplies the muscles in its neighborhood, namely, the pectineus, psoas, adductors, gracilis, and obturator externus, anastomosing with the obturator artery; the other, the *descending*, passes down to behind the adductor brevis to supply it and the adductor magnus; the continuation of the artery, called the *transverse*, will be seen in the dissection of the back of the thigh, between the adductor magnus and the quadratus femoris. This latter sometimes gives off a small branch to the hip joint, which runs through the notch in the acetabulum to the ligamentum teres; it afterward inosculates with the sciatic, the external circumflex and superior perforating arteries, forming the *crucial anastomosis*.

The *external circumflex artery* comes off from the outer side of the profunda, runs transversely outward beneath the sartorius and rectus, between the branches of the anterior crural nerve, and then subdivides into three sets of branches, ascending, transverse and descending. The *ascending* run up to the outer side of the

[1] The point at which the profunda is given off below the crural arch varies very much, even in the two limbs of the same body. We have measured it in 19 bodies, or 38 femoral arteries. It varied from half an inch to 3 inches. In 22 cases the profunda came off between 1½ and 2 inches; in 9 this distance was exceeded; in 7 this distance was less.

ilium, beneath the tensor fasciæ and gluteus medius, supply these muscles, and inosculate with the terminal branches of the gluteal and deep circumflex iliac arteries. The *transverse* pass directly outward over the crureus, then enter the vastus externus, and get between the muscle and the femur. They inosculate with the sciatic, the internal circumflex, the gluteal, and the perforating

FIG. 151.

1. Crural arch.
2. Internal iliac.
3. Superficial femoral.
4. Profunda.
5. Internal circumflex.
6. External circumflex.



7. First perforating.
8. Second ditto.
9. Third ditto.
10. Gluteal.
11. Obturator.
12. Sciatic.
13. Anastomotica magna.

PLAN OF THE INOSCULATIONS OF THE CIRCUMFLEX ARTERIES.

arteries. The *descending*, two or more in number, of considera size, run down between the rectus and crureus, and sup both these muscles: one branch, larger than the rest, runs do in the substance of the vastus externus, along with the nerve that muscle, and inosculates with the superior articular branc of the popliteal.

The *perforating branches* of the profunda are so named becau they pass through the adductors to supply the hamstring muscl

There are generally four. The *first* passes between the pectineus and the adductor brevis, then pierces the adductor magnus, and communicates with the internal and external circumflex, the sciatic and perforating arteries. The *second*, the largest, passes through the tendons of the adductor brevis and magnus, divides into an ascending and descending branch, which anastomose respectively with the first and third perforating arteries. It usually furnishes the nutrient artery of the femur. The *third*, given off below the adductor brevis, passes through the tendon of the adductor magnus. The *fourth*, or terminal branch, passes through the tendon of the adductor magnus, and supplies the hamstring muscles, and inosculates with the perforating and articular arteries. They not only supply the hamstring muscles—namely, the biceps, semitendinosus, and semimembranosus—but the vastus externus, and even the gluteus maximus. The perforating arteries inosculate with one another, with the internal and external circumflex, and with the sciatic arteries.

Muscular branches, from four to seven in number, are distributed by the superficial femoral to the sartorius and the vastus internus.

The *anastomotica magna* arises from the femoral artery just before it leaves its tendinous canal. It emerges through the canal, and runs in front of the tendon of the adductor magnus, in company with the long saphenous nerve to the inner side of the knee. Here it divides into two branches: one, the *superficial*, accompanies the saphenous nerve beneath the sartorius, and is subsequently distributed to the skin; the other, the *deep*, enters the vastus internus, ramifies over the capsule, and communicates with the other articular arteries.[1]

Arterial Inosculations.

If the common femoral were tied *above* the origin of the profunda, how would the circulation be carried on? The gluteal, the ilio-lumbar, and the circumflex iliac communicate with the ascending branch of the external circumflex; the obturator and sciatic communicate with the internal circumflex (see Fig. 151); the arteria comes nervi ischiatici com-

[1] In its course down the thigh the femoral artery gives off a branch of considerable size for the supply of the vastus internus. We may trace this branch through the substance of the vastus down to the patella, where it joins the network of vessels on the surface of that bone.

municates with branches from the lower perforating and popliteal arteries. Again, how is the circulation maintained when the superficial femoral is tied *below* the profunda? The descending branch of the external circumflex and the perforating branches of the profunda communicate with the articular branches of the popliteal and the tibial recurrent.[1]

Anterior Crural Nerve. The anterior crural nerve is the largest branch of the lumbar plexus (p. 518). It comes from the third and fourth lumbar nerves, also by a small fasciculus from the second. It passes beneath the crural arch, lying in the groove between the iliacus and psoas, about a quarter of an inch to the outer side of the artery, and soon divides into branches, some of which are cutaneous, but the greater number supply the extensor muscles of the thigh. The cutaneous branches, already described (p. 630), and the long saphenous nerve, are given off from the *superficial* part of the trunk; the muscular from the *deep* part.

The *long saphenous nerve*, the largest of the cutaneous branches, descends close to the outer side of the femoral artery, and enters the tendinous canal with it in the middle third of the thigh. In the canal it crosses over the artery to its inner side. The nerve leaves the artery just before it becomes popliteal, and then runs in company with the anastomotica magna to the inner side of the knee, where it becomes superficial, between the gracilis and the sartorius. In the middle third of the thigh it gives off a small branch which communicates beneath the fascia lata with the internal cutaneous and obturator nerves; and lower down another branch is distributed to the skin over the patella. Its further relations will be seen in the dissection of the leg and foot.

The *muscular branches* are to be traced to the sartorius, rectus, crureus, and subcrureus; the branch to the vastus externus accompanies the descending branch of the external circumflex artery, and sends a filament to the knee joint; that to the vastus internus runs parallel with, but external to, the long saphenous nerve, and supplies filaments to the knee joint. One branch, often two, passes under the femoral artery and vein to enter the anterior surface of the pectineus.

[1] Read the account of the dissection of an aneurismal limb by Sir A. Cooper, *Med.-Chir. Trans.*, vol. ii, 1811.

The *obturator nerve*, also a branch of the lumbar plexus, arising from the second, third and fourth lumbar nerves (p. 519), supplies the adductor muscles. It enters the thigh through the upper part of the obturator foramen above the corresponding artery, and immediately divides into two branches, of which one passes in front of, the other behind, the adductor brevis. The *anterior* branch subdivides for the supply of the gracilis, the adductor longus, and sometimes the adductor brevis and pectineus; it, moreover, sends a filament to the hip joint; another to the femoral artery; and a third forms a plexiform communication at the lower border of the adductor longus with the internal cutaneous and long saphenous nerves. The *posterior* branch supplies the obturator externus, the adductor brevis and magnus. In some bodies you can trace a filament of this nerve through the notch of the acetabulum into the hip joint, and another, which runs near the popliteal artery, into the back part of the knee joint. We have frequently seen cutaneous branches from the obturator on the inner side of the thigh. This is interesting practically, since it helps to explain the pain often felt on the inner side of the knee in disease of the hip joint.

The *accessory obturator nerve*, when present, comes either from the obturator nerve or from the third and fourth lumbar nerves. Descending, it runs between the horizontal ramus of the os pubis and the pectineus, and supplies a branch to this muscle, also a filament of communication to the anterior branch of the obturator, and a third branch to the hip joint.

The *obturator artery*, after passing through the foramen, divides into two branches, an *internal* and an *external*, which form a circle round the obturator membrane. These supply the external obturator and adductors of the thigh, and inosculate with the internal circumflex artery (p. 655). The latter branch sometimes gives off the small artery to the ligamentum teres of the hip joint.

DISSECTION OF THE FRONT OF THE LEG.

Surface Marking. About an inch below the patella is the prominent tubercle of the tibia, to which the ligamentum patellæ is attached; on each side of this is a depression, filled with more or less fat. About the same distance below the outer tuber-

osity of the tibia is the head of the fibula, situated far back, and to it can be traced the tense tendon of the biceps. The crest of the tibia is easily felt in front, commencing above at the outer tuberosity, and passing down nearly vertically, gradually inclining to the inner side, so that it is continuous below with the front of the internal malleolus. Internal to the crest is the subcutaneous internal surface of the tibia, and externally is the interval between the tibia and the fibula, which is filled up by the extensor muscles. The lower fourth of the fibula is subcutaneous, ending in a well-marked prominence, the external malleolus, which, it should be observed, descends lower than the internal malleolus. The student should notice well the tendons which surround the ankle joint, which are easily recognizable under the skin; thus, behind, the tendo-Achillis stands out prominently, having a deep depression on each side; on the outer side, the tendons of the peronei, longus and brevis, are felt, the latter being the anterior. Running round the inner ankle we can only feel the tibialis posticus close to the tibia, and next to it the flexor longus digitorum; in front of the ankle, but bound down by the anterior annular ligament, can be felt, from within outward, the strong tendon of the tibialis anticus, the extensor proprius hallucis, and the long extensor of the toes with the peroneus tertius.

The foot should be turned inward, and fixed in this position. An incision must be made from the knee, down the front of the leg, over the ankle, along the top of the foot to the great toe; a second, at right angles to the first, on either side of the ankle; a third, across the bases of the toes. Reflect the skin from the front and sides of the leg and foot.

Cutaneous Veins and Nerves. Having traced the internal saphena vein (p. 630) to the inner side of the knee, follow it down the inner side of the leg, in front of the inner ankle[1] to the dorsum of the foot. On the dorsum of the foot notice that the principal veins form an arch, with the convexity forward, as on the back of the hand. This arch receives the veins from the toes. From the inner side of the arch the *internal saphena* originates; from the outer side, the external saphena. The latter vein commences

[1] The French commonly bleed from the internal saphena vein, as it crosses over the inner ankle, this being a convenient and safe place for venesection.

on the outer side of the arch on the dorsum, runs behind the external ankle, along the outer border of the tendo-Achillis, up the back of the calf of the leg, between the two heads of the gastrocnemius, and finally pierces the deep fascia at the lower part of the popliteal space, to join the popliteal vein. The external saphenous nerve accompanies this vein, as the long saphenous nerve does the internal saphena vein.

Long Saphenous Nerve. The skin on the inner side of the leg is supplied by the long or internal saphenous nerve (p. 658). It becomes subcutaneous on the inner side of the knee, between the gracilis and sartorius. Here it meets the saphena vein, and accompanies it down the leg, distributing its branches on either side, till it is finally lost on the inner side of the foot and the great toe. The largest branch curves round the inner side of the knee, just below the patella, to supply the skin in this situation. It pierces the sartorius close to the knee, and forms, with branches from the internal, middle, and external cutaneous nerves, the *plexus patellæ*.

The *internal cutaneous nerve* supplies the skin of the upper and inner aspect of the leg, and joins the internal saphenous nerve.

The skin on the front and outer parts of the upper half of the leg is supplied by *cutaneous branches* from the *external popliteal* or *peroneal* nerve; the skin of the lower half by its *external cutaneous branches*, as follows:—

External Cutaneous Branch of the Peroneal Nerve. This branch of the peroneal nerve comes through the fascia about the lower third of the outer side of the leg; and, descending over the front of the ankle, divides into two. Trace them and you will find that the *inner* and smaller supplies the inner side of the great toe, and the contiguous sides of the second and third toes; toward its termination it communicates with the long saphenous and anterior tibial nerves. The *outer* distributes branches to the outer side of the third toe, both sides of the fourth and the inner side of the fifth toe, and joins the short or external saphenous nerve.

The outside of the little toe is supplied by the external saphenous nerve, which runs behind the outer ankle with the corresponding nerve.

The contiguous sides of the great and second toes are supplied by the termination of the anterior tibial nerve.[1]

Muscular Fascia and Annular Ligaments.

This is remarkably thick and strong. Besides its general purpose of forming sheaths for the muscles, and straps for the tendons, it gives origin, as in the forearm, to muscular fibres, so that it cannot be removed near the knee without leaving the muscles ragged. The fascia, continuous above with the fascia lata, is attached to the head of the tibia and the fibula; it is connected on the inner side with the expanded tendons of the sartorius, gracilis, and semi-tendinosus; on the outer side with that of the biceps; consequently, when these muscles act, it is rendered tense. Following it down the leg, you find that it is attached to the edge of the tibia, and that it becomes stronger as it approaches the ankle, to form the ligaments which confine the tendons in this situation. Of these ligaments, called *annular*, there are three, as follows:—

a. The *anterior annular ligament* extends obliquely across the front of the ankle joint, and confines the extensor tendons of the ankle and toes. It consists of two converging straps—one oblique, the other horizontal, which join, and are continued on as a common band, like the letter ≺ placed transversely; the upper or oblique binds down the tendons in front of the lower end of the tibia; the lower or oblique the tendons which lie over the tarsus. The common band is attached to the external malleolus, cuboid and os calcis; it is continued horizontally inward for a short distance, and in front of the ankle splits into two fasciculi: the upper or oblique ascends to be attached to the tibia; the lower or horizontal passes inward to be attached to the internal malleolus, the scaphoid, and the internal cuneiform. Beneath the upper fasciculus, enclosed in two synovial sheaths, run the tibialis anticus on the inner side, and the extensor longus digitorum and peroneus tertius on the outer side; the extensor proprius hallucis and the anterior tibial vessels lying behind the ligament, but not having any synovial sheath. Beneath the lower fasciculus are three synovial sheaths—an inner one for the tibialis anticus, a middle one for the extensor proprius hallucis, and an outer one for the

[1] Such is the most common distribution of the nerves to the upper surface of the toes. But deviations from this arrangement are frequent.

xtensor longus digitorum and peroneus tertius. It is the strain f this ligament which occasions the pain in sprains of the ankle.

b. The *external annular ligament* extends from the outer nalleolus to the os calcis, and confines the tendons of the peronei nuscles, which are enclosed in a common synovial sheath.

c. The *internal annular ligament* is a strong fasciculus of ill-lefined fibrous tissue, which extends from the inner malleolus to he os calcis, where it becomes continuous with the plantar fascia nd the tendinous origin of the abductor hallucis. It binds down he flexor tendons of the foot and toes, and, as these pass round he inner ankle, it forms three compartments, each lined with a separate synovial sheath—one each for the tibialis posticus, the lexor longus digitorum, and the flexor longus hallucis.

Remove the fascia, leaving enough of the annular ligaments to etain the tendons in their places.

Muscles on the Front of the Leg. The muscles on the front of the leg are: (1) the tibialis anticus; (2) the extensor longus digi-orum and peroneus tertius; (3) the extensor proprius hallucis.

Tibialis Anticus. The tibialis anticus *arises* by fleshy fibres from the external tuberosity and the upper two-thirds f the outer side of the shaft of the tibia, from the interosseous nembrane, from the fascia which covers it, and from the inter-nuscular septum which separates it from the extensor longus ligitorum. About the lower third of the leg the fibres terminate on strong flat tendon, which descends obliquely over the front of the nkle, through the innermost compartment of the anterior annular igament, to the inner side of the foot; here it becomes a little roader, and is *inserted* into the internal cuneiform bone and the ase of the metatarsal bone of the great toe. The synovial mem-rane, which lines the sheath of the tendon beneath the anterior nnular ligament, accompanies it to within an inch of its insertion; onsequently, it is opened when the tendon is divided for club-oot. The action of this muscle is to draw the foot upward nd inward.[1] When the foot is the fixed point, it assists in alancing the body at the ankle. Its nerve comes from the nterior tibial.

[1] It is generally necessary to divide this tendon in the distortion of the foot inward alled talipes varus.

Extensor Longus Digitorum. This muscle lies along the fibular side of the preceding. It *arises* from the external tuberosity of the tibia, from the upper three-fourths of the anterior surface of the shaft of the fibula, from the interosseous membrane, from the fascia of the leg and the intermuscular septa. Its fibres terminate in a penniform manner upon a long tendon, situated on the inner side of the muscle; this tendon descends in front of the ankle and divides into four slips, which pass to the four outer toes. They diverge from each other, and are *inserted* into the toes thus: On the base of the first phalanx, each tendon (except that of the little toe) is joined on its outer side by the corresponding tendon of the extensor brevis, and a little further on by a fibrous expansion from the interosseous and lumbrical muscles. The united tendons then expand, cover the dorsal surface of the first phalanx, and at the articulation between this and the second phalanx split into three fasciculi; the middle one is inserted into the base of the second phalanx; the two lateral ones, running on and reuniting, are inserted into the base of the third phalanx. Its nerve comes from the anterior tibial.

Immediately below the ankle the anterior annular ligament forms a pulley through which the tendon of this muscle plays. It is like a sling, of which the two ends are attached to the os calcis, while the loop serves to confine the tendon. The play of the tendon is facilitated by a synovial membrane, which is prolonged for a short distance along each of its four divisions. Besides its chief action, this muscle extends the ankle joint.[1]

Peroneus Tertius. This is a portion of the preceding. Its fibres *arise* from the lower fourth of the anterior surface of the shaft of the fibula, the interosseous membrane, and the intermuscular septum between it and the peroneus brevis, and terminate on their tendon, like barbs on a quill. The tendon passes through the same synovial pulley with the long extensor of the toes, and, expanding considerably, is *inserted* into the tarsal end of the metatarsal bone of the little toe. It is not always present. It is supplied by a branch of the anterior tibial nerve.

[1] There is often a large bursa between the tendon of the extensor longus digitorum and the outer end of the astragalus. This bursa sometimes communicates with the joint of the head of the astragalus.

his muscle extends the foot and draws the outer border of the ot upward.

The peroneus tertius and the tibialis anticus are important uscles in progression. They raise the toes and foot from the ound. Those who have lost the use of the muscles are obliged drag the foot along the ground, or to swing the entire limb itward in walking.

tensor Proprius Hallucis. This muscle lies partly concealed between the tibialis anticus and the extensor longus digitorum. *arises* from rather more than the middle third of the anterior rface of the fibula, and from the interosseous membrane. The res terminate in a penniform manner on the tendon, which runs er the ankle, between the tendons of the tibialis anticus and the tensor longus digitorum, along the top of the foot to the great e, where it is *inserted* into the base of the last phalanx. It has special pulley beneath the horizontal portion of the anterior nular ligament, lined by a synovial membrane, which accom- nies it as far as the metatarsal bone of the great toe. It is pplied by the anterior tibial, a branch of the peroneal nerve.

Now examine the course, relations and branches of the ante- or tibial artery. Since it lies deeply between the muscles, it is ecessary to separate them from one another; this is easily done proceeding from the ankle toward the knee.

Course and Relations of the Anterior Tibial Artery. The *anterior tibial artery* is one of the two branches into which the popliteal divides at the lower border of the popliteus. It comes at first horizontally forward, about 1¼ inch below the ad of the fibula, between the two heads of the tibialis posticus, ove the interosseous membrane, and then descends, lying, in ther more than the first half of its course, upon the interosseous embrane, afterward along the front of the tibia. It runs be- ath the anterior annular ligament, over the front of the ankle, here it takes the name of the dorsal artery of the foot. Thus, line drawn from the head of the fibula to the interval between e first and second metatarsal bones would nearly indicate its urse. In the upper third of the leg it lies deeply between the ialis anticus and the extensor longus digitorum; in the lower o-thirds, between the tibialis anticus and the extensor proprius

hallucis. In *front of the ankle* the artery is crossed by the extensor proprius hallucis, and lies between the tendon of this muscle and the inner tendon of the extensor longus digitorum.

The artery is accompanied by the anterior tibial nerve (a branch of the peroneal), which runs for some distance upon its fibular side, then in front of it, and lower down is again situated on its outer side. It is accompanied by two veins, one on each side, which communicate at intervals by cross branches.

The branches of the anterior tibial are as follows:—

a. The *recurrent tibial* branch ascends close by the outer side of the head of the tibia, through the tibialis anticus, to the front of the knee joint, where it inosculates with the other articular arteries derived from the popliteal, and with the anastomotica magna.

b. Muscular branches, in its course down the leg, and others which pierce the interosseous membrane, and communicate posteriorly with branches of the posterior tibial and peroneal arteries.

c. The *malleolar* branches, *external* and *internal*, ramify over the ankle; the *external*, descending beneath the tendon of the extensor longus digitorum and peroneus tertius, ramifies on the external malleolus, inosculating with the anterior peroneal and the tarsal arteries; the *internal* passes beneath the extensor proprius hallucis and the tibialis anticus, and anastomoses with the posterior tibial, with its internal calcanean branch, and with the internal plantar artery. They supply the joint, the articular ends of the bones, and the sheaths of the tendons around them.

Extensor Brevis Digitorum. This muscle is situated on the dorsum of the foot, beneath the long extensor tendons of the toes. It *arises* from the outer part of the os calcis, from the external calcaneo-astragaloid ligament, and from the anterior annular ligament. The fibres run obliquely over the foot, and terminate in four tendons, which pass forward to the four inner toes. The inner one and the largest is *inserted* by an expanded tendon into the base of the first phalanx of the great toe; the others *join* the fibular side of the long extensor tendons to be inserted with them into the second and ungual phalanges. The tendon to the great toe crosses over the dorsal artery of the foot. It is supplied by a branch of the anterior tibial nerve.

orsal Artery of the Foot. This artery, the continuation of the anterior tibial, runs over the instep to the back of the erval between the first and second metatarsal bones, where divides into two branches: one, the dorsalis hallucis, runs ng the dorsal aspect and the first interosseous space; the ıer, the communicating, sinks into the sole and joins the deep ntar arch. On the dorsum of the foot the artery *lies upon* astragalus, the scaphoid, and the internal cuneiform bones, arated, however, from them, by their dorsal ligaments; *in front*, artery has the skin, superficial and deep fasciæ, the venous h across the dorsum, and the innermost tendon of the short ensor of the toes; on its *outer side*, it has the extensor longus itorum and the anterior tibial nerve; on its *inner side*, the ensor proprius hallucis. The dorsal artery gives off the follow- branches:—

a. The *tarsal* branch arises near the scaphoid bone, passes ward in an arched direction beneath the extensor brevis digi- um toward the outside of the foot, supplies the bones and nts of the tarsus, and inosculates with the external malleolar, peroneal, the metatarsal, and the external plantar arteries.

b. The *metatarsal* branch generally runs toward the outside of foot, in front of and parallel with the tarsal artery, beneath short extensor tendons, near the bases of the metatarsal nes, and gives off the three outer *dorsal interosseous arteries*. ese pass forward over the corresponding interosseous muscles, pply them and then subdivide to supply the contiguous sides the upper surfaces of the toes. The outer interosseous branch, ddition to giving off a branch to the fourth interosseous space, es off a small branch to the outer side of the little toe. They eive, at the back of each interosseous space, the posterior per- ating branches of the plantar arch, and at the front of each erosseous space they receive the anterior perforating branches m the plantar digital arteries.

c. The *dorsalis hallucis* is, strictly speaking, the artery of the t interosseous space. It is the continuation of the dorsal ery of the foot, after it has given off the communicating branch the sole, and runs forward to supply *digital branches* to the es of the great toe and the inner side of the second toe.

Peronei Muscles. These muscles are situated on the outer side of the fibula, and are named, respectively, peroneus longus and brevis.

Peroneus Longus. This *arises* from the head and outer surface of the fibula along its upper two-thirds, from the deep fascia which covers it, and the intermuscular septa. The fibres terminate in a penniform manner upon a tendon, which runs with the peroneus brevis, in a groove behind the external malleolus, then along the outer side of the os calcis, and, lastly, through a groove on the under surface of the os cuboides, deep into the sole. It crosses the sole obliquely forward and inward, and is *inserted* into the tarsal end of the metatarsal bone of the great toe, and usually into the internal cuneiform bone. In its course through these several bony grooves the tendon is confined by a fibrous sheath, lined by a synovial membrane. In removing the metatarsal bone of the great toe, if possible, leave the attachment of this tendon, which is usually inserted by means of a sesamoid bone. Its nerve comes from the peroneal.

Peroneus Brevis. This muscle lies beneath the preceding. It *arises* from the lower two-thirds of the outer surface of the fibula, internal to the preceding muscle, and from the intermuscular septa. It terminates on a tendon which runs behind the external malleolus, through the same sheath with the peroneus longus, then proceeds along the outside of the foot, and is *inserted* into the dorsal surface of the tarsal end of the metatarsal bone of the little toe. On the outside of the os calcis there is a ridge which separates the tendons of the peronei. Each has a distinct sheath. The short tendon runs above, the long one below the ridge. Its nerve is from the musculo-cutaneous branch of the peroneal nerve.

The *action* of the peronei is to raise the outer side of the foot.[1] This movement regulates the bearing of the foot in progression, so as to throw the principal part of the weight on the ball of the great toe. Its action is well exemplified in skating. Again, supposing the fixed point to be at the foot, they tend to prevent the body from falling on the opposite side, as when we balance ourselves on one leg.

[1] In distortion of the foot outward, called talipes valgus, it is generally necessary to divide the tendons of the peronei.

Peroneal or External Popliteal Nerve.

Near the inner side of the tendon of the biceps flexor of the leg, is a large nerve, the *external popliteal* or *peroneal*, a branch of the great sciatic. By reflecting the upper part of the peroneus longus, you will find that this nerve runs round the outer side of the fibula immediately below its head, and, piercing the origin of the peroneus longus, divides into two main branches—the anterior tibial and the musculo-cutaneous nerves. It gives off several branches, as follows: 1. *Articular branches*, two in number, to the knee joint, which pass in with the external articular arteries, and a third, which accompanies the tibial recurrent artery. 2. *Cutaneous branches*, two or more, supply the skin on the back and outer aspect of the leg, and one, the *communicans peronei*, which joins the external saphenous to supply the dorsal aspect of the outer side of the little toe; this will be seen later on in the dissection of the back of the leg. 3. The *anterior tibial*, which accompanies the corresponding artery and supplies the muscles between which it runs —namely, the tibialis anticus, extensor longus digitorum, extensor proprius hallucis and peroneus tertius; also, the extensor brevis digitorum. 4. The *musculo-cutaneous* (p. 661), which comes through the fascia between the peroneus longus and the extensor longus digitorum. 5. *Branches*, which supply the peronei, longus and brevis, muscles.

If, then, the peroneal nerve were divided in the popliteal space, the result would be paralysis of the tibialis anticus, the extensors of the toes, long and short, and all the peronei.

DISSECTION OF THE GLUTEAL REGION.

The body having been placed on its face, the pelvis is to be raised to such a height, by blocks beneath it, that the lower extremities hang down over the end of the table. Then rotate the thighs inward as much as possible, and cross them.

Surface Marking.

The bony prominences are very marked and prominent; they are, on the outer side, the great trochanter, and on the inner, the posterior superior spine of the ilium, the spines of the sacral vertebræ, which are continued on to the coccyx. Between the latter bone and the great trochanter is the rounded tuberosity of the ischium, which in the erect po-

sition is covered by the gluteus maximus, but is uncovered by it when the femur is flexed. Notice a transverse curved fold of the skin extending from the coccyx to the base of the great trochanter, which does not, as it might be thought, correspond with the lower border of the gluteus maximus, which is much lower. This fact is important to bear in mind in operations for stretching the great sciatic nerve, for the relief of sciatica or other affections of the nerve or its branches. The buttock is convex toward its inner part, and in health presents on its outer side a hollow behind the great trochanter, which usually becomes lost in hip-joint disease. The back of the thigh is convex, and toward its lower part it presents a flattening, corresponding to the lozenge-shaped hollow of the popliteal space, so that the tendons and muscles forming its boundaries can be easily distinguished.

The incision through the skin should commence at the coccyx, and be continued in a semicircular direction along the crest of the ilium. Another incision should be made from the coccyx downward and outward for about six inches below the great trochanter. In reflecting the skin, notice the thick cushion which the subcutaneous adipose tissue forms over the tuberosity of the ischium, and the peculiar manner in which the fat is enclosed in meshes formed by dense connective tissue. A large bursa is often formed between this cushion and the bone.

Cutaneous Nerves. These are derived from the following sources: *Branches* from the posterior divisions of the *first* and *second lumbar nerves* descend over the crest of the ilium, near the origin of the erector spinæ, to supply the skin over the gluteus maximus as far as the great trochanter (Fig. 84, p. 389).

Branches from the posterior branches of the *three upper sacral nerves* pass downward and outward to supply for a short distance the integument over the sacrum and coccyx.

The *lateral branch* of the *twelfth dorsal nerve* descends vertically over the crest of the ilium, near to its anterior part, and supplies the integument of the front of the gluteal region.

The *iliac branch* of the *ilio-hypogastric nerve* passes over the crest of the ilium, between the posterior branches of the lumbar nerves and the preceding nerve, and supplies for a short distance the skin of the buttock.

The *external cutaneous nerve* distributes cutaneous branches to the lower and outer part of the buttock over the great trochanter.

Cutaneous branches from the *lesser sciatic nerve* proceed upward from beneath the lower border of the gluteus maximus to supply the skin over the lower part of this muscle.

Gluteal Muscles. Three powerful muscles are situated in the region of the buttock, one above the other, named, according to their size, the gluteus maximus, medius, and minimus. The fascia covering the gluteus maximus is comparatively thin, posteriorly, where it is attached to the sacrum, coccyx and ilium; but anteriorly it is very dense and glistening, and gives origin to the fibres of the gluteus medius, and lower down becomes continuous with the fascia lata.

Gluteus Maximus. This is the largest muscle of the body, and is covered by a fascia, which sends prolongations inward deeply between the muscular bundles. Its great size is characteristic of man, in reference to his erect position. Its texture is thick and course. It *arises* from the posterior fifth of the crest of the ilium, and from the rough surface below it, from the lower part of the sacrum, the coccyx, and the great sacro-sciatic ligament. The fibres descend obliquely forward, and are *inserted* thus: The anterior two-thirds terminate on a strong, broad aponeurosis which plays over the great trochanter, and joins the fascia lata on the outside of the thigh (p. 632); the remaining third is inserted into the femur, along the gluteal ridge leading from the linea aspera to the base of the great trochanter.

This muscle *extends* the thigh bone upon the pelvis, and is therefore one of those most concerned in raising the body from the sitting to the erect position, and in maintaining it erect. It propels the body in walking, running, or leaping, and rotates the thigh outward. It is supplied with blood by the gluteal and sciatic arteries; with nerves from the lesser sciatic, and the sacral plexus.

What is Seen Beneath the Gluteus Maximus The gluteus maximus should be reflected from its origin. The best way is to begin at the front border, which overlaps the gluteus medius. The

dissection is difficult, and he who undertakes it for the first time is almost sure to injure the subjacent parts. The numerous vessels which enter its under surface must be divided before the muscle can be reflected. This having been accomplished, the following objects will be exposed:—

The muscle covering the ilium is the gluteus medius. At the posterior border of this are the several objects which emerge from the pelvis through the great sciatic notch—namely, the pyriformis muscle, above which is the trunk of the gluteal vessels and nerve, and below which are the greater and lesser sciatic nerves, the arteria comes nervi ischiatici, the long pudendal nerve, the sciatic vessels, the pudic vessels and nerve, the nerve to the obturator internus and the coccygeus. Coming through the lesser sciatic notch, is the tendon of the obturator internus, and attached to it are the gemelli muscles, one above, the other below it. Extending from the tuber ischii transversely outward to the great trochanter is the quadratus femoris, and below this is seen the upper part of the adductor magnus. The origins of the semi-membranosus, biceps, semi-tendinosus, and of the adductor magnus, from the tuber ischii, are also seen; as well as the great sacro-sciatic ligament, which passes upward to the sacrum, and is pierced by the coccygeal branch of the sciatic artery. The great trochanter is exposed, together with a small portion of the vastus externus; and where the tendon of the gluteus maximus plays over the trochanter major, there is a large bursa, simple or multilocular. Lastly, the side of the sacrum, the coccyx, part of the crest of the ilium, the tuberosity of the ischium, are brought into view.

Gluteus Medius. This muscle, covered behind by the gluteus maximus, and in front by the fascia lata, *arises* from the surface of the ilium, between the crest and the upper curved line; also from the strong fascia which covers it toward the front. The fibres converge to a tendon, which is *inserted* into the oblique line on the upper and outer surface of the great trochanter; some of the anterior fibres—in immediate connection with the tensor fasciæ—terminate on the aponeurosis of the thigh. Between its insertion and the bone is a bursa.

Reflect the gluteus medius to see the third gluteal muscle.

The line of separation between them is marked by a large branch of the gluteal artery.

Gluteus Minimus. This muscle arises from the surface of the ilium below the upper and lower curved lines, and as far back as the margin of the great sacro-sciatic notch. Its fibres pass over the capsule of the hip joint, and converge to a tendon which is *inserted* into a depression on the front part of the great trochanter, a bursa being interposed. This muscle and the preceding are supplied by the superior gluteal nerve, a branch of the lumbo-sacral cord. The chief action of this and the preceding muscle is to assist in balancing the pelvis steadily on the thigh, as when we are standing on one leg; with the fixed point at the ilium, they are abductors of the thigh. The anterior fibres of the gluteus medius co-operate with the gluteus minimus and the tensor fasciæ in *rotating the thigh inward.*

Gluteal Vessels and Nerves. The *gluteal artery* is the largest branch of the internal iliac (p. 559). Emerging from the pelvis through the great sciatic foramen between the pyriformis and the gluteus medius, it divides into two large branches for the supply of the gluteal muscles. Of these, the *superficial* proceeds forward between the gluteus maximus and medius, both of which they supply, and eventually anastomose with the posterior sacral and sciatic arteries; the other, the *deep*, after a short course, divides into two branches: one, the superior, curves forward along the origin of the gluteus minimus, toward the anterior part of the ilium, to anastomose with the ascending branches of the external circumflex and the circumflex iliac arteries; the other, the inferior branch, crosses obliquely over the gluteus minimus toward the insertion of this muscle, and anastomoses with the external and internal circumflex arteries.

The nerve which accompanies the gluteal artery is the *superior gluteal nerve*, a branch of the lumbo-sacral cord. It passes out above the pyriformis, and divides into two branches, a superior and an inferior branch; the *superior branch* accompanies the corresponding branch of the gluteal artery, and supplies the gluteus medius and minimus; the *inferior branch* accompanies the inferior branch of the gluteal artery, and distributes filaments to the gluteus medius and minimus, and the tensor fasciæ femoris.

In some subjects it sends a branch to the gluteus maximus; but this muscle is chiefly supplied by the lesser sciatic nerve.

A surgeon ought to be able to cut down and tie the gluteal artery as it emerges from the pelvis. The following is the best rule[1] for finding it:—

Draw a line from the posterior superior spine of the ilium to the trochanter major, rotated inward. The junction of the upper with the middle third of this line lies over the artery as it emerges from the upper border of the great sciatic notch.

Now examine the series of muscles which rotate the thigh outward, namely, the pyriformis, the obturator internus, the gemelli, the quadratus femoris, and the obturator externus.

Pyriformis. This muscle lies immediately below and parallel with the lower fibres of the gluteus medius. It *arises* within the pelvis by three fleshy fasciculi from the second, third and fourth segments of the front surface of the sacrum between the foramina for the sacral nerves, from the margin of the great sacro-sciatic notch, and from the great sacro-sciatic ligament. The fibres, passing horizontally outward, converge to a tendon, which is *inserted* into the upper border of the great trochanter. Its nerve comes from the sacral plexus. Its action is that of an abductor and an external rotator of the femur; and, if the femur be the fixed point, it steadies the pelvis on the femur, and when the pelvis has been drawn backward it will bring it forward.

Obturator Internus. This muscle, of which little more than the tendon can be seen at present, *arises* within the cavity of the pelvis, from the inner surface of the ischium, bounded posteriorly by the margin of the great sacro-sciatic notch and the articular surface for the sacrum, and superiorly by the brim of the true pelvis; from the obturator membrane, and the obturator fascia; in front, from the inner surface of the descending ramus of the os pubis and the ascending ramus of the ischium. The fibres are directed backward and outward, and terminate on four tendons which converge toward the lesser sacro-sciatic notch; pass through the foramen of the same name at nearly right angles, as round a pulley, and then unite into a single tendon to be inserted into the

[1] The operation of tying the gluteal artery was first performed by John Bell. See his *Principles of Surgery*, vol. i, p. 421.

top of the great trochanter, in front of the pyriformis. Divide the tendon about three inches from its insertion, to see the four tendons which play over the smooth cartilaginous surface on the inner side of the tuberosity of the ischium. There is a large synovial bursa to diminish friction. The nerve to this muscle comes from the sacral plexus (sometimes from the pudic) within the pelvis; it emerges from the great sacro-sciatic foramen, winds round the spine of the ischium, and re-enters the pelvis through the lesser sacro-sciatic foramen to supply the muscle on its inner surface. The action of this muscle is to rotate the femur outward; but, in the sitting position, it loses this action, and becomes an abductor of the thigh. Between the capsule of the hip joint and the tendon, a synovial bursa is commonly found, which not infrequently communicates with the bursa placed between the tendons and the tuberosity of the ischium.

Gemelli. These small muscles are accessory to the obturator internus, and are situated, one above, the other below it. The *gemellus superior*, the smaller of the two, and occasionally absent, *arises* from the outer surface of the spine of the ischium; the *gemellus inferior* from the upper and back part of the tuberosity of the ischium. Their fibres, attached to the tendon of the obturator internus, are *inserted* with it into the upper border of the great trochanter. The nerves to these muscles come from the sacral plexus; that to the superior gemellus from the lower part of the plexus; that to the inferior gemellus comes out through the great sacro-sciatic foramen, passes beneath the superior gemellus and obturator internus to enter the muscle on its deep aspect; this branch also distributes a filament to the quadratus femoris, and another to the hip joint.

Quadratus Femoris. This quadrilateral muscle *arises* from the ridge on the outer border of the tuber ischii. Its fibres run horizontally outward, and are *inserted* into the back of the great trochanter, into the greater part of the linea quadrati. The lower border of the quadratus femoris runs parallel with the upper edge of the adductor magnus; in fact, it lies on the same plane. Between these muscles is generally seen a terminal branch of the internal circumflex artery. Its nerve, as previously described, comes from the sacral plexus, and enters its deep surface.

Obturator Externus. To see this muscle, reflect the quadratus femoris from its origin. It *arises* from the outer surface of the body of the os pubis, from the front surface of the rami of the os pubis and ischium, which form the inner border of the obturator foramen, from the inner two-thirds of the outer surface of the obturator membrane, and from the tendinous arch over the obturator vessels. Its fibres converge to a tendon which runs

FIG. 152.



12. N. of pyriformis.
13. N. of gemellus superior.
14. N. of gemellus inferior.
15. N. of quadratus femoris.
16. N. of gluteus maximus.
17. Long pudendal n.
18. Cutaneous n. of the buttock.
19. N. of the long head of the biceps.
20. N. of semi-tendinosus.
21. N. of semi-membranosus.
22. N. of short head of the biceps.

1, 2, 3, 4, 5. Sacral nn.
6. Superior gluteal n.
7. Great sciatic n.
8. Lesser sciatic n.
9. Pudic n.
10. N. of obturator internus.
11. N. of levator ani.

PLAN OF THE SACRAL PLEXUS AND BRANCHES.

horizontally outward over a groove on the ischium, and, running across the back of the hip joint, is *inserted* into the deepest part of the trochanteric fossa of the femur. Its nerve is a branch of the posterior division of the obturator nerve. This muscle has *in front* of it the adductor longus and brevis, the pectineus, the psoas and iliacus, the neck of the femur and the capsular ligament; *above* it are the capsular ligament and the inferior gemellus; *below* it

re the adductor magnus and quadratus femoris; *behind* it are the bturator membrane and the quadratus femoris.

Great Sciatic Nerve. This large nerve, formed by the union of the last lumbar and the four upper sacral nerves (Fig. 52), is the largest nerve in the body, being three-quarters of an ch in breadth, and supplies all the flexor muscles of the lower xtremity and the extensors of the foot.

Emerging from the pelvis through the great sacro-sciatic ramen below the pyriformis, it descends over the external rotator uscles of the thigh, along the interval between the tuber ischii nd the great trochanter, but rather nearer to the former; so that,

FIG. 153.

1. Gluteus medius.
2. Pyriformis.
3. Lesser sciatic nerve.
4. Obturator internus, with the two gemelli.
5. Coccygeus.
6. Great sciatic nerve.
7. Quadratus femoris.
8. Gluteus maximus.
9. The semitendinosus and biceps.
10. Adductor magnus.

DEEP MUSCLES OF THE GLUTEAL REGION.

n the sitting position, the nerve is protected from pressure by this ony prominence. The nerve does not descend quite perpendicularly, but rather obliquely forward upon the adductor magnus, arallel with the great sacro-sciatic ligament, and below the middle f the thigh divides into the internal popliteal and the peroneal or external popliteal). It is accompanied by a branch of the ciatic artery, called the *comes nervi ischiatici*.[1] The nerve dis-

[1] The arteria comes nervi ischiatici runs generally by the side of the nerve, but ometimes in the centre of it. This artery becomes one of the chief channels by hich the blood reaches the lower limb after ligature of the femoral. See in the Iuseum of the Royal College of Surgeons a preparation in which the femoral was ed by John Hunter fifty years before the man's death.

tributes branches to the hamstring muscles and the adductor magnus, and sends two or more small branches to the hip joint which pierce the posterior part of the capsular ligament.

Small Sciatic Nerve. This comes from the lower part of the sacral plexus. It leaves the pelvis below the pyriformis, with the great sciatic nerve, but on the inner side of it, and in company with the sciatic artery. It descends behind the gluteus maximus, and becomes cutaneous at its lower border. The muscular branches which it gives off are one or more—*inferior gluteal*—which enter the under surface of the gluteus maximus near its lower border. All its other branches are cutaneous, and are divided into an ascending and internal group; the *ascending branches* turn round the lower border of the gluteus maximus, and supply the skin of the buttock; the *internal branches* supply the skin on the inner and posterior aspect of the thigh in its upper part, and one branch, larger than the rest, called the *inferior pudendal*, turns inward toward the perineum to supply the skin of that region and the scrotum, communicating with the inferior hæmorrhoidal and superficial perineal nerves. The continued trunk runs down the back of the thigh beneath the muscular fascia, as low as the upper part of the calf with the external saphenous vein, supplying the skin all the way down, and communicates with the short saphenous nerve.

Sciatic Artery. This, one of the terminal branches of the internal iliac, courses along the inner aspect of the sacral plexus and pyriformis, behind the pudic artery, while this vessel is still within the pelvis. It emerges from the pelvis between the pyriformis and coccygeus, and is then seen in the gluteal region coming out between the pyriformis and superior gemellus. It then descends between the tuber ischii and the great trochanter, along the inner side of the great sciatic nerve. It gives off: (1) within the pelvis, branches to the muscles which form the muscular floor of the pelvis, to the rectum, the bladder, prostate and vesiculæ seminales; (2) external to the pelvis, it gives off: *a*, a *coccygeal branch*, which runs inward through the great sacro-sciatic ligament, then ramifies in the gluteus maximus, and on the back of the coccyx; *b*, the *comes nervi ischiatici*, which accompanies the great sciatic nerve for a short distance, and then

enters its substance; *c*, the *inferior gluteal branches*, which enter the gluteus maximus; *d*, *articular branches*, which pierce the posterior capsule of the hip joint; *e*, *muscular branches* to the several external rotators and the hamstring muscles, and which inosculate with the external and internal circumflex, gluteal, obturator, and first perforating arteries.

Pudic Artery and Nerve. The course of this artery and nerve has been fully described (p. 561). Observe now that they pass over the spine of the ischium, accompanied by the nerve to the obturator internus, and that in a thin subject it is possible to

FIG. 154.



1. Gluteal artery and nerve.
2. Pudic artery and nerve, and nerve to obturator internus.
3. Great sacro-sciatic nerve.
4. Sciatic artery.
5. Internal circumflex artery.
6. The first perforating artery.

THE ARTERIES OF THE GLUTEAL REGION.

compress the artery against the spine. The rule for finding it is this: rotate the foot inward, and draw a line from the top of the great trochanter to the base of the coccyx; the junction of the inner with the outer two-thirds gives the situation of the artery.[1]

Popliteal Space; its Boundaries. It is advisable to examine the popliteal space at this stage of the dissection, in order that the various parts may be carefully made out with as little disturbance as possible of their mutual relations.

[1] Mr. Travers succeeded in arresting hemorrhage from a sloughing ulcer of the glans penis by pressing the pudic artery with a cork against the spine of the ischium.

Surface Marking. The popliteal space is a lozenge-shaped hollow at the back of the knee-joint, extending as high as the junction of the middle with the lower third of the femur, and as low as the upper sixth of the tibia. The hollow is most apparent when the knee is flexed, as then the tendinous boundaries stand out in bold relief; it is almost lost when the leg is extended. The tendon on the outer side is that of the biceps, diverging to the head of the fibula; on the outer side, and below, are the plantaris and outer head of the gastrocnemius, which are not well defined; on the inner side, above, we can feel three tendons in the following order from within outward—the semitendinosus, the semimembranosus, and the gracilis; below, on their inner side, is the inner head of the gastrocnemius. The upper angle of this space is formed by the diverging biceps and semitendinosus; the lower angle by the converging heads of the gastrocnemius. Passing from above downward in the middle of the space, and in the following order, are the internal popliteal nerve, the popliteal vein and artery; and along the inner border of the biceps can be felt the external popliteal nerve. Filling up the hollow is a quantity of soft fat, with some lymphatic glands, and on the bone rest the articular arteries.

Dissection. A vertical incision must be made along the middle of the ham, extending from six inches above to three inches below the knee; transverse incisions should be made at each extremity of the vertical, so that the skin may be conveniently reflected. In doing so, care must be taken to preserve the cutaneous branch of the lesser sciatic nerve, which descends over the space to the back of the leg.

The muscular fascia covering the space is very strong, and strengthened by numerous transverse fibres. It is pierced by the posterior saphena vein, which passes in to join the popliteal vein.

The fascia having been reflected, the muscles and tendons constituting the boundaries of the popliteal space are to be cleaned. The boundaries of the space can now be seen to be formed, as before stated, above, by the divergence of the hamstring muscles to reach their respective insertions; below, by the converging heads of the gastrocnemius; its shape is, therefore, that of a lozenge. Above, it is bounded on the inner side by the semiten-

dinosus, semimembranosus, gracilis and sartorius; on the outer side, by the biceps; below, it is bounded, on the inner side by the internal head of the gastrocnemius; on the outer, by the external head of this muscle and the plantaris.

The space is occupied by a quantity of fat, which permits the easy flexion of the knee; and in this fat are found the popliteal vessels and nerves, in the following order: nearest to the surface

FIG. 155.



LEFT POPLITEAL SPACE.

are the nerves; the artery lies close to the bone, the vein being superficial to the artery (Fig. 155).

Great Sciatic Nerve. Along the outer border of the semimembranosus, and covered by the long head of the biceps, is the great sciatic nerve, which, after giving off branches to the three great flexor muscles and the adductor magnus, divides, about the lower third of the thigh (higher or lower in

different subjects), into two large nerves—the peroneal or external popliteal and the internal popliteal.

The *peroneal nerve* runs close by the inner side of the tendon of the biceps,[1] and subsequently in the groove between this muscle and the outer head of the gastrocnemius, toward the head of the fibula. As it passes round the joint it gives off *two articular* branches to the outer side of the knee, which accompany the external superior and inferior articular arteries, and a *recurrent articular* branch, which runs with the recurrent tibial artery to the front of the knee. It supplies also two or three *cutaneous branches* to the posterior and outer surfaces of the leg, as far as its middle third.

The *communicans peronei* (Fig. 155) is a small branch given off as the nerve passes over the gastrocnemius; it crosses the outer head of this muscle and joins the external saphenous, which runs down the back of the calf, and behind the external malleolus, to supply the outer side of the foot and little toe.

Below the head of the fibula we have already traced the division of the peroneal into the anterior tibial, and the musculo-cutaneous nerves (p. 669).

The *internal popliteal nerve*, the larger of the two divisions of the great sciatic nerve, accompanies the popliteal artery, and, at the lower border of the popliteus, it is continued under the name of the posterior tibial. The nerve in the popliteal space lies superficial to and rather external to the artery, and gives off four or five *muscular branches* which supply the two heads of the gastrocnemius, the plantaris, the soleus, and the popliteus; *three articular* branches, two accompanying the internal superior and inferior articular arteries, the third piercing the back of the capsule accompanied by the azygos artery; and the *short* or *external saphenous*, which descends in the groove between the two heads of the gastrocnemius, is joined about the middle of the leg by the communicans peronei, and then, running down behind the outer malleolus in company with the external saphena vein, is distributed to the outer side of the foot and the little toe. The con-

[1] The nerve is, therefore, very liable to be injured in the operation of dividing the outer hamstring. In the diagram, the nerve is not near enough to the tendon, their connections having been severed.

tinuation of the internal popliteal nerve, as posterior tibial, supplies all the flexor muscles on the back of the leg and the sole of the foot.

Popliteal Vessels. By clearing out all the fat, we observe that the popliteal vessels enter the ham through an aperture in the adductor magnus, and descend close to the back part of the femur, and the back of the knee joint. At first they are partially overlapped (in muscular subjects) by the semimembranosus; indeed, the outer border of this muscle is a good guide to the artery in the operation of tying it. The *popliteal artery* lies upon the triangular surface at the back of the lower third of the femur; then, upon the ligamentum posticum Winslowii; and, lastly, upon the popliteus, at the lower border of which it divides into the anterior and posterior tibial.

Superficial to the artery are the semimembranosus, a considerable amount of fat, the gastrocnemius, the plantaris, the soleus, the popliteal vein, and the internal popliteal nerve; *internally*, it has the semimembranosus, the internal condyle of the femur, and the inner head of the gastrocnemius; *externally*, it has the biceps, the external condyle, the outer head of the gastrocnemius, and the plantaris.

The artery gives off the *external* and *internal superior articular arteries;* lower down, the *external* and *internal inferior articular arteries*, the *superior* and *inferior muscular branches*, the *azygos*, and *cutaneous branches*. The description of these branches of the popliteal will be deferred till later, until the muscles of the calf have been reflected.

The articular branches which come from the popliteal are given off at right angles to that vessel; and besides these it gives off the *sural*, which supply the muscles of the calf, and the azygos artery; close to the vessel is the articular branch of the obturator nerve which supplies the knee joint.

The *popliteal vein* lies superficial to the artery, and rather to its outer side. It receives the short saphena vein. Its coats are remarkably thick, and on transverse section resemble those of an artery of similar size.

Lymphatic Glands. Two or more lymphatic glands are situated one on each side of the artery. They deserve atten-

tion, because, when enlarged, their close proximity to the artery may communicate a pulsation which might be mistaken for an aneurism.

DISSECTION OF THE BACK OF THE THIGH.

Dissection. The incision should be continued along the remainder of the back of the thigh, and the skin reflected. The fat should be removed, and the cutaneous branches derived from the external and internal cutaneous nerves and the small sciatic should be carefully sought out.

Cutaneous Nerves and Veins. The skin on the middle of the back of the thigh is supplied by the small sciatic nerve, which runs down beneath the deep fascia as far as the middle third; then pierces it, and runs down as far as the middle third of the calf, distributing branches on each side. On the outer side, a few cutaneous branches from the *posterior division* of the *external cutaneous nerve* supply the skin as far as the middle third; on the inner side are small branches from the small sciatic and the internal cutaneous nerves as low as the knee joint.

The subcutaneous veins at the back of the thigh are very small; here they would be liable to pressure. But near the popliteal space there is a vein, called the *external* or *short saphena*. It comes up the back of the calf, and joins the popliteal vein after perforating the strong fascia covering the space.

Muscular Fascia. Respecting this, remark that its fibres run chiefly in a transverse direction, that it becomes stronger as it passes over the popliteal space, and that here it is connected with the tendons on either side. Remove it, to examine the powerful muscles which bend the leg, called the hamstrings.

Hamstring Muscles. There are three of these, and all arise by strong tendons from the tuber ischii. One, the biceps, passes downward and outward, to be inserted into the head of the fibula; the other two—namely, the semitendinosus and semimembranosus—descend inward and are inserted into the tibia. The divergence of these muscles toward their respective insertions occasions the space termed the *popliteal*, which is occupied by soft fat, the popliteal vessels, nerves, and lymphatic glands.

Biceps. This muscle has two origins, a long and a short. The *long head arises*, by a strong tendon, from the back part of the tuber ischii in common with the semitendinosus; the *short head*, by fleshy fibres, from the outer lip of the linea aspera of the femur between the vastus externus and the adductor magnus, and from the external intermuscular septum; this origin begins at the linea aspera, just below the insertion of the gluteus maximus, and continues nearly down to the external condyle. It joins the long head of the muscle, and both terminate on a common tendon, which is *inserted* into the outer side of the head of the fibula, by two portions separated by the external lateral ligament of the knee joint. It also gives off a strong expansion to the fascia of the leg. The tendon covers part of the external lateral ligament of the knee joint, and a small bursa intervenes.

The biceps is not only a flexor of the leg, but rotates the leg, when bent, outward. It is the muscle which in chronic disease of the knee dislocates the leg outward and backward, and at the same time rotates it outward. Each head of the biceps is supplied by the great sciatic nerve. The short head is sometimes supplied by the peroneal.

Semitendinosus. This *arises*, in common with the biceps, from the back part of the tuber ischii, by muscular fibres, and also from the inner border of the tendon of the biceps, for about three inches. The muscle passes downward and inward, and terminates in the middle of the thigh, in a long round tendon, which rests upon the semimembranosus, and is *inserted* into the upper part of the inner surface of the tibia by an expanded tendon, below the tendon of the gracilis, and behind that of the sartorius. Like them, it plays over the internal lateral ligament of the knee, and is provided with a bursa. Its nerve comes from the great sciatic.

The semitendinosus sends off from the lower border of its tendon a very strong fascia to cover the leg, which is attached along the inner edge of the tibia. The middle of the muscle is usually intersected by an oblique tendinous line.

Semimembranosus. This muscle *arises* from the upper and outer facet on the back of the tuber ischii, above and

times high up, occasionally lower down than usual. The further course of this nerve has already been described (p. 681).

Deferring the course, relations, and branches of the popliteal artery till this vessel is exposed throughout its whole course, pass on now to the dissection of the calf.

DISSECTION OF THE BACK OF THE LEG.

Surface Marking. The back of the leg gradually narrows from above downward so as to form a long cone; the upper half is convex and fleshy, corresponding to the gastrocnemius and the soleus muscles; the lower half suddenly diminishes, so that the posterior borders of the tibia and fibula can be easily felt extending to their respective malleoli. In a well-developed subject with not much fat the two heads of the gastrocnemius can be seen through the skin, the inner head being the broader and lower of the two; the tendon into which they are inserted (tendo-Achillis) rapidly narrows to be attached to the posterior and upper part of the tuberosity of the os calcis. On the inner and outer sides of the lower part of this tendon there is a well-defined vertical groove; bounded laterally by the tibia and fibula. In cases of synovial disease of the ankle joint these grooves are lost, so that instead of a depression there is a convexity. In these grooves can be felt the tendons passing round the ankle behind the malleoli; on the inner side are the tibialis posticus, flexor longus digitorum, and flexor longus hallucis; on the outer side are the peronei longus and brevis.

Continue the incision down the centre of the calf to the heel, where a transverse incision must also be made. The skin should now be reflected, taking care of the subcutaneous veins and nerves.

Short or Posterior Saphena Vein. The large vein seen in the middle of the back of the leg is called the *short* or *posterior saphena*. It commences on the outer side of the dorsum of the foot, ascends behind the outer ankle, where it has a communication with the deep veins, and then runs up the calf between the two bellies of the gastrocnemius, receiving numerous veins in its course, and being guarded by several valves. It eventually passes through the muscular fascia, and joins the popliteal vein.

ıtaneous Nerves. The back of the leg is supplied by cutaneous nerves; in the middle, above, by the small sciatic :rve, and below, by the short or external saphenous; on the ıter side, by the communicans peronei; and on the inner side, ' branches from the internal saphenous nerve.

hort or External phenous Nerve. The *short saphenous* nerve[1] is derived from the internal popliteal nerve (Fig. 155), and passes ›wn beneath the deep fascia between the two heads of the gas- ɔcnemius to the middle of the calf, where it pierces the fascia. ere it is joined by a branch from the peroneal nerve (*communicans ronei*); it then descends with the short saphena vein, usually ı its fibular side, and is finally distributed to the outer side of e foot and the little toe.

To expose the muscles of the calf, reflect the muscular fascia / incisions corresponding to those made through the skin.

Muscles of the Calf. The great flexor muscle of the foot consists of two portions: the superficial one, called the gas- ɔcnemius, *arises* from the lower end of the femur; the deep one, lled the soleus, *arises* from the tibia and fibula. They are tached to one thick tendon, called the tendo-Achillis, which is *serted* into the os calcis.

Gastrocnemius. This muscle arises by two strong tendinous heads from the back of the condyles of the femur ig. 155). The inner head is the larger, longer and more mus- lar, and *arises* from a depression at the upper and back part of e internal condyle, and, for a short distance, by fleshy fibres ›m the line leading from the linea aspera to the internal condyle; e outer head from the back and upper part of the external con- le above the popliteus, and also from the line leading to the ıea aspera. The two parts of the muscle descend, distinct from ch other, and form the two bellies of the calf, of which the inner rather the lower. Both *terminate*, rather below the middle of e leg, on the broad commencement of the tendo-Achillis.

The gastrocnemius should be divided transversely near its tachment to the tendo-Achillis, and reflected upward from the

[1] This nerve is sometimes called the *communicans poplitei*, and does not take name of short saphenous nerve till its junction with the *communicans peronei* . 682).

subjacent soleus, as high as its origin. By this proceeding you observe that the contiguous surfaces of the muscles are covered by a glistening tendon, which receives the insertion of their fibres, and transmits their collected force to the tendo-Achillis.

Observe also the large sural vessels and nerves (branches of the popliteal) which enter the mesial aspect of each head of the muscle. To facilitate the play of the inner tendon over the condyle, there is a bursa, which generally communicates with the knee joint; and in the substance of the outer tendon is commonly found a small piece of fibro-cartilage. Lastly, between the gastrocnemius and soleus is the tendon of the plantaris.

Plantaris. This small muscle[1] arises from the rough line leading from the linea aspera to the outer condyle of the femur, and from the posterior ligament of the knee joint. It descends close to the inner side of the outer head of the gastrocnemius, having a fleshy belly for about two inches, and terminates a little below the knee, in a long, thin tendon, which can be traced down the inner border of the tendo-Achillis to the calcaneum. This muscle is occasionally absent. Its nerve comes from the internal popliteal.

Soleus. This muscle arises by tendinous fibres from the head and from the upper third of the posterior surface of the fibula, from the oblique ridge on the back of the tibia,[2] from about the middle third of the inner border of this bone, and from an aponeurotic arch thrown over the posterior tibial vessels. The muscular fibres bulge out beyond the gastrocnemius, and terminate on a broad tendon, which, gradually contracting, forms a constituent part of the tendo-Achillis. The muscle lies upon the flexor longus digitorum, the tibialis posticus, the flexor longus hallucis, and the posterior tibial vessels and nerve. The soleus is supplied with blood by several branches from the posterior tibial; also by a large branch from the peroneal. Its nerve

[1] This is the representative of the palmaris longus of the forearm. In man it is lost on the calcaneum, but in monkeys, who have prehensile feet, it is the proper tensor muscle of the plantar fascia. It is remarkably strong in bears and plantigrade mammals.

[2] The tibial and fibular origins of the soleus constitute what some anatomists describe as the two heads of the muscle. Between them descend the popliteal vessels, protected by a tendinous arch.

comes from the internal popliteal and enters the top of the muscle. This is an important muscle in a surgical point of view, for two reasons—(1) by reflecting its tibial origin, we can reach the posterior tibial artery; (2) by reflecting its fibular origin we can reach the peroneal.

The *tendo-Achillis*, the common tendon of the gastrocnemius and soleus, begins about the middle of the leg, and is at first of considerable breadth, but it gradually contracts and becomes thicker as it descends. The narrowest part of it is about one inch and a half above the heel; here, therefore, it can be most conveniently and safely divided, for the relief of club-foot. There is no risk of injuring the deeper-seated parts, because they are separated from the tendon by a quantity of fat. Its insertion is into the under and back part of the tuberosity of the os calcis. The tendon previously expands a little; between it and the bone is a bursa of considerable size.

The *action* of the gastrocnemius and soleus is to raise the body on the toes. Since the gastrocnemius passes over two joints, it has the power (like the rectus) of extending the one while it bends the other, and it is, therefore, admirably adapted to the purpose of walking. For instance, by first extending the foot it raises the body; and then, by bending the knee, it transmits the weight from one leg to the other. Supposing the fixed point to be at the heel, the gastrocnemius is also concerned in keeping the body erect, for it keeps the tibia and fibula perpendicular on the foot, and thus counteracts the tendency of the body to fall forward.

The tendo-Achillis, in pointing the toes, acts upon a lever of the *first* order. The fulcrum is at the ankle joint, F (Fig. 157); the resistance, W, at the toes; the power at the heel, P. All the conditions are those of a lever of the first order. The power and the weight act in the *same* direction on *opposite* sides of the fulcrum. In raising the body on tiptoe, the tendo-Achillis acts as a lever of the second order, the fulcrum being then at the ball of the great toe, and the weight of the body at the ankle.

FIG. 157.



Course and Relations of the Popliteal Artery.

After passing through the opening in the tendon of the adductor magnus, the femoral artery takes the name of *popliteal*. It descends nearly perpendicularly behind the knee joint, between the origins of the gastrocnemius, as far as the lower border of the popliteus, where it divides into the anterior and posterior tibial. In its descent it *lies*, first, upon the lower part of the femur, and here it is slightly overlapped by the semimembranosus; next, it lies upon the posterior ligament of the knee joint; and, lastly, upon the popliteus. At its lower part the artery is covered, at first, by the semimembranosus, the popliteal vein, the internal popliteal nerve, and a considerable amount of fat; then, lower down, by the gastrocnemius and soleus, and it is crossed by the plantaris. The vein closely accompanies the artery, and is situated superficially with regard to it, and rather to its outer side in the first part of its course. The internal popliteal nerve runs also in a similar direction with the vein, but is still more superficial and to the outer side (Fig. 155). The vessels and the nerve are surrounded by fat, and one or two lymphatic glands are generally found in the immediate neighborhood of the artery, just above the joint.

The branches of the popliteal artery are—the *articular*, the *sural*, and the *cutaneous*.

There are five *articular branches* for the supply of the knee joint and the articular ends of the bones; the two *superior*, *external* and *internal*, run, one above each condyle, close to the bone; the two *inferior*, *external* and *internal*, run below the joint.

1. The *superior external articular* artery runs above the external condyle, passes beneath the biceps, and through the intermuscular septum; it then divides into a superficial and a deep branch; the superficial supplies the vastus externus, and then forms part of the patellar arterial plexus; the deep branch keeps close to the femur and supplies the knee, forming an arch above the joint, with a branch from the anastomotica magna.

2. The *superior internal articular* artery runs above the internal condyle, under the tendon of the adductor magnus and vastus internus, and divides into two branches, a superficial

and a deep, which take a corresponding course to those on the outer side.

3. The *inferior external articular* artery runs under the gastrocnemius, over the popliteus, then, passing beneath the external lateral ligament and the tendon of the biceps, it reaches the patella, where it breaks up into branches anastomosing with the other articular arteries, and the recurrent branch of the anterior tibial artery.

4. The *inferior internal articular* artery runs between the tuberosity of the tibia and the internal lateral ligament, and supplies the inner and anterior part of the joint.

5. The *azygos* artery is given off from the deep aspect of the popliteal, pierces the ligamentum posticum Winslowii, to supply the crucial ligaments and the synovial membrane.

The several articular arteries form, over the front and sides of the joint, a network of vessels, which anastomose, superiorly, with the descending branch of the external circumflex and the anastomotica magna; inferiorly, with the anterior tibial recurrent; and also among themselves. It is mainly through these channels that the collateral circulation is established in the leg after ligature of the superficial femoral.

The *sural arteries* proceed one to each head of the gastrocnemius, and are proportionate in size to the muscle; one or two branches are distributed to the soleus. These arteries are accompanied by branches of the internal popliteal nerve for the supply of the muscles.

The *superior muscular* branches supply the vasti and hamstring muscles, and inosculate with the perforating and articular arteries.

Cutaneous arteries pass down between the gastrocnemius and the skin, supplying the integument of the calf, as far as the middle of the leg.

Popliteal Vein. This vein is formed by the junction of the venæ comites of the anterior and posterior tibial arteries, and is situated superficial to the artery. It crosses obliquely from the inner to the outer side of the artery, and is continued upward as the femoral. It receives in the popliteal space the short saphena, the articular and sural veins. It is usually provided with four valves.

The insertion of the tendon of the semimembranosus into the head of the tibia, and its several connections (described p. 686), should now be fully examined.

Popliteus. This triangular muscle *arises* within the capsule of the knee joint, from a depression on the outside of the external condyle, and from the posterior ligament of the knee, by a thick tendon, which runs beneath the external lateral ligament and the tendon of the biceps. The muscular fibres gradually spread out, and are *inserted* into the triangular surface of the tibia above the soleal ridge on the bone, and into the aponeurotic expansion covering the muscle. It is supplied by a branch of the internal popliteal nerve which enters its deep surface. Its action is to flex the leg, and then to rotate the tibia inward. The tendon plays over the articulation between the tibia and fibula; and a bursa intervenes, which generally communicates by a wide opening with the knee joint. The tendinous origin grooves the external semilunar cartilage, and has an investment from the synovial membrane of the knee.

Reflect the soleus from its origin, and remove it from the deep-seated muscles, observing at the same time the numerous arteries which enter its under surface. This done, notice the deep fascia which separates the superficial and the deep layers of muscles and which binds down the deep muscles. It is attached to the margins of the tibia and the fibula on either side, increases in strength toward the ankle, and forms an *internal annular ligament* which confines the tendons and the vessels and nerves in their passage into the sole of the foot.

The deep fascia should now be removed from the deep muscles which must be cleaned in the course of their muscular fibres.

Deep Muscles on the Back of the Leg. There are three: the flexor longus digitorum on the tibial side; the flexor longus hallucis on the fibular; the tibialis posticus upon the interosseous membrane, between and beneath them both.

Flexor Longus Digitorum. This *arises* from the posterior surface of the tibia, commencing below the popliteus, and extending to within four inches of the lower end of the bone, also from the fascial septum between it and the tibialis posticus. The fibres terminate on a tendon which is placed on the tibial side of

the muscle, and it runs through a groove behind the inner ankle, lined by a special synovial sheath. It enters the sole and divides into four tendons, which are inserted into the ungual phalanges of the four outer toes. It is supplied by the posterior tibial nerve.

Flexor Longus Hallucis. This powerful muscle *arises* from the lower two-thirds of the posterior surface of the fibula, from the septum between it and the peronei, from the lower part of the interosseous membrane, and from the aponeurosis over the tibialis posticus. The fibres terminate on a tendon which runs through a groove on the back of the lower extremity of the tibia, then through the groove on the back of the astragalus; thence it passes through another groove on the under aspect of the sustentaculum tali, and is *inserted* into the ungual phalanx of the great toe. The chief action of this muscle is to raise the body on the tip of the great toe. It is essential to the propulsion of the body in walking. It is supplied by the posterior tibial nerve.

Tibialis Posticus. This is so concealed between the two preceding muscles that it cannot be properly examined without reflecting them. It *arises* by two processes from the interosseous membrane, between which the anterior tibial artery passes forward, from the opposite surfaces of the tibia and fibula for about their middle three-fifths, and from the aponeurosis covering it. In the lower part of the leg it passes beneath the tibia and the flexor longus digitorum. Its muscular fibres terminate on a tendon which comes into view a short distance above the inner ankle, and, running through the same groove with the tendon of the flexor longus digitorum, enters the sole, and is *inserted* into the scaphoid and internal cuneiform bones, and by fibrous prolongations into most of the tarsal and metatarsal bones. Its action is to bend and turn the foot inward. It is supplied by the posterior tibial nerve. The precise situation of the tendon of the tibialis posticus is interesting, surgically, because the tendon has to be divided for the relief of talipes varus. It lies close to and parallel with the inner edge of the tibia, so that this is the guide to it. It is necessary to relax the tendon while the knife is introduced between the tendon and the bone. Its synovial sheath commences about 1½ inch above the end of the internal malleolus, and is consequently opened in the operation.

Attention should now be directed to the *internal* or *posterior annular ligament*, which binds down the tendons behind the inner ankle.

It is attached to the internal malleolus and the inner border of the os calcis. It is continuous, above, with the deep fascia of the leg, below, with the plantar fascia and the abductor hallucis. Beneath it are a series of compartments through which pass the tendons of the deep-seated muscles of the leg and the vessels into the sole of the foot. The relative positions of the structures passing under this ligament, proceeding from within outward, are, the tendons of the tibialis posticus, and the flexor longus digitorum; the posterior tibial artery accompanied by its venæ comites; the posterior tibial nerve; and, lastly, the tendon of the flexor longus hallucis. Each of the tendons is lined by a separate synovial membrane, as they pass behind the inner ankle.

Course and Relations of the Posterior Tibial Artery. This artery is one of the branches into which the popliteal divides at the lower border of the popliteus. It descends between the superficial and the deep muscles at the back of the leg, to the interval between the internal malleolus and the os calcis, and, entering the sole, divides beneath the abductor hallucis into the external and internal plantar arteries. It lies, first, for a short distance, *upon* the tibialis posticus, then, on the flexor longus digitorum; but behind the ankle it is in contact with the tibia, so that here it can be felt beating, and effectually compressed; and lastly, on the back of the ankle joint. In the upper part of its course it runs nearly midway between the bones, and is *covered* by the gastrocnemius and soleus, and is crossed obliquely from within outward by the posterior tibial nerve. To tie the artery, therefore, in this situation, is difficult; but in the lower part of its course it gradually approaches the inner border of the tibia, from which, generally speaking, it is not more than half or three-quarters of an inch distant. Here, being comparatively superficial, it may easily be tied. Immediately behind the internal malleolus, it lies between the tendons of the flexor longus digitorum on the inner side, and the flexor longus hallucis on the outer. It has two venæ comites, which communicate at intervals. The posterior tibial nerve which accompanies the artery is at its upper third

its *inner side*, then crosses over it, and for the lower two-thirds of its course lies *external* to the artery. Its branches are as follows:—

a. Numerous *muscular* branches to the soleus, the peronei, and the deep muscles.

b. The *peroneal* is a branch of considerable size—often as large as the posterior tibial. Arising about an inch below the division of the popliteal, it descends close to the interosseous border of the fibula, and then over the articulation between the tibia and fibula to the outer part of the os calcis, where it inosculates with the malleolar and plantar arteries. All down the leg it is imbedded among the muscles, being covered, first, by the soleus, afterward, by the flexor longus hallucis, by the fibres of which it is more or less surrounded. To both these muscles, to the latter especially, it sends numerous branches, and just above the ankle it gives off a constant one, the *anterior peroneal*, which passes through the interosseous membrane to the under aspect of the peroneus tertius, then runs in front of the inferior tibio-fibular articulation, and inosculates with the other malleolar and tarsal arteries. The peroneal supplies the *nutrient* artery of the fibula.

c. The *communicating artery* passes transversely across the interosseous membrane, about an inch above the os calcis. It runs under the tendon of the flexor longus hallucis, and anastomoses with the peroneal artery.

d. The *internal calcanean* arteries, several in number, run down, ramifying over the posterior and inner aspect of the tendo-Achillis and os calcis; they anastomose with the internal malleolar and peroneal arteries.

e. The *nutrient artery* to the tibia enters the bone about an inch below the oblique line, the foramen through which it passes being directed away from the knee joint.

The *posterior tibial veins*, formed by the junction of the external and internal plantar veins, accompany the artery as *venæ comites*, and, after receiving the peroneal veins, join with the veins corresponding to the anterior tibial artery, at the lower border of the popliteus, to form the popliteal vein.

Posterior Tibial Nerve. This is the continuation of the popliteal. It descends close to its corresponding artery, and,

behind the inner ankle, divides into the external and internal plantar nerves. In the first part of its course the nerve lies superficial to the artery, and rather to its inner side; but lower down the nerve crosses the artery, and passes to its outer side, and lies to the inner border of the tendo-Achillis. It supplies branches to the three deep-seated muscles, the branch to the flexor longus hallucis accompanying the peroneal artery; and a cutaneous branch, *calcaneo-plantar*, which pierces the internal annular ligament, and supplies the skin of the heel and the inner side of the sole of the foot.

THE DISSECTION OF THE SOLE OF THE FOOT.

Surface Marking. The skin of the sole is remarkably thick, especially over the os calcis and the heads of the metatarsal bones. The sole of the foot is convex and narrow behind, but gradually increases in breadth forward to the clefts of the toes. The inner arch of the foot extending from the inner tuberosity of the os calcis to the distal end of the first metatarsal bone, is well marked in a well-developed foot; along this arch can be distinguished the sustentaculum tali; about an inch in front of this is the prominent tubercle of the scaphoid, and another inch in front of this we can feel the articulation between the internal cuneiform and the base of the metatarsal bone of the great toe. The outer arch of the foot is neither so deep nor so long as the inner; it extends from the external tubercle of the os calcis to the base of the metatarsal bone of the little toe, and along it we can recognize the peroneal tubercle of the os calcis and the base of the fifth metatarsal bone. The transverse arch, between the heads of the metatarsal bones, is not well marked through the skin, for it is obscured by the tense plantar fascia. The plantar fascia can be made tense by extending the foot, when its narrowest part is seen to be about one inch in front of the os calcis. The course of the external plantar artery is indicated by a line drawn from the internal tubercle of the os calcis to the base of the fifth metatarsal bone; that of the internal plantar artery by a line drawn from the same tuberosity to the first interosseous space.

Dissection. The foot should be firmly fixed to a block, with the sole directed toward the dissector, and the

toes either fastened by strings or nailed to the block, so as to put the plantar fascia on the stretch. Make a perpendicular incision down the middle of the sole; another transverse one across the foot at the clefts of the toes, and continue the perpendicular incision along the middle of the toes to their terminations. Reflect the skin, and notice the peculiar structure of the subcutaneous tissue. It is composed of globular masses of fat, separated by strong, fibrous septa, and forms elastic pads, especially marked at the heel, and at the ball of the great and the little toes; these being the points which form the tripod supporting the arch of the foot.

In removing the subcutaneous tissue from the ball of the great and the little toes, we often meet with bursæ, simple or multilocular. They are generally placed between the skin and sesamoid bones, and have remarkably thick walls. Frequently an artery and nerve can be traced running directly through one of these sacs, which explains the acute pain produced by their inflammation.

Cutaneous Nerves. In the fat the student must make out the *cutaneous branch* of the posterior tibial nerve, which supplies the skin of the sole of the foot and the heel; the remainder of the sole is supplied by small branches of the plantar nerves which come through the fascia, as in the palm of the hand.

Plantar Fascia. This is a remarkably dense, white and glistening fascia. It extends from the under and back part of the os calcis to the distal extremities of the metatarsal bones. It is divided into a strong central and two lateral less dense portions; from which prolongations pass deeply inward, separating the lateral from the central muscles. The *middle portion*, covering the flexor brevis digitorum, is narrow behind, and, as it passes forward toward the toes, is spread out, and strengthened by transverse fibrous bands. The *inner portion* is comparatively thin, and surrounds the abductor hallucis, becoming continuous posteriorly with the internal annular ligament. The *outer portion* is thicker than the inner, especially as it passes forward to be attached to the proximal end of the fifth metatarsal bone. It covers the abductor minimi digiti. Both the inner and the outer portions are continuous with the fascia of the dorsum of the foot round the inner and outer borders of the

foot, and with the central portion of the plantar fascia toward the centre of the sole.

Near the distal ends of the metatarsal bones the central part divides into five portions; each of these subdivides into two slips, which embrace the corresponding flexor tendons, and are attached to the metatarsal bones and their connecting ligaments. Between the primary divisions of the fascia—that is, in a line between the toes—are seen the digital vessels and nerves. This arrangement is in all respects like that in the palm.

In the interdigital folds of the skin there are also ligamentous fibres, which run from one side of the foot to the other, and answer the same purpose as those in the hand (p. 375).

The plantar fascia must be partially removed to examine the muscles. Toward the os calcis its removal is not accomplished without some difficulty, since the muscles arise from it.

Superficial Muscles. After the removal of the fascia three muscles are exposed. All arise from the os calcis and the fascia, and proceed forward to the toes.[1] The central one is the flexor brevis digitorum, the one on the inner side is the abductor hallucis, and the outer one is the abductor minimi digiti.

Abductor Hallucis. This muscle *arises* from the internal tubercle of the os calcis, from the plantar fascia, from the internal annular ligament, and from the intermuscular septum between it and the flexor brevis digitorum. Its origin arches over the plantar muscles and nerves in their passage to the sole. The fibres run along the inner side of the sole, and terminate on a tendon, which is *inserted*, with the inner tendon of the flex brevis hallucis, into the inner side of the base of the first phala of the great toe, through the medium of the internal sesam bone. Its nerve comes from the internal plantar.

Abductor Minimi Digiti. This muscle lies on the outer border of t foot, and has a very strong *origin* from the und surface of the os calcis, from its external tubercle, from the plant fascia, and from the external intermuscular septum between and the flexor brevis digitorum. Some of its fibres terminate

[1] They are separated from each other by strong perpendicular partitions—int muscular septa—which pass in from the plantar fascia.

a tendon which is inserted into the proximal end of the metatarsal bone of the little toe; but the greater part run on to a tendon which is *inserted*, with the flexor brevis minimi digiti, into the outer side of the first phalanx of the little toe. It is supplied by the external plantar nerve.

Flexor Brevis Digitorum. This muscle *arises* from the under surface of the os calcis, between the two preceding, from the deep surface of the plantar fascia and the intermuscular septa. It passes forward and divides into four tendons, which run super-

FIG. 158.

MUSCLES, VESSELS AND NERVES OF THE SOLE OF THE RIGHT FOOT, AFTER REFLECTION OF THE FLEXOR BREVIS DIGITORUM.

ficial to those of the long flexor. Cut open the sheath which contains them; follow them on to the toes, to see that each bifurcates over the first phalanx, to allow the long tendon to pass through; then the two slips, reuniting, are inserted into the sides of the second phalanx. The same arrangement prevails in the fingers. It is supplied by the internal plantar nerve.

The three superficial muscles should now be reflected, by sawing off about half an inch of the os calcis, and then turning it downward with the muscles attached to it. This done, we bring

into view the plantar vessels and nerves, and the second layer of muscles—*i. e.*, the long flexor tendon of the great toe, that of the other toes, and the flexor accessorius.

Tendon of the Flexor Longus Digitorum.

Musculus Accessorius.

Tracing this tendon into the sole, you find that an accessory muscle is attached to it. The *flexor accessorius arises* by muscular fibres from the inner concave side of the os calcis and the calcaneo-scaphoid ligament, and by tendinous fibres from the outer side in front of the external tubercle, and from the long plantar ligament. Its fibres run straight forward, and are *inserted* into the fibular side of the upper surface of the tendon, so that their action is not only to assist in bending the toes, but to make the common tendon pull in a straight line toward the heel; which, from its oblique direction, it could not do without the accessory muscle. The common tendon then divides into four, one for each of the four outer toes. These run in the same sheath with the short tendons, and, after passing through their divisions, are inserted into the bases of the ungual phalanges. Respecting the manner in which the tendons are confined by fibrous sheaths, and lubricated by a synovial lining, what was said of the fingers (p. 383) applies equally to the toes. The flexor accessorius is supplied by the external plantar nerve.

Lumbricales.

These four little muscles are placed between the long flexor tendons. Each, excepting the most internal, which is attached only to the inner side of the tendon going to the second toe, *arises* from the adjacent sides of two tendons, proceeds forward, and then, sinking between the toes, terminates in an aponeurosis which passes round the inner side of the four outer toes, and *joins* the extensor tendon on the dorsum of the first phalanges of the toes. Concerning their use, refer to p. 385. The two outer lumbricales are supplied by the external, the two inner by the internal, plantar nerve.

Now trace the long flexor tendon of the great toe. From the groove in the astragalus it runs along the groove in the lesser tuberosity of the os calcis, above, that is nearer to the bones than the tendon of the flexor longus digitorum, between the two heads of the flexor brevis hallucis, and then straight to the base of the last phalanx. It crosses the long flexor tendon of the toes, and

the two tendons are connected by an oblique slip; so that we cannot bend the other toes without the great toe.

Plantar Arteries. The posterior tibial artery, having entered the sole between the origins of the abductor hallucis, divides into the external and internal plantar arteries.

The *internal plantar artery* is smaller than the external plantar artery; it passes forward between the abductor hallucis and the flexor brevis digitorum to the base of the great toe, and then is continued along the inner side of that toe, where it terminates in small inosculations with the digital arteries. Its chief use is to supply the muscles between which it runs.

The *external plantar* is the principal artery of the sole, and alone forms the plantar arch (Fig. 159). It runs obliquely outward across the sole, toward the base of the fifth metatarsal bone; then, sinking deeply, it bends inward across the bases of the metatarsal bones, and inosculates with the communicating branch of the dorsalis pedis in the first interosseous space. At first it lies between the os calcis and the abductor hallucis; it then passes between the flexor brevis digitorum and the flexor accessorius; still continuing its course forward, it is placed between the flexor brevis digitorum and the flexor brevis minimi digiti, covered only with skin, fat and plantar fascia; and, lastly, it lies deep beneath the flexor tendons, the lumbricales, the adductor hallucis, upon the interossei muscles. Deeply seated as it appears to be, that part of its curve near the fifth metatarsal bone lies immediately beneath the fascia.

FIG. 159.

1. Internal plantar artery.
2. External ditto.

The external plantar sends two or three *internal calcanean* branches to the skin of the heel—one round the outer edge of the foot, which anastomoses with the tarsal and metatarsal branches of the arteria dorsalis pedis; and also some cutaneous branches which emerge between

the adjacent borders of the flexor brevis minimi digiti and the flexor brevis digitorum. It also gives off—

The *digital arteries*, four in number, which arise from the deepest part of the arch. They supply both sides of the fifth, fourth, third and the outer side of the second toes; and, running forward along the interossei, divide at the cleft of the toes into two branches, which supply the contiguous sides of the adjacent toes. At the point of division the digital arteries send upward, through the front part of the three outer interosseous spaces, small branches, *anterior perforating*, which anastomose, on the dorsum of the foot, with the interosseous arteries.

The *posterior perforating* are three branches which perforate the back part of the three outer interosseous spaces, and inosculate with the dorsal interosseous arteries at each end of the spaces.

The *digital artery*, supplying the inner side of the great toe and the adjacent sides of the great and second toes, comes from the communicating branch of the dorsal artery of the foot, which pierces the back of the first interosseous space to get to the sole of the foot. It is joined here by a branch from the external plantar artery.

Plantar Nerves. The posterior tibial nerve divides, like the artery, into an external and internal plantar.

The *internal plantar nerve* is the larger, and runs with its corresponding artery along the inner side of the foot between the abductor hallucis and the flexor brevis digitorum; in this part of its course it distributes *cutaneous branches* to supply the skin of the sole, *muscular branches* to the two above-named muscles, and *articular branches* to the joints of the tarsus and metatarsus. It then gives off *four digital branches* which supply the three inner toes and a half, like the median in the palm; the *first* digital branch runs along the inner side of the great toe to its tip, and in its passage gives off a branch to the flexor brevis hallucis; the *second* divides into two branches, one of which supplies the inner lumbricalis, and the other, the contiguous borders of the great and second toes; the *third* sends a filament to the second lumbrical, and then bifurcates for the supply of the adjacent sides of the second and third toes; the *fourth*, after receiving a communicating filament from the external plantar nerve, is distributed to the contiguous

sides of the third and fourth toes. These digital nerves send off small branches to supply the dorsum of the toes at the last phalanges.

The *external plantar nerve* passes obliquely forward and outward with the artery of the same name, passing between the flexor accessorius and the flexor brevis digitorum, to the former of which it sends a muscular branch; it then runs along the inner border of the abductor minimi digiti, supplies it, and then divides into two branches, a superficial and a deep.

The *superficial branch* sends one *digital branch*, which supplies the flexor brevis minimi digiti, the plantar and dorsal interossei of the fourth space, and the outer side of the little toe; and another and larger *digital branch* which supplies the contiguous sides of the fourth and fifth nerves, and sends a communicating filament to the outer digital branch of the internal plantar nerve.

The *deep branch* accompanies the plantar arch deep into the sole of the foot, beneath the adductor hallucis, and furnishes branches to the plantar and dorsal interossei of all the interosseous spaces except the fourth, the adductor hallucis, the transversalis pedis, and the two outer lumbricales.

Third Layer of Muscles. Having traced the principal vessels and nerves, divide them with the flexor tendons near the os calcis, and turn them down toward the toes, to expose the deep muscles in the sole. These are, the flexor brevis and adductor hallucis, the flexor brevis minimi digiti, and the transversalis pedis.

Flexor Brevis Hallucis. This muscle *arises* by a pointed tendon from the cuboid bone, and from the fibrous prolongation of the tibialis posticus into the external cuneiform. It proceeds along the metatarsal bone of the great toe, and divides into two portions, which run one on each side of the long flexor tendon, and are *inserted* by tendons into the sides of the first phalanx of the great toe. The inner tendon is inseparably connected with the abductor hallucis, the outer with the adductor hallucis. In each tendon there is a sesamoid bone. These bones not only increase the strength of the muscle, but, both together, form a pulley for the free play of the long flexor tendon; so that in walking the tendon is not pressed upon. Its nerve comes from the internal plantar (Fig. 160, *k*).

Adductor Hallucis. This very powerful muscle *arises* from the bases of the second, third and fourth metatarsal bones, and from the sheath of the peroneus longus. Passing obliquely forward and inward across the foot, it is *inserted* through the external sesamoid bone into the outer side of the base of the first phalanx of the great toe, together with the inner head of the flexor

FIG. 160.

a. Abductor hallucis.
b. Flexor brevis digitorum.
c. Abductor minimi digiti.
d. Tendon of peroneus longus.
e, *n*. Flexor longus hallucis.
f. Tendon of flexor longus digitorum.
g. Flexus accessorius.
h. Flexor brevis minimi digiti.
i. Adductor hallucis.
k. Flexor brevis hallucis.
l. Interossei.
m. Transversalis pedis.
o. Tibialis posticus.

VIEW OF THE THIRD LAYER OF MUSCLES OF THE FOOT.

brevis. This muscle greatly contributes to support the arch of the foot. Like the adductor of the thumb, it should be considered as an interosseous muscle. Its nerve is derived from the external plantar (Fig. 160, *i*).

Flexor Brevis Minimi Digiti. This little muscle rests on the fifth metatarsal bone, and *arises* from the base of the fifth meta-

tarsal bone and the sheath of the peroneus longus; it proceeds forward along the bone, and is *inserted* into the outer side of the base of the first phalanx of the little toe. It is supplied by the external plantar nerve (Fig. 160, *h*).

Transversalis Pedis. This slender muscle runs transversely across the distal ends of the metatarsal bones. It *arises* by little fleshy slips from the inferior metatarso-phalangeal ligaments of the three outer toes, and the transverse ligament of the metatarsus, and is *inserted* into the outer side of the first phalanx of the great toe with the adductor hallucis, of which it ought to be considered a part. Its nerve comes from the external plantar (Fig. 160, *m*).

The fourth layer of muscles consist of the interossei.

Interossei. These muscles are arranged nearly like those in the hand. They occupy the intervals between the metatarsal bones, and are seven in number, four being on the dorsal aspect of the foot, three on the plantar. The four *dorsal interossei arise*, each by two heads, from the contiguous sides of the metatarsal bones, and are *inserted* into the bases of the first phalanges, and into the aponeurosis of the extensor communis digitorum on the dorsum of the toes. The first is inserted into the inner side of the second toe; the remaining three into the outer sides of the second, third and fourth. The *plantar interossei*, three in number, *arise* from the inner sides and under surfaces of the third, fourth and fifth metatarsal bones, and are *inserted* respectively into the inner sides of the bases of the first phalanges of the third, fourth and fifth toes, and into the aponeurosis of the common extensor tendon.

The use of the interosseous muscles is to draw the toes to or from each other, and they do the one or the other according to the side of the phalanx on which they act. Now, if we draw a longitudinal line through the second toe, we find that all the dorsal muscles draw *from* that line, and the plantar *toward* it. This is the key to the action of them all. A more detailed account of these muscles is given in the dissection of the hand (p. 419). Between the tendons of the interossei, that is, between the distal ends of the metatarsal bones, there are bursæ which facilitate movement. They sometimes become enlarged and occasion

Attention should now be directed to the *internal* or *posterior annular ligament*, which binds down the tendons behind the inner ankle.

It is attached to the internal malleolus and the inner border of the os calcis. It is continuous, above, with the deep fascia of the leg, below, with the plantar fascia and the abductor hallucis. Beneath it are a series of compartments through which pass the tendons of the deep-seated muscles of the leg and the vessels into the sole of the foot. The relative positions of the structures passing under this ligament, proceeding from within outward, are, the tendons of the tibialis posticus, and the flexor longus digitorum; the posterior tibial artery accompanied by its venæ comites; the posterior tibial nerve; and, lastly, the tendon of the flexor longus hallucis. Each of the tendons is lined by a separate synovial membrane, as they pass behind the inner ankle.

Course and Relations of the Posterior Tibial Artery. This artery is one of the branches into which the popliteal divides at the lower border of the popliteus. It descends between the superficial and the deep muscles at the back of the leg, to the interval between the internal malleolus and the os calcis, and, entering the sole, divides beneath the abductor hallucis into the external and internal plantar arteries. It lies, first, for a short distance, *upon* the tibialis posticus, then, on the flexor longus digitorum; but behind the ankle it is in contact with the tibia, so that here it can be felt beating, and effectually compressed; and lastly, on the back of the ankle joint. In the upper part of its course it runs nearly midway between the bones, and is *covered* by the gastrocnemius and soleus, and is crossed obliquely from within outward by the posterior tibial nerve. To tie the artery, therefore, in this situation, is difficult; but in the lower part of its course it gradually approaches the inner border of the tibia, from which, generally speaking, it is not more than half or three-quarters of an inch distant. Here, being comparatively superficial, it may easily be tied. Immediately behind the internal malleolus, it lies between the tendons of the flexor longus digitorum on the inner side, and the flexor longus hallucis on the outer. It has two venæ comites, which communicate at intervals. The posterior tibial nerve which accompanies the artery is at its upper third on

its *inner side*, then crosses over it, and for the lower two-thirds of its course lies *external* to the artery. Its branches are as follows:—

a. Numerous *muscular* branches to the soleus, the peronei, and the deep muscles.

b. The *peroneal* is a branch of considerable size—often as large as the posterior tibial. Arising about an inch below the division of the popliteal, it descends close to the interosseous border of the fibula, and then over the articulation between the tibia and fibula to the outer part of the os calcis, where it inosculates with the malleolar and plantar arteries. All down the leg it is imbedded among the muscles, being covered, first, by the soleus, afterward, by the flexor longus hallucis, by the fibres of which it is more or less surrounded. To both these muscles, to the latter especially, it sends numerous branches, and just above the ankle it gives off a constant one, the *anterior peroneal*, which passes through the interosseous membrane to the under aspect of the peroneus tertius, then runs in front of the inferior tibio-fibular articulation, and inosculates with the other malleolar and tarsal arteries. The peroneal supplies the *nutrient* artery of the fibula.

c. The *communicating artery* passes transversely across the interosseous membrane, about an inch above the os calcis. It runs under the tendon of the flexor longus hallucis, and anastomoses with the peroneal artery.

d. The *internal calcanean* arteries, several in number, run down, ramifying over the posterior and inner aspect of the tendo-Achillis and os calcis; they anastomose with the internal malleolar and peroneal arteries.

e. The *nutrient artery* to the tibia enters the bone about an inch below the oblique line, the foramen through which it passes being directed away from the knee joint.

The *posterior tibial veins*, formed by the junction of the external and internal plantar veins, accompany the artery as *venæ comites*, and, after receiving the peroneal veins, join with the veins corresponding to the anterior tibial artery, at the lower border of the popliteus, to form the popliteal vein.

Posterior Tibial Nerve. This is the continuation of the popliteal. It descends close to its corresponding artery, and,

behind the inner ankle, divides into the external and internal plantar nerves. In the first part of its course the nerve lies superficial to the artery, and rather to its inner side; but lower down the nerve crosses the artery, and passes to its outer side, and lies to the inner border of the tendo-Achillis. It supplies branches to the three deep-seated muscles, the branch to the flexor longus hallucis accompanying the peroneal artery; and a cutaneous branch, *calcaneo-plantar*, which pierces the internal annular ligament, and supplies the skin of the heel and the inner side of the sole of the foot.

THE DISSECTION OF THE SOLE OF THE FOOT.

Surface Marking. The skin of the sole is remarkably thick, especially over the os calcis and the heads of the metatarsal bones. The sole of the foot is convex and narrow behind, but gradually increases in breadth forward to the clefts of the toes. The inner arch of the foot extending from the inner tuberosity of the os calcis to the distal end of the first metatarsal bone, is well marked in a well-developed foot; along this arch can be distinguished the sustentaculum tali; about an inch in front of this is the prominent tubercle of the scaphoid, and another inch in front of this we can feel the articulation between the internal cuneiform and the base of the metatarsal bone of the great toe. The outer arch of the foot is neither so deep nor so long as the inner; it extends from the external tubercle of the os calcis to the base of the metatarsal bone of the little toe, and along it we can recognize the peroneal tubercle of the os calcis and the base of the fifth metatarsal bone. The transverse arch, between the heads of the metatarsal bones, is not well marked through the skin, for it is obscured by the tense plantar fascia. The plantar fascia can be made tense by extending the foot, when its narrowest part is seen to be about one inch in front of the os calcis. The course of the external plantar artery is indicated by a line drawn from the internal tubercle of the os calcis to the base of the fifth metatarsal bone; that of the internal plantar artery by a line drawn from the same tuberosity to the first interosseous space.

Dissection. The foot should be firmly fixed to a block, with the sole directed toward the dissector, and the

toes either fastened by strings or nailed to the block, so as to put the plantar fascia on the stretch. Make a perpendicular incision down the middle of the sole; another transverse one across the foot at the clefts of the toes, and continue the perpendicular incision along the middle of the toes to their terminations. Reflect the skin, and notice the peculiar structure of the subcutaneous tissue. It is composed of globular masses of fat, separated by strong, fibrous septa, and forms elastic pads, especially marked at the heel, and at the ball of the great and the little toes; these being the points which form the tripod supporting the arch of the foot.

In removing the subcutaneous tissue from the ball of the great and the little toes, we often meet with bursæ, simple or multilocular. They are generally placed between the skin and sesamoid bones, and have remarkably thick walls. Frequently an artery and nerve can be traced running directly through one of these sacs, which explains the acute pain produced by their inflammation.

Cutaneous Nerves. In the fat the student must make out the *cutaneous branch* of the posterior tibial nerve, which supplies the skin of the sole of the foot and the heel; the remainder of the sole is supplied by small branches of the plantar nerves which come through the fascia, as in the palm of the hand.

Plantar Fascia. This is a remarkably dense, white and glistening fascia. It extends from the under and back part of the os calcis to the distal extremities of the metatarsal bones. It is divided into a strong central and two lateral less dense portions; from which prolongations pass deeply inward, separating the lateral from the central muscles. The *middle portion*, covering the flexor brevis digitorum, is narrow behind, and, as it passes forward toward the toes, is spread out, and strengthened by transverse fibrous bands. The *inner portion* is comparatively thin, and surrounds the abductor hallucis, becoming continuous posteriorly with the internal annular ligament. The *outer portion* is thicker than the inner, especially as it passes forward to be attached to the proximal end of the fifth metatarsal bone. It covers the abductor minimi digiti. Both the inner and the outer portions are continuous with the fascia of the dorsum of the foot round the inner and outer borders of the

foot, and with the central portion of the plantar fascia toward the centre of the sole.

Near the distal ends of the metatarsal bones the central part divides into five portions; each of these subdivides into two slips, which embrace the corresponding flexor tendons, and are attached to the metatarsal bones and their connecting ligaments. Between the primary divisions of the fascia—that is, in a line between the toes—are seen the digital vessels and nerves. This arrangement is in all respects like that in the palm.

In the interdigital folds of the skin there are also ligamentous fibres, which run from one side of the foot to the other, and answer the same purpose as those in the hand (p. 375).

The plantar fascia must be partially removed to examine the muscles. Toward the os calcis its removal is not accomplished without some difficulty, since the muscles arise from it.

Superficial Muscles. After the removal of the fascia three muscles are exposed. All arise from the os calcis and the fascia, and proceed forward to the toes.[1] The central one is the flexor brevis digitorum, the one on the inner side is the abductor hallucis, and the outer one is the abductor minimi digiti.

Abductor Hallucis. This muscle *arises* from the internal tubercle of the os calcis, from the plantar fascia, from the internal annular ligament, and from the intermuscular septum between it and the flexor brevis digitorum. Its origin arches over the plantar muscles and nerves in their passage to the sole. The fibres run along the inner side of the sole, and terminate on a tendon, which is *inserted*, with the inner tendon of the flexor brevis hallucis, into the inner side of the base of the first phalanx of the great toe, through the medium of the internal sesamoid bone. Its nerve comes from the internal plantar.

Abductor Minimi Digiti. This muscle lies on the outer border of the foot, and has a very strong *origin* from the under surface of the os calcis, from its external tubercle, from the plantar fascia, and from the external intermuscular septum between it and the flexor brevis digitorum. Some of its fibres terminate on

[1] They are separated from each other by strong perpendicular partitions—intermuscular septa—which pass in from the plantar fascia.

a tendon which is inserted into the proximal end of the metatarsal bone of the little toe; but the greater part run on to a tendon which is *inserted*, with the flexor brevis minimi digiti, into the outer side of the first phalanx of the little toe. It is supplied by the external plantar nerve.

Flexor Brevis Digitorum. This muscle *arises* from the under surface of the os calcis, between the two preceding, from the deep surface of the plantar fascia and the intermuscular septa. It passes forward and divides into four tendons, which run super-

FIG. 158.

1. Abductor minimi digiti.
2. Flexor accessorius.
3. Abductor hallucis.
4. External plantar artery and nerve.
5. Tendon of flexor longus hallucis.
6. Internal plantar artery and nerve.
8. Flexor brevis minimi digiti.

9. Lumbricales.
10. Internal plantar nerve.
11. Tendons of the flexor brevis digitorum bifurcating, for the passage of the tendons of the flexor longus digitorum.

MUSCLES, VESSELS AND NERVES OF THE SOLE OF THE RIGHT FOOT, AFTER REFLECTION OF THE FLEXOR BREVIS DIGITORUM.

ficial to those of the long flexor. Cut open the sheath which contains them; follow them on to the toes, to see that each bifurcates over the first phalanx, to allow the long tendon to pass through; then the two slips, reuniting, are inserted into the sides of the second phalanx. The same arrangement prevails in the fingers. It is supplied by the internal plantar nerve.

The three superficial muscles should now be reflected, by sawing off about half an inch of the os calcis, and then turning it downward with the muscles attached to it. This done, we bring

into view the plantar vessels and nerves, and the second layer of muscles—*i. e.*, the long flexor tendon of the great toe, that of the other toes, and the flexor accessorius.

Tendon of the Flexor Longus Digitorum.

Tracing this tendon into the sole, you find that an accessory muscle is attached to it. The *flexor accessorius arises* by muscular fibres from the inner

Musculus Accessorius.

concave side of the os calcis and the calcaneo-scaphoid ligament, and by tendinous fibres from the outer side in front of the external tubercle, and from the long plantar ligament. Its fibres run straight forward, and are *inserted* into the fibular side of the upper surface of the tendon, so that their action is not only to assist in bending the toes, but to make the common tendon pull in a straight line toward the heel; which, from its oblique direction, it could not do without the accessory muscle. The common tendon then divides into four, one for each of the four outer toes. These run in the same sheath with the short tendons, and, after passing through their divisions, are inserted into the bases of the ungual phalanges. Respecting the manner in which the tendons are confined by fibrous sheaths, and lubricated by a synovial lining, what was said of the fingers (p. 383) applies equally to the toes. The flexor accessorius is supplied by the external plantar nerve.

Lumbricales.

These four little muscles are placed between the long flexor tendons. Each, excepting the most internal, which is attached only to the inner side of the tendon going to the second toe, *arises* from the adjacent sides of two tendons, proceeds forward, and then, sinking between the toes, terminates in an aponeurosis which passes round the inner side of the four outer toes, and *joins* the extensor tendon on the dorsum of the first phalanges of the toes. Concerning their use, refer to p. 385. The two outer lumbricales are supplied by the external, the two inner by the internal, plantar nerve.

Now trace the long flexor tendon of the great toe. From the groove in the astragalus it runs along the groove in the lesser tuberosity of the os calcis, above, that is nearer to the bones than the tendon of the flexor longus digitorum, between the two heads of the flexor brevis hallucis, and then straight to the base of the last phalanx. It crosses the long flexor tendon of the toes, and

› tendons are connected by an oblique slip; so that we bend the other toes without the great toe.

· Arteries. The posterior tibial artery, having entered the sole between the origins of the abductor hallucis, into the external and internal plantar arteries.

internal plantar artery is smaller than the external artery; it passes forward between the abductor hal- nd the flexor brevis digitorum to the base of the great ıd then is continued along the inner side of that toe, it terminates in small inosculations with the digital arte- .ts chief use is to supply the muscles between which it

external plantar is the principal artery of the sole, and alone the plantar arch (Fig. 159). It runs obliquely outward the sole, toward the base of the fifth metatarsal bone; then, ; deeply, it bends inward across the ›f the metatarsal bones, and inoscu- ith the communicating branch of the s pedis in the first interosseous space. : it lies between the os calcis and the or hallucis; it then passes between cor brevis digitorum and the flexor ›rius; still continuing its course for- t is placed between the flexor brevis um and the flexor brevis minimi :overed only with skin, fat and plantar and, lastly, it lies deep beneath the endons, the lumbricales, the adductor s, upon the interossei muscles. Deeply as it appears to be, that part of its ıear the fifth metatarsal bone lies im- ely beneath the fascia.

FIG. 159.

1. Internal plantar artery.
2. External ditto.

external plantar sends two or three *l calcancan* branches to the skin of :l—one round the outer edge of the hich anastomoses with the tarsal and rsal branches of the arteria dorsalis and also some cutaneous branches which emerge between

the adjacent borders of the flexor brevis minimi digiti and the flexor brevis digitorum. It also gives off—

The *digital arteries*, four in number, which arise from the deepest part of the arch. They supply both sides of the fifth, fourth, third and the outer side of the second toes; and, running forward along the interossei, divide at the cleft of the toes into two branches, which supply the contiguous sides of the adjacent toes. At the point of division the digital arteries send upward, through the front part of the three outer interosseous spaces, small branches, *anterior perforating*, which anastomose, on the dorsum of the foot, with the interosseous arteries.

The *posterior perforating* are three branches which perforate the back part of the three outer interosseous spaces, and inosculate with the dorsal interosseous arteries at each end of the spaces.

The *digital artery*, supplying the inner side of the great toe and the adjacent sides of the great and second toes, comes from the communicating branch of the dorsal artery of the foot, which pierces the back of the first interosseous space to get to the sole of the foot. It is joined here by a branch from the external plantar artery.

Plantar Nerves. The posterior tibial nerve divides, like the artery, into an external and internal plantar.

The *internal plantar nerve* is the larger, and runs with its corresponding artery along the inner side of the foot between the abductor hallucis and the flexor brevis digitorum; in this part of its course it distributes *cutaneous branches* to supply the skin of the sole, *muscular branches* to the two above-named muscles, and *articular branches* to the joints of the tarsus and metatarsus. It then gives off *four digital branches* which supply the three inner toes and a half, like the median in the palm; the *first* digital branch runs along the inner side of the great toe to its tip, and in its passage gives off a branch to the flexor brevis hallucis; the *second* divides into two branches, one of which supplies the inner lumbricalis, and the other, the contiguous borders of the great and second toes; the *third* sends a filament to the second lumbrical, and then bifurcates for the supply of the adjacent sides of the second and third toes; the *fourth*, after receiving a communicating filament from the external plantar nerve, is distributed to the contiguous

sides of the third and fourth toes. These digital nerves send off small branches to supply the dorsum of the toes at the last phalanges.

The *external plantar nerve* passes obliquely forward and outward with the artery of the same name, passing between the flexor accessorius and the flexor brevis digitorum, to the former of which it sends a muscular branch; it then runs along the inner border of the abductor minimi digiti, supplies it, and then divides into two branches, a superficial and a deep.

The *superficial branch* sends one *digital branch*, which supplies the flexor brevis minimi digiti, the plantar and dorsal interossei of the fourth space, and the outer side of the little toe; and another and larger *digital branch* which supplies the contiguous sides of the fourth and fifth nerves, and sends a communicating filament to the outer digital branch of the internal plantar nerve.

The *deep branch* accompanies the plantar arch deep into the sole of the foot, beneath the adductor hallucis, and furnishes branches to the plantar and dorsal interossei of all the interosseous spaces except the fourth, the adductor hallucis, the transversalis pedis, and the two outer lumbricales.

Third Layer of Muscles. Having traced the principal vessels and nerves, divide them with the flexor tendons near the os calcis, and turn them down toward the toes, to expose the deep muscles in the sole. These are, the flexor brevis and adductor hallucis, the flexor brevis minimi digiti, and the transversalis pedis.

Flexor Brevis Hallucis. This muscle *arises* by a pointed tendon from the cuboid bone, and from the fibrous prolongation of the tibialis posticus into the external cuneiform. It proceeds along the metatarsal bone of the great toe, and divides into two portions, which run one on each side of the long flexor tendon, and are *inserted* by tendons into the sides of the first phalanx of the great toe. The inner tendon is inseparably connected with the abductor hallucis, the outer with the adductor hallucis. In each tendon there is a sesamoid bone. These bones not only increase the strength of the muscle, but, both together, form a pulley for the free play of the long flexor tendon; so that in walking the tendon is not pressed upon. Its nerve comes from the internal plantar (Fig. 160, *k*).

Adductor Hallucis. This very powerful muscle *arises* from the bases of the second, third and fourth metatarsal bones, and from the sheath of the peroneus longus. Passing obliquely forward and inward across the foot, it is *inserted* through the external sesamoid bone into the outer side of the base of the first phalanx of the great toe, together with the inner head of the flexor

FIG. 160.

a. Abductor hallucis.
b. Flexor brevis digitorum.
c. Abductor minimi digiti.
d. Tendon of peroneus longus.
e, n. Flexor longus hallucis.
f. Tendon of flexor longus digitorum.
g. Flexus accessorius.
h. Flexor brevis minimi digiti.
i. Adductor hallucis.
k. Flexor brevis hallucis.
l. Interossei.
m. Transversalis pedis.
o. Tibialis postica.

VIEW OF THE THIRD LAYER OF MUSCLES OF THE FOOT.

brevis. This muscle greatly contributes to support the arch of the foot. Like the adductor of the thumb, it should be considered as an interosseous muscle. Its nerve is derived from the external plantar (Fig. 160, *i*).

Flexor Brevis Minimi Digiti. This little muscle rests on the fifth metatarsal bone, and *arises* from the base of the fifth meta-

tarsal bone and the sheath of the peroneus longus; it proceeds forward along the bone, and is *inserted* into the outer side of the base of the first phalanx of the little toe. It is supplied by the external plantar nerve (Fig. 160, *h*).

Transversalis Pedis. This slender muscle runs transversely across the distal ends of the metatarsal bones. It *arises* by little fleshy slips from the inferior metatarso-phalangeal ligaments of the three outer toes, and the transverse ligament of the metatarsus, and is *inserted* into the outer side of the first phalanx of the great toe with the adductor hallucis, of which it ought to be considered a part. Its nerve comes from the external plantar (Fig. 160, *m*).

The fourth layer of muscles consist of the interossei.

Interossei. These muscles are arranged nearly like those in the hand. They occupy the intervals between the metatarsal bones, and are seven in number, four being on the dorsal aspect of the foot, three on the plantar. The four *dorsal interossei arise*, each by two heads, from the contiguous sides of the metatarsal bones, and are *inserted* into the bases of the first phalanges, and into the aponeurosis of the extensor communis digitorum on the dorsum of the toes. The first is inserted into the inner side of the second toe; the remaining three into the outer sides of the second, third and fourth. The *plantar interossei*, three in number, *arise* from the inner sides and under surfaces of the third, fourth and fifth metatarsal bones, and are *inserted* respectively into the inner sides of the bases of the first phalanges of the third, fourth and fifth toes, and into the aponeurosis of the common extensor tendon.

The use of the interosseous muscles is to draw the toes to or from each other, and they do the one or the other according to the side of the phalanx on which they act. Now, if we draw a longitudinal line through the second toe, we find that all the dorsal muscles draw *from* that line, and the plantar *toward* it. This is the key to the action of them all. A more detailed account of these muscles is given in the dissection of the hand (p. 419). Between the tendons of the interossei, that is, between the distal ends of the metatarsal bones, there are bursæ which facilitate movement. They sometimes become enlarged and occasion

painful swellings between the roots of the toes. The flexor brevis minimi digiti, the transversalis pedis, and all the interossei, are supplied by the external plantar nerve.

Now trace the tendons of the peroneus longus and tibialis posticus. The *tendon* of the *peroneus longus* is the deepest in the sole. It runs through a groove in the cuboid bone, obliquely across the sole, toward its insertion into the outer side of the base of the metatarsal bone of the great toe, and into the internal cuneiform bone; not infrequently it has a fasciculus of attachment into the second metatarsal bone. It is confined in a strong fibrous sheath, lined throughout by synovial membrane.

The *tendon* of the *tibialis posticus* may be traced over the internal lateral ligament of the ankle, and thence under the head of the astragalus to the tuberosity of the scaphoid, and the internal cuneiform bones. Prolongations are sent off to the cuneiform bones, to the cuboid, to the sustentaculum tali, and to the bases of the second, third, and fourth metatarsal bones. Observe that the tendon contributes to support the head of the astragalus, and that for this purpose it often contains a sesamoid bone. This is one of the many provisions for the maintenance of the arch of the foot.

DISSECTION OF THE LIGAMENTS.

Ligaments of the Pelvis with the Fifth Lumbar Vertebra.

The sacrum is united to the last lumbar vertebra in the same manner as one vertebra is to another—viz., by the prolongation of the anterior and posterior common ligaments, the intervertebral fibro-cartilage, the ligamenta subflava, supra- and interspinous ligaments, and the capsular ligaments. The student should, therefore, refer to the description of the ligaments of the spine (p. 314).

The *ilio-lumbar ligament* is very strong, and extends directly outward from the tip of the transverse process of the last lumbar vertebra to the crest of the ilium (Fig. 162).

The *lumbo-sacral ligament* varies much in its extent and attachment, and passes from the anterior and lower border of the transverse process of the fifth lumbar vertebra to the lateral part of the base of the sacrum; the fibres as they descend obliquely

outward become frayed out, joining in part the anterior sacro-iliac ligament.

Ligaments of the Sacrum and Coccyx.

The sacrum is connected with the coccyx by means of an anterior and a posterior sacro-coccygeal ligament and by an intervertebral fibro-cartilage.

The *posterior sacro-coccygeal ligament* is a flattened fasciculus of fibres extending from the lower margin of the sacral canal to the posterior surface of the coccyx; this ligament closes in the inferior termination of the sacral canal.

The *anterior sacro-coccygeal ligament* is a thin band of fibres passing along the front of the sacrum to the coccyx.

The *intervertebral disc* is a thin layer of fibro-cartilage, firm in the centre, thinner laterally and in front and behind, with occasionally a synovial membrane. Laterally, there are some irregular strands of fibres, the *lateral ligaments*, which extend from the lower lateral part of the sacrum to the transverse process of the coccyx.

The segments of the coccyx are in early life separated by interposed fibro-cartilages, which subsequently ossify; they have in front and behind a continuation of the anterior and posterior common ligaments.

The innominate bones are connected to each other in front, constituting the *symphysis pubis;* posteriorly, to the sacrum, forming the *sacro-iliac symphysis.*

Sacro-Iliac Articulation.

This articulation is an example of that form of amphiarthrodial joints where the surfaces are covered with fibro-cartilage, with an incomplete synovial membrane. The articulation is formed between the auricular surfaces of the lateral portions of the sacrum and ilium. The anterior part of the bones forming this articulation is incrusted with articular cartilage, of which the shape is like that of the ear. Later in life these two surfaces are more or less connected by thin interarticular transverse fibres, so that the interval between them is very irregular, and frequently contains yellow, viscid material. In front of the articulation there is the anterior sacro-iliac ligament, and behind, the posterior iliac ligament.

The *anterior sacro-iliac ligament* consists of thin ligamentous fibres passing in front of the sacrum and ilium.

The *posterior sacro-iliac ligament* is composed of fibres much stronger and more marked, which pass behind the articulation. It consists of two portions: the upper, or *horizontal*, extends from the upper two transverse tubercles of the sacrum, and is attached to the rough surface of the ilium above the auricular surface; the lower, or oblique, is a well-marked fasciculus of fibres, the *oblique sacro-iliac ligament*, passing from the posterior superior spine to the third segment of the sacrum.

Sacro-sciatic Ligaments. These are two strong ligaments passing from the sacrum to the ischium.

FIG. 161.



The *great sacro-sciatic ligament* is triangular and thick; but narrower in the middle than at either extremity. Its base is broad, and is attached to the posterior inferior iliac spine, and to the sides of the sacrum and coccyx; rapidly narrowing, it descends obliquely outward toward the tuberosity of the ischium, where it again expands to be attached to the inner margin of this bone. This attached portion is continued upward for some distance as a prolongation, the *falciform process*, into the inner margin of the ramus of the ischium, where it becomes continuous with the obturator fascia, forming a protection for the pudic vessels and nerve.

The *lesser sacro-sciatic ligament* lies in front of the preceding

ligament, and, like it, is triangular, though smaller and shorter. It passes from the sides of the sacrum and coccyx to the spine of the ischium, where it narrows considerably. The attachments of the sacro-sciatic ligaments to the sacrum and coccyx are more or less blended, and they not only serve to connect the bones, but also, from their great breadth, contribute to diminish the lower aperture of the pelvis.

Pubic Symphysis. This is formed by the union of the pubic bones, in front, by means of an interposed piece of fibro-cartilage. It is an amphiarthrodial articulation, and is secured by the following ligaments:—

The *anterior pubic ligament* consists of several layers of irregular superficial fibres which run obliquely and decussate with each other, and of deeper fibres which pass transversely across from one bone to the other, and are connected with the fibro-cartilage.

The *posterior pubic ligament* consists of fibres, less distinct than the anterior, which connect the two pubic bones posteriorly.

The *superior pubic ligament* passes across the upper surface of the pubic bones.

The *subpubic ligament* is very strong, and extends between the rami of the pubic bones, beneath the fibro-cartilage with which it is blended; it rounds off the pubic arch, and is situated between the two layers of the triangular ligament.

The *intermediate fibro-cartilage* is composed of two layers of cartilage, each attached to the inner border of the body of the os pubis by a number of nipple-like processes fitting into corresponding depressions on the bony surface. Between these cartilaginous plates there is a thick stratum of fibrous and fibro-elastic tissue. In the middle line at the upper and back part is usually a smooth cavity lined with epithelium. The cartilage acts as a buffer, and breaks the force of shocks passing through the pelvic arch.

Ligaments of the Hip Joint. This joint is secured by the form of the bones, and by the strength of the powerful muscles which surround it. Although an enarthrodial or ball-and-socket joint, its range of motion is somewhat limited; the disposition of its ligaments restricts its range of motion to those directions only which are most consistent with the maintenance of the erect attitude and the requirements of this part of the skeleton.

The ligaments of the hip joint are, the capsular, the ilio-femoral, the ligamentum teres, the cotyloid and the transverse.

Capsular Ligament. The *capsular ligament* is attached above to the circumference of the acetabulum, a little external to the margin, also to the transverse ligament, and by a few fibres to the outer margin of the obturator foramen; below, to the anterior intertrochanteric ridge in front; above, to the root of the great trochanter, and to the middle of the neck behind, about half an inch above the posterior intertrochanteric ridge. The anterior and upper part of the capsular ligament is very thick and strong, composed chiefly of longitudinal fibres with a few deeply seated circular fibres, which are concealed by the superficial longitudinal bands. The posterior aspect of the capsular ligament is represented by a few sparsely scattered fibres. The front part of the ligament is rendered exceedingly strong by several accessory ligaments, one of which, called the *ilio-femoral ligament*, extends from the anterior inferior iliac spine, and from a depression above the acetabulum, and then divides like the two arms of the inverted letter Λ: one, the inner and vertical, passes to the base of the lesser trochanter; the outer, to the upper part of the anterior intertrochanteric line. In addition, there is at the lower and back part a broad ligament, the *ischio-capsular ligament*, whose fibres extend from the ischium to the inner part of the joint, close to the lesser trochanter; and a third accessory ligament, the *pubo-femoral*, consists of the thin fibres converging from the ilio-pectineal eminence, and the margin of the obturator foramen to the front and inner part of the capsular ligament.

The ilio-femoral ligament is very strong, and serves as a strap to prevent the femur being extended beyond a certain point, and limits rotation inward and outward.

The capsule is in relation, in front, with the iliacus and psoas muscles, from which it is separated by a synovial bursa. This bursa occasionally communicates by a rounded aperture with the synovial cavity of the hip joint.

Open the capsule to ascertain its great thickness in front and its strong attachment to the bones. This exposes the cotyloid ligament and the ligamentum teres.

Ligamentum Teres.

The *ligamentum teres* is exposed by drawing the head of the femur out of the socket. This ligament is somewhat flat and triangular. Its base, which is bifid, is attached, below, to the borders of the notch in the acetabulum, where it becomes continuous with the transverse ligament; its apex, to the fossa in the head of the femur. To prevent pressure on it, and to allow free room for its play, there is a gap at the bottom of the acetabulum. This gap is not

FIG. 162.



VERTICAL SECTION THROUGH THE HIP.

crusted with cartilage like the rest of the socket, but is occupied by soft fat. The ligamentum teres is surrounded by the synovial membrane. An artery runs up with it to the head of the femur. It is a branch of the obturator, and enters the acetabulum through the notch at the lower part.

The chief use of the ligamentum teres is to assist in steadying the pelvis on the thigh in the erect position. In this position, the ligament is vertical and quite tight (Fig. 162); it therefore pre-

vents the pelvis from rolling toward the opposite side, or the thigh from being adducted beyond a certain point. Another purpose served by this ligament is to limit rotation of the thigh, both inward and outward.

Cotyloid Ligament. The *cotyloid ligament* is an annular piece of fibro-cartilage which is attached all round the margin of the acetabulum. Its circumference is thicker than its free margin, which is very thin, so that on a transverse section the cartilage is triangular. Both its surfaces are covered with synovial membrane, and its attachment to the margin of the acetabulum is effected by oblique fibres passing from without inward, and interlacing in all directions at an acute angle. The ligament is thicker above and behind than elsewhere; it thus deepens the socket, and embraces the head of the femur like a sucker. It extends over the notch at the lower part of the acetabulum, being attached to a ligament, the *transverse*, which passes

Transverse Ligament. across the notch, and thus converts it into a foramen. Beneath the transverse ligament some of the vessels and nerves pass into the joint to supply it.

Synovial Membrane. The *synovial membrane* extends from the cartilaginous border of the head, round the neck as far as the attachment of the capsular ligament, on the inner surface of which it is reflected as far as the margin of the acetabulum. Thence it passes over the superficial surface of the cotyloid ligament, curves round its inner sharp border so as to line its deeper surface; it then covers the osseous surface of the acetabulum, and is finally continued as a tubular sheath over the ligamentum teres to the head of the femur. The synovial membrane, it will be seen, extends down to the base of the neck of the femur in *front*, but only two-thirds *behind*. Between the bottom of the acetabulum, as far as the cotyloid notch and the synovial membrane, is a collection of fat and connective tissue called the *gland of Havers*.

The ligaments of the hip joint are so arranged that, when we stand "at ease," the pelvis is spontaneously thrown into a position in which its range of motion is the most restricted; for the accessory ligaments of the capsule prevent it from being extended beyond a straight line, and the ligamentum teres prevents its roll-

ing toward the opposite side. This arrangement economizes muscular force in balancing the trunk.

The atmospheric pressure is, of itself, sufficient to keep the limb suspended from the pelvis, supposing all muscles and ligaments to be divided. When fluid is effused into the hip-joint, the bones are no longer maintained in accurate contact; and it sometimes happens that the head of the femur escapes from its cavity, giving rise to a spontaneous dislocation.

The *movements at the hip joint* are those of flexion, extension, abduction, adduction, rotation, and circumduction.

The *flexors* are, the ilio-psoas, the sartorius, the pectineus, the adductor longus and brevis, the gluteus medius and minimus.

The *extensors* are, the gluteus maximus, the biceps, semitendinosus, and the semimembranosus.

The *abductors* are, the upper fibres of the gluteus maximus, the gluteus medius and minimus, the pyriformis, and when the joint is flexed, the obturator internus and the two gemelli.

The *adductors* are, the three adductors, the pectineus, the gracilis, and the sartorius.

The *external rotators* are, the ilio-psoas, the three adductors, the pectineus, the gluteus maximus, the posterior fibres of the gluteus medius, the obturator externus and internus, the gemelli, the quadratus femoris, the pyriformis, and the sartorius.

The *internal rotators* are, the tensor fasciæ femoris, the anterior fibres of the gluteus medius, and the gluteus minimus.

Circumduction is effected by the successive action of the different muscles in the order of their attachment into the femur.

The muscles in immediate relation with the hip joint are, in front, the iliacus and psoas; on the outer side, the reflected tendon of the rectus, the gluteus minimus; behind, the pyriformis, gemellus superior, obturator internus, gemellus inferior, obturator externus, and quadratus femoris; on the inner side, the pectineus and obturator externus.

Ligaments of the Knee joint. The knee joint is a ginglymus or a hinge joint, and the bones entering into its formation are, above, the condyles of the femur, below, the head of the tibia, and in front, the patella. Looking at the skeleton, one would

suppose that it was very insecure; but this insecurity is only apparent, the joint being surrounded by powerful ligaments, and a thick capsule formed by the tendons of the muscles which act upon it.

First examine the tendons concerned in the protection of the knee joint. In front is the ligamentum patellæ; on each side are the tendons of the vasti; on the outer side, in addition, it is strengthened by the strong ilio-tibial band; on the inner side there are also the tendons of the sartorius and gracilis; at the back of the joint are the tendons of the gastrocnemius and plantaris, with the semimembranosus and semitendinosus, in addition, on its inner part, and the tendons of the popliteus and biceps on its outer part. It deserves to be mentioned that the weakest part of the articulation is near the tendon of the popliteus, which arises *within* the joint; here, therefore, pus or fluid formed in the popliteal space may make its way into the joint, or vice versâ.

The ligaments of the joint may be divided into those outside the joint and those within it.

Those *outside* the joint are, the ligamentum patellæ, the internal lateral, the two external lateral, the posterior ligament, and the capsular ligament.

Those *within* the joint are, the two crucial, the two semilunar fibro-cartilages, the transverse, the coronary, the ligamenta alaria, and the ligamentum mucosum.

Ligamentum Patellæ. The *ligamentum patellæ* is a strong, thick, ligamentous band, about three inches long, extending from the lower border of the patella to the tubercle of the tibia. Beneath it is found a considerable amount of fat, which separates the ligament from the joint, and in fat people forms a prominent elastic mass on each side of the ligament. There is a synovial bursa between the ligamentum patellæ and the tuberosity of the tibia.

Internal Lateral Ligament. This is a broad, flat band, which extends from the inner condyle of the femur to the inner tuberosity and the inner aspect of the shaft of the tibia (Fig. 163). A few of the deeper fibres are attached to the inner semilunar cartilage, and serve to keep it in place. The inferior internal articular

artery and part of the tendon of the semimembranosus pass underneath this ligament. In the several motions of the joint there is a certain amount of friction between the ligament and the head of the tibia, and consequently a small bursa is interposed.

FIG. 163.

DIAGRAM OF THE SEMILUNAR CARTILAGES AND LATERAL LIGAMENTS OF THE KNEE.

1. Internal lateral ligament.
2. External ditto.

External Lateral Ligaments. The *long external lateral ligament* is a strong round band which extends from the outer condyle of the femur to the outer part of the head of the fibula. This ligament separates the two divisions of the tendinous insertion of the biceps. Beneath it pass the tendon of the popliteus and the inferior external articular artery.

The *short external lateral ligament* is situated posterior to and runs parallel with the preceding ligament; it passes from the posterior and outer part of the condyle of the femur to the tip of the styloid process of the fibula; the tendon of the popliteus also lies beneath it.

Posterior Ligament. This, which is generally called the *ligamentum posticum Winslowii*, covers the whole of the posterior surface of the knee joint, and consists of two portions, one formed by a broad, flat band of vertical fibres passing from the posterior surface of the femur between and above the condyles to the posterior part of the tuberosity of the tibia; the other consists of an oblique tendinous expansion from the semimembranosus (p. 685), which passes upward and outward from the internal tuberosity of the tibia to the back of the outer condyle of the femur. It is pierced by numerous blood vessels to supply the knee joint, chiefly by the azygos artery, and a branch from the obturator nerve (p. 659). This ligament not only closes and protects the joint behind, but prevents its extension beyond the perpendicular.

Capsular Ligament. The *capsular ligament* occupies the intervals between the other ligaments, and so completes

the fibrous investment of the joint. It is materially strengthened by fibrous expansions from many of the tendons in connection with the articulation.

The joint should be opened above the patella. Observe the great extent of the fold which the synovial membrane forms above this bone.[1] It allows the free play of the bone over the lower part of the femur. The fold extends higher above the inner than the outer condyle, which accounts for the form of the swelling produced by effusion into the joint.

Crucial Ligaments. The *crucial ligaments*, so named because they cross like the letter X, extend from the mesial side of each condyle to the head of the tibia.

The *anterior* or *external ligament*, the smaller, ascends from the inner part of the fossa in front of the spine of the tibia, backward and outward, to the inner and back part of the external condyle. It is attached to the tibia close to the anterior termination of the external semilunar cartilage.

The *posterior* or *internal ligament*, best seen from behind, extends from the back of the fossa behind the spine of the tibia, and from the posterior termination of the external semilunar cartilage, upward, forward and inward, to the front of the inner condyle. The direction of this ligament is more vertical than the anterior one.

Interarticular or Semilunar Fibro-cartilages. Between the condyles and the articular surfaces of the tibia are two incomplete rings of fibro-cartilage, shaped like the letter C. They serve to deepen the articular surfaces of the tibia; their mobility and flexibility enable them to adapt themselves to the condyles in the several movements of the joint; they distribute pressure over a greater surface and break shocks. They are thickest at the circumference, and gradually shelve off to a thin margin; thus they fit in between the bones, and adapt a convex surface to a flat one, as shown in Fig. 163. Their form is suited to the condyles, the inner being oval, the outer circular, and the synovial membrane covers both surfaces of the cartilages.

The *external semilunar fibro-cartilage* is nearly a circular ring

[1] In performing operations near the knee, the joint should always be bent, in order to draw the synovial fold as much as possible out of the way.

of fibro-cartilage, its two extremities being firmly attached to the fossæ, one in front of and the other behind the spine of the tibia; they are enclosed by the two extremities of the internal cartilage.

On its outer border it presents a groove for the tendon of the popliteus; its anterior border gives off a transverse fibrous fasciculus, the transverse ligament, which passes across to be connected with the anterior border of the internal cartilage. The anterior extremity of the fibro-cartilage is connected with the anterior crucial ligament; the posterior is attached partly into the outer side of the inner tuberosity in front and behind the posterior crucial ligament, and partly into the anterior crucial ligament.[1]

The *internal semilunar fibro-cartilage* forms about two-thirds of an oval ring, and is narrower in front than behind. Its anterior extremity is pointed, and is attached to the tibia internal to the anterior crucial ligament; its posterior extremity to the pit behind the spine immediately in front of the posterior crucial ligament.

The *transverse ligament*, already alluded to, is a thin fibrous fasciculus in front of the anterior crucial ligament, and connects the anterior borders of the semilunar fibro-cartilages.

The *coronary ligaments* are two ligaments which connect the circumference of the two semilunar cartilages to the borders of the tibial tuberosities. The external ligament is the weaker of the two, so that the external cartilage is the more movable.

Synovial Membrane.

The synovial membrane is very extensive, the most extensive in the body. It lines the posterior surface of the quadriceps tendon, and the aponeuroses of the vasti, and is reflected on to the femur a variable distance above the incrusting cartilage; traced from the femur, we find that it lines the inner surface of the capsular ligament as far as the circumference of the tibia; thence it is reflected over the upper surfaces of the semilunar cartilages, round their inner concave margins to get to their under surfaces, from which the membrane passes to cover the

[1] Of the two cartilages the external has the greater freedom of motion, because in rotation of the knee the outer side of the tibia moves more than the inner. Consequently, itis not in any way connected to the external lateral ligament; so far from this, it is separated from it by the tendon of the popliteus, of which the play is facilitated by a bursa communicating freely with the joint. For this reason the external cartilage is more liable to dislocation than the internal.

articular surface of the head of the tibia. It forms tubular prolongations round the crucial ligaments, and below the patella a slender band of the synovial membrane proceeds backward to the space between the condyles, and is called the *ligamentum mucosum*. Two lateral folds, extending from the sides of the mucous ligament, pass upward and outward to the sides of the patella; these are termed the *ligamenta alaria*. These are not true ligaments, but merely remnants of the partition which, in the early stage of the joint's growth, divided it into two equal portions.

Outside the synovial membrane there is always fat; especially under the ligamentum patellæ. Its use is to fill up vacuities, and to mould itself to the several movements of the joint.

The movements which the knee joint permits are those of flexion and extension, together with rotation outward and inward. In order completely to master its various movements, the student should examine the movements first as between the femur and the patella, and then as between the condyles of the femur and the articular surface of the tibia crowned by its two semilunar cartilages.

The articular surface of the patella glides upon the femoral condyles in extension and in flexion. If this surface of the patella be examined, it will be seen that each lateral facet is subdivided by two indistinct transverse ridges into three very shallow transverse zones; each of these zones rests upon a definite part of the trochlear surface of the femur in the different stages of extension or flexion; thus, in extreme extension, the lower zones of the patella rest upon the upper border of the trochlear surface; in mid-flexion, the middle zones alone rest on the femur; and in nearly extreme flexion, the upper zones lie in the lower part of the femoral condyles. In addition to the six shallow facets just described, there is a seventh, which is seen on the inner margin.

The respective points of the attachment of the ligaments are such that, when the joint is extended, all the ligaments are tight, to prevent extension beyond the perpendicular; thus muscular force is economized. But when the joint is bent the ligaments are relaxed, enough to admit a slight rotatory movement of the tibia.

This movement is more free outward than inward, and is

effected, not by rotation of the tibia on its own axis, but by rotation of the outer head round the inner. Rotation outward is produced by the biceps; rotation inward by the popliteus and semimembranosus.

FIG. 164.

CRUCIAL LIGAMENTS OF THE KNEE.

The crucial ligaments, though placed inside the joint, answer the same purposes as the coronoid process and the olecranon of the elbow. They make the tibia *slide* properly forward and backward. In extension, the anterior crucial ligament is tight, as are also the lateral ligaments; in flexion, the posterior ligament becomes tight and consequently limits flexion. They also conjointly limit excessive rotation. They not only prevent dislocation in front or behind, but they prevent lateral displacement, since they cross each other like braces, as shown in Fig. 164.

Superior Tibio-fibular Articulation. This is an arthrodial or gliding joint, and is formed by the flat oval surfaces of the upper part of the tibia and fibula. It is secured by an anterior and a posterior tibio-fibular ligament.

The *anterior superior tibio-fibular ligament* is a strong, flat ligament, whose fibres pass obliquely downward and outward from the external tuberosity of the tibia to the head of the fibula.

The *posterior superior ligament* passes in the same direction as the anterior, only being placed behind the joint.

The *synovial membrane* occasionally communicates with that of the knee joint.

Interosseous Membrane. The contiguous borders of the tibia and fibula are connected by the interosseous membrane. The purpose of it is to afford additional surface for the attachment of muscles. Its fibres pass chiefly downward and outward from the tibia to the fibula, but a few fibres cross, like the letter X. The anterior tibial artery comes forward above the interosseous membrane, through an oval space about an inch below the head of the fibula. Lower down there is an aperture for the anterior peroneal artery. It is, moreover, pierced here and there by small blood vessels.

Inferior Tibio-fibular Articulation.

The lower extremities of the tibia and fibula are firmly connected, for it is essential to the security of the ankle joint that there should be little or no movement between the two bones.

The *anterior inferior ligament* passes between the adjacent borders of the two bones; it is narrow above but broader below, and consists of oblique fibres which pass downward and outward.

The *posterior inferior ligament* is stronger and narrower than the anterior, and its fibres pass horizontally from the outer malleolus to the posterior border of the tibia, above the articular surface.

FIG. 165.



1. Plantar fascia.
2. Calcaneo-scaphoid ligament which supports the head of the astragalus.
3. Internal lateral ligament, called, from its shape, deltoid.

The *transverse ligament* is the lower part of the preceding, and may be usually recognized as a distinct narrow fasciculus.

The *inferior interosseous ligament* consists of strong, short fibres connecting the contiguous surfaces of the two bones, and continuous above with the interosseous membrane.

The *synovial membrane* of this joint is an extension upward of that of the ankle joint.

Ankle Joint.

From the form of the bones, it is obvious that the ankle is a ginglymus or hinge joint; consequently, its security depends upon the great strength of its lateral ligaments. The hinge, however, is not so perfect but that it

admits of a slight rotatory motion, of which the centre is on the fibular side, and therefore the reverse of that in the case of the knee.

The ligaments of the ankle joint comprise the anterior, the internal and the external lateral ligaments.

The *anterior ligament* is a thin, loose, membranous ligament, attached above to the tibia, and below to the astragalus, in front of their articular surfaces, and is sufficiently loose to permit the necessary range of motion.

The *internal lateral ligament*, sometimes called, from its shape, *deltoid*, is exceedingly thick and strong, and compensates for the comparative shortness of the internal malleolus (Fig. 165). The great strength of it is proved by the fact that, in dislocation of the ankle inward, the summit of the malleolus is more often broken off than the ligament torn. The *superficial portion* of this ligament is attached above to the margin of the internal malleolus, and passing downward radiates to be inserted into the scaphoid, the inner side of the astragalus, the sustentaculum tali, and the inferior calcaneo-scaphoid ligament; the *deeper portion*, thick and strong, passes from the tip of the malleolus to the astragalus, close to its articular border.

FIG. 166.

DIAGRAM OF THE EXTERNAL LATERAL LIGAMENT.

1. Anterior part.
2. Posterior part.
3. Middle part.
4. Interosseous ligament between the astragalus and os calcis.

The *external lateral ligament* consists of three distinct fasciculi, an anterior, a posterior, and a middle (Fig. 166). The *anterior fasciculus* passes from the front of the tip of the external malleolus, nearly horizontally forward and inward to the astragalus in front of its malleolar articular surface. The *middle fasciculus*, round and long, passes obliquely downward and backward to the outer surface of the os calcis. The *posterior fasciculus* passes from the posterior part of the external malleolus, nearly horizontally outward, to the back of the astragalus below its upper articular surface.

Besides flexion and extension, the ankle joint admits of a slight lateral movement, only permitted in the extended state, for the better direction of our steps. In adaptation to this movement the internal malleolus is shorter than the outer; it is not so tightly confined by its ligaments, and its articular surface is part of a cylinder.

Open the joint to see that the breadth of the articular surfaces of the bones is greater in front than behind. The object of this is to render the astragalus less liable to be dislocated backward. Whenever this happens, the astragalus must of necessity become firmly locked between the malleoli.

Ligaments Connecting the Bones of the Tarsus.

The astragalus is the keystone of the arch of the foot, and supports the whole weight of the body. It articulates with the os calcis and the os scaphoides in such a manner as to permit the abduction and adduction of the foot, so useful in the direction of our steps.

Astragalo-calcanean Ligaments.

The astragalus articulates with the os calcis by two surfaces separated by the deep interosseous groove, of which the posterior is concave and the anterior convex. The articulations are strengthened by the three ligaments, the external and the posterior astragalo-calcanean, and the interosseous.

The *external astragalo-calcanean ligament* is a short, quadrilateral fasciculus, passing from the outer surface of the astragalus, in front of the anterior fasciculus of the external lateral ligament, almost directly downward to the outer surface of the os calcis.

The *posterior astragalo-calcanean ligament* is a short, oblique band, which passes from the posterior border of the astragalus to the upper border of the os calcis.

The *interosseous ligament*, a very thick, strong band of fibres which descends vertically in the interosseous canal, and is the principal bond of union between the two bones.

Astragalo-scaphoid Ligament.

The anterior surface of the astragalus is broadly convex, fitting into the concave surface of the scaphoid bone. Superiorly the dorsal surfaces of the two bones are connected by a broad, membranous ligament, *astragalo-*

scaphoid, which passes obliquely across, blending externally with the external calcaneo-scaphoid, and below with the inferior calcaneo-scaphoid ligament.

Calcaneo-scaphoid Ligaments. In the skeleton the head of the astragalus articulates in front with the scaphoid, but the lower part of it is unsupported. This interval is bridged over by a very strong and slightly elastic ligament, which extends from the os calcis to the scaphoid (Fig. 167); this is the inferior calcaneo-scaphoid ligament.

The *inferior calcaneo-scaphoid ligament* is thick and strong, and passes horizontally forward and inward from the sustentaculum tali to the plantar surface of the scaphoid, where it is connected with the tendon of the tibialis posticus, and, superiorly, with the astragalo-scaphoid ligament. Thus the os calcis, scaphoid and this ligament form a complete socket for the head of the astragalus; it is this joint, chiefly, which permits the abduction and adduction of the foot. In chronic disease of the ankle joint, leading to much impairment of movement at the joint, the motion at the astragalo-scaphoid articulation is so great as to take the place of the ankle joint. This ligament being slightly elastic, allows the keystone of the arch (the astragalus) a play, which is of great service in preventing concussion of the body. Whenever this ligament yields, the head of the astragalus falls, and the individual becomes gradually flat-footed. In this yielding of the ligament the head of the astragalus not only falls, but becomes also rotated inward.

The *superior calcaneo-scaphoid ligament* is short and triangular, lying in the hollow between the outer part of the astragalus and the os calcis; it passes forward and upward from the ridge on the anterior and outer part of the os calcis to the outer side of the scaphoid.

Calcaneo-cuboid Articulation. The os calcis articulates with the os cuboides nearly on a line with the joint between the astragalus and the scaphoid. The bones are connected together, on the dorsum, by the superior and internal calcaneo-cuboid ligaments, and on the plantar aspect by the long and short calcaneo-cuboid ligaments.

The *superior calcaneo-cuboid ligament* is a short quadrilateral

FIG. 167.

1. Calcaneo-scaphoid ligament.
2. Calcaneo-cuboid ligament.

band of fibres passing from the upper part of the dorsal aspect of the os calcis to the back and upper part of the os cuboides.

The *internal calcaneous-cuboid ligament* connects the front part of the ridge of the os calcis to the dorsal and inner part of the cuboid. It is sometimes called the *interosseous ligament*, and is closely associated with the superior calcaneo-scaphoid ligament.

The *long calcaneo-cuboid ligament*, a broad, long and strong band of ligamentous fibres, is the more superficial of the two inferior calcaneo-cuboid ligaments. It is known as the *long plantar ligament*, and is attached to the under surface of the os calcis in front of the tuberosities, as far as the anterior tubercle; it passes forward to the plantar aspect of the cuboid, being attached to the ridge, while some of its fibres extend to the bases of the second, third and fourth metatarsal bones, and complete the canal for the tendon of the peroneus longus.

The *short calcaneo-cuboid ligament*, deeper than the former, is seen somewhat on its inner aspect, and is separated from it by some fat and connective tissue. It is very broad, and passes from the front of the tubercle of the os calcis, for about an inch in breadth, to the inner and posterior surface of the cuboid, behind the ridge.

The articulations between the cuboid and the scaphoid bones behind, and the three cuneiform bones in front, are maintained by dorsal, plantar and interosseous ligaments.

The *dorsal* and the *plantar ligaments* consist of parallel fasciculi passing between the contiguous borders of the respective bones on their dorsal and plantar aspects.

The *interosseous ligaments*, four in number, are composed of transverse fibres (Fig. 168) connecting the rough non-articular surfaces of the contiguous bones; the first one is between the scaphoid and the cuboid; the second connects the internal and middle cuneiform cones; the third is between the middle and

external cuneiform bones; and the fourth between the external cuneiform and the cuboid bones.

Though there is very little motion between any two bones the collective amount is such that the foot is enabled to adapt itself accurately to the ground; pressure is more equally distributed, and consequently there is a firmer basis for the support of the body. Being composed, moreover, of several pieces, each of which possesses a certain elasticity, the foot gains a general springiness and strength which could not have resulted from a single bone.

FIG. 168.



INTEROSSEOUS LIGAMENTS OF THE WEDGE-BONES.

Tarso-metatarsal Joints. The tarsus articulates with the metatarsus in an oblique line which inclines backward on its outer side. This line is interrupted at the joint of the middle cuneiform bone and the second metatarsal bone. Here there is a deep recess, so that the base of this metatarsal bone is wedged in between the internal and external cuneiform bones.

These joints are maintained in position, above, by the dorsal tarso-metatarsal ligaments, and below, by the plantar ligaments. Interosseous ligaments also pass between the wedge bones, maintaining them in their normal positions.

Synovial Membranes of the Tarsus. Exclusive of the ankle joint and the phalanges of the toes, the bones of the foot are provided with six distinct synovial membranes; namely—

1. Between the posterior articular surface of the os calcis and that of the astragalus.

2. Between the head of the astragalus and the scaphoid, and between the anterior articular surface of the astragalus and os calcis.

3. Between the os calcis and the os cuboides.

4. Between the inner cuneiform bone and the metatarsal bone of the great toe.

5. Between the scaphoid and the three cuneiform bones, and between these and the adjoining bones (the great toe excepted).

6. Between the os cuboides and the fourth and fifth metatarsal bones.

The tarso-metatarsal articulations are arthrodial joints.

Inter-Metatarsal Articulations. The metatarsal bones are connected at their proximal and distal ends by *dorsal* and *plantar ligaments;* those at the proximal extremities are very strong, and are supplemented by interosseous ligaments, as in the metacarpus (p. 436). The movement between the proximal ends of the inter-metatarsal articulations is arthrodial or gliding.

The distal extremities of the metatarsal bones are united by a *transverse metatarsal ligament;* this extends from the great toe to the little toe on their plantar surfaces.

FIG. 169.

DIAGRAM OF THE ARTICULATIONS OF THE TARSUS AND THE TARSO-METATARSUS.

1. Posterior calcaneo-astragaloid synovial cavity.
2. Calcaneo-scaphoid synovial cavity.
3. Calcaneo-cuboid synovial cavity.
4. Synovial cavity between metatarsal bone of great toe and internal cuneiform bone.
5. Common scapho-cuneiform, intercuneiform, and metatarso-cuneiform synovial cavity.
6. Cubo-metatarsal synovial cavity.

Metatarso-phalangeal Articulations. These are connected by a *plantar* and *two lateral ligaments;* the dorsal ligament being formed by the expansion of the tendon of the extensor longus digitorum. The movements which take place between the articulations are those of flexion, extension, abduction and adduction.

Phalangeal Articulations. These articulations have the same kind of ligaments as the preceding, and the movements are also nearly identical.

DISSECTION OF THE BRAIN.

Membranes of the Brain. Before passing on to the examination of the brain, the student should study the arrangement, the structure and the uses of the three membranes by which the brain is enveloped.

The most external one, the *dura mater*, has been described (p. 24). The second, or intermediate one, is a serous membrane, termed the *arachnoid;* the third, the *pia mater*, is a vascular layer, and is in contact with the encephalon.

Arachnoid Membrane. The *arachnoid membrane*,[1] the second investment, constitutes the smooth, polished membrane covering the surface of the brain, and is exposed after the removal of the dura mater. This tunic was formerly considered by anatomists to be an example of an ordinary serous membrane, and was described as consisting of two layers—an external or parietal, which lined the inner surface of the dura mater, and an internal or visceral, which was reflected over the brain.

It is now regarded as consisting of one layer only, viz., the one which envelops the brain; the under aspect of the dura mater being covered only with a layer of flattened epithelium cells. The cavity which was formerly described as the cavity of the arachnoid, is now called the *subdural space*, and contains a very limited amount of fluid.

The arachnoid membrane is a colorless and transparent layer, and is spread uniformly over the surface of the brain, from which it is separated by the pia mater. It does not, like the pia mater, dip down into the furrows between the convolutions of the brain, and it is more or less connected with the pia mater by delicate connective tissue, the *subarachnoid*. On account of its extreme tenuity, and its close adhesion to the pia mater, the two membranes cannot be readily separated; but there are places, especially at the base of the brain, termed *subarachnoid spaces*, where the arachnoid membrane can be seen distinct from the subjacent pia mater. The two membranes can be artificially separated by blowing air beneath the arachnoid with a blowpipe.

[1] ἀράχνη, a spider's web; εἶδος, form.

Subarachnoid Spaces and Fluid. Wherever the arachnoid membrane is separated from the pia mater, a serous fluid (*cerebro-spinal*) intervenes, contained in the meshes of a very delicate areolar tissue. The spaces between these membranes are termed *subarachnoid*, and are very manifest in some places. For instance, there is one well-marked space in the longitudinal fissure, where the arachnoid does not descend to the bottom, but passes across the edge of the falx cerebri, a little above the corpus callosum. At the base of the brain there are two of considerable size: one, the *anterior subarachnoid space*, is situated between the anterior border of the pons Varolii, the middle and the anterior cerebral lobes; the other, the *posterior subarachnoid space*, is placed between the cerebellar hemispheres and the medulla oblongata. The fluid in the subarachnoid space communicates with the fluid of the general ventricular cavities of the brain through an aperture (*foramen of Magendie*) in the fourth ventricle, close to its lower boundary; and also through an opening, on each side, behind the glosso-pharyngeal nerves. In the spinal cord, also, there is a considerable interval, containing fluid, between the arachnoid and the pia mater. The purpose of this fluid is, not only to fill up space, as fat does in other parts, but mechanically to protect the nerve centres from the violent shocks and vibrations to which they would otherwise be liable.

The brain, therefore, may be said to be supported in a fluid, which insinuates itself into all the inequalities of the surface, and surrounds in fluid sheaths all the nerves as far as the foramina through which they pass. This fluid sometimes escapes through the ear, in cases of fracture through the base of the skull, involving the meatus auditorius internus and the petrous portion of the temporal bone.

The arachnoid is supplied with filaments from the motor root of the fifth, the facial and the spinal accessory nerves.

The *cerebro-spinal fluid* varies in amount from two drachms to two ounces. It is a clear, limpid fluid, slightly alkaline, containing 98.5 parts of water, and 1.5 parts of solid matter. The cerebro-spinal fluid of the encephalon and that of the spinal cord communicate.

Pia Mater. This, the immediate investing membrane of the brain, is extremely vascular, and composed

of a minute network of blood vessels held together by delicate connective tissue. It covers the cerebral surface, and dips into the fissures between the convolutions, forming a double layer. From its internal surface numerous vessels pass off at right angles into the substance of the brain.[1] The pia mater sends a prolongation through the transverse fissure into the lateral and third ventricles, forming the *velum interpositum* and the *choroid plexuses*, and also another along the roof of the fourth ventricle, forming the *tela choroidea inferior*. Upon the surface of the cerebellum, the pia mater is thinner, not so vascular, and only sends prolongations down the larger sulci; on the pons Varolii and the medulla, the membrane is more fibrous and much less vascular than elsewhere.

The pia mater is supplied with nerves by the third, fifth, sixth, facial, glosso-pharyngeal, pneumogastric, spinal accessory, and sympathetic nerves, which chiefly accompany the blood vessels forming the pia mater.

Arteries of the Brain. The brain is supplied with blood by the two internal carotid and the two vertebral arteries.

Internal Carotid. This artery enters the skull through the carotid canal in the temporal bone, and ascends very tortuously, by the side of the body of the sphenoid, along the inner wall of the cavernous sinus. It appears on the inner side of the anterior clinoid process, and after giving off the ophthalmic, divides into an anterior and middle cerebral, posterior communicating and anterior choroid arteries.

a. The *anterior cerebral artery* is given off from the internal carotid at the inner end of the fissure of Sylvius. It passes forward and inward to reach the longitudinal fissure between the hemispheres, curves round the front part of the corpus callosum, then runs backward over its upper surface (under the name of the *artery of the corpus callosum*), and terminates in branches which anastomose with the posterior cerebral arteries. The anterior cerebral arteries of opposite sides run side by side, and supply the olfactory lobes, the optic nerves, the frontal lobes, the anterior perforated spaces, and the corpus callosum. At the

[1] Owing to these vessels, the pia mater, when placed in water, presents a flocculent, woolly appearance, and hence it is sometimes called *tomentum cerebri*.

base of the brain, as they enter the longitudinal fissure, they are connected by a short transverse branch, called the *anterior communicating artery* (Fig. 170).

b. The *middle cerebral artery*, the largest branch of the internal carotid, runs outward deeply within the fissure of Sylvius, and

FIG. 170.



divides into many branches distributed to the anterior and middle lobes. Near its origin it gives off a number of small arteries, which pierce the locus perforatus anticus to supply the corpus striatum.

c. The *posterior communicating artery*, unequal in size, usually, on the two sides, proceeds directly backward to join the posterior

cerebral; thus establishing at the base of the brain the free arterial inosculation called the *circle of Willis*.

d. The *anterior choroid artery*, a small branch of the internal carotid, arises external to the posterior communicating artery. It runs backward, and enters the fissure at the bottom of the middle horn of the lateral ventricle, to terminate in the choroid plexus of that cavity. It supplies, in addition, the hippocampus major and the corpus fimbriatum.

Vertebral Artery. This artery, a branch of the subclavian in the first part of its course, enters the foramen in the transverse process of the sixth cervical vertebra, and ascends through the transverse processes of the cervical vertebræ. It then winds backward along the arch of the atlas, and enters the skull through the foramen by perforating the posterior occipito-atlantal ligament and the dura mater. It then curves around the medulla oblongata between the hypoglossal nerve and the anterior root of the first cervical nerve. At the lower border of the pons Varolii the two arteries unite to form a single trunk, the *basilar*, which is lodged in the groove on the middle of the pons, and bifurcates, at its upper border, into the posterior cerebral arteries.

Each vertebral artery, before joining its fellow, gives off:—

a. *Lateral spinal branches*, which enter the spinal canal, to supply the spinal cord and its membranes, and the bodies of the cervical vertebræ.

b. *Muscular branches* to the deep muscles of the neck, which anastomose with the occipital and deep cervical arteries.

c. A *posterior meningeal branch*, distributed to the posterior fossa of the skull.

d. *Anterior* and *posterior spinal arteries*, which are given off immediately before the vertebral arteries join to form the basilar, run along the median fissures of the front and the back surfaces of the spinal cord, and anastomose with the spinal branches of the ascending cervical arteries.

e. The *posterior inferior cerebellar artery*, sometimes a branch of the basilar, but more frequently of the vertebral, passes backward between the spinal accessory and the pneumogastric nerves, to the under surface of the cerebellum. It divides into two branches: an outer, which ramifies on the lower surface of the cerebellum

as far as its outer border; and an inner, which passes to the vallecula between the two hemispheres, and supplies branches to the fourth ventricle.

The *basilar artery*, formed by the junction of the two vertebral, in its course along the pons, gives off on each side:—

a. *Transverse* branches which pass outward on the pons: one, the *internal auditory*, enters the meatus auditorius internus with the auditory nerve, to be distributed to the internal ear on each side.

b. The *anterior inferior cerebellar*, which supplies the front part of the lower surface of the cerebellum, and anastomoses with the other cerebellar arteries.

c. The *superior cerebellar arteries*, given off near the bifurcation of the basilar, are distributed to the upper surface of the cerebellum, and anastomose with the inferior cerebellar; branches are supplied to the pineal body, the valve of Vieussens, and the velum interpositum.

d. The *posterior cerebral arteries* are the two terminal branches into which the basilar artery divides. They run outward and backward, in front of the third cranial nerve, and wind round the crura cerebri to the under surface of the posterior cerebral lobes, where they divide into numerous branches for the supply of the brain, anastomosing with the anterior and middle cerebral arteries. Shortly after their origins they receive the two posterior communicating arteries from the internal carotids. Each gives off small branches to the posterior perforated space, and also the small *posterior choroid artery*, which, passing beneath the posterior border of the corpus callosum and fornix, is distributed to the velum interpositum and choroid plexus.

Circle of Willis. This important arterial inosculation (Fig. 170) takes place between the branches of the two internal carotid and the two vertebral arteries. It is formed, laterally, by the two anterior cerebral, the two internal carotid, and the two posterior communicating arteries; in front, it is completed by the anterior communicating artery; behind, by the two posterior cerebral. The tortuosity of the large arteries before they enter the brain serves to mitigate the force of the heart's action; and the circle of Willis provides a free supply of blood from other vessels,

in case any accidental circumstance should stop the flow of blood through any of the more direct channels.[1]

Peculiarities of the Cerebral Circulation. Besides the circle of Willis, there are other peculiarities relating to the circulation of the blood in the brain, namely, the length and tortuosity of the four great arteries as they enter the skull; their passage through tortuous bony canals; the spreading of their ramifications in a very delicate membrane, the pia mater, before they enter the substance of the brain; the minuteness of the capillaries, and the extreme thinness of their walls; the formation of the venous sinuses (p. 27), which do not accompany the arteries; the chordæ Willisii in the superior longitudinal sinus; the absence of valves in the sinuses; and the confluence of no less than six sinuses, forming the *torcular Herophili*, at the internal occipital protuberance.

General Division of the Brain. The mass of nervous substance contained within the cranial cavity, comprised under the common term brain (*encephalon*), is divided into four parts. The *cerebrum* forms the largest portion, and occupies the whole of the upper part of the cranial cavity; its base resting on the anterior and middle fossæ and the tentorium cerebelli. It is connected with the pons Varolii by two white nerve masses, the *crura cerebri*, and with the cerebellum by two white cords, the *crura cerebelli*. The *cerebellum*, or small brain, occupies the space between the tentorium cerebelli and the inferior occipital fossæ. The *pons Varolii* is the quadrilateral mass of white fibres which rests upon the basilar process of the occipital bone. The *medulla oblongata* is the portion below the pons, which is continuous below with the spinal cord and rests upon the lower part of the basilar process of the occipital bone.

[1] In many of the long-necked herbivorous quadrupeds a provision has been made in the disposition of the internal carotid arteries, for the purpose of equalizing the force of the blood supplied to the brain. The arteries, as they enter the skull, divide into several branches, which again unite and form a remarkable network of arteries called by Galen, who first described it, the "*rete mirabile*." The object of this evidently is to moderate the rapidity with which the blood would otherwise enter the cranium in the different positions of the head, and thus preserve the brain from those sudden influxions to which it would under other circumstances be continually exposed.

The result of a large number of observations shows that the weight of the brain averages in males 49½ oz., and in females about 44 oz.; although it has been known to weigh as much as 64 oz., as in the case of Cuvier's brain, and as little as 23 oz., in the case of an idiot's brain.

The following are the weights of the various portions of the encephalon, which have been carefully collated by Dr. J. Reid.[1]

	MALE.		FEMALE.	
	oz.	drs.	oz.	drs.
Cerebrum	43	15¾	38	12
Cerebellum	5	4	4	12¼
Pons and medulla oblongata . . .		15¾	1	0¼
Total	50	3½	44	8½

The average specific gravity of the brain is about 1.036; that of the white matter being 1.040, and that of the gray 1.034.

The relative proportion of the amount of white to gray matter is 64 per cent. of white substance to 36 per cent. of gray matter.

The weight of the encephalon varies greatly in different subjects, and although its weight seems to bear some proportionate relation to the intellectual power, yet in many instances there appears to be no such definite relation.

The brain weight gradually increases up to the age of forty, when it attains its maximum; after this period the weight decreases at the rate of one ounce for every additional ten years of life.

MEDULLA OBLONGATA.

The *medulla oblongata* is that part of the cerebro-spinal axis which is placed below the pons Varolii, and is continuous with the spinal cord on a level with the upper border of the atlas. It is slightly pyramidal in shape, with the broad part above. It lies on the basilar groove of the occipital bone, and descends obliquely backward through the foramen magnum. Its posterior surface is received into the fossa (vallecula) between the hemispheres of the cerebellum. It is about an inch and a quarter in length, three

[1] Reid, *London and Edinburgh Monthly Journal of Medical Science*, April, 1843.

quarters of an inch at its broadest part, and half an inch in thickness.

In front and behind, the medulla is marked by a median fissure, the *anterior* and *posterior median fissures*, which are the continuations of the median fissures of the spinal cord. The anterior ends, below the pons Varolii, in a cul-de-sac, termed the *foramen cæcum*, and is occupied by a process of pia mater. The posterior runs along the floor of the fourth ventricle, as a shallow median groove.

FIG. 171.



C. Crus cerebri.
P. V. Pons Varolii.
P. Anterior pyramid.
O. Olive.
R. Restiform tract or body.

1. Gasserian ganglion.
2. Motor root of the fifth n.
3. Third n.
4. Arciform fibres.
5. Sensitive root of the fifth n.
6. Sixth n.
7. The seventh and eighth nerves.
8. The ninth, tenth and eleventh nerves.
9. Twelfth or hypoglossal n.

DIAGRAM OF THE FRONT SURFACE OF THE MEDULLA OBLONGATA.

The surface of the medulla is marked out on each side into four longitudinal columns, which receive the following names, from before backward: the *anterior pyramids*, the *lateral tracts* and the *olivary bodies*, the *restiform bodies* and the *posterior pyramids*.

The *anterior pyramids* are two columns of white matter, narrow below, but increasing gradually in breadth as they ascend toward the pons. At this part they become constricted, and may be traced through the pons into the crura cerebri. The fibres of which they are in the main composed are derived from the anterior columns of the spinal cord, and consist, therefore, of motor fibres. On separating the pyramids about an inch below the pons, bundles of

nerves are seen decussating across the anterior fissure (Fig. 171). This decussation, which consists of three or four bundles on each side, involves only the inner fibres of the pyramid; the outer fibres ascend through the pons without crossing. The decussating fibres are the continuations upward of the deep fibres of the lateral pyramidal tract and the lateral columns of the cord, which here come forward to the surface, and push aside the anterior pyramids. Thus it will be seen that the thickness of the anterior pyramids is mainly due to these decussating bundles; that the fibres on the outer side of the pyramids are those continued upward from the anterior column of the cord; and that the inner part is made up of the fibres of the lateral tract of the opposite side.

This decussation explains the phenomenon of *cross paralysis*, *i. e.* when one side of the brain is injured, the loss of motion is manifested on the opposite side of the body.[1]

The *olivary bodies* are the two oval eminences situated on the outer side of the upper part of the anterior pyramids, from which they are separated by a shallow depression. They do not ascend quite as high as the pons, for there is a deep groove between them. They consist externally of white matter; and at their lower part some white fibres may be observed arching round from the anterior median fissure, constituting the *arciform fibres* of Rolando. In the groove between the anterior pyramids and these bodies are seen the fasciculi of the hypoglossal nerve emerging from the medulla, and in the groove behind the olive emerge the roots of the glosso-pharyngeal, the pneumogastric and spinal accessory nerves.

The *lateral tracts*, situated on the outer side of the anterior pyramids, are the continuations upward of the lateral columns of the spinal cord. As described, the greater number of their fibres pass across the anterior median fissure to the opposite anterior pyramid; but some ascend as far as the lower border of the olivary body, where they divide, a few passing in front of, but the larger part passing behind, this body in their course to the brain.

The *restiform bodies* are the broadest and thickest of the

[1] The phenomenon of *cross paralysis of sensation* is explained by the fact, made out by Brown-Séquard, that the paths of sensory impressions cross each other in the gray matter of the cord.

[2] From *restis*, a rope.

columns of the medulla. They are situated to the outer side and behind the lateral tracts and the olivary bodies. They are the continuations upward of the posterior columns of the spinal cord, and as they ascend, they diverge and pass into the cerebellum, constituting its inferior peduncles (Fig. 172). Owing to this divergence, the gray matter of the medulla is exposed, so that the floor of the fourth ventricle (of which the restiform bodies and the posterior pyramids assist in forming the lateral boundaries) is mainly composed of gray matter. Near the commencement of each restiform body there is easily recognizable an eminence

FIG. 172.



DIAGRAM OF THE FOURTH VENTRICLE AND RESTIFORM BODIES.

1. Thalamus opticus.
2. Nates and testes, or corpora quadrigemina.
3. Origin of fourth nerve.
4. Processus a cerebello ad testes.
5. Restiform bodies diverging.
6. Origins of eighth or auditory nerve.

interposed between the restiform body and the groove which separates it from the lateral tract; this eminence, narrow at first, is known as the *funiculus of Rolando*, but it soon enlarges to form a considerable prominence, called the *tubercle of Rolando*. The restiform body on a transverse section is wedge-shaped, and hence it is sometimes called the *funiculus cuneatus;* near the level where the two bodies diverge, each presents an enlargement, known as the *cuneate tubercle*. The restiform bodies

consist, as before said, of white fibres derived from the posterior and lateral columns of the cord; in its interior is some gray matter continuous with that in the posterior part of the cord.

The *posterior pyramids* (*funiculi graciles*) are two slender white columns on each side of the posterior median fissure, and are the continuations upward of the posterior median columns of the spinal cord. 'Ascending, they diverge, and thus form the apex of the fourth ventricle. At their point of separation the posterior pyramids enlarge, and form the *clava;* after which they rapidly taper off, and run up on the inner side of the restiform bodies, which, however, they soon leave, and are continued upward into the cerebrum, along the floor of the fourth ventricle.

Emerging from the anterior median fissure may be noticed some superficial transverse white fibres, which cross over the anterior pyramids and the lower extremity of the olivary bodies; these are known as the *arciform fibres* of Rolando (Fig. 171), and they pass upward to join the restiform body. They are probably connected with white fibres which run horizontally, constituting an imperfect septum between the two halves of the medulla, and may be seen when a longitudinal section is carefully made through its middle. The majority of these septal fibres enter the olivary bodies, and then emerging through the gray matter of the corpus dentatum, become continuous with the fibres of the restiform bodies and lateral tracts; others pass out from the posterior fissure and wind round the restiform bodies. These latter fibres are the *transverse striæ*, seen on the floor of the fourth ventricle, some of which form the roots of the auditory nerves.

The lower portion of the fourth ventricle can now be seen, bounded laterally by the diverging posterior pyramids and the restiform bodies, the floor being formed by the gray matter of the medulla. As only part, however, can be examined in the present stage, the full consideration of the fourth ventricle is deferred to a later stage in the dissection of the brain (p. 784).

Minute Structure of the Medulla Oblongata. These are among the most complicated parts of the central nervous system. They contain white and gray matter intermixed. The white matter consists, in part, of the continuation upward of the longitudinal fibres of the cord; in part, of horizontal fibres.

Anterior Pyramids. The anterior columns of the cord (Fig. 173), having reached the lower part of the medulla oblongata, are not continued straight through it, but diverge from each other, being reinforced by the deep fibres of the lateral columns, which here cross over and form the decussation of the anterior pyramids. In their further progress the fibres of the anterior columns are arranged thus: some of them run up and form the outer portion of their own pyramid; some of them pass deeply beneath the pyramid to constitute the longitudinal fibres of the reticularis alba; these fibres are continued upward into the cerebrum; some ascend beneath the olive to join the restiform body and thence to the cerebellum; another and larger fasciculus passes upward and, after embracing the olive, reunites to form a single bundle; this, joined by fibres (*olivary fasciculus*) from the olive, ascends under the name of the *fillet of Reil*, over the superior crus of the cerebellum, to pass to the corpora quadrigemina and the cerebral hemispheres (Fig. 173).

Lateral Tracts. The *lateral columns* on reaching the medulla are disposed of in three ways, as follows: some of its fibres, the *internal*, come forward between the diverging anterior columns, decussate in the middle line, and form part of the pyramid of the opposite side; others, the *external*, ascend with the restiform body (or tract) to the cerebellum; a third set, the *middle*, ascend beneath the olivary body, along the floor of the fourth ventricle (concealed by its superficial gray matter) as the *fasciculus teres*, and are continued along the upper part of the crus cerebri into the corpora quadrigemina and optic thalamus.

Olivary Bodies. The *olivary bodies* are composed externally of white matter; but if a transverse section be made into them, their interior presents an undulating line of yellowish-brown color, called, from its zigzag shape, the *corpus dentatum* or the *nucleus of the olivary body*. It forms an interrupted circle, incomplete at its upper and inner side, so that it allows the entrance of a bundle of white fibres—*olivary peduncle*—from the medulla behind the anterior pyramid. The peduncular fibres pass in various directions; some pass into the gray matter; some pass through it and join the restiform body, under the name of the internal arcuate fibres; and some, passing between the fibres

over the olivary body, come to the surface and curve round it, forming the external arcuate fibres. Two other isolated gray nuclei may be recognized in this transverse section, one on the inner side of, and the other behind the corpus dentatum; these, called the *accessory olivary nuclei*, are linear in shape, and are about a line in length; the root fibres of the hypoglossal nerve pass between the inner nucleus and the corpus dentatum, to emerge between the anterior pyramid and the olivary body.

Restiform Bodies. The posterior columns are continued upward, under the name of the *restiform bodies*, along the back of the medulla. At the apex of the fourth ventricle they diverge from each other and receive bundles of accessory fibres

FIG. 173.

1. Anterior column.
2. Lateral column.
3. Posterior column.
4. Posterior median column of the spinal cord.
5. Anterior pyramid.
6. Restiform body.
7. Posterior pyram'd.
8. Fasciculus teres.
9. Inferior crus of the cerebellum.
10. To the corpora quadrigemina.
11. Crus cerebri.

DIAGRAM OF THE COURSE OF THE FIBRES THROUGH THE MEDULLA OBLONGATA.

from the anterior pyramids and the lateral tracts, and are continued partly into the cerebellum, forming its *inferior crura*, and partly as the *fasciculi teretes* along the floor of the fourth ventricle into the cerebrum.

Posterior Pyramids. The *posterior pyramids* are the continuations upward of the posterior median columns of the spinal cord; and, like the restiform body, each divides into two fasciculi, one of which ascends and helps to form the inferior crus of the cerebellum, the other runs up with the *fasciculus teres*.

Septum or Raphé. The horizontal fibres in the medulla oblongata were first accurately described by Stilling and

Rolando, and subsequently by Clarke and others. Some of them form a *septum* and divide the medulla oblongata into symmetrical halves; some run longitudinally and others obliquely. The fibres which run from the dorsal to the ventral surface of the medulla —*fibræ rectæ*—pass outward, in front, from the anterior median fissure, and are continuous with the superficial arched fibres of Rolando; while behind they are continuous with the fibres of the fasciculi graciles and their gray nuclei. The longitudinal fibres run vertically and are derived from the arcuate fibres and the fibræ rectæ, which have altered their direction. The oblique fibres consist of the deep arcuate fibres, which enter or come out of the septum, and then pass to the outer part of the medulla. Some fibres, again, are transverse commissural fibres, and pass into the opposite anterior pyramid or the formatio reticularis,[1] and eventually assume a longitudinal direction.

Gray Matter of the Medulla.

Although in the lower part of the medulla the gray matter in its interior is arranged in the same way as it is in the spinal cord, yet owing to the course of the decussating fibres of the lateral tracts which break up the anterior gray cornua in front, and to the diverging restiform bodies and posterior pyramids behind, the gray matter in the upper two-thirds of the medulla becomes altered both in appearance and in arrangement.

The gray matter of the anterior cornua soon loses its characteristic arrangement, as it becomes detached from the main part of the gray matter by the passage through it of bundles of the white fibres of the medulla. At the lower part of the olivary body it is pushed backward and outward, in consequence of the increase of the anterior pyramids and the olivary bodies, so as to lie in the lateral part of the medulla, just in front of the posterior horn, and constitutes a gray nucleus called the *nucleus lateralis*. The larger portion of the gray matter of the anterior cornu is intersected by white fibres and becomes broken up into an interlacement of fibres and nerve cells, forming the *formatio reticularis*.

The gray matter of the posterior cornua becomes largely in-

[1] The *formatio reticularis* comprises the anterior and lateral parts of the medulla, and consists of intersecting fibres which cross each other at right angles, some being longitudinal and some transverse in their direction.

creased in the upper part of the medulla. At first the posterior cornua are pushed outward by the restiform bodies, and subsequently by the posterior pyramids, so that they lie at right angles to the posterior median fissure. At the same time the caput cornu posterioris enlarges, and later on forms a prominent gray mass, the *tubercle of Rolando*, which becomes separated from the chief part of the gray matter by being intersected by transverse and longitudinal white fibres. Internal to the tubercle of Rolando there are situated masses of gray matter in the restiform bodies and the posterior pyramids, constituting their nuclei. Higher up on a level with the middle of the olivary body we find, internal to the gray matter of the posterior pyramid (nucleus gracilis), the nuclei of origin of the two portions of the pneumogastric nerve, and of the hypoglossal nerve, the latter being nearest the middle line.

The central canal gradually approaches the posterior surface of the medulla, and above the middle of it opens out to form the median groove in the floor of the fourth ventricle. On each side of the groove we find a collection of the gray matter which enlarges in the upper part and constitutes the *nucleus teres*.

The nuclei of the origins of the spinal accessory, the pneumogastric and the glosso-pharyngeal nerves will be described subsequently.

PONS VAROLII.

The *pons Varolii*, or *tuber annulare*, is the convex eminence of transverse white fibres (Fig. 171), which is situated at the base of the brain immediately above the medulla oblongata. It rests upon the basilar groove of the occipital and the sphenoid bones, and in its antero-posterior diameter measures rather more than an inch. Its *upper margin* is convex and well defined, and arches over the crura cerebri; the *lower*, also well defined, is nearly straight, being separated from the medulla by a transverse groove. *Laterally*, the pons becomes narrower, in consequence of its transverse fibres being more closely aggregated; these enter the anterior and under part of the cerebellum, constituting its *middle peduncle*. Along the middle runs a shallow groove, broader in front than behind, which lodges the basilar artery. If the pia mater be

emoved, we observe how the superficial fibres pass transversely, o connect the two hemispheres of the cerebellum. Throughout he mammalia the size of the pons bears a direct ratio to the legree of development of the lateral lobes of the cerebellum; herefore it is larger in man than in any other animal.[1]

The pons consists of four layers of alternating transverse and longitudinal white fibres, intermingled with more or less gray matter, which is chiefly found on its upper surface, where it forms part of the floor of the fourth ventricle.

The *superficial layer* of white fibres is transverse, connecting the cerebellar hemispheres; the middle fibres pass transversely across, the inferior ascend slightly, while the superior pass backward and outward to enter the cerebellum.

The *second layer* consists of longitudinal fibres which are the continuation of the fibres of the medulla oblongata in their passage to the cerebrum. It is mainly composed of fibres derived from the anterior pyramids which pass up to form the superficial fibres, *crusta*, of the crura cerebri.

The *third layer* is formed of transverse fibres, which from their peculiar arrangement take the name of *trapezium;* the fibres in their course outward pass round in front of the superior olivary nuclei, then across the fasciculi of the facial nerves, and lastly in front of the ascending roots of the fifth nerves to enter the middle peduncle of the cerebellum.

The *deepest* and *uppermost layer* is composed of longitudinal nerve fibres; those derived from the olivary fasciculi divide into two bundles, one ascending to the corpora quadrigemina, the other passing to the cerebrum; and those derived from the lateral and posterior columns of the cord, which, with a fasciculus from the fillet, form the fasciculi teretes in the floor of the fourth ventricle, and pass upward to form the *tegmenta*, or deeper portions of the crura cerebri.

The gray matter is chiefly aggregated at the posterior part of the pons, and varies in thickness in different sections. Thus a section through the middle of the pons will show the following nuclei of gray matter, beginning from the central groove and

[1] Birds, reptiles and fishes have no pons, as there are no lateral lobes to the cerebellum.

passing outward: a small intermediate portion of the facial nerve, the large nucleus of the sixth, the facial nerve, the large superior nucleus of the auditory nerve; while below we notice the superior olivary nucleus, the nucleus of the facial nerve, and externally the gray substance of Rolando, enclosing the ascending root of the fifth nerve.

The pons, like the medulla oblongata, has an imperfect median septum, composed of horizontal fibres, some of which at the anterior border surround the crura cerebri.

THE CEREBRUM.

The *cerebrum* in man is so much more developed than the other parts of the encephalon that it completely overlies them, and forms by far the largest portion. It is oval in form, and convex on its external aspect. It is divided in the middle line into two symmetrical parts, termed the *right* and *left hemispheres*, by the deep *longitudinal fissure*, which is occupied by the falx cerebri (p. 25).[1] The cerebrum is composed of numerous parts—viz., of certain internal ganglionic masses, the corpora striata, optic thalami, and corpora quadrigemina; of commissural white fibres, the fornix, corpus callosum, and the commissures of the third ventricle; of the pineal and pituitary bodies; and, lastly, of the two lateral hemispheres, which overlie and conceal the parts previously mentioned.

FIG. 174.



DIAGRAM OF THE GENERAL DIVISIONS OF THE BRAIN.

1, 2, 3. Anterior, middle and posterior lobes of the cerebrum.
4. Cerebellum.
5. Pons Varolii.
6. Medulla oblongata.

The cerebrum rests upon the anterior and middle fossæ of the base of the skull and the tentorium cerebelli. There are three surfaces to each hemisphere: an *external* or convex; an

[1] Examples are occasionally met with, where the longitudinal fissure is not exactly in the middle line, the consequence of which want of symmetry is, that one hemisphere is larger than the other. Bichat (*Recherches physiologiques sur la Vie et la Mort*, Paris, 1829) was of opinion that this anomaly exercised a deleterious influence on the intellect. It is remarkable that the examination of his own brain after death went to prove the error of his theory.

inner or median; and an *inferior*, interrupted by the fissure of Sylvius.

By widely separating the two hemispheres at the longitudinal fissure (the brain being in its natural position), we discover that they are connected in the middle by the transverse white commissure called the *corpus callosum*. In front of and behind this mass the longitudinal fissure extends to the base of the brain.

The cerebral hemispheres, viewed from above, form an ovoid mass, broader in front than behind, and convex, to correspond with the vault of the calvarium. Their surface is mapped out by tortuous eminences termed *convolutions* (*gyri*), separated from each other by deep furrows (*sulci*), which extend to a variable depth. Many of the sulci are occupied by large veins in their course to the sinuses; others are filled with subarachnoid fluid. The convolutions are folds of the brain, and their outer surface consists of gray matter, so that the extent of the gray substance is thus very largely increased; the gray matter here is called the *cortical substance*. The interior of the convolutions consists of white nerve substance. The convolutions are not symmetrical on both sides, although they follow a somewhat similar arrangement. The number, arrangement and depth of the convolutions vary somewhat in different individuals, and, to a certain extent, may be considered an index of the degree of intelligence.[1]

The depth of the sulci between the convolutions varies in different brains, from an inch to half an inch; hence it follows that two brains of equal size may be very unequal in point of extent of surface for the gray matter, and therefore in amount of intellectual capacity. Under the microscope the cortical layer is seen to consist of four layers—two of gray alternating with two of white—the external layer being always white. These layers are not equally thick in all situations, and in some parts six layers

[1] Those who wish to investigate the cerebral convolutions in their simplest form in the lower classes of mammalia, and to trace them through their successive development and arrangement into groups as we ascend to the higher classes, should consult Leuret, *Anatomie comparée du Système Nerveux considérée dans ses Rapports avec l'Intelligence*, Paris 1839; also Foville, *Traité de l'Anat. du Système Nerveux* etc., Paris, 1844. The convolutions of the human brain have been described by Ecker, *On the Convolutions of the Human Brain*, 1873; and by Turner, *The Convolutions of the Human Brain topographically considered*, Edin. 1866.

may be demonstrated, owing to the interpolation of a layer of white substance in the outer stratum; these are chiefly seen near the corpus callosum and in the occipital lobe.[1]

Some of the sulci, from their depth, regularity, and early period of development, are termed the *primary* or *interlobular fissures*, and

FIG. 175.

VIEW OF THE CONVOLUTIONS AND FISSURES OF THE EXTERNAL SURFACE OF THE BRAIN (LEFT SIDE).

A. Fissure of Rolando.
B. Fissure of Sylvius.
C. Inter-parietal fissure.
P.O. Parieto-occipital fissure.
c.m. Calloso-marginal fissure.
F. Frontal lobe.
P. Parietal lobe.
O. Occipital lobe.
T.S. Temporo-sphenoidal lobe.
A.F. Ascending frontal convolution.
A.P. Ascending parietal convolution.
f_1, f_2, f_3. Superior, middle, and inferior frontal convolutions, separated by the superior and inferior frontal sulci.
p_1, p_2. Superior and inferior parietal convolutions, separated by the inter-parietal fissure.
o_1, o_2, o_3. Superior, middle, and inferior occipital convolutions, separated by the occipital fissures.
t_1, t_2, t_3. Superior, middle and inferior temporo-sphenoidal convolutions, separated by the superior and inferior temporo-sphenoidal fissures.

map out the surface of the cerebrum into five lobes. Of these sulci there are three: the *fissure of Sylvius*, the *fissure of Rolando* or *central fissure*, and the *parieto-occipital fissure* (Fig. 175).

[1] For an account of these laminæ, see Lockhart Clarke, *Proceedings of the Royal Society*, 1863.

The *fissure of Sylvius* is seen on the base of the cerebrum, where it receives the lesser wing of the sphenoid bone. It begins, outside the locus perforatus anticus, as a deep triangular depression—*vallecula Sylvii*, and then curves outward as a deep cleft to the external surface of the cerebrum; it divides into two rami, an ascending or vertical, about an inch in length, and a posterior or horizontal ramus, which passes backward and slightly upward, and ends at the posterior third of the cerebrum by a bifid extremity. Within the fissure, near its commencement, a series of convolutions may be seen deeply placed; these are called the *gyri operti*, or the *island of Reil*. In the fork between the two rami of the Sylvian fissure are several convolutions, which have been termed by Broca the *operculum of the insula*.

The *fissure of Rolando* or *central fissure* (Fig. 175, A) runs obliquely over the outer convex surface of the hemisphere. It commences close to the longitudinal fissure, about its middle, from which it is separated by the marginal convolution. It then runs obliquely downward and forward, and terminates a little above the fork of the Sylvian fissure. As seen in Fig. 176, the two fissures form a V-shape, failing to be joined at the angle. The fissure is formed, in early fœtal life, by a large vein, which subsequently atrophies, and is rarely bridged over.

The *parieto-occipital fissure* (Fig. 178, P.O) is seen on the median surface of the hemisphere, toward its posterior part. It begins as a deep cleft on the median surface, about half an inch behind the corpus callosum, then ascends nearly vertically, and ends on the external aspect of the cerebrum, about an inch beyond the longitudinal fissure; this latter portion taking the name of the *external parieto-occipital fissure*.

The fissure of Sylvius is first seen about the middle of the third month of fœtal life, and is caused by the extension backward, and folding upon itself, of the *mantle;* the fissure of Rolando begins to be developed about the fifth month; and the parieto-occipital fissure between the third and fourth month of uterine life.

The primary fissures form the boundaries of the various lobes of which each hemisphere is composed.

The *frontal lobe* is that part of the front portion of the cerebrum,

bounded behind by the fissure of Rolando, below by the anterior part of the fissure of Sylvius, and on the median plane by the calloso-marginal fissure. Its inferior part rests on the anterior fossa, and is called the *orbital surface*, while its convex external surface is called the *frontal surface*.

The convolutions on its frontal surface are four in number: The *ascending frontal convolution* (Fig. 175, A.F) which bounds,

FIG. 176.



CONVOLUTIONS OF THE UPPER SURFACE OF THE BRAIN.

A. Fissure of Rolando.
A.F. Ascending frontal convolution.
A.P. Ascending parietal convolution.
S.M. Supra-marginal convolution, below which is the angular gyrus.
P.O.*f.* Parieto-occipital fissure.
F. Frontal lobe.
P. Parietal lobe.
O. Occipital lobe.
f_1, f_2, f_3. Superior, middle and inferior frontal convolutions.
P_1, P_2. Superior and inferior parietal convolutions.
O_1, O_2, O_3. Superior, middle and inferior occipital convolutions.
C. m. f. Calloso-marginal fissure.

in front, the fissure of Rolando, and is usually connected above and below the fissure with the ascending parietal convolution; the union below forming the *operculum*, the union above part of the *paracentral lobule;* the upper and the lower union of these

two convolutions thus shut off the fissure of Rolando from joining the fissure of Sylvius below and the longitudinal fissure above.

The *superior*, *middle* and *inferior frontal convolutions* (Fig. 175, f_1, f_2, f_3) course nearly horizontally; the *superior* runs along the margin of the longitudinal fissure, the *inferior* along the lower border of the lobe, arching over the ascending ramus of the Sylvian fissure; the *middle* is placed between these. In front of the ascending frontal convolution is a vertical fissure, *præcentral sulcus*, which prevents the frontal convolutions joining the former convolution.

The sulci which map out the frontal convolutions are two, the *superior* and *inferior frontal sulci*.

On the *orbital surface* there is a deep sulcus, the *tri-radiate sulcus*, whose rami pass forward, outward and backward, mapping this surface out into internal, anterior and posterior orbital convolutions (Fig. 177, *Tr. s*).

On this surface also we find the *olfactory lobe*, lodged in a deep cleft, the *olfactory sulcus* (Fig. 177, *Olf. s*), and although it will be described under the cranial nerves, it is strictly a cerebral lobe, for it is developed as a distinct outgrowth from the anterior cerebral vesicle; moreover, in early fœtal life it possesses a cavity continuous with the general ventricular cavity of the brain.

The *parietal lobe* (Fig. 175, P) is placed between the fissure of Rolando and the external parieto-occipital fissure, and is bounded below by the horizontal limb of the fissure of Sylvius, and by a line continued from it to meet one passing down from the parieto-occipital fissure.

The convolutions of the parietal lobe are three: The *ascending parietal convolution* is bounded in front by the fissure of Rolando, and runs parallel with the ascending frontal convolution in front, which it joins above and below the fissure.

The *superior parietal convolution* (Fig. 175, *p*) is placed above the interparietal fissure, and courses horizontally backward; posteriorly it runs beneath the parieto-occipital fissure, and is connected with the superior occipital convolution; this is known as the *first annectent convolution;* this convolution is also seen on the median surface of the cerebrum. The *inferior parietal con-*

volution lies below the interparietal fissure, and consists of two portions: one, the ***supra-marginal***, lies in front of the posterior ramus of the Sylvian fissure; the other, the ***angular*** gyrus, lies behind the fissure of Sylvius, and bends over the termination of the parallel sulcus; the angular convolution is connected behind with the occipital convolutions by the ***second*** and ***third annectent convolutions.***

The ***interparietal fissure*** ascends at first nearly vertically, and

FIG. 177.



CONVOLUTIONS OF THE BASE OF THE CEREBRUM.

Olf.s. Olfactory sulcus.
Tr.s. Tri-radiate sulcus.
U. Uncinate convolution.
Cal.f. Calcarine fissure.
T_1, T_2, T_3. Superior, middle, and inferior temporo-sphenoidal convolutions.

then runs horizontally backward from the ascending parietal convolution.

The *occipital lobe* (Fig. 175, O) consists of the posterior part of the hemisphere behind, bounded in front by the external parieto-occipital fissure, and by a line continued from it.

The convolutions on the external surface are three :—

The ***superior***, ***middle***, and ***inferior occipital convolutions*** (Fig. 175,

o_1, o_2, o_3) run nearly horizontally backward, and are separated by the superior and inferior occipital fissures; anteriorly these convolutions are continuous with the parietal and temporo-sphenoidal convolutions, through the four *annectent gyri*. The first three have been previously described; the lowest or fourth connects the inferior occipital with the inferior temporo-sphenoidal convolution.

There is usually a small, shallow, vertical fissure, the *transverse occipital*, which passes down behind the external parieto-occipital fissure.

On the median plane is the *cuneate lobule*, which forms part of the occipital lobe, and will be described further on.

The *temporo-sphenoidol lobe* is bounded in front and above by the fissure of Sylvius and its horizontal ramus, and forms that part of the hemisphere which occupies the middle cerebral fossa.

The convolutions of this lobe are three:—

The *superior temporo-sphenoidal* (Fig. 177 T_1,) is bounded above by the horizontal ramus of the Sylvian fissure, and below by the parallel fissure; it is continuous behind with supra-marginal and angular gyri. The *middle temporo-sphenoidal* is joined behind to the angular gyrus and to the middle occipital convolution through the third annectent gyrus; the *inferior* is connected with the inferior occipital convolution through the fourth annectent gyrus.

There are three fissures running from before backward, the *superior temporo-sphenoidal* or *parallel*, the *middle*, and the *inferior*.

The convolutions and fissures of the median and tentorial surfaces can only be properly examined by making an antero-posterior vertical section of the brain through the longitudinal fissure. As this would spoil the brain for future demonstration, the student is recommended to examine one in which this section has been already done.

The *fissures* to be examined on this surface are the callosomarginal, the internal parieto-occipital, the calcarine, the collateral, and the hippocampal or dentate.

The *convolutions* are the præcuneus, the cuneate, the uncinate, the marginal, the gyrus fornicatus, and the dentate.

The *calloso-marginal fissure* commences beneath the rostrum of the corpus callosum, curves in front and round the genu, and then runs nearly parallel with the anterior two-thirds of the corpus callosum; then, changing its direction, it ascends obliquely and terminates on the external aspect of the hemisphere, where it forms a deep notch immediately behind the fissure of Rolando.

The *internal parieto-occipital fissure* passes nearly vertically downward, and joins the calcarine fissure at an acute angle.

FIG. 178.



CONVOLUTIONS AND FISSURES OF THE MEDIAN AND TENTORIAL SURFACES OF RIGHT HEMISPHERE.

C.C. Corpus callosum.
A.F. Ascending frontal convolution.
A.P. Ascending parietal convolution.
P.O. Parieto-occipital fissure.
P.C. Præcuneus or quadrate lobe.
C. Cuneus.
c. Calcarine fissure.
Cl.f. Collateral fissure.
G.F. Gyrus fornicatus.
cm. Calloso-marginal fissure.
fi. Superior frontal or marginal convolution.
O.T. Optic thalamus.
P. Pituitary body.
dc. Dentate convolution.
to_4. Gyrus occipito-temporalis lateralis.
to_5. Gyrus occipito-temporalis medialis.

The *calcarine fissure* begins close to the posterior border of the cerebrum, and then, running nearly horizontally forward, is joined by the preceding fissure; it terminates a little below the posterior border of the corpus callosum. It corresponds with the hippocampus minor in the posterior horn of the lateral ventricle.

The *collateral fissure* (Fig. 178, *Cl.f*) is situated below the calcarine fissure, and runs parallel with it. It separates the

superior and inferior occipito-temporal convolutions, and causes the eminentia collateralis in the descending horn of the lateral ventricle.

The *hippocampal fissure* takes its origin in the posterior part of the gyrus fornicatus, and, passing downward and forward, it ends by forming the notch in the uncinate gyrus. It corresponds to the hippocampus major.

The *præcuneus*, or *quadrate lobule* Fig. 178, P.C), is bounded in front by the calloso-marginal fissure, behind by the internal parieto-occipital fissure. It consists of numerous convolutions, and belongs to the parietal lobe.

The *cuneate lobule* (Fig. 178, C) is triangular in shape, and is situated between the converging internal parieto-occipital and calcarine fissures. It is part of the superior occipital convolution.

The *gyrus fornicatus* (Fig. 178, G.F) begins in front of the locus perforatus anticus, and winds round the corpus callosum, keeping close to its upper surface. It curves round its posterior free border, and is continuous below with the uncinate gyrus. Between this convolution and the corpus callosum is a well-marked furrow, which sometimes takes the name of the *ventricle of the corpus callosum*. The part of the convolution that forms the boundary of this ventricle is termed the *labium cerebri*.

The *dentate convolution* (*fascia dentata*, Fig. 178, *dc*) is the gray convolution lying in the dentate fissure, and takes its name from the notched appearance it presents, owing to the arrangement of the choroid arteries as they pass in through the fissure into the descending horn of the lateral ventricle.

The *uncinate gyrus* (Fig. 177, U) is the anterior part of the superior occipito-temporal convolution, which ends in a peculiar hook-like process at the front of the temporo-sphenoidal lobe.

The *inferior occipito-temporal convolution* (Fig. 178, to_4) is situated between the collateral and the inferior temporo-sphenoidal fissures, and runs backward to the posterior border of the cerebrum.

The *marginal convolution* skirts the longitudinal fissure from the locus perforatus anticus, as far as the termination of the calloso-marginal fissure. It is frequently indented by secondary furrows, and usually is joined to the gyrus fornicatus.

The *island of Reil*, or the *central lobe*, lies deeply in the fissure of Sylvius, not far from its commencement. It is triangular in shape, the apex being close to the anterior perforated spot, and from it radiate outward five or six short convolutions (*gyri operti*), which are separated from the operculum by a deep fissure. In the normal position of the brain it forms the floor of the lenticular nucleus of the corpus striatum. It appears very early in fœtal life, and is at first very prominent, but subsequently becomes closed in by the increasing development of the temporo-sphenoidal lobe.

Nomenclature of the Parts at the Base of the Brain.

The several objects seen at the base of the brain should now be examined, proceeding in order from the front (Fig. 170, p. 732). In this description the cerebral nerves are omitted. These will be examined hereafter.

In front we notice the triangular frontal lobes, separated from each other by the longitudinal fissure, and bounded behind by the fissure of Sylvius.

In the middle line, dividing the frontal lobes, is the longitudinal fissure. By gently separating these lobes, we expose the corpus callosum, or the great transverse commissure which connects the two hemispheres of the cerebrum. Continued backward and outward on each side, from the corpus callosum to the fissure of Sylvius, is a white band, the peduncle of the corpus callosum. Extending from the corpus callosum to the optic commissure is a thin gray layer, the lamina cinerea. Between the frontal and temporo-sphenoidal lobes is the fissure of Sylvius, which lodges the middle cerebral artery. The optic commissure, formed by the union of the two optic tracts, is seen in the middle line behind the lamina cinerea. At the root of the fissure of Sylvius is the locus perforatus anticus.[1] Immediately behind the optic commissure is a slight prominence of gray matter, the tuber cinereum; from this descends a conical tube of reddish color, the infundibulum, to the apex of which is attached the pituitary body. Behind the tuber cinereum are two round white bodies, the corpora albicantia. Posterior to these is the locus perforatus posticus,

[1] Called *perforatus* from its being perforated by a number of blood vessels for the supply of the corpus striatum.

which is bounded behind by the pons, and laterally by the two diverging crura cerebri, two round cords of white substance, which emerge from the anterior border of the pons. Winding round the outer side of each crus is a soft, white band, the optic tract.

Examine now in detail the various objects above enumerated, most of which are shown in Fig. 170.

The *longitudinal fissure* is visible in front, where it separates the two frontal lobes, and, by lifting up the cerebellum, it can be seen behind, dividing the temporo-sphenoidal lobes. It can be more satisfactorily examined later on.

The *lamina cinerea* is a thin layer of gray substance, which runs backward from the termination of the corpus callosum, and passes above the optic commissure to be connected with the tuber cinereum. Laterally it is continuous with the gray matter of the two anterior perforated spots. If the lamina be torn, which is very easily done, an opening is made into the anterior part of the floor of the third ventricle.

The *olfactory lobe* lies in its own sulcus on the orbital surface, nearer its mesial aspect.

The *optic commissure* is placed immediately behind the lamina cinerea. It is formed by the junction in the middle line of the two optic tracts. From it the two optic nerves can be traced, running forward and outward.

The *locus perforatus anticus* is a shallow, triangular depression, placed to the inner side of the commencement of the fissure of Sylvius. It is bounded in front by the two diverging white roots of the olfactory lobe, and behind by the optic tract. It is composed partly of gray substance, and is continuous with the lamina cinerea on the inner side. Crossing it is seen a broad, white band, the *peduncle of the corpus callosum*. This space is pierced by a number of small apertures for the transmission of small vessels to the corpus striatum; hence its name.

The *tuber cinereum* (Fig. 170, p. 732) is a prominence of gray matter immediately behind the optic commissure, and in front of the corpora albicantia. It forms part of the floor of the third ventricle, and from it a conical tube of reddish color, the *infundibulum*, descends to the posterior lobe of the pituitary body. There

is a large collection of gray matter on the outer side of the tuber cinereum, and internal to the optic tract, called the *basal optic ganglion*, from which fibres pass to the corresponding optic tract.

The *pituitary body* occupies the sella turcica, is of a reddish-brown color, and consists of two lobes. Of its two lobes the anterior and larger is concave posteriorly, to receive the posterior lobe, and weighs from five to ten grains. The two lobes consist of different structure, and differ in their development; the *posterior* is developed downward from the third ventricle, and is hollow; subsequently there is a large increase of connective-tissue structure, and blood vessels in it, so that the cavity is usually obliterated. The *anterior* is darker, and is surrounded by a connective-tissue capsule; on section it resembles in structure the thyroid gland, being composed of reticular tissue, with numerous cavities filled with nucleated cells and granular matter; it is originally developed as a prolongation from the ectoderm of the buccal cavity, from which it soon becomes isolated.

The *corpora albicantia* (*mammillaria*) are two round, white bodies, situated behind the tuber cinereum. Each is formed by the curl upon itself of the anterior crus of the fornix, called the *bulb of the fornix*, which then turns backward and upward to end in the optic thalamus. They contain within them some gray matter, and up to the seventh month of fœtal life they form one mass.

The *locus perforatus posticus* (*pons Tarini*) is a depression of gray matter placed between the diverging crura cerebri and behind the corpora albicantia. Its surface is penetrated by small vessels which supply the optic thalami. From its gray substance some white fibres emerge and turn round over the crura cerebri to enter the white medullary portion of the cerebellum.

The *crura cerebri* (Fig. 170) are the two rounded masses of white matter which emerge from the anterior border of the pons Varolii, and then pass forward and outward to enter the anterior and inner aspect of the temporo-sphenoidal lobes. Each is about three-quarters of an inch long, and is rather broader in front than behind. On the inner side the third nerve is seen emerging from a groove (*oculo-motor*) in the crus, which marks the division of the crus into two portions, an upper (dorsal) and larger, called the

tegmentum, and a lower or ventral, called the *crusta*. The optic tract curves round the anterior part of each crus, and is adherent to it by its anterior border.

Structure of the Crura Cerebri. These are composed of longitudinal fibres, derived from the pyramids, from part of the lateral and restiform columns of the medulla, and from the gray matter in the pons Varolii. If one of the crura be divided longitudi-

FIG. 179.



DIAGRAM OF THE COURSE OF THE FIBRES THROUGH THE MEDULLA AND PONS.

nally, there is found in the middle of it a layer of dark-colored nerve-substance, called *locus niger*, which separates the crus into an upper and lower stratum of fibres. The lower stratum (*crusta*) is tough and coarse, and consists of the continuation of the fibres proceeding from the pyramid and the pons. The upper stratum (*tegmentum*) is much softer and finer in texture; it is composed of the fibres proceeding from the lateral and restiform columns;

also from the superior crus of the cerebellum. Tracing the fibres of the crus cerebri into the cerebral hemisphere, we find that its lower fibres ascend chiefly through the corpora striata, its upper fibres through the thalami optici. In passing through these ganglia, the crus receives a large addition to its fibres; these branch out widely toward all parts of the hemisphere, in order to reach the cortical substance on the surface.

Origin of the Cerebral Nerves. The *cerebral nerves* are given off in pairs, named the first, second, third, etc., according

FIG. 180.



DIAGRAM OF THE ORIGINS OF THE OLFACTORY AND OPTIC NERVES.

to the order in which they appear, beginning from the front. There are twelve pairs. Some are nerves of special sense—as the olfactory, the optic, the auditory; others are nerves of common sensation—as the larger root of the fifth, the glosso-pharyngeal, and the pneumogastric; others, again, are nerves of motion—as the third, the fourth, the smaller root of the fifth, the sixth, the facial, the spinal accessory, and the hypoglossal.

First Pair, or Olfactory Nerves. These (Fig. 180, 1) are, from their early development, outgrowths from the cerebral lobes, and

not, strictly speaking, nerves. The nerve is triangular on section, the apex of the triangle being lodged in a straight furrow (*olfactory sulcus*) in the orbital surface of the frontal lobe. It proceeds straight forward, and terminates in the olfactory bulb, which lies on the cribriform plate of the ethmoid bone.

The olfactory lobe is oval, of a reddish-gray color, and very soft consistence, owing to the large amount of gray matter contained in it. It gives off from its under surface about twenty branches, which pass through the foramina in the cribriform plate.[1] For description of these, see p. 296.

The nerve arises by three roots—an outer and an inner, composed of white matter, and a middle, composed of gray (Fig. 180).

The *outer root* passes backward and outward, as a thin, white line, along the outer side of the locus perforatus anticus, to the commencement of the fissure of Sylvius. Its deeper origin has been traced to a nucleus of gray matter in the anterior part of the temporo-sphenoidal lobe.

The *inner root* passes backward and inward to the posterior extremity of the internal convolution of the frontal lobe, and thence may be traced to the gyrus fornicatus.

The *middle* or *gray root* arises from the gray matter of the sulcus in which the nerve is lodged, and from the gray matter of the locus perforatus anticus in the fork between the two white roots; this is called the *tuber olfactorium*. It contains white fibres in its interior, which have been traced to the corpus striatum.

Second Pair, or Optic.

The *optic tracts* arise from the anterior lobes (*nates*) of the corpora quadrigemina, the corpora geniculata, and the posterior part of the optic thalami (Fig. 180).

[1] The olfactory nerve and its ganglion, as stated above, are integral parts (the prosencephalic lobe) of the brain. What in human anatomy is called the origin of the nerve is, in point of fact, the crus of the olfactory lobe, and is in every way homologous to the crus cerebri or cerebelli. In proof of this, look at the enormous size and connections of the crus in animals which have very acute sense of smell. Throughout the vertebrate kingdom there is a strict ratio between the sense of smell and the development of the olfactory lobes. Again, in many animals, these lobes are actually larger than the cerebral, and contain in their interior a cavity which communicates with the lateral ventricles. According to Tiedemann, this cavity exists even in the human fœtus at an early period.

They wind round the crura cerebri, with which they are connected by their anterior borders, and after receiving some fibres from the basal optic nucleus (p. 758), join in the middle line to form the *optic commissure*. This commissure rests upon the sphenoid bone in front of the sella turcica; and from it each optic nerve, invested by its fibrous sheath, passes through the optic foramen into the orbit, and terminates in the retina.

At the commissure some of the nerve fibres cross from one side to the other. The decussation affects only the middle fibres of the nerve; the outer fibres pass from one optic tract to the optic nerve of the same side; the inner fibres pass from one optic tract round to the optic tract of the opposite side; while in front of the commissure are fibres which pass from one optic nerve to its fellow (p. 767).

Third Pair, or Motores Oculorum.

The apparent origin of the *third nerve* is from the inner side of the crus cerebri, immediately in front of the pons. Some of its roots, however, pass through the locus niger and the tegmentum of the crus, to reach a nucleus of large yellow cells beneath the iter a tertio ad quartum ventriculum, extending forward as far as the posterior commissure, and behind as far as the nucleus of the fourth nerve (see below). It runs forward through the cavernous sinus, and, passing through the sphenoidal fissure in two divisions, supplies all the muscles of the orbit except the superior oblique and the external rectus.

Fourth Pair, or Trochlear Nerves.

The *fourth nerve* has its deep origin from a nucleus of gray matter in the floor of the aqueduct of Sylvius, beneath the corpora quadrigemina, and almost continuous superiorly with the yellow nucleus of the third nerve. The nerve fibres then run backward, upward and inward in the lateral wall of the Sylvian aqueduct, and reach the anterior part of the valve of Vieussens, where they cross over to the opposite side. The nerve then emerges from the valve of Vieussens close to the middle line, and winding round the crus cerebri, enters the orbit through the sphenoidal fissure and supplies the superior oblique.

Fifth Pair, or Trigeminal Nerves.

The *fifth nerve* is the largest of all the cranial nerves, and consists of two roots, a larger or sensory, and a smaller or motor. It has its

apparent origin from the outer side of the pons Varolii, and a few of the transverse fibres of this body separate the two roots of the fifth. The *motor* or *smaller root* consists of fibres which take origin from an oval gray nucleus (*motor nucleus*) situated in the front part of the floor of the fourth ventricle, internal to its lateral boundary; in their passage forward the fibres are joined by filaments from the descending root of the fifth, which arise from the gray matter in the lateral wall of the aqueduct of Sylvius, beneath the anterior lobes of the corpora quadrigemina. It also receives some fibres from the raphé. The *sensory* and *larger root* arises by fibres having their origin chiefly in the *superior sensory nucleus*, which is situated external to the motor nucleus, and partly by fibres known as the ascending fibres, which may be traced far down in the medulla from a mass of nerve cells in connection with the gray tubercle of Rolando and its upward prolongation. The two divisions of the nerve proceed forward over the apex of the petrous portion of the temporal bone; here is developed, upon the sensory root, the Gasserian ganglion. The root then divides into three branches—the *ophthalmic*, which passes through the sphenoidal fissure; the *superior maxillary*, which passes through the foramen rotundum; the *inferior maxillary*, which passes through the foramen ovale. They all confer common sensation upon the parts they supply, which comprise the entire face and sides of the head. The small motor root passes beneath the ganglion, with which it has no connection, and accompanies the inferior maxillary division, to be distributed to the muscles of mastication.

Sixth Pair, or Abducentes. The *sixth nerve* emerges from the transverse groove between the pons and the anterior pyramid (p. 732), with both of which it is connected. Its deep origin can be traced to an oval gray mass of nerve cells in the fasciculus teres in the floor of the fourth ventricle, close to the median groove and in front of the transverse striæ. The nerve fibres pass downward from their origin through the pons, parallel with the septum, and emerge from the transverse groove, as before stated. It leaves the skull through the sphenoidal fissure, and passing between the two heads of the external rectus, is distributed to this muscle.

Seventh Pair, or Facial Nerves. The *facial nerve*, or *portio dura* (p. 732), has its apparent origin from the groove between the pons and the restiform tract, and behind the olivary body. Its deep origin may be traced to an elongated mass of gray substance, placed deeply in the floor of the fourth ventricle, between the motor nucleus of the fifth and the transverse striæ. From this origin its fibres run upward, backward and inward to the floor of the fourth ventricle, and wind round the nucleus of the sixth, so as to course superficial to it in the fasciculus teres. The nerve then makes a sharp bend upon itself, and passes downward and outward through the pons between the superior olivary

FIG. 181.

a. Superior peduncles of the cerebellum.
b. Restiform tracts.
c. Posterior pyramids.
d. Fasciculus teres; external to it is the superior fovea.
e. Striæ acusticæ.
f. Fasciculus teres.
g. Tuberculum acusticum.
h. Inferior fovea.
i. Cerebellum.

VIEW OF THE FLOOR OF THE FOURTH VENTRICLE.

nucleus and the ascending root of the fifth nerve. A small separate fasciculus of this nerve—*pars intermedia*—lies between it and the auditory nerve, and forms connections with both; it arises from the lateral column of the cord. The nerve enters the meatus auditorius internus. For the further description of the portia dura, see p. 61.

Eighth Pair, or Auditory Nerves. The *auditory nerve* emerges from the same groove as the preceding nerve, and is situated immediately beneath it, being separated from it only by the pars intermedia. Its deep origin is principally from the *inner auditory nucleus*, situated in the floor of the fourth ventricle, under the tuberculum acusticum; this nucleus extends from beneath the

acoustic tubercle to the middle of the anterior half of the floor, passing beneath the transverse striæ; on its inner side, below, is the vagal nucleus, on its outer side is the restiform body; from this nucleus the fibres pass outward, and, on curving round the restiform body, are joined by some filaments from the transverse striæ. A few of the filaments of the auditory nerve come from another nucleus situated in front of the medullary striæ and external to the preceding nucleus, and which gets larger as it passes upward. These two bundles unite, and the nerve passes outward and enters the meatus auditorius internus in company with the portio dura. It divides at the bottom of the meatus into cochlear and vestibular branches, which are distributed to the internal ear.

Ninth Pair, or Glosso-Pharyngeal Nerves. The *glosso-pharyngeal nerve* arises, apparently by several filaments, from the restiform body below the auditory nerve. Its deep origin is from a nucleus in the inferior fovea of the fourth ventricle, continuous behind with the vagal nucleus, and covered in front by the inner auditory nucleus. The glosso-pharyngeal nerve passes through the middle compartment of the foramen jugulare, and is distributed to the mucous membrane of the pharynx and back of the tongue (p. 284).

Tenth Pair, or Pneumogastric Nerves. The *pneumogastric nerve* arises from a gray nucleus (divided into two by a bundle of white fibres), which is placed between the glosso-pharyngeal nucleus in front and the spinal accessory nucleus behind, in the inferior fovea. The fibres, about twelve in number, pass through the medulla, and emerge from the restiform body, below the glosso-pharyngeal, and join to form a single nerve. This passes through the foramen jugulare, separated from the preceding by a septum of dura mater, and is distributed to the pharynx, larynx, the heart and lungs, the œsophagus and stomach.

Eleventh Pair, or Spinal Accessory Nerves. The *spinal accessory nerve* is composed of two parts: an upper or accessory portion, which arises from the medulla below the vagus; and a lower or spinal portion, which arises from the spinal cord. The accessory fibres may be traced to the gray nucleus, which is connected in

front with the vagal nucleus, and lies close to the median sulcus of the fourth ventricle, extending to the apex of the calamus scriptorius and along the side of the central canal; the spinal portion may be traced below to the tractus intermedio-lateralis and anterior cornu, and above the posterior cornu, arising by slender filaments as low down as the fifth or sixth cervical vertebra. The spinal portion ascends behind the ligamentum denticulatum, through the foramen magnum, into the skull, and joins the accessory part. The nervus accessorius then passes through the foramen jugulare with the two preceding nerves; its accessory portion joins the pneumogastric nerve, and its spinal portion supplies the sterno-mastoid and the trapezius. The course and distribution of this nerve have been described in the dissection of the head and neck.

The Twelfth, or Hypoglossal Nerve.

The *hypoglossal nerve* arises by several filaments from the medulla, which emerge from the groove between the anterior pyramid and the olivary body. Its fibres may be traced to a long gray nucleus, which forms an eminence in the floor of the fourth ventricle, in front and to the inner side of the vagal nucleus. The filaments are collected into two fasciculi, which pierce the dura mater through two apertures and join in the anterior condylar foramen; it is distributed to the muscles of the tongue and the depressor muscles of the os hyoides and larynx.

Dissection of the Brain.

The brain should now be laid on its base. We first notice a median fissure, separating the cerebrum into two symmetrical hemispheres: this is the *longitudinal fissure*. By gently separating the hemispheres we see that the fissure extends in front and behind to the base, but that in the middle there is at the bottom a white band of nerve substance, which is the great transverse commissure of the cerebrum, and termed the *corpus callosum*, upon which are seen the two anterior cerebral arteries.

Slice off the hemispheres to about half an inch above the level of the corpus callosum. The cut white surface presents an oval appearance, and is called the *centrum ovale minus*. The white substance is surrounded by a tortuous layer of gray matter about one-eighth of an inch in thickness. This gray substance consists

of four layers—two of gray alternating with two of white, the most external layer being white. In some places, chiefly at the base of the brain, six layers have been demonstrated. The white substance is spotted with red dots (*puncta vasculosa*); these are due to the escape of blood from the divided vessels. The corpus callosum is now seen to be overlaid on each side by the gyrus fornicatus; the border is termed the *labium cerebri*, and the space between the gyrus and the corpus callosum is called the *ventricle of the corpus callosum.*

Now slice off the hemisphere down to the level of the corpus

FIG. 182.



UPPER SURFACE OF CORPUS CALLOSUM.

1, 1. Lineæ transversæ.
2. Raphé.
3, 3. Anterior cerebral a.

FIG. 183.



DIAGRAM OF LAMINA CINEREA.

1, 1. Peduncles of corpus callosum.
2. Lamina cinerea.
3. Commissure of optic nerves.

callosum, when a section is made of the white substance, called the *centrum ovale majus.* The corpus callosum is now well exposed.

Corpus Callosum. This stratum of white substance, consisting of transverse commissural fibres, is the chief connecting medium between the two hemispheres, and is called the *great transverse commissure* of the cerebrum; and moreover, on each side forms the roof of the lateral ventricles. Its surface is

slightly arched from before backward; it is about four inches long and one inch in its greatest breadth, which is behind. It is rather nearer to the front than to the back part of the brain, and it is thicker at the ends than in the middle, and thicker behind than in front. A shallow groove, called the *raphé*, runs along the middle of its upper surface (Fig. 182); in a fresh brain, two longitudinal white tracts, named *striæ longitudinales*, or the *nerves of Lancisi*, run parallel to it; and external to these again are two

FIG. 184.



VERTICAL SECTION THROUGH THE CORPUS CALLOSUM, AND PARTS BELOW.

other longitudinal fibres, *striæ longitudinales laterales*. The surface of the corpus callosum is marked by transverse lines which indicate the course of its fibres; these are the *lineæ transversæ* of the old anatomists. The anterior cerebral arteries proceed along the surface of the corpus callosum to the back of the brain.

The anterior part of the corpus callosum turns downward and backward, forming a bend called its *genu*. The inferior part of this bend—*rostrum*—becomes gradually thinner and narrower, and terminates in two peduncles, which diverge from each other,

and are lost, one in each fissure of Sylvius. Between these crura is placed the lamina cinerea (Fig. 180). The posterior part of the corpus callosum terminates in a thick, round border—the *splenium*—which is free, and beneath it the pia mater enters the interior of the ventricles. A satisfactory view cannot be obtained of the arch formed by the corpus callosum, of its terminations in front and behind, and of the relative thickness of its different parts, without making a perpendicular section through a fresh brain, as shown in the preceding figure.[1]

Connected with the under surface of the posterior part of the corpus callosum is the fornix, which separates from it in front, the two structures being connected by a vertical septum—the *septum lucidum* (Fig. 184).

Lateral Ventricles. A longitudinal incision should be made on each side through the corpus callosum, about half an inch from its median raphé. Care must be taken not to cut too near the middle line, in order to preserve the delicate partition which descends from the under surface of the corpus callosum, and separates the ventricles from each other. Two cavities, called the *lateral ventricles*, will thus be exposed, one in each cerebral hemisphere, and they should afterward be laid open throughout their whole extent. Their general form should be first examined; then the several objects seen in them.

The *lateral ventricles* are two serous cavities, one in each hemisphere of the brain. They are occasioned by the enlargement and folding backward of the cerebral lobes over the other parts of the central nervous axis. They contain a serous fluid, which, even in a healthy brain, sometimes exists in considerable quantity; when greatly in excess it constitutes one form of the disease, termed hydrocephalus. The ventricles are lined with ciliated epithelium, laid upon a layer of neuroglia (*ependyma*); a term which has been applied to that peculiarly delicate connective tissue found throughout the brain and spinal cord.

The ventricles are crescentic in shape, with their backs toward

[1] The corpus callosum is more or less developed in all mammalia, but is absent in birds, reptiles and fish. It has been absent in the human subject without any particular mental deficiency. See cases recorded by Reil, *Archiv. für die Phys.*, t. xi, and Wenzel, *De penitiori Struct. Cereb.*, p. 302.

each other. Each consists of a central part or *body*, and three horns or *cornua*, *anterior*, *middle* and *posterior*, which extend, respectively, into the frontal, temporo-sphenoidal and occipital lobes. The body, situated in the middle of the hemisphere, is triangular in shape, and is separated from its fellow by the septum lucidum. Its roof is formed by the corpus callosum (Fig. 186, 1);

FIG. 185.



VIEW OF LATERAL VENTRICLES FROM ABOVE, AFTER THE REMOVAL OF THE CORPUS CALLOSUM.

1. Corpus striatum.
2. Optic thalamus.
3. Tænia semicircularis.
4. Fornix.
5. Hippocampus minor.
6. Hippocampus major, with the eminentia collateralis behind it.
7. The corpus callosum (cut through).
8. Fifth ventricle
9. Pes Hippocampi.
10. Choroid plexus.

internally, it is bounded by the septum lucidum (Fig. 184); on the floor, beginning from the front, are seen the *corpus striatum*, the *tænia semicircularis*, the *optic thalamus*, the *choroid plexus*, and the *corpus fimbriatum* of the *fornix* (Fig. 185).

The *anterior horn* extends into the frontal lobe, and as it passes forward it diverges slightly from its fellow of the opposite side. It is triangular in shape, its roof and anterior wall are formed by

the corpus callosum, and it curves round the anterior extremity of the corpus striatum.

The *posterior horn* can be traced into the occipital lobe, where it passes at first backward and outward, and then, narrowing to a point, converges toward its fellow. Its roof is formed by the fibres of the corpus callosum as they pass backward and outward from the splenium; on its floor are seen, on the inner side, an eminence, the *hippocampus minor*, and external to it a triangular flat surface, called the *pes accessorius* or *eminentia collateralis*.[1]

FIG. 186.



TRANSVERSE VERTICAL SECTION THROUGH THE BRAIN.

1. Corpus callosum.
2. Lateral ventricle.
3. Third ventricle.
4. Corpus striatum.
5. Thalamus opticus.
6. Corpus mammillare.
7. Choroid plexus.
8. Fornix.
9. Pituitary gland.

The *middle* or *descending horn* runs into the temporo-sphenoidal lobe, descends toward the base of the brain, making a curve, at first backward and outward, then downward and forward, and lastly inward; the initial letters of which make the memorial word "*bodfi*." Its roof is formed by the fibres of the corpus callosum, partly by the posterior narrow extremity of the corpus

[1] The posterior horns are not always equally developed in both hemispheres, and sometimes they are absent in one or both.

In the carnivora, ruminantia, solipeda, pachydermata and rodentia, the lateral ventricles are prolonged into the largely-developed olfactory lobes. This is the case in the human fœtus only at an early period.

striatum, with the tænia semicircularis, and the rounded extremity of the optic thalamus. On its floor are the *hippocampus major*, a large, rounded, white eminence which follows the curve of the cornu; the *pes hippocampi*, the expanded paw-like extremity of the former; the *eminentia collateralis* on the outer side of the hippocampus major, and part of which is seen in the posterior horn; the *corpus fimbriatum* of the fornix, attached to the anterior concave border of the hippocampus major; the *fascia dentata*, a crimped edge of gray matter under the corpus fimbriatum; the *choroid* plexus; and the outer part of the *transverse fissure*.

The various structures seen in the body and horns of the lateral ventricle will be described later on, when they are fully exposed.

Appearance on Perpendicular Section.

If a vertical transverse section is made across the middle of the brain, the lateral ventricles would appear as represented in Fig. 186. Observe that the roof and the floor are almost in actual contact, unless separated by ventricular fluid. Together with the third or middle ventricle, their shape slightly resembles the letter T. Such a section shows well the radiating fibres of the corpus callosum, the fornix, and the velum interpositum beneath it; also the beginning of the transverse fissure at the base of the brain, between the crus cerebri and the temporo-sphenoidal lobe.

If the corpus callosum be slightly raised, a thin vertical median septum, *septum lucidum*, will be seen, extending from the under aspect of this body to the under surface of the fornix.

Septum Lucidum.

This is a thin and almost translucent partition which descends vertically in the middle line from the under surface of the corpus callosum, and separates the anterior part of the lateral ventricles from each other. It is attached above to the corpus callosum, below to the reflected part of the corpus callosum and fornix (Fig. 184). It is not of equal depth throughout. Its broadest part is in front and corresponds with the knee of the corpus callosum. It becomes narrower behind, tapering to a thin point, where the corpus callosum and the fornix become continuous. The septum consists of two layers, which enclose a space called the *fifth ventricle* or the *ventricle of the septum* (Fig. 185). Each layer consists of gray matter inside and of white matter outside; the former representing the cortical, the

latter the medullary substance of the brain. The cavity is not lined with epithelium, as is the case with those ventricles developed from the cerebral vesicles.[1]

Cut transversely through the corpus callosum about its middle, with the septum lucidum, and turn forward the anterior half. In this way the ventricle of the septum will be exposed. By turning back the posterior half of the corpus callosum a view is obtained of the fornix. This proceeding requires care, or the fornix will be reflected also, since these two arches of nerve substance are here so closely connected.

FIG. 187.



DIAGRAM OF THE FORNIX.

(The arrow is passed through the foramen of Monro.)

1, 1. Corpora striata.
2, 2. Thalami optici.
3, 3. Anterior crura of fornix bending down to join the corpora mammillaria.
4, 4. Posterior crura of the fornix joining the hippocampi.
5, 5. Choroid plexus.
6, 6. Hippocampi majores.
7. Corpus callosum cut through.
8. Ventricle of septum lucidum.

Fornix. The *fornix* is a layer of white matter, extending in the form of an arch (whence its name) from before backward, beneath the corpus callosum. It is the great longitudinal white commissure, and lies over the velum interpositum (Fig. 187). Viewed from above, it is triangular with the base backward, and is called the *body;* from its anterior

[1] The development of the septum lucidum commences about the fifth month of fœtal life, and proceeds from before backward, *pari passu* with the corpus callosum and the fornix. It is developed from the lower part of the great longitudinal fissure, but becomes shut off from it in the process of development.

narrow part are given off the *two anterior crura*, and from its posterior and outer part the *two posterior crura*.

The *body* is the broad triangular part with the narrow portion in front. The posterior broad part is connected with the corpus callosum; in front of this it arches downward, so as to leave the corpus callosum, to which, however, it is still connected by the septum lucidum. Its lateral free edges rest on the choroid plexuses, and are seen on the floor of the lateral ventricles.

The *anterior pillars* or *crura* proceed from the front narrow part of the body, one on each side of the mesial line. As they pass forward the crura diverge slightly, and descend through a mass of gray matter in the sides of the third ventricle toward the base of the brain, where, making a sudden bend upon themselves, they form the corpora mammillaria, from which they may be traced backward and upward, each to the anterior nucleus of the optic thalamus of its own side. As they descend, the anterior crura are joined by the peduncles of the pineal body, by the tænia semicircularis, and by fibres from the septum lucidum. Immediately behind and below the anterior crura is a triangular passage, through which the choroid plexuses of opposite sides are continuous with each other. This aperture is called the *foramen of Monro*. Strictly speaking, it is not a foramen, but only an interval caused by the anterior crus arching over the groove between the corpus striatum and optic thalamus on each side; it establishes a communication between the two lateral and third ventricles, and is in shape like the letter Y, the passage from each lateral ventricle passing downward and inward, and meeting below, to be continued as a single passage for a short distance before opening into the third ventricle.

The *posterior pillars* or *crura* are continued downward and outward from the thickened free borders of the body of the fornix, and are at first connected to the under surface of the corpus callosum. Each leaves the body at the posterior and outer angle as a thin, flat, white band, resting on the choroid plexus and the pulvinar of the optic thalamus, and curving downward and outward, becomes intimately connected with the concave border of the hippocampus major as far down as the pes hippocampi, gradually tapering to a point at its termination. The free border

of the posterior crus is known as the *taniæ hippocampi*, or the *corpus fimbriatum;* and on raising this up we expose an indented layer of gray matter, the *fascia dentata*, which is the free border of the cortical substance of the cerebrum.[1]

The fornix should now be cut through transversely, and its two portions reflected backward and forward respectively. On the under surface of the posterior portion are seen fibres, passing transversely, belonging to the corpus callosum, and forming what is termed the *lyra*.

Between the fornix and the upper surface of the cerebellum is the *transverse fissure*, or *fissure of Bichat*, through which the pia mater enters the ventricles. The fissure extends from the middle downward on each side to the base of the brain, as far as the end of the descending horn. It is of a horse-shoe shape, with the concavity directed forward. The upper boundary of that part of the transverse fissure which extends into the middle horn is sometimes called the *free margin of the hemisphere*.

The contents of the lateral ventricles should now be examined more in detail.

Corpus Striatum. The *corpus striatum* is so called because, when cut into, it presents alternate layers of a white and gray substance. It is a large ovoid mass of gray substance, part of which forms an eminence in the body of the lateral ventricle (the intraventricular portion), but the larger part (extraventricular portion) is imbedded in the white substance of the cerebrum. The *intraventricular portion*, called the *nucleus caudatus*, is pear-shaped, broad in front, and when traced backward is found to taper gradually to a point on the outside of the optic thalamus (Fig. 188). Its surface is of pinkish-gray color, and is crossed by numerous small veins (*venæ corporis striati*), which open into the venæ Galeni. When a horizontal cut is made into it, it shows a thin layer of white substance covering a mass of gray streaked with white. The *extraventricular portion* or *nucleus lenticularis*, can only be seen on a horizontal section being made outward; the section reveals a biconvex mass of gray matter, separated from the

[1] The fornix and septum lucidum are absent in fish; they are merely rudimentary in reptiles and birds; but all mammalia have them in greater or less perfection, according to the degree of development of the cerebral hemispheres.

nucleus caudatus by a broad band of white substance, the *internal capsule*, and corresponds with the island of Reil. Running parallel with the outer border of the nucleus lenticularis, but separated from it by a thin layer of white substance, the *external capsule*, is a wavy streak of gray matter, the *claustrum*, of variable thickness. Outside the claustrum is another layer of white

FIG. 188.



VIEW OF THE LATERAL VENTRICLES AND VELUM INTERPOSITUM, AFTER REFLECTING THE FORNIX.

1. Anterior horn.
2. Corpus striatum.
3. Tænia semicircularis.
4. Optic thalamus.
5. Velum interpositum, with the venæ Galeni.
6. Lyra.
7. The posterior half of the fornix turned backward.
8. Hippocampus minor.
9. Hippocampus major.
10. Eminentia collateralis.
11. Fifth ventricle.
12. Choroid plexus.

matter, and then we see the indented convolutions of the island of Reil. If a vertical transverse section be made through the nucleus lenticularis, it appears triangular and intersected by two white lines, which divide it into three parallel gray bands. Beneath the lenticular nucleus is a mass of gray matter, called the *nucleus amygdalæ*, which causes an elevation at the apex of the roof of the middle horn.

Tænia Semicircularis. The *tænia semicircularis*, or *stria terminalis*, is a narrow, semi-transparent band of longitudinal white fibres, which lies in the groove between the corpus striatum and the optic thalamus (Fig. 188). In front, it is connected with the anterior crus of the fornix, and descends with it to the corpus mammillare; it passes backward and outward, and behind it is lost in the white substance of the middle horn of the lateral ventricle. Several veins from the corpus striatum pass underneath the tænia semicircularis to join the venæ Galeni. The upper surface of the tænia is firmer in structure than its deeper part, and is called the *horny band* of Tarinus.

Hippocampus Major. The *hippocampus major* is an elongated convex eminence of gray matter, covered with white, and is situated in the posterior part of the descending horn. It extends to the bottom of the horn, following its curve, where it becomes somewhat expanded and indented on the surface, so as to resemble the paw of an animal, whence its name, *pes hippocampi*. Attached to the front concave border of the hippocampus is the posterior crus of the fornix. It corresponds to the hippocampal fissure, which itself is filled with gray matter, which forms the fascia dentata.

Hippocampus Minor. The *hippocampus minor*, called also *calcar avis* and *ergot*, is a rounded eminence, smaller than the preceding, occupying the inner curved wall of the posterior horn. It consists of white matter externally, and corresponds to the calcarine fissure. Between the hippocampus major and minor is a triangular smooth surface, called the *pes accessorius* or *eminentia collateralis*, and is found in the posterior and the descending horns. This corresponds to the collateral fissure.

Velum Interpositum and Choroid Plexus. The *velum interpositum*, which supports the fornix, should now be examined. This is a layer of pia mater, which penetrates into the ventricles through the transverse fissure, beneath the posterior border of the corpus callosum, as shown in Fig. 184. The shape of this vascular membrane is like that of the fornix, and its borders project beneath that body and form the red convoluted fringes called the *choroid plexuses*. These plexuses consist almost entirely of tortuous ramifications of minute blood vessels, and are covered with vas-

cular villi. The villi themselves are covered with large spheroidal epithelial cells. In front the plexuses communicate with each other through the foramen of Monro; behind, they descend into the middle horns of the lateral ventricles, and become continuous with the pia mater at the base of the brain. From the under surface of the velum two small vascular processes are prolonged into the third ventricle, forming the *choroid plexuses* of that cavity. Along the centre of the velum run two large veins, called *venæ Galeni*, which return the blood from the ventricles into the straight sinus.

Venæ Galeni.

The velum interpositum, with the choroid plexuses, must now be removed to expose the following structures shown in diagram (p. 781): 1. A full view of the *optic thalamus*. 2. Between the optic thalami is the *third ventricle*, a deep vertical fissure, situated in the middle line. 3. Behind the fissure is the *pineal body*, a vascular structure, about the size of a pea. From this body may be traced forward two slender white cords, called its *peduncles*, or *striæ pineales*—one along the inner side of each optic thalamus. 4. Passing transversely across the third ventricle are *three commissures*—anterior, middle and posterior—connecting the opposite sides of the brain. 5. Immediately behind the pineal body are four elevations, two on each side, called the *corpora quadrigemina*, or *nates* and *testes*. 6. These bodies are connected with the cerebellum by two bands, one on each side, termed the *processus a cerebello ad cerebrum*. 7. Between these cords extends a thin layer of gray substance, called the *valve of Vieussens*, beneath which lies the fourth ventricle.

Thalamus Opticus.

This, called also the *posterior cerebral ganglion*, is the convex oval elevation seen immediately behind the corpus striatum and tænia semicircularis. Superficially it is covered with a thin layer of white, but internally it is composed of gray substance. The under surface rests upon the tegmentum of the crus cerebri, and forms part of the roof of the middle horn of the lateral ventricle; externally it is bounded by a broad band of white substance derived from the crusta, which forms the internal capsule, already described. Externally, the optic thalamus is bounded by the tænia semicircularis; superficially, it is covered by the choroid plexus and the fornix; inter-

nally, it forms the lateral boundary of the third ventricle, and has, running along it, the peduncle of the pineal body; posteriorly, it overlaps the sides of the corpora quadrigemina and forms a prominence in the roof of the middle horn, where it receives the crus cerebri. The upper surface of the thalamus is divided into two portions by an oblique shallow groove, passing from before backward; the anterior and outer portion forms a prominent convex surface, called the *anterior tubercle*, which is covered with the epithelium of the lateral ventricle; the posterior and inner portion is pointed in front, and posteriorly enlarges to form a prominent rounded eminence, the *posterior tubercle* or *pulvinar*, and is not lined with epithelium.[1] Beneath the posterior part of the thalamus are two small oval eminences, termed the *corpora geniculata*, *internum* and *externum*. These consist of small accumulations of gray matter, beneath the white; the outer one being situated external to and above the internal, and to the outer side of one of the roots of the optic tract (Fig. 180). From each of these bodies proceeds a white band to join the root just referred to, and from the junction of these three roots (*brachia*) the optic tract has its commencement. A narrow band of white substance connects the external one with the nates, and a similar band connects the internal one with the testes.[2]

Third Ventricle. The *third ventricle* is the long, narrow fissure between the optic thalami, and reaches down to the base of the brain. Its *roof* is formed by the fornix and the velum interpositum, the under aspect of which is lined by the epithelium covering the general ventricular cavities, and is reflected from the velum and choroid plexuses on to the optic thalami; the *floor*, which increases in depth in front, is formed by certain parts at the base of the brain, found within the interpeduncular space—viz.: the locus perforatus posticus, corpora mammillaria, tuber cinereum, infundibulum, and lamina cinerea, all of which are best seen in a vertical section, as shown on page 768. In *front*, it is bounded by the anterior crura of the fornix and

[1] There is a triangular depression between the pulvinar and the peduncle of the pineal body, which has received the name of the *trigonum habenulæ*.

[2] These bands are faintly marked in man, but are more apparent in the lower animals.

the anterior commissure; *laterally*, by the optic thalami and the peduncles of the pineal body; *behind*, by the posterior commissure and the *iter a tertio ad quartum ventriculum*, which is a long canal beneath the corpora quadrigemina, connecting the third with the fourth ventricle.

Commissures. Passing across the third ventricle are seen three commissures, the anterior, middle and posterior. The *middle commissure* may be seen by gently separating the optic thalami, and is about half an inch in breadth. This is composed entirely of gray substance, and in most brains, owing to its softness, is generally torn before it can be examined.[1] The *anterior commissure* is a round white cord, which lies immediately in front of the anterior crura of the fornix, and connects the corpora striata. This commissure may be traced on each side, through the corpora striata, below the nuclei lenticulares, extending backward far into the temporo-sphenoidal lobes. Situated immediately in front of, and rather below the pineal body, is another thin, round, white cord, called the *posterior commissure*. Its fibres pass into the substance of the hemispheres and connect the optic thalami. Its fibres are derived from the fillet which comes from the tegmentum of the crus cerebri.

Middle.

Anterior.

Posterior.

The third ventricle communicates with the lateral ventricles by the two openings of the foramina of Monro, with the fourth ventricle through the iter a tertio ad quartum ventriculum, and in front of its floor by a conical cavity, *iter ad infundibulum*, with the infundibulum.

The third ventricle is covered with an epithelial lining continuous with that of the lateral ventricles through the foramina of Monro; after covering the walls of the third ventricle it lines the aqueduct of Sylvius, to pass to the fourth ventricle.

Pineal Body, or Gland. The *pineal body* (conarium) is a very vascular oval body, situated immediately in front of the

[1] The soft commissure does not appear to be a very essential constituent part of the brain. It is not found before the ninth month of fœtal life, and in some instances, according to our observation, is never developed. Wenzel states that it is absent in about one out of seven subjects (*De penitiori Struct. Cerebri Hom. et Brut.* Tübingen, 1812).

corpora quadrigemina (Fig. 189). It is about the size of a cherry-stone, and is firmly connected with the under surface of the velum, and is apt to be separated from its normal position when that membrane is reflected. It is connected to the cerebrum by two white crura, the *peduncles of the pineal body*, which extend forward, one on the inner side of each optic thalamus along their upper margin, and terminate by joining the anterior crura of

FIG. 189.



the fornix. The peduncles join together behind, in front of the pineal body, and are connected with the front of the posterior commissure.

The pineal body consists of numerous small follicles filled with cells, which are separated by connective tissue; so that in structure it much resembles that of the anterior lobe of the pituitary body. In its interior it contains, beside some viscid fluid, more

or less gritty particles (*acervulus cerebri*), consisting of phosphate and carbonate of lime and phosphate of magnesia and ammonia. Besides the calcareous particles, these follicles contain corpora amylacea; and, when abundant, this sabulous matter is found on the peduncles of the pineal body.

The pineal body is larger in the female than in the male subject, and is largest of all in the child. It is found in all mammalia, birds and reptiles, in the same typical position, but its functions are entirely unknown.

Corpora Quadrigemina.

The *corpora quadrigemina* are four round eminences, situated two on each side, behind the pineal body, and are separated from each other by a crucial depression. Though white on their surface, they contain gray matter in their interior, for the purpose of giving origin to the optic tract. Laterally they are continued outward as two convex white cords, the *anterior* and *posterior brachia*. The anterior brachium passes between the corpora geniculata, and is continued on into the optic tract, of which it may be considered its direct root; the posterior brachium passes forward and outward, and is lost beneath the corpus geniculatum internum. They are situated above the iter a tertio ad quartum ventriculum. The anterior pair are called the *nates*, and are larger and darker than the posterior pair, which take the name of *testes*. A more appropriate term for these bodies would be the optic lobes.[1]

The corpora quadrigemina are developed very early in fœtal life, and at first are only two in number, one on each side of the mesial line; but about the seventh month a transverse groove is apparent, thus mapping out the four bodies.[2]

[1] Eminences homologous to the corpora quadrigemina are found in all vertebrate animals; they are the mesocephalic lobes; they always give origin to the optic nerves, and their size bears a direct relation to the power of sight. They are relatively smaller in man than in any other animal. In birds there are only two eminences, and these are very large, especially in those far-seeing birds which fly high, as the eagle, falcon, vulture, etc., who require acute sight to discern their prey at a distance.

[2] On making a transverse vertical section through the nates, we find that there is a superficial thin layer of white fibres (stratum zonale); beneath this is a crescentic layer of gray matter (stratum cinereum); deeper than this is a thick, biconvex mass of gray matter, with nerve filaments and nerve cells (stratum opticum); and lowest of all is an arched layer of white nerve fibres derived from the fillet (stratum lemnisci).

Processus a Cerebello ad Cerebrum.

By gently drawing back the overlapping cerebellum, two broad white cords are seen, which pass backward, diverging from each other, from the optic thalami and the corpora quadrigemina to the cerebellum (Fig. 189). These are the *processus a cerebello ad cerebrum*, or *superior peduncles of the cerebellum*. They connect the cerebrum and cerebellum, and rest upon the crura cerebri. Below they pass to the inferior vermiform process and to the white matter within the corpus dentatum.

Valve of Vieussens.

The triangular space between the superior peduncles is occupied by a thin layer of gray matter, which covers over the anterior part of the fourth ventricle. This layer is called the *valve of Vieussens*, or the *anterior* or *superior medullary velum;* it is narrow in front and broad posteriorly, where it is connected with the central portion of the cerebellum. Along the mesial line of its upper surface there is an irregular ridge, the *frænulum*, which becomes lost toward its lower part; the lower part is overlapped by a corrugated lobule of gray matter from the anterior part of the cerebellum, and is called the *linguetta laminosa*.

Iter a Tertio ad Quartum Ventriculum, or Aqueduct of Sylvius.

The third ventricle is connected with the fourth by a canal, large enough to admit a probe, which runs downward and backward beneath the posterior commissure and the corpora quadrigemina. It is about half an inch in length, and its shape varies in different parts of its course; in the lower being T-shaped, and in the upper part shield-shaped, on transverse section. In its walls is a large amount of gray matter, in which are the nuclei of origin of the third, fourth and upper part of the fifth cranial nerves. It is lined with ciliated columnar epithelium. This passage, together with the third and fourth ventricles, are persistent parts of the central canal, which in early fœtal life extended down the middle of the cerebro-spinal axis. It subsequently becomes much encroached upon by the large increase of gray substance in the process of development.

Fourth Ventricle.

The fourth ventricle is the space situated between the cerebellum behind and the posterior surface of the medulla oblongata and pons Varolii in front. It is

the dilated portion of the primordial canal alluded to in the last paragraph. If viewed in a vertical section, as represented in the diagram (Fig. 184), it appears triangular, with its base forward; but if seen from behind, it is a lozenge-shaped space, the long axis being antero-posterior (Fig. 190).

The *upper wall* or *roof* of the fourth ventricle is formed by the valve of Vieussens, and by the front of the inferior vermiform process, with the two amygdalæ; *laterally* it is bounded, in front by the processus a cerebello ad cerebrum, and behind by the diverging posterior pyramids and restiform bodies; *below*, by the continuation of the arachnoid membrane on to the posterior surface of the spinal cord, in which there is an aperture called the

FIG. 190.

FLOOR OF THE FOURTH VENTRICLE.

foramen of Magendie; in front its floor is formed by the medulla oblongata and pons Varolii. The pia mater is prolonged for a short distance into the lower part of the cavity, and forms the *choroid plexus* of the *fourth ventricle*.

The anterior wall is diamond-shaped, pointed above and below, while laterally the space broadens out into an angular point, between the cerebellum and the medulla, called the *lateral recess*. Below, the ventricle is bounded by the restiform bodies and posterior pyramids diverging like the branches of the letter V to form the *inferior peduncles* of the *cerebellum;* the divergence of these cords, with the median furrow, was called by the older anatomists

the *calamus scriptorius*, from its fancied resemblance to a writing pen. At the termination of the posterior pyramid there is a slight overhanging thickening, turning over the restiform body at the lateral recess, of which it forms the lateral boundary; it is called the *ligula* or *tænia*. We find also a similar thickening, partly of the lining membrane and partly of nerve matter, arching over the apex of the calamus scriptorius, known as the *obex*.[1]

The *floor*, formed by the posterior surface of the medulla and pons, is marked by a median groove passing from the apex of the calamus scriptorius to the iter. It is divided into two portions, a lower and an upper, by some transverse white fibres called the *striæ acusticæ* or *medullares*, which emerge from the median groove and pass outward over the inferior cerebellar peduncle to join part of the roots of the auditory nerves. The lower part of the floor, on each side, is mapped out into three surfaces by a triangular depression, *fovea inferior* (Fig. 190, *h*), having its apex at the transverse striæ, and its base below at the posterior pyramids. On the outer side of the fovea, there is a convex triangular surface, with its base upward (Fig. 190, *g*), called the *tuberculum acusticum;* on the inner side of the fovea, and bounded internally by the median groove, is the rounded triangular surface which marks the commencement of the *fasciculus teres*. Toward the base of the inferior fovea there is a dark surface of gray matter called the *ala cinerea*, which becomes raised into an eminence (*eminentia cinerea*).

The upper part of the fourth ventricle is that portion between the acoustic striæ and the iter a tertio ad quartum ventriculum. The median groove is still continued upward, although it becomes fainter, and on each side of it is the parallel rounded eminence, the fasciculus teres. Outside this fasciculus is a triangular depression, the *fovea superior;* and passing upward under cover of the superior cerebellar peduncle, we notice a depression of gray substance called the *locus cæruleus*.

The lower part of the fourth ventricle is developed from the

[1] *Obex*, a bar.

metencephalic, the upper part from the epencephalic, portion of the posterior primary vesicle.[1]

Structure of the Cerebrum.

The white substance of the cerebrum consists of medullated fibres, which are, as a rule, smaller than those in the spinal cord. The general arrangement of the fibres may be classified under three heads: 1. The diverging, or peduncular fibres. 2. The transverse commissural fibres; and 3. The longitudinal fibres.

The *diverging* or *peduncular fibres* are derived partly from the crusta, and partly from the tegmentum of the crus cerebri. Those from the crusta pass forward and outward between the nucleus caudatus and nucleus lenticularis with the internal capsule, and in front of these ganglia the fibres radiate outward in all directions, called the *corona radiata*. Most of these fibres pass indirectly to the cortical portion of the cerebrum; some proceed direct to the cortex, through the gray ganglionic structure, among which are the *pyramidal tract*, passing to the gray matter in the neighborhood of the fissure of Rolando, and the *direct sensory tract*, to the cortex of the occipital lobe. The fibres from the tegmentum are joined by others from the processus a cerebello ad cerebrum, and the corpora quadrigemina, and pass under the optic thalamus, and probably through this body, and radiate outward, joining the corona radiata to proceed to the temporo-sphenoidal, post-parietal and occipital lobes.

The *transverse commissural fibres* connect the two hemispheres, and are the corpus callosum, the anterior and posterior commissures.

The *longitudinal fibres* consist of the fornix, the striæ longitudinales of the corpus callosum, the tæniæ semicirculares, the gyrus fornicatus, the gyrus uncinatus, and the peduncles of the pineal body.

THE CEREBELLUM.

The *cerebellum* is that portion of the encephalon situated in the occipital fossa, beneath the posterior lobes of the cerebrum, from

[1] Tiedemann proposed to call the fourth ventricle the first, because in the fœtus it is formed sooner than any of the others; because it exists in all vertebrated animals, whereas the lateral ventricles are absent in all osseous fishes; and because the ventricle of the septum lucidum is absent in all fishes, in reptiles, and in birds.

which it is separated by the tentorium. It measures in its transverse diameter from three and a half to four inches, in its antero-posterior diameter two to two and a half inches, and two inches in its vertical diameter. Its form is ellipsoidal, with the long axis transverse. When the arachnoid membrane and the pia mater are removed, it is noticed that its surface is darker, and not arranged in tortuous convolutions like those of the cerebrum. It is covered externally with gray matter, and consists of a multitude of thin laminæ disposed in a series of nearly parallel concentric curves, with the concavity forward. By a little dissection it is easy to separate some of the laminæ from each other, and to

FIG. 191.



SUPERIOR SURFACE OF THE CEREBELLUM.

i c s. Incisura cerebelli anterior.
i c p. Incisura cerebelli posterior.
a s or *l q*. The anterior superior or quadrate lobe.
p s. The posterior superior lobe.
p i. The posterior inferior lobe.
h f. The great horizontal fissure.
l c. The lobulus centralis.
m c. Monticulus cerebelli.
c s. Commissura simplex.

see that the intervening fissures increase in depth from the centre toward the circumference.

The cerebellum consists of two *lateral hemispheres* united by an intermediate portion, the *vermiform process*, the upper aspect of which takes the name of the *superior vermiform process*, the inferior that of the *inferior vermiform process*. Comparative anatomy proves that this is the fundamental part of the cerebellum, the lateral masses not being developed in the vertebrate series until after the birds. In man the lateral masses form by far the largest part of the cerebellum.

The two hemispheres on their under aspect are convex and

separated from each other by a deep fossa, the *vallecula;* and, behind, this is continued so as to form a deep notch between the posterior borders of the cerebellum.

The upper surface is separated from the lower by a deep fissure named the *great horizontal*, which extends along the free border of each hemisphere.

Upper Surface. The *upper surface* of the cerebellum slopes on each side, having a ridge along the middle line, called the *superior vermiform process.* This process presents three eminences, an anterior, middle and posterior, which are named respectively, the *lobulus centralis*, the *monticulus cerebelli*, and the *commissura simplex.* The hemispheres are separated posteriorly by a deep notch, the *incisura cerebelli posterior*, which receives the falx cerebelli; and anteriorly by a broader notch, the *incisura cerebelli anterior*, which lodges the pons Varolii. On this surface of the cerebellum are two lobes, one of which, the *quadrate*, is situated on its external and anterior aspect; the other, the *posterior* or *crescentic*, is placed along its posterior border.

Under Surface. On the *under surface* of the cerebellum, its division into two hemispheres is clearly perceptible. The deep furrow between them is called the *vallecula.* The front part of it is occupied by the medulla oblongata. To examine the surface of the valley, the medulla must be raised, and the hemispheres separated from each other. Along the middle line of the vallecula is the *inferior vermiform process*, which is the under surface of the original part of the cerebellum. Traced forward, this process terminates in the *nodule*, which projects into the fourth ventricle, and is called the *laminated tubercle* of Malacarne; traced backward, it ends in a small conical projection, called the *pyramid;* between these is a tongue-like body, called the *uvula*, which is connected with the adjacent amygdalæ by an indented gray ridge, named the *furrowed band.* Passing from the nodule to the flocculus is a thin, valve-like fold of white matter, which together take the name of the *posterior* or *inferior medullary velum.* Its anterior crescentic margin is free, and its posterior is attached to the furrowed band.[1] To see this satisfactorily, the tonsils must be carefully separated from each other.

[1] These are sometimes called the *valves of Tarini.*

Each hemisphere presents on its under surface certain secondary lobes, to which different names have been applied (Fig. 192). That portion which immediately overlies the side of the vallecula is called the *tonsil* (*amygdala*); this is connected with the uvula by an indented layer of gray matter, called the furrowed band. At the anterior part of each hemisphere, near the middle line, is a little lobe, named the *flocculus* or *subpeduncular lobe*.

In addition to the amygdalæ and flocculi, already mentioned, other lobes have been described on the under surface of the cerebellum. Thus, there is the *digastric lobe*, situated external to the

FIG. 192.



INFERIOR SURFACE OF THE CEREBELLUM.

A. The amygdala.
Bi. The biventral lobe.
G. The slender lobe.
Pi. The posterior inferior lobe.
h f. The great horizontal fissure.
fl. The flocculus.
n. The nodule
u. The uvula
p. The pyramid
} situated in the vallecula.

amygdala; and behind this are successively the *slender* and the *posterior inferior lobes* (Fig. 192).

Peduncles of the Cerebellum. The cerebellum is connected with the cerebro-spinal axis by three *peduncles* or *crura*—a superior, middle and inferior. With the medulla oblongata it is connected by means of the restiform tracts; these are called the *processus a cerebello ad medullam*, or its *inferior peduncles;* with the cerebrum it is connected by means of the *processus c cerebello ad cerebrum;* these are called its *superior peduncles*. The lateral portions of the pons constitute its *middle peduncles*.

Internal Structure. To examine the internal structure of the cerebellum, a longitudinal section must be made through the thickest part of one of its hemispheres. There is then seen in the centre a large nucleus of white substance, from which branches radiate into the gray substance in all directions, and upon which the gray cortical substance is deposited (*laminæ*).

The *laminæ*, about twelve in number, have branches from them at right angles, secondary laminæ; and, from these again, tertiary laminæ. This racemose arrangement of the white matter in the substance of the gray has been likened to the branches of a tree deprived of its leaves, and is generally known as the *arbor vitæ*.

Corpus Dentatum. In the centre of the white substance of each hemisphere is a nucleus of gray matter, the *corpus dentatum*, consisting of a zigzag line, of yellowish-gray color, incomplete at its upper and inner part, and enclosing within it some white substance. From its centre white fibres may be traced to the superior cerebellar peduncles and the valve of Vieussens. It is displayed either by a vertical or by a horizontal section.

Functions. Respecting the function of the cerebellum, the deductions derived from comparative anatomy and physiological experiments render it probable that it is the co-ordinator of muscular movements—*e. g.*, in walking, flying, and swimming.

The encephalon is originally developed from three primary vesicles, from which the following parts are, in the later stages, severally developed:—

1. Anterior vesicle	Prosencephalon	Cerebral hemispheres, corpora striata, corpus callosum, fornix, lateral ventricles, olfactory lobe.
	Thalamencephalon	Optic thalami, pineal body, pituitary body, third ventricle, optic nerve.
2. Middle vesicle	Mesencephalon	Corpora quadrigemina, crura cerebri, aqueduct of Sylvius, optic nerve.
3. Posterior vesicle	Epencephalon	Cerebellum, pons Varolii, front part of the fourth ventricle.
	Metencephalon	Medulla oblongata, posterior part of the fourth ventricle, auditory nerve.

he component parts of the encephalon begin to be developed lifferent periods of fœtal life, and the ages at which they rally begin to appear are given as follows :—

POSTERIOR VESICLE.

	Part.	*Month.*
ncephalon :	Medulla oblongata	Third
	Restiform bodies	Third to fourth
	Anterior pyramids	Fifth
	Olivary bodies	Sixth
	Striæ acusticæ	After birth
ıcephalon :	Cerebellum	Second, end of
	Inf. cerebellar ped.	Third
	Middle "	Fourth
	Corpus dentatum	Fourth
	Superior cereb. ped.	Fifth
	Valve of Vieussens	Fifth
	Lobes of cerebellum	Fifth
	Folia	Sixth
	Flocculus	Seventh
	Post. medullary velum	Seventh
	Amygdalæ	Eighth

MIDDLE VESICLE.

ncephalon :	Corpora quadrigemina	Fourth
	Fillet	Fourth
	Corpora quadrig.:	
	Vertical groove	Sixth
	Transverse	Seventh

ANTERIOR VESICLE.

lamencephalon :	Optic thalami	Second to third
	Anterior commissure	Third
	Posterior commissure	Third, end of
	Pineal body	Third to fourth
	Optic tracts	Third to fourth
	Peduncles of pineal body	Third to fourth
	Middle commissure	Ninth ?
encephalon :	Island of Reil	Earliest of all
	Corpora striata	Third
	Corpus callosum	Third, end of
	Fornix	Fourth to fifth

ANTERIOR VESICLE (*Continued*).

Prosencephalon:	Sulci, primitive:	
	Fissure of Sylvius	Middle of third
	Parieto-occipital	Third
	Dentate	Third
	Calcarine	Third
	Sulci, secondary:	
	Rolando	Fifth to sixth
	Parallel	Fifth
	Interparietal	Sixth, end of
	Calloso-marginal	Sixth, end of
	Collateral	Sixth, end of
	Frontal	Seventh
	Hippocampus major	Fourth to fifth
	Convolutions	Fourth to fifth
	Convolutions develop rapidly	Seventh to eighth
	Septum lucidum	Fifth

The cerebral hemispheres enlarge at first slowly, but later they develop much more rapidly; the extent to which they reach backward in the various periods of intra-uterine life is as follows:—

To the optic thalami at third month,
To the corpora quadrigemina at fourth month,
To the greater part of cerebellum at sixth month,
To the posterior border of cerebellum at seventh month.

DISSECTION OF THE SPINAL CORD.

To examine, in situ, the spinal cord covered with its membranes, the arches of the vertebræ must be sawn through, and removed. It is then noticed that the cord does not occupy the whole cavity of the spinal canal. The dura mater does not adhere to the vertebræ, and does not form their internal periosteum, as in the skull. Between the bones and this membrane, a space intervenes, which is filled with a soft, reddish-looking fat, with watery cellular tissue, and the ramifications of a plexus of veins.

Spinal System of Veins. The spine is remarkable for the number of large and tortuous veins which ramify about it, inside and outside the vertebral canal (Fig. 193). They are:—

1. The *dorsi-spinal* or *posterior external veins*, which form a tortuous plexus outside the spinous, transverse, and articular processes, and the arches of the vertebræ; they communicate with corresponding veins above and below, and they send off branches, which pass through the ligamenta subflava and intervertebral foramina, and end in the plexus inside the vertebral canal. They join the vertebral veins in the cervical region, the intercostal in the dorsal, and the lumbar and sacral veins below.

2. The *veins of the bodies of the vertebræ* (*venæ basis vertebrarum*), emerge from the backs of the bodies, and empty themselves into the transverse vein connecting the two anterior longitudinal spinal veins.

3. The *anterior longitudinal spinal veins*, two in number, one on each side, are very large tortuous veins, which extend along

FIG. 193.

1. Anterior external veins.
2. Dorsi-spinal veins.
3. Posterior longitudinal spinal veins.
4. Anterior longitudinal spinal veins.

DIAGRAM OF THE SPINAL VEINS. (VERTICAL SECTION.)

the whole length of the spinal canal. They communicate by transverse branches, passing beneath the posterior common ligament, opposite the body of each vertebra, where they receive the venæ basis vertebrarum. They are larger in the dorsal and lumbar regions, and communicate externally with the vertebral, the intercostal, the lumbar and the sacral veins.

4. The *posterior longitudinal spinal veins*, like the anterior, run along the whole length of the spinal canal. They form a tortuous venous plexus, situated inside the vertebral arches, and communicate in front with the anterior longitudinal veins by cross branches at frequent intervals, and externally with the vertebral, intercostal, lumbar and sacral veins by branches through the intervertebral foramina.

The anterior and posterior longitudinal spinal veins are situated between the spinal canal and the dura mater of the spinal cord, and are called the *meningo-rachidian veins.*

5. The *medulli-spinal* or *proper veins of the spinal cord* lie within the dura mater. They form a fine plexiform arrangement of veins over both surfaces of the cord, and can with difficulty be injected from the other spinal veins. This complicated system of veins discharges itself through the intervertebral foramina in the several regions of the spine, as follows: In the cervical, into the vertebral veins; in the dorsal, into the intercostal veins; in the lumbar, into the lumbar veins. None are provided with valves;

FIG. 194.

1. Anterior external veins.
2. Dorsi-spinal veins.
3. Posterior longitudinal spinal veins.

4. Anterior longitudinal spinal veins.
5. Internal veins of the body of the vertebra.
6. Lateral veins.

DIAGRAM OF THE SPINAL VEINS. (TRANSVERSE SECTION.)

hence they are liable to become congested in diseases of the spine.

The membranes of the spinal cord, though the same in number and continuous with those of the brain, differ from them in certain respects, and require separate notice.

Dura Mater. The *dura mater* of the cord is a tough, fibrous membrane, like that of the brain, but does not adhere to the bones, being separated from them by fat, loose areolar tissue, and the plexus of veins described above. Moreover, such adhesion would impede the free movements of the vertebræ upon each other. It is attached firmly above to the margin of the foramen magnum, and by slender tissue to the posterior common ligament, and may be traced downward, as a sheath, as

ar as the second bone of the sacrum, from which it is prolonged as a fibrous cord to the coccyx, where it becomes continuous with the periosteum. It forms a complete canal or bag (*theca*), which surrounds loosely the spinal cord, and is relatively larger in the cervical and lumbar regions than in the dorsal. On each side are two openings in the dura mater for the anterior and posterior roots of the spinal nerves, and the membrane is prolonged over the trunk of each spinal nerve. These prolongations accompany the nerves only as far as the intervertebral foramina, and are there blended with the periosteum. The inner surface of the dura mater is covered with a layer of polygonal cells, so that it is smooth and secerning; this was formerly described as the parietal layer of the arachnoid membrane.

Cut through the nerves which proceed from the spinal cord on each side, and remove the cord with the dura mater entire. Then lay it flat on the table and slit up the dura mater along the middle of the front of the cord, to examine the arachnoid membrane.

It will be seen that the functions of the dura mater of the cord are not identical with those of encephalon, since it does not form an internal periosteum to the bones of the spinal canal; nor does it send in partitions to support the cord; and it does not split to form venous sinuses.

Arachnoid Membrane. The *arachnoid membrane of the cord* is a continuation from that of the brain, and is reflected over the spinal nerves as they pass from the cord to the apertures in the dura mater. This membrane invests the cord, and is in contact by its superficial aspect with the dura mater, there being an interval between them called the *sub-dural space*, although in some situations they are more or less connected by connective-tissue bands. On its deeper surface it is in contact with the pia mater, but is loosely connected with it by delicate areolar tissue, so that there is a considerable interval between them (*subarachnoid space*), which is occupied by a transparent watery fluid (*cerebro-spinal fluid*) contained in the meshes of the subarachnoid tissue. The separation between the arachnoid and the pia mater varies in different parts, and is greatest in the lowest part of the cord.

Cerebro-Spinal Fluid. This *cerebro-spinal fluid* cannot be demonstrated unless the cord be examined very soon after death,

and before the removal of the brain.[1] The nerves proceeding from the cord are loosely surrounded by a sheath of the arachnoid; but this only accompanies them as far as the dura mater, where the two are continuous. The cerebro-spinal fluid of the cord communicates with that of the brain, and also with the general ventricular cavity through an aperture in the lower boundary of the fourth ventricle, called the *foramen of Magendie*.

Pia Mater.

The pia mater to the cord is the protecting membrane which immediately invests it. It is very different in structure from that of the brain, since it does not constitute a membrane in which the arteries break up, but serves rather to support and strengthen the cord; consequently, it is much less vascular, more fibrous in its structure, and more adherent to the substance of the cord. It sends down thin folds into the anterior and posterior median fissures of the cord, and is prolonged upon the spinal nerves, forming their investing membrane or "neurilemma."

Along the anterior median fissure may be traced a well-marked fibrous band, formed by the pia mater, which has been named the *linea splendens*.

Below the level of the second lumbar vertebra, the pia mater is continued as a slender filament, called the *filum terminale*, or *central ligament*, which runs down in the middle of the bundle of nerves into which the spinal cord breaks up. About the level of

[1] The existence and situation of the cerebro-spinal fluid were first discovered by Haller *Element. Phys.* vol. iv, p. 87, and subsequently more minutely investigated by Magendie *Recherches Phys. et Cliniques sur le Liquide Cephalo rachidien*, in-4°, avec atlas: Paris, 1842. This physiologist has shown that if, during life, the arches of the vertebræ are removed in a horse, dog, or other animal, and the dura mater of the cord punctured, there issue jets of a fluid which had previously made the sheath tense. The fluid communicates, through the fourth ventricle, with that in the general ventricular cavity. The collective amount of the fluid varies from 1 to 2 oz. or more. It can be made to flow from the brain into the cord, or *vice versâ*. This is proved by experiments on animals, and by that pathological condition of the spine in children termed spina bifida. In the later instance, coughing and crying make the tumor swell; showing that the fluid is forced into it from the ventricles. Again, if pressure be made on the tumor with one hand and the fontanelles of the child examined with the other, in proportion as the spinal swelling decreases so is the brain felt to swell up, accompanied by symptoms resulting from pressure on the nervous axis generally. See some remarks very much to the point, by Sir George Burrows, *On Diseases of the Cerebral Circulation* p. 50, 1846.

the third sacral vertebra it becomes continuous with the dura mater of the cord, and is then prolonged as far as the base of the coccyx. The spine of the third sacral vertebra marks the level to which the cerebro-spinal fluid descends in the vertebral canal. It is supplied with nerves from the sympathetic and from the posterior roots of the spinal nerves.

Ligamentum Denticulatum.

From each side of the cord along its whole length there runs a fibrous band, *ligamentum denticulatum*, which gives off a series of processes to steady and support the cord. They are triangular, their bases being attached to the cord, and their points to the inside of the dura mater (Fig. 195). There are from eighteen to twenty-two of them on each side, and they lie between the anterior and posterior roots of the spinal nerves. The first process passes between the vertebral artery and the hypoglossal nerve; the last is found at the termination of the cord. It is composed of fibrous tissue, and is covered with nucleated cells continuous with the arachnoid membrane.[1]

FIG. 195.



DIAGRAM OF THE LIGAMENTUM DENTICULATUM.

1. Dura mater.
2, 2, 2. Ligamentum denticulatum.

SPINAL CORD.

The *spinal cord* (medulla spinalis) is that part of the cerebro-spinal axis contained in the vertebral canal, and is enclosed within a sheath of dura mater (*theca vertebralis*), which is separated from the canal by a plexus of veins and connective tissue. It is the continuation of the medulla oblongata, and extends from the foramen magnum down to the lower border of the first lumbar vertebra, where it terminates in a conical point, *conus medullaris*, after having given off a large bundle of nerves, termed the *cauda equina*, for the supply of the lower limbs.

It is from fifteen to eighteen inches in length, and is about an ounce and a half in weight. Its lower extremity from the conus medullaris is continued downward as a thin, silvery cord, the *filum terminale*, which descends along the posterior aspect of the cauda

[1] *Vide* Axel Key and Retzius; Max Schultze's *Archives*, 1873.

equina. It passes down within the sheath as far as the second sacral vertebra, and then, piercing the dura mater, becomes attached to the periosteum of the canal at the back of the coccyx.[1] In its upper part, the filum terminale contains some gray nerve substance. It is a prolongation of the pia mater of the cord, and in many subjects there is a continuation of the central canal of the cord in its upper half.

The cord is not of uniform dimensions throughout. It presents a considerable enlargement in the lower part of the cervical region; another in the lower part of the dorsal, from which proceed the large nerves to the upper and lower limbs, respectively. The upper or cervical enlargement, which is the larger, extends from the third cervical to the first dorsal vertebra, and is largest at the sixth cervical vertebra; the lower, or lumbar, extends from the twelfth dorsal vertebra, and is largest opposite the last dorsal vertebra.[2]

The cord is divided into two symmetrical halves by a median longitudinal fissure in front and behind (Fig. 196). The *anterior fissure* is the more distinct and wider, and penetrates about one-third of the substance of the cord; deeper in the lower than in the upper part of the cord. It contains a fold of pia mater, with many blood vessels for the supply of its interior. At the bottom of this fissure is a transverse layer of white substance, named the anterior white commissure, connecting the two anterior halves of the cord. The *posterior fissure* is much less apparent than the anterior, and is better marked in the upper and the lower parts of the cord. It does not contain a fold of pia mater, but contains a thin septum of connective tissue with blood vessels. It can be traced to a greater depth than the anterior, and reaches down as far as the *posterior gray commissure* of the cord.

Besides the anterior and posterior fissures, is another superficial lateral groove, from which the posterior roots of the spinal

[1] The explanation of this is, that, at an early period of fœtal life the length of the cord corresponds with that of the vertebral canal; but after the third month the lumbar and sacral vertebra grow away from the cord, in accordance with the more active development of the lower limbs. See Tiedemann, *Anatomie und Bildungsgeschichte des Gehirns im Fœtus des Menschen, etc.* Nüremberg, 1816.

[2] In very early fœtal life these enlargements do not exist, and only make their appearance with the development of the extremities.

nerves emerge; this is termed the *postero-lateral groove* (Fig. 196). This leads down to the posterior horn of the gray matter in the interior of the cord. There is sometimes described an *antero-lateral groove*, corresponding to the line whence the anterior roots of the spinal nerves emerge; but this is not really a groove, although it serves to map out each half of the cord into three longitudinal portions, a posterior, a lateral and an anterior column. On each side of the posterior median fissure, in the cervical region, is a slender column, called the *posterior median column*, which is separated from the posterior column by a shallow furrow.[1] The anterior column is continuous with the anterior pyramid; the lateral column with the lateral tract of the

FIG. 196.



DIAGRAM OF A TRANSVERSE SECTION THROUGH THE SPINAL CORD AND ITS MEMBRANES.

1. Dura mater.
2. Arachnoid membrane.
3. Ganglion on posterior root of spinal nerve.
4. Anterior root of spinal nerve,
5, 5. Seat of sub-arachnoid fluid.
4. Posterior branch of spinal nerve.
7. Anterior branch of spinal nerve.

medulla; the posterior column with the restiform body; and the posterior median column with the posterior pyramid.

Internal Structure. A transverse section through the cord (Fig. 196) shows that, externally, it is composed of white nerve substance, and that its interior contains gray matter, arranged in the form of two crescents, with their backs to each other. Each crescent is placed in its corresponding half of the cord, and is connected with its fellow across the centre by a portion called the *posterior* or *gray commissure*. The *posterior horns* are long and narrow, and extend to the postero-lateral groove, where they are connected with the posterior roots of the spinal nerves. At their extremities they taper to a point, the *apex cornu*

[1] The posterior median column is said by Foville to be present along the whole length of the cord.

posterioris, and near their bases they present a constriction, the *cervix cornu*, beyond which they slightly enlarge to form the *caput cornu*. The outline of the gray matter of the posterior horns at their commencement is indefinite and frayed out, which is especially noticeable in the cervical region, and is called the *processus reticularis;* at their apices the gray matter is semi-transparent in appearance, and hence is known as the *substantia gelatinosa*. In the centre of the concavity of the posterior horn is a rounded projection, most marked in the upper dorsal region, termed the *tractus intermedio-lateralis*, the continuation of which has been traced upward through the medulla oblongata. The *anterior horns* are short and thick, and come forward toward the attachment of the anterior roots of the nerves, but do not reach the surface. Separating the gray commissure from the anterior median fissure is the *anterior* or *white commissure*.

On making transverse sections through different regions of the spinal cord, the gray substance is seen to vary in shape and in amount; in the cervical region the anterior cornua are thick and short, the posterior are long and slender; in the dorsal, the anterior and posterior cornua are both thin; in the lumbar, the anterior and posterior cornua are large and broad; in the lower part of the cord the gray matter is arranged in a central mass.

Running along the centre of the cord in its whole length is a minute canal, the *central canal*, just visible to the naked eye. Below, in the conus medullaris, it ends in a dilated cul-de-sac, of the shape of the letter T; above, it opens out at the calamus scriptorius into the fourth ventricle. It is lined with cylindrical ciliated epithelium. This central canal is interesting, as it is the remains of the cavity formed by the spinal cord at the earliest period of its development.[1]

Spinal Nerves. Thirty-one pairs of nerves arise from the spinal cord, namely, eight in the cervical region, twelve in the dorsal, five in the lumbar, five in the sacral, and one in the coccygeal. Each nerve is formed by the junction of two series of roots, one from the front, which is the *motor root*, the other from the back of the cord, which is the *sensory* and *larger root*. The two roots pierce the dura mater separately and then con-

[1] The central canal is well seen in fishes, birds, and reptiles.

verge to the corresponding intervertebral foramen to form a single nerve, composed of motor and sensory filaments.

The filaments composing the *posterior roots* are finer, but their fasciculi are thicker and more numerous than the anterior.[1] They proceed from the postero-lateral fissure, and previous to their union with the anterior roots are collected into two bundles which pass through a ganglion. The *ganglion* is of an oval form, bilobate on its external extremity, and lies in the intervertebral foramen; each fasciculus of the posterior root enters the corresponding lobe of the ganglion. The ganglia of the first and second cervical nerves are placed upon the arches of the atlas and axis respectively; the ganglia of the sacral and coccygeal nerves are situated within the spinal canal. The *anterior roots* arise from the antero-lateral column, are smaller than the posterior roots, but like them divide into two fasciculi as they approach the ganglion on the posterior root.[2]

The compound nerve formed by the junction of the two roots, external to the ganglion of the posterior, divides, outside the intervertebral foramen, into an anterior and a posterior (dorsal) branch. (See diagram, p. 799.)

The fibres of the anterior roots, after entering the antero-lateral column, pass through the white fibres to enter the gray matter; here they pass in all directions, the larger number passing upward and downward, and some decussating with corresponding filaments of the opposite side, through the anterior white commissure; some also pass upward vertically through the lateral column. The fibres of the posterior roots enter the caput cornu posterioris through the postero-lateral groove, and then pass upward and downward in the gray matter, chiefly of the posterior

[1] This does not apply to the first cervical nerve, in which the anterior root exceeds the posterior in size.

[2] The researches of Blandin, *Anat. descript.*, t. ii, p. 648, 1838, have led him to establish the following relation between the respective size of the anterior and posterior roots of the nerves in the several regions of the spine :—

The posterior roots are to the anterior in the			cervical region	::	2	:	1
"	"	"	dorsal "	::	1	:	1
"	"	"	lumbar and sacral	::	1½	:	1

This relation quite accords with the greater delicacy of the sense of touch in the upper extremity.

horn, but some curve round to enter the anterior horn, while others pass to the opposite side through the posterior gray commissure.

Variation in the Length of the Roots.

The direction and length of the roots of the nerves vary in the different regions of the spine, owing to the respective parts of the cord from which they arise not being opposite to the foramina through which the nerves leave the spinal canal. In the upper part of the cervical region, the origins of the nerves and their point of exit are nearly on the same level; therefore the roots proceed transversely, and are very short. Lower down, however, the obliquity and length of the roots gradually increase, so that the roots of the lower dorsal nerves are at least a vertebra higher than the foramina through which they emerge. Again, since the cord itself terminates at the lower border of the first lumbar vertebra, the lumbar and sacral nerves must descend from it almost perpendicularly through the lower part of the spinal canal. To this bundle of nerves the old anatomists have given the name of *cauda equina*, from its resemblance to a horse's tail.

Cauda Equina.

To sum up briefly, it appears that the spinal cord consists of two symmetrical halves, separated in front and behind by a deep median fissure; that the two halves are connected at the bottom of the anterior fissure by an anterior or white commissure, at the bottom of the posterior fissure by the posterior or gray commissure; that each half of the cord is divided into three tracts or columns of longitudinal white nerve fibres—an anterior, a lateral and a posterior—the boundaries between them being the respective lines of origin of the roots of the spinal nerves; that the interior of the cord contains gray matter disposed in the form of two crescents, placed with their convexities toward each other, and connected by a transverse bar of gray matter, which constitutes the posterior commissure.

Blood Vessels of the Cord.

The cord is supplied with blood by: 1. The *anterior spinal* artery, which commences at the medulla oblongata by a branch from the vertebral of each side, and then runs down the middle of the front of the cord. Other branches are derived from the vertebral, ascending cervical, inter-

costal and lumbar arteries, which pass through the intervertebral foramina, and assist in keeping up the size of this anterior artery. 2. The *posterior spinal* arteries, which proceed also from the vertebral, intercostal and lumbar arteries, and ramify somewhat irregularly on the back of the cord.

On the posterior part of the bodies of the vertebræ, the spinal arteries of opposite sides communicate by numerous transverse branches along the entire length of the spine, thus resembling the arrangement of its venous plexuses.

Functions of the Spinal Cord. The spinal cord performs at least three functions: 1. It is the general conductor of impressions to and from the brain. 2. It transfers impressions. 3. It is a centre of reflex action. *Sensory* impressions are conducted by the posterior roots of the spinal nerves to the cord, and are thence transmitted to the brain through the posterior columns and the gray matter of the cord. These impressions do not run up on the same side, for the fibres, immediately on entering the gray matter, cross over to the opposite side to reach the brain; so that if the posterior column of the *right* side be divided, the *left* leg, and not the right, would be deprived of sensation. *Motor* impulses are conveyed along the antero-lateral columns and the gray matter in them, and carry the commands of the will from the brain to the muscles. The crossing of the motor fibres takes place in the medulla oblongata, at the decussation of the anterior pyramids, so that they run in the corresponding half of the cord as far as their point of decussation. Division, therefore, of one-half of the cord below this point causes paralysis of motion on the same side of the body. The cord is, moreover, concerned in the conduction of impressions to and from the *vaso-motor centre* of the medulla oblongata, which determines the varying conditions of the blood vessels. The cord also transfers impressions; this is more manifest in disease than in health; a well-marked example of transference is, that pain is felt at the knee in cases of disease of the hip joint. The spinal cord has probably no power of originating impressions—in other words, it is *not automatic*.

DISSECTION OF THE EYE.

Since the eye in the human subject cannot be obtained sufficiently fresh for anatomical purposes, the student should examine the eye of the sheep, bullock, or pig. The conjunctiva should be removed, together with the loose connective tissue which unites it to the sclerotic coat.

Conjunctiva.

The *conjunctiva* is the mucous membrane which covers the ocular surface of the eyelids and the anterior part of the globe. It presents different characters in the various situations over which it is reflected, and is described as the palpebral, the sclerotic and the corneal portions.

The *palpebral portion* is thicker than the other portions, is very vascular, and is provided with fine papillæ abundantly supplied with nerves.[1] As described p. 47, it is continuous with the mucous membrane of the lachrymal sac through the canaliculi, and lines the Meibomian glands and the ducts of the lachrymal gland. The columnar epithelium which lines the palpebral conjunctiva becomes at the margins of the eyelids more stratified and continuous with the flattened cells of the skin of the eyelid. It forms at the inner canthus a reduplicated fold, the *plica semilunaris*, and is reflected from the eyelids to the globe, its angle of reflection being called the *fornix conjunctivæ*, and the folds into which this is thrown are termed the *superior* and *inferior palpebral folds*.

The *sclerotic conjunctiva* is loosely attached by submucous tissue to the sclerotic coat, so as not to impede the movements of the globe. It is thinner, and has no papillæ. It is transparent and nearly colorless, except when inflamed; it then becomes intensely vascular, and of a bright scarlet color. The arteries are derived from the lachrymal and palpebral branches of the ophthalmic artery, and at the circumference of the cornea they form capillary loops which anastomose with each other. The lymphatics are well marked in the palpebral and sclerotic portions of the conjunctiva, and at the margin of the cornea they rapidly diminish in size and become connected with the cell spaces in the cornea. An abundant supply of nerves is distributed to the membrane;

[1] These papillæ were first described by Eble, *Ueber den Bau und die Krankheiten der Bindehaut des Auges.*

their arrangement is in the form of plexuses as far as the margin of the cornea, where they terminate in "end-bulbs," described by Krause, resembling in many respects the central portion of Pacinian corpuscles.

The *corneal conjunctiva* is composed chiefly of epithelium arranged in layers, the deepest of which consists of columnar cells resting by their bases on the substantia propria of the cornea; superficial to these are two or three layers of polygonal cells, the deepest of which, called the *fingered cells of Cleland*, interdigitate with the columnar cells; and on the surface there is a layer of flattened squamous epithelial cells. This portion of the conjunctiva cannot be separated by dissection in recent eyes, but it possesses the same acute sensibility as the rest of the conjunctiva. Changes produced by inflammation of the conjunctiva often involve the cornea and render its texture thick and opaque.[1]

The eyeball is imbedded in a large quantity of fat and delicate connective tissue; and surrounding it in its posterior three-fourths is a serous membrane, the *capsule of Tenon*, which allows of its free movement in the orbit.

The axes of the two eyeballs are nearly parallel with each other; thus they do not correspond with the axes of the orbits, nor of the optic nerves, which enter the globes on their nasal aspect.

The human eye is nearly spherical, and consists of segments of two spheres: a large posterior one, which corresponds with the sclerotic, and a small anterior one, with the cornea. The antero-posterior and vertical diameters of the globe are equal, the transverse exceeding these by less than half a line. The convexity of the cornea varies in different persons and at different periods of life; this is one cause of the varying degrees of near and far sight.

Coats and Humors of the Eye. The globe is composed of three concentric coats, arranged one within the other, which enclose cer-

[1] The facts of comparative anatomy confirm this view. In the serpent tribe, which annually shed the skin, the front of the cornea comes off with the rest of the external surface of the body. In the eel the surface of the cornea is often drawn off in the process of skinning. In some species of rodents which burrow under the ground, like the mole, the eye is covered with hair, like other parts.

tain transparent structures for the transmission of light. The external coat, consisting of the *sclerotic* and *cornea*, is fibrous, thick, and strong. The second coat, consisting of the *choroid*, the *iris*, and the *ciliary processes*, is composed of blood vessels, muscular tissue and pigment cells, and is very dark in color. The third coat, called the *retina*, consists of the expansion of the optic nerve for the reception of the impression of the waves of light. The bulk of the interior is filled with a transparent humor, called the *vitreous body*. Embedded in the front of this, and just behind the pupil, is the crystalline lens, for the purpose of concentrating the rays of light. In front of the lens is placed a movable curtain, called the *iris*, to regulate the amount of light which shall be admitted through a central aperture, the *pupil*, to the fundus of the eye. The space in which the iris is suspended is filled with a fluid, termed the *aqueous humor*.

Sclerotic Coat. The *sclerotic coat* is the white, tough, protecting coat of the eye which serves to maintain the form of the globe.[1] It covers the posterior five-sixths of the globe, the remaining anterior one-sixth being completed by the cornea. It is of dense white color, except in front, where the tendons of the recti and obliqui are inserted into it. The thickest part of the sclerotic coat is at the back of the globe (Fig. 198); the thinnest is a short distance behind the cornea.[2] The back of the sclerotic is perforated by the optic nerve, which enters it about one-tenth of an inch on the inner or nasal side of the axis of vision. The sheath of the optic nerve becomes continuous with the sclerotic, where it perforates this coat. The optic nerve at its entrance into the sclerotic is much constricted, and instead of passing through a single aperture in this coat, it enters it through a number of minute apertures, so that this membrane forms a porous lamina, called the *lamina cribrosa*. In the centre of the

[1] The sclerotic coat of the eye in fishes is of extraordinary thickness and density; and in birds this coat is further strengthened by a circle of bony plates, fourteen or fifteen in number, arranged in a series round the margin of the cornea. Similar plates are found in some of the reptiles, and particularly in the fossil ichthyosauri and plesiosauri.

[2] The greatest thickness posteriorly is about the $\frac{1}{20}$ of an inch; its thinnest in front is about the $\frac{1}{40}$ of an inch.

lamina cribrosa is an opening (*porus opticus*), larger than the rest, which transmits the arteria centralis retinæ. Around the optic nerve the sclerotic is pierced by the ciliary arteries, veins and nerves, for the supply of the choroid and iris. About a quarter of an inch from the cornea the sclerotic receives the insertions of the recti muscles; here also it transmits the anterior ciliary arteries, which run forward along the tendons of these muscles, and form a vascular ring around the circumference of the cornea (Fig. 197).

FIG. 197.

INSERTION OF THE RECTI MUSCLES WITH ANTERIOR CILIARY ARTERIES.

Structure. The sclerotic is composed of connective tissue arranged in bundles, which run, some longitudinally, some transversely, and are intermingled with fine elastic fibres. The longitudinal fibres are the most external and abundant. Under the microscope numerous connective-tissue corpuscles may be seen filling cell spaces, similar to those in the cornea but not so abundant, and containing pigment granules. The inner surface of the sclerotic is of a dark brownish color, due to the presence of a thin layer of connective tissue, *lamina fusca*, in which are found pigment cells. This surface of the sclerotic is grooved for the passage of the ciliary nerves, which run forward in a sort of lymph space; and it is, moreover, connected by filamentous tissue with the subjacent choroid coat.

Cornea. The *cornea* is the brilliant translucent coat which forms about the anterior one-sixth of the globe. It is nearly circular in shape, its diameter being nearly half an inch and its thickness about $\frac{1}{15}$ of an inch. The curve of the cornea, forms part of a smaller circle than that of the sclerotic, so that it projects further forward, varying in this respect in different eyes, and at different ages of life. It is firmly connected at its margin to the sclerotic, with the fibres of which it is continuous. The margin of the sclerotic is beveled on the inside; that of the cornea on the outside, so that the former overlaps the latter (Fig. 198).

To examine the cornea, it should be removed with the sclerotic coat. This should be done under water, by making a circular cut with scissors, about a quarter of an inch from the margin of the cornea. With a little care it will be easy to remove the outer coat of the eye without injuring the dark choroid coat, the ciliary muscle, or the iris. In the loose, brown-colored, connective tissue between the sclerotic and the choroid, are the ciliary nerves passing forward to the iris; their white color makes them very conspicuous on the dark ground.

Structure.

The cornea consists of four layers, which are not all composed of the same kind of tissue; they

FIG. 198.



DIAGRAM OF A VERTICAL SECTION OF THE EYE.

1. Anterior chamber filled with aqueous humor.
2. Posterior chamber. 3. Canal of Petit.

are, from without inward, the conjunctiva, the substantia propria or cornea proper, the posterior elastic lamina, and the epithelial lining.

The *conjunctiva* is the most superficial layer, and consists of several strata of epithelial cells; the deeper ones are columnar and placed vertically; the next consist of several layers of polygonal cells, and the most superficial ones are flattened, scaly epithelium cells, with well-marked nuclei.

The *cornea proper* or *substantia propria* consists of translucent

connective tissue, upon which the thickness and strength of the cornea mainly depend. The fibres are arranged in laminæ, about sixty in number. Those composing a lamina are arranged in a parallel direction, but the fibres of each layer cross at right angles those of each succeeding layer. The lamellæ are connected together by filaments passing from one to another, so that they are not perfectly separate from each other. The cornea proper in the recent subject presents no trace of structure, but it is only after death, by means of lenses and reagents, that these lamellæ can be satisfactorily demonstrated. Between the lamellæ are irregularly branched spaces, called the *cell spaces of the cornea*, in which are lodged the *corneal corpuscles*, having outstanding processes, which communicate freely with each other in their own plane, and also with those of the planes on either side. These corpuscles correspond in shape to the spaces within which they lie.[1] In inflammation of the cornea they undergo considerable changes. Immediately below the conjunctiva, the cornea proper presents a different appearance to that of the main thickness of this layer, so that this has been described by some anatomists under the name of the *anterior elastic lamina* of Bowman. It presents, however, a definite fibrillar structure, similar to that of the corner proper, but is destitute of the corneal corpuscles and cells. The greatest thickness of the cornea proper is about $\frac{1}{25}$ of an inch, and that of the anterior elastic lamina about $\frac{1}{1500}$ of an inch.

The *posterior elastic lamina*, called also the *membrane of Descemet* or *Demours*, is translucent, elastic and brittle, and may be easily separated from the preceding lamina. It consists of a perfectly structureless lamina, which, when peeled off, has a remarkable tendency to curl, with the attached surface innermost. It is unaffected by boiling, or by the action of acids or alkalies, and is from $\frac{1}{2000}$ to $\frac{1}{3000}$ of an inch in thickness. At the junction of this lamina with the sclerotic, on its inner surface, it spreads out

[1] If fluid be injected very gently into the cornea proper, there may be demonstrated a system of canals, called *Recklinghausen's canals*, which are the communications between the corneal corpuscles; but if the fluid be injected more forcibly, it passes in the course of the fibres composing the various laminæ of the cornea, which gives the appearance of a number of varicose and enlarged tubes crossing each other at right angles; these are termed *Bowman's corneal tubes*.

into a number of radiating tooth-like processes, the *ligamentum pectinatum iridis*, which are attached to the front of the circumference of the iris and to the sclerotic and choroid coats. The processes alone are covered with epithelial cells, and the intervals between the processes form small spaces, the *spaces of Fontana*, which communicate freely with the fluid of the aqueous chamber. In the sclerotic coat, close to its junction with the cornea, is situated a small oval canal, lined with epithelium, termed the *sinus circularis iridis* or *canal of Schlemm* (Fig. 198). Although, by some, it is considered a lymph-space, it is probably a venous sinus, for it can be injected from the arteries; but it has probably some free communication with the fluid of the anterior chamber, as this fluid passes readily from the chamber into this sinus.[1]

The *epithelial lining* consists of a single layer of polygonal nucleated cells, and lines the inner surface of the posterior elastic lamina. They resemble those which line serous membranes generally.

Arteries and nerves.—In the healthy condition the cornea contains no blood vessels, except at its circumference, where they form loops. Nor have any *lymphatics* been demonstrated in it. Its *nerves*, which are numerous—forty to forty-five in number—are derived from the ciliary nerves, and may be traced forward to the fibrous portion of the cornea, where they lose their dark outline and become transparent, forming a fine plexus—the *primary plexus*. This gives off minute filaments which ramify beneath the epithelium, constituting the *secondary* or *sub-epithelial plexus*. From this very minute varicose fibres run between the epithelial cells, forming the *intra-epithelial plexus*. Besides these plexuses, filaments are given off from the primary plexus to supply the cornea proper, and the filaments are said by some to be continuous with the anastomosing processes of the cell spaces.

Choroid Coat.

After the removal of the sclerotic coat and the cornea, which constitute the first tunic, we expose the second tunic, consisting of the choroid, the iris and ciliary processes in front, and of the ciliary ligament and the ciliary muscle.

[1] For further information on this point consult Leber, *Archiv. f. Opth.*, 1878; Heisrath, *Archiv. f. Opth.*, xxvi; and Schwalbe, Graefe and Saemisch's *Handbook*, 1874.

The *choroid* is the soft and flocculent tunic of the eye, recognized by its dark-brown color and great vascularity. It covers the posterior five-sixths of the globe, and is thickest posteriorly, where there is a circular aperture in it for the passage of the optic nerve. In front, the choroid passes beneath the ciliary muscle and ligament with which it is connected, and then extends forward, terminating in a series of plaited folds, called the *ciliary processes*. It is connected with the sclerotic by delicate connective tissue, the *lamina fusca*, through which the ciliary vessels and nerves pass forward (Fig. 199) to the iris. Its inner surface is smooth, and is in contact with the retina, and when detached from it presents a layer of hexagonal pigmented cells, which are now recognized as forming a part of the retinal coat and as part of which it is originally developed.

Structure. Under the microscope the choroid is seen to consist of two layers, an external and an internal. The choroid is covered externally by a connective-tissue layer, similar to the lamina fusca of the sclerotic, and known as the *lamina supra-choroidea;* it consists of connective tissue intermingled with elastic tissue, and imbedded in the meshwork are pigment cells and lymphoid cells. The contiguous surfaces of the lamina fusca and the lamina supra-choroidea are lined with squamous epithelium, having between them a more or less complete lymph space, which is continuous with that of Tenon's capsule through the apertures in the sclerotic, through which the ciliary vessels and nerves pass.

The *external layer* consists of the larger branches of the blood vessels; the arteries (short ciliary) running forward between the veins, previous to dipping down to form the internal layer. The veins are arranged with great regularity in drooping branches (*venæ vorticosæ*) like the weeping willow (Fig 199), and converge to four or five nearly equidistant trunks, which, after running backward for a short distance, perforate the slerotic not far from the entrance of the optic nerve, and empty themselves into the ophthalmic vein. Between the veins there are interspersed numerous stellate pigment cells which anastomose with neighboring cells; on the inner side of this layer the cells are absent.

The *internal layer* is formed by the capillaries of the ciliary

arteries, and is called the *tunica Ruyschiana*, after the Dutch anatomist, Ruysch. The capillaries branch off from the choroid vessels in a radiating manner, and form the most delicate vascular network found in any tissue. It extends forward as far as the retina, where the intervals become larger and the vessels freely communicate with those of the ciliary processes. This tissue has on its inner surface a transparent membrane, the *membrane of Bruch*, which rests on the pigmentary layer of the retina.

Ciliary Processes. The *ciliary processes* are the plaited folds formed by the anterior part of the choroid, and may be best seen when the globe has been divided by a trans-

FIG. 199.



SCLEROTIC COAT REMOVED TO SHOW THE CHOROID, CILIARY MUSCLE AND NERVES.

verse vertical section into an anterior and a posterior half, the vitreous humor being left undisturbed. They are black, and consist of from sixty to seventy radiating folds arranged in a circle about three lines broad. These processes consist of longer and shorter folds, the former being the more numerous, and in the proportion of three to one of the latter. The longer fold is about $\frac{1}{10}$ of an inch in length; the smaller about $\frac{1}{20}$ of an inch. One of the longer processes is seen in the diagram. The processes fit into corresponding folds of the suspensory ligament of the lens, and their free ends project for a short distance into the posterior chamber. The circumference of the processes is

attached to the ciliary ligament; their inner ends are free and rest upon the circumference of the lens.

Structure. The vascular supply of the ciliary processes is most abundant and resembles in the main that of the choroid, except that the plexus is coarser, with its meshes arranged longitudinally. The *arteries* come chiefly from the anterior ciliary, and from the front vessels of the choroid; and after breaking up into a fine plexus, they form loops which arch backward, to end in the smaller veins. Their dark color is due to several layers of pigmented cells, which disappear, however, at the free end of the processes.

Ciliary Muscle. The *ciliary muscle* consists of unstriped muscular fibres, and forms a muscular zone at the front of the choroid, close to the junction of the sclerotic with the cornea. It arises by a thin tendon from the sclerotic close to the cornea, and near the spaces of Fontana. Thence some of its fibres radiate backward, forming the *meridional* or *radiating fibres*, and are lost in the choroid behind the ciliary processes; some of the fibres form a circular muscle around the outer circumference of the iris, the *circular ciliary muscle*, which was formerly described as the ciliary ligament. Its action is to accommodate the eye to objects at various distances by compressing the lens and increasing the convexity of its anterior surface.[1]

Iris. The *iris* is the contractile and colored curtain suspended in the clear fluid which fills the space between the cornea and the lens. The iris divides this space into two unequal parts, called the *anterior* and *posterior chambers* (Fig. 198); these communicate with each other through a circular aperture in the centre of the iris, called the *pupil*, which is situated a little to the inner side of the iris.[2]

The circumference of the iris is nearly circular, and is immovably connected with the choroid, the ciliary muscle, and through the ligamentum pectinatum with the cornea. The diameter of

[1] Sir P. Crampton has noticed that this muscle is well developed in birds. In them its muscles are of the striped variety, as are the circular fibres of the iris.

[2] The size and shape of the pupil vary in different animals. In the bullock, sheep, horse, etc., it is oblong; in carnivorous quadrupeds it is often a mere vertical slit during the day, but dilates into a large circle at night.

the iris is about half an inch, and that of the pupil in man varies from the $\frac{1}{10}$ to the $\frac{1}{3}$ of an inch.

The color of the iris varies in different subjects, and gives the peculiar tint and brilliancy to the eye. The coloring matter or pigment is contained in minute cells, *pigment cells*, lining the anterior and posterior surfaces, of the iris, the posterior taking the name of *uvea*, from its grape-like color. Pigmented cells are also found in the substance of the iris.

The use of the iris is to regulate the amount of light which shall be admitted into the eye; for this purpose its inner circumference is capable of dilating and contracting according to circumstances, while its outer circumference is immovably attached.

Structure. When the iris is laid under water, and viewed with a low magnifying power, it is seen to be composed of fine fibres converging from all sides toward the pupil; many of them unite and form arches, leaving elongated interspaces, which are most marked toward the middle of the iris.

In front of the iris is a thin layer of polyhedral cells, which is continuous with that covering the membrane of Descemet, but the cells are smaller and more granular.

The *stroma* consists of connective tissue and cells. The fibres of the connective tissue are arranged longitudinally and circularly; the longitudinal fibres radiate from the circumference toward the pupil, and between them are contained the blood vessels and nerves; the circular fibres are found at the circumference of the iris. Intermingled in the meshes of this connective tissue and throughout its whole thickness are numerous and various-shaped pigment cells having anastomosing processes, like those of the choroid. They are chiefly found in the uvea, which is continuous with the pigmented layer of the retina. Upon the disposition of these pigment cells depend the color of the iris; in dark eyes, the pigment cells are scattered throughout the thickness of the stroma; in light eyes, only on its posterior surface.

The *muscular tissue* is of the unstriped kind, and is arranged partly in a radiating, partly in a circular manner. The circular fibres, the *sphincter*, well marked, are collected on the posterior aspect of the pupillary margin, where they form a ring about $\frac{1}{10}$

of an inch in width; at the free margin of the iris they form a thick bundle, but become more frayed out toward the circumference.[1] The radiating fibres, the *dilatator*, converge toward the pupil, where they form arches and blend with the circular fibres.

The *pigment*, as before described, is found in varying thickness and position, differing according to the color of the iris.

The *arteries* of the iris are derived from the two long and the anterior ciliary arteries. The *long ciliary arteries* perforate the sclerotic coat on each side of the optic nerve, and then run forward between this tunic and the choroid to the ciliary muscle at the outer circumference of the iris. Each artery divides into an upper and a lower branch, which form with each other and the anterior ciliary arteries a vascular circle (*circulus major*); from this circle numerous small branches pass inward and form another circle (*circulus minor*) of anastomosis, which terminates in the veins of the iris. The *anterior ciliary arteries*, five or six in number, are derived from the muscular and lachrymal branches of the ophthalmic artery, and ramify on the tendons of the recti muscles (p. 807), where they pierce the sclerotic about the $\frac{1}{12}$ of an inch behind the margin of the cornea. These vessels supply the ciliary processes and iris, and join the circulus major; it is from their enlargement that the red zone round the cornea is produced in inflammation of the iris.

The *veins* follow the same arrangement as the arteries, and communicate, as stated (p. 810), with the canal of Schlemm.

The *nerves of the iris* come from the nasal branch of the ophthalmic nerve, and by twelve to fifteen branches from the lenticular ganglion. They pierce the sclerotic around the entrance of the optic nerve, and run forward between the sclerotic and the choroid as far as the ciliary muscle. On the choroid they form a gangliated plexus which lies in connection with and among the blood vessels. In the ciliary muscle the nerves form another plexus, from which numerous non-medullated fibrils are given off to terminate in the muscular tissue of the iris. The sphincter iridis is supplied through the motor root of the lenticular gan-

[1] The circular fibres of the iris in the bird are of the striped variety, and discernible without difficulty.

glion, which is derived from the third nerve; the dilatator iridis is supplied by the sympathetic system.

Membrana Pupillaris. Until the seventh or eighth month of fœtal life the pupil is closed by a transparent vascular membrane, the *membrana pupillaris*, so that the anterior and posterior chambers are divided from each other by this membrane. Its vessels, derived from those of the iris and capsule of the lens, are arranged in loops which converge toward the centre of the membrane without joining each other. About the eighth month this membrane becomes gradually absorbed, so that at birth it is completely lost.

Retina. To obtain a view of the *retina*, the choroid coat must be carefully removed while the eye is under water; this should be done with the forceps and scissors on a fresh eye. When the choroid is thus removed, there will be seen on its inner surface a layer of pigmented cells, which has been already referred to as really the external layer of the retina, and not in any way part of the choroid coat. The optic nerve, having entered the interior of the globe through the sclerotic and the choroid, expands into the delicate nerve layer, called the retina, which forms the third tunic of the globe. The retina is in contact, externally, with the choroid; internally, with the hyaloid membrane, which separates it from the vitreous; and it extends forward nearly to the posterior margin of the ciliary processes, where it terminates in a thin serrated border—the *ora serrata;* from this border a thin membrane—*pars ciliaris retinæ*—destitute of nerve fibres, is continued forward to the tips of the ciliary processes, and thence to the posterior surface of the iris.

In passing through the coats of the eye, the optic nerve becomes gradually constricted and reduced to one-half of its diameter; here it presents a round disc, called the *porus opticus*, in the centre of which may be seen the arteria centralis retinæ. At this point, too, the nerve substance projects slightly into the interior of the globe, forming a little prominence, to which the term *colliculus nervi optici* has been applied. This prominence is remarkable, in that it is insensible to the rays of light, and is hence called the "blind spot."

The retina when fresh is nearly transparent, but soon it becomes

of a pink milky tint. Precisely opposite the pupil, in the centre of the axis of vision, there is an oval, yellow spot, *macula lutea*, in the retina, about $\frac{1}{25}$ of an inch in diameter, having a depression, *fovea centralis*, in the centre, and fading off gradually at the edges.

Here vision is most perfect, so that it might be called the "spot of light." This central spot was believed by its discoverer, Sömmering, to be a perforation; but it is now ascertained to be due to the pigmentary layer of the retina showing through it. These appearances are lost soon after death, and are replaced by a minute fold, into which the retina gathers itself, reaching from the centre of the spot to the prominence of the optic nerve.[1]

Structure. Although to the naked eye the retina appears a simple, soft, semi-transparent membrane, yet when examined under the microscope it is found to be most minutely and elaborately organized. It varies in thickness from the $\frac{1}{50}$ to the $\frac{1}{100}$ of an inch, being thickest behind, and gradually diminishes toward the front. It consists of eight layers, through which may be traced a considerable amount of extremely delicate connective tissue (*fibres of Müller*), which constitutes a sustentacular tissue for the various strata, and is said to form for them two more or less continuous boundary layers, termed *membranæ limitantes*, *interna* and *externa*, and which are classed by some anatomists as two additional layers. The layers of the retina are as follows, beginning from within :—

1. The layer of nerve fibres.
2. The ganglionic layer.
3. The inner molecular layer.
4. The inner nuclear layer.
5. The outer molecular layer.
6. The outer nuclear layer.
7. The layer of rods and cones.
8. The pigmentary layer.

The membranæ limitantes are situated as follows: the internal stratum lies on the inner surface of the layer of nerve fibres; the external, between the outer nuclear layer and the layer of rods and cones.

[1] In birds the retina has throughout the yellowish color seen only at one part in the human eye.

1. The *layer of nerve fibres* (Fig. 200, 1) is composed of the spreading out of the optic nerve fibres, and of connective tissue cells. The nerve fibres, consisting only of the axis cylinders, run forward as a continuous layer to the ora serrata, partly arranged in bundles and partly in plexuses, and become connected with

FIG. 200.



DIAGRAM OF THE VARIOUS LAYERS OF THE RETINA. (AFTER SCHWALBE.)

the nerve cells of the next layer. The fibres are almost absent on the yellow spot.

2. The *ganglionic layer* (Fig. 200, 2) is a stratum of spheroidal nerve cells; from the deeper part of each cell there is given off a single elongated process, which passes obliquely into the nerve-fibre layer, with which it becomes continuous; from the outer side of the cell two or more processes are given off, which branch

dichotomously and become at first imbedded and then lost in the inner molecular layer. The ganglionic cells, which in the greater part of the stratum form a single layer, are at the yellow spot arranged eight or ten deep, and in its neighborhood two or three deep.

3. The *inner molecular layer* (Fig. 200, 3) is a granular stratum of considerable thickness which exhibits, under high powers, a reticular structure, having small interstices filled probably with lymph. In it are found the processes of the nerve cells of the preceding layer, which pass outward for a considerable distance, some varicose filaments which pass inward from the next layer; and some Müllerian fibres which pass through this layer. Other cells, like those found in the nerve-fibre and ganglionic layers, are also found in this stratum, chiefly on its surfaces.

4. The *inner nuclear layer* (Fig. 200, 4) contains three kinds of cells, and some fibres which belong to the Müllerian or connective tissue fibres of the retina. The first kind consists of oval bipolar cells placed longitudinally, and having a distinct nucleus and nucleolus; from the inner extremities of these cells there extend long, varicose, thin processes, which pass vertically downward, without division, into the inner molecular layer, and are presumably connected with the processes of the ganglionic layer and thence with the nerve fibres; from the outer extremities of the cells pass processes, thicker than the ones just described and not varicose, which pass to the next layer and there break up into numerous filamentous processes. The second kind of cells are small, granular, protoplasmic cells, which are confined to the deeper part of this layer; and the third kind, similar in their appearance, are disposed here and there in the most external stratum of the inner nuclear layer. The course which the Müllerian fibres take through this layer will be described later on.

5. The outer *molecular* or *internuclear layer* (Fig. 200, 5) resembles in most respects the inner molecular layer, but is much thinner. It contains, however, numerous flattened, branched cells, having well-marked nuclei and nucleoli, and whose fine branching processes exhibit varicosities in their course, resembling nerve fibrils. Whether these are nerve fibres or only the fibres of the sustentacular tissue is at present not determined; but

this layer, as well as those already described, is developed in the same manner as those of the brain, so that probably these fibres are nerve fibrils.

6. The *outer nuclear layer* (Fig. 200, 6) consists of a thick stratum of nucleated cells, having outward and inward prolongations, which may be recognized as connected respectively with the rods and cones of the next layer. The *rod granules* are the most numerous, and each presents an oval cell, which has a well-marked transverse striation, due to the highly refracting substance being crossed by discs of a less refracting medium. There are usually two, one on each side of the middle of the cell. From this enlargement one varicose filament passes inward and becomes connected with the outer molecular layer by a dilatation, from which numerous filaments pass inward; the other extremity is thicker, not varicose, and passes outward toward the membrana limitans externa, where it becomes somewhat expanded, and then becomes continuous with a rod. The *cone granules* are fewer, and each has an oval nucleated cell, which presents no transverse striation characteristic of the rod granule. The cell is situated close to the membrana limitans externa, and rests upon a thick cone fibre, much thicker than a rod fibre, which enlarges as it approaches the outer molecular layer, upon which it rests by a pyramidal base. From this base numerous fine processes are given off into the molecular layer; the outer extremity is very short and broad, and supports the base of a cone.

7. The *layer of rods and cones*, *bacillary layer*, or *Jacob's membrane* (Fig. 200, 7) is composed of minute cylindrical elements, arranged at right angles to the surface of the retina. The *rods*, the more numerous, are tapering processes running through the whole thickness of this layer, and, externally, are embedded to a greater or less depth in the pigmented layer, so that when viewed from without they have the appearance of mosaic pavement made up of round segments. Among the rods are intermingled numerous shorter, flask-shaped bodies called *cones*, which do not extend through more than half the thickness of this layer. Their outer extremities taper off toward the choroid; their inner or broad ends, like the rods, rest upon the membrana limitans externa, and thence are connected with the outer nuclear layer.

Each rod and cone consists of two segments of equal lengths; the inner, in the case of the cones, very broad and bottle-shaped, of the rods only slightly bulged; the outer, fine and tapering off. The two segments vary in their microscopic appearance and in relations to reagents; the outer segments of both have a transverse striation, and break up in the direction of this striation; the inner segments are composed externally of longitudinal fibrillæ, internally of finely granular homogeneous substance continuous with the rod or cone fibre. The inner segments are deeply stained by carmine, iodine, etc.; the outer segments not by the same reagents, but are by osmic acid. The rods are absent at the yellow spot.

8. The *pigmentary layer* (Fig. 200, 8) is usually described as forming part of the choroid coat, but it should, both developmentally and physiologically, be included as one of the layers of the retina. It consists of a single layer of hexagonal nucleated cells filled with pigment granules, which are most numerous toward the margins of the cells. The surface of the cells which look toward the choroid is smooth and destitute of pigment granules, and it is here that the nucleus is situated; the surface toward Jacob's membrane is filled with pigment, which is not well defined, but runs down among the rods, so that their outer part is imbedded among the pigment cells. The use of the pigment is to absorb the rays of light which pass through the retina, and thus prevent their being reflected. It serves the same purpose as the black paint with which the inside of optical instruments is darkened. Albinoes, in whom this layer has little or no pigment, are, consequently, dazzled by daylight and see better in the dusk.[1]

The *sustentacular tissue* (*Müllerian fibres*) is a tissue which runs through the greater thickness of the retina, beginning at the so-called membrana limitans interna, and ending at the membrana limitans externa; but in neither of these two situations does it

[1] In many of the nocturnal carnivorous quadrupeds, the inner surface of the choroid at the bottom of the eye presents a brilliant color and metallic lustre. It is called the *tapetum*. By reflecting the rays of light a second time through the retina, it probably enables the animal to see better in the dusk. It is the cause of the well-known glare of the eyes of cats and other animals; and the great breadth of the luminous appearance arises from the dilatation of the pupil.

form a continuous layer, so that it cannot be classed under the layers forming the retina. These fibres are probably of the nature of a delicate connective tissue, which serves to sustain the various layers and their constituent elements. Each fibre begins by a broad, conical base, on the deeper aspect of the layer of nerve fibres (the bases of these fibres being more or less in connection with each other); it then passes through the layers of nerve fibres and ganglionic cells, and consequently, the inner molecular layer, gradually diminishing in thickness; on reaching the inner nuclear layer it gives off thin filamentous processes which support the structures of this stratum, presenting here a lateral bulge with a well-marked nucleus. After passing through the outer molecular layer it reaches the outer nuclear layer, and then breaks up into filaments which join with fibrils from other Müllerian fibres, thus enclosing and supporting the cells and their prolongations of this layer. These filamentous offsets reach as far as the bases of the rods and cones, forming a bed on which they rest; this is described as the membrana limitans externa.

The *structure* of the *macula lutea* and *fovea centralis*.—In the *macula lutea*, the nerve fibres do not form a continuous layer; the ganglionic layer consists of cells six to eight deep; there are no rods; the cones are longer and narrower than elsewhere; and the outer nuclear layer has only cone fibres. In the *fovea centralis* there are no rods, and the cones are longer than in the macula; and all the other layers are much thinned. At the margin of the fovea most of the layers are thicker than elsewhere.

The *structure* of the *ora serrata* is much less complex than the other parts of the retina; the layer of rods and cones disappear, the former first of all; the ganglionic and nerve-fibre layers become thin and then cease; the inner molecular layer loses much of its granular appearance, and is largely formed of sustentacular tissue, and then abruptly ceases; the inner and outer nuclear layers become thinner, and then gradually merge into a single layer, which is continued on to the *pars ciliaris* as a single stratum of columnar epithelial cells.

The *arteria centralis retinæ*, after emerging through the porus opticus, divides into two branches—an *upper* and a *lower*—which then form a delicate network of blood vessels throughout the

nerve-fibre layer, penetrating as far as the inner nuclear layer, beyond which no capillaries can be traced. After maceration in water, the nervous substance can be removed with a camel's-hair brush, and then in an injected eye the network formed by the vessels can be distinctly seen. The arteries of the retina do not communicate directly with the choroidal vessels.

Aqueous Humor. The *aqueous humor* consists of a few drops of an alkaline, clear, watery fluid, which fills the space between the cornea and the lens.[1] The iris lies in it, and divides the space into two chambers of unequal size—an anterior and a posterior. The *posterior* is much the smaller of the two; indeed, the iris rests on the capsule of the lens, so that, strictly speaking, there is no interval between the opposed surfaces, except a triangular interval bounded by the attachment of the iris, the ciliary processes, and the zone of Zinn. This accounts for the frequent adhesions which take place during inflammation of the iris, between the iris and the capsule of the crystalline lens.[2] A delicate layer of epithelium covers the posterior surface of the cornea, but nothing like a continuous membrane can be demonstrated on the iris or the capsule of the lens. The *anterior* chamber is remarkable for the rapidity with which it absorbs and secretes; as is proved, in the one case, by the speedy removal of extravasated blood; in the other, by the rapid reappearance of the aqueous humor after the extraction of a cataract.

The Vitreous Body and the Hyaloid Membrane. The vitreous body is a transparent, gelatinous-looking substance, which fills up nearly four-fifths of the interior of the globe (p. 808). It can be easily separated from the retina, except at the optic disc; in front it presents a deep depression, in which the crystalline lens is imbedded. It is surrounded, except in front, by a delicate, transparent membrane—the *hyaloid membrane*—which forms a capsule for the vitreous body, and is sufficiently strong to keep it in shape after the stronger tunics of the eye have been removed.

When the vitreous humor has been hardened in chromic acid

[1] The solid constituent is mainly composed of chloride of sodium.

[2] Some anatomists describe the anterior chamber as lined by a serous membrane, called the membrane of the aqueous humor.

third of the circumference. Each of these portions is composed of numerous concentric layers, arranged one within the other, like the coats of an onion. If any single layer be examined with the microscope, it is seen to be composed of fibres about $\frac{1}{3000}$ of an inch in thickness, running in a curved direction, and connected together by finely serrated edges. On a transverse section the lens fibres are found to be hexagonal prisms, with very little connecting substance. Between the front of the lens and its capsule is a layer of flattened cells with well-marked eccentric nuclei. The beautiful dovetailing of the fibres of the lens was first pointed out by Sir David Brewster; and to see it in perfection, one ought to examine the lens of the codfish.

The function of the lens is to bring the rays of light to a focus upon the retina.[1]

DISSECTION OF THE ORGAN OF HEARING.

The parts constituting the organ of hearing should be examined in the following order: (1) the outer cartilage or pinna: (2) the meatus auditorius externus; which leads to (3) the tympanum or middle ear; and (4) the labyrinth or internal ear, comprising the vestibule, cochlea and semicircular canals, which contain the distribution of the auditory nerve.

Pinna.

The *pinna* or *auricle* consists of yellow fibro-cartilage covered with integument, and is irregularly concave, to receive the undulations of sound. It is unevenly oval, and presents on its external aspect numerous eminences and hollows, which have received the following names: The circumferential folded border is called the *helix;* the ridge within it, the *antihelix;* between these is a curved groove, called the *fossa of the helix.* The antihelix bifurcates toward the front, and encloses the *fossa of the antihelix* (*fossa scaphoidea*). The conical eminence in front of the meatus is termed the *tragus*, on which some hairs are usually found. Behind the tragus, and separated from it by a deep notch (*incisura intertragica*), is the *antitragus.* The *lobule* is the soft, pendulous part placed below the concha, and consists of fat and fibrous tissue. The deep hollow which

[1] The lens contains about 60 per cent. of water, and 30 per cent. of albuminoids.

collects the vibrations of sound, and conveys them into the external meatus, is termed the *concha*.

Structure. The pinna is composed of yellow fibro-cartilage, with some fat and connective tissue, covered with integument, and attached to neighboring parts, partly by fibrous tissue and partly by muscles.

The *skin* is very thin, intimately adherent to the subjacent cartilage, and provided with numerous sebaceous glands, found chiefly in the scaphoid fossa and the concha.

The *cartilage* is a single, uneven plate of fibro-cartilage, which presents all the irregularities of the external ear. The cartilage is incomplete, for there is a deficiency behind the tragus, at the bottom of the concha, which is filled up with fibrous tissue. It has a tubular prolongation inward, which forms the external part of the meatus auditorius externus. The cartilage presents several fissures (*fissures of Santorini*) at the anterior part of the tubular prolongation, which are completed by fibrous tissue. In the front part of the pinna, where the helix makes its first bend, is a conical projection of cartilage, termed the *process of the helix*. The lobule, attached to the lower part of the pinna, is a rounded projection formed of fat and connective tissue; it is this which enlarges with age and obesity of the subject.

Ligaments. The ligaments are: the *anterior ligament*, broad and strong, which passes from the process of the helix to the root of the zygoma; the *posterior ligament*, which extends from the cranial surface of the concha to the mastoid process of the temporal bone. There are also *intrinsic ligaments* which bridge over and fill up the deficiencies in the pinna.

Muscles of the Pinna. The muscles which move the cartilage of the ear as a whole have been described (page 19). Other small muscles extend from one part of the cartilage to another; but they are so indistinct that, unless the subject be very muscular, it is difficult to make them out. The following six—four on the front of the auricle and two behind it—are usually described:—

(*a*) The *musculus major helicis* runs vertically along the front margin of the helix; it arises below from the process of the helix, and is inserted into the curve of the helix as it passes backward.

(*b*) The *musculus minor helicis*, an oblique muscle, lies over that part of the helix which is connected with the concha.

(*c*) The *musculus tragicus* lies vertically over the outer surface of the tragus.

(*d*) The *musculus antitragicus* passes transversely from the antitragus to the lower part of the tail-like process of the helix behind the lobule.

(*e*) The *transversus auriculæ* is on the cranial aspect of the pinna; it passes nearly transversely from the back of the concha to the prominence corresponding to the fossa of the helix.

(*f*) The *obliquus auris* extends vertically from the cranial aspect of the concha to the convexity below it.

The *arteries* of the pinna are derived from the posterior auricular, and from the auricular branches of the temporal and occipital. The *veins* empty themselves into the temporal vein. The *nerves* are furnished by the great auricular branch of the superficial cervical plexus, the auriculo-temporal branch of the inferior maxillary, the posterior auricular branch of the facial, and the auricular branch of the pneumogastric.

Meatus Auditorius Externus. This oval passage leads down to the membrana tympani, and conveys the vibrations of sound to the tympanum. It is about an inch and a quarter in length; its external opening is longest in its vertical direction; its termination is broadest in its transverse. The canal inclines at first upward and forward, and then curves a little downward.[1] Its floor, owing to the oblique direction of the membrana tympani, is a little longer than the roof. It is not of equal calibre throughout, the narrowest part being about the middle; hence the difficulty of extracting foreign bodies which have passed to the bottom of the canal. It is formed, partly by a tubular continuation of the cartilage of the pinna, partly by an osseous canal in the temporal bone.

The *cartilaginous portion* is about half an inch long, and is

[1] To obtain a correct knowledge of the length and dimensions of the meatus, sections should be made through it in different directions, or a cast be taken of it in plaster-of-Paris.

firmly connected to the osseous portion. The cartilage is incomplete at the upper and back part, and the interval is filled in with fibrous tissue.

The *osseous portion*, about three-quarters of an inch in length, is narrower than the cartilaginous portion, and is curved forward and inward. Its outer extremity is rough, for the attachment of the cartilage; its inner presents a narrow groove, except at the upper part, for the insertion of the membrana tympani. The lower and anterior wall of the osseous portion is formed by a semicircular plate of bone, the *tympanic plate*, the outer border of which is thickened and is termed the *external auditory process*.

The *skin* and the cuticle are continued down the passage, and becoming gradually thinner, form a cul-de-sac over the membrana tympani. The outer portion is furnished with hairs and ceruminous glands, which secrete the cerumen or wax, and are only found over the cartilaginous portion of the canal.

Its *arteries* are derived from the posterior auricular, the internal maxillary and the temporal, all branches of the external carotid artery. Its *nerves* come from the auriculo-temporal branch of the inferior maxillary nerve.

Tympanum.

The *tympanum*, or middle ear, is an irregular cavity in the petrous part of the temporal bone, having on its outer side the membrana tympani; on its inner side the labyrinth; behind, the mastoid cells; in front, the carotid canal; below, the wall of the jugular fossa. It is rather less than half an inch in its long diameter; from $\frac{1}{8}$ to $\frac{1}{12}$ of an inch between its outer and inner boundaries; and a quarter of an inch in its vertical direction. It is lined with mucous membrane and filled with air, which is freely admitted through the Eustachian tube; so that atmospheric pressure is equal on both sides of the membrane. A chain of small bones, the *ossicles*, retained in their position by ligaments and acted upon by muscles, passes across it. The use of these bones is to communicate the vibrations of the membrana tympani to the labyrinth. For this purpose one end of the chain is attached to the membrane, the other to the fenestra ovalis. The tympanum is bounded by a floor, a roof, an outer, an inner, an anterior and a posterior wall.

Its *roof* is formed by a thin plate of bone corresponding with

the anterior surface of the pars petrosa, which separates the tympanum from the cranial cavity.

The *floor*, which is narrow, is formed by a thin plate, corresponding to the jugular fossa beneath; it is perforated in front by a small aperture for Jacobson's nerve.

Its *outer wall* is formed mainly by the membrana tympani, and by a ring of bone which affords attachment to it; the latter is pierced by the fissura Glaseri (which gives passage to the processus gracilis of the malleus, the laxator tympani, and the tympanic branch of the internal maxillary artery), by the foramen chordæ posterius, through which the chorda tympani enters the tympanum, and by the foramen chordæ anterius, which is the commencement of the canal of Huguier, for the exit of the chorda tympani nerve.

The *inner wall* is vertical and uneven, and presents the following objects, beginning from above; 1. A horizontal *ridge*, indicating the line of the aquæductus Fallopii; 2. The *fenestra ovalis*, a reniform opening, nearly horizontal, which leads into the vestibule, but is closed in the recent state by a membrane, to which is attached the base of the stapes; 3. Below and in front of the fenestra ovalis is a convex bony prominence, the *promontory*; it is occasioned by the first turn of the cochlea, and is marked by vertical grooves, in which lie the branches of the tympanic plexus of nerves; 4. Below and behind this is the *fenestra rotunda*, which lies at the bottom of a conical depression and is overhung by a projection of bone, so that it cannot be seen, except when viewed obliquely; it leads to the scala tympani of the cochlea, but is closed in the recent state by membrane; 5. Immediately behind the fenestra ovalis is a small conical eminence, named the *pyramid*, in the summit of which is a small aperture, from which the tendon of the stapedius emerges; within the pyramid at its base is a small aperture which leads to the aquæductus Fallopii, and transmits a special filament from the facial nerve to the stapedius.

The *posterior wall* presents three or four openings, one of them large, which lead to the mastoid cells, and convey air into them from the tympanum. The mucous membrane of the tympanum is continued into the mastoid cells through these openings.

The *anterior wall* is pierced by an aperture for the transmission of a small artery from the internal carotid. Into this wall open the *Eustachian tube*, and (in the dry bone) the canal for the *tensor tympani*, which are separated from each other by a bony septum, the *processus cochleariformis*. The *Eustachian tube* is partly osseous, partly cartilaginous; the *cartilaginous portion* has been described, p. 255; the *osseous portion*, about half an inch in length, opens into the lowest part of the anterior wall, and is lined with mucous membrane continuous behind with that of the tympanum and in front with that of the pharynx. The *canal for the tensor tympani* terminates in the anterior wall above the Eustachian tube, as a conical projection, in the apex of which is a small aperture for the tensor muscle; this projection is frequently called the *anterior pyramid*.

Lastly, a nerve called the chorda tympani (a branch of the facial) runs in an arched direction from the back to the front of the tympanum, and is covered with mucous membrane.

Membrana Tympani. The *membrana tympani* is a thin, semi transparent, oval disc, which completely closes the bottom of the meatus auditorius externus. Its transverse diameter slightly exceeds its vertical, and its circumference is set in a bony groove, so that it is stretched, somewhat like the parchment of a drum, on the outer wall of the tympanum.[1] Its plane is not vertical, but slants from above downward, forming, with the floor of the meatus, an angle of 55°. It is slightly conical, the apex being directed inward toward the tympanum, and between its layers is inserted the handle of the malleus which runs downward and forward to a little below the centre.

Structure. It is formed of three layers; an outer, formed by an extremely thin layer of true skin; an inner, by the mucous membrane of the tympanum; and a middle fibrous layer; most of the fibres radiate from the attachment of the tip of the handle of the malleus in a bowed direction, so that the membrane is not a strict cone; other fibres are annular, forming a circumferential ring close to the osseous ring; these stretch over a notch in the upper part of the ring (*notch of Rivini*) so that the

[1] The transverse diameter of the membrane is 0.37 inch; its vertical diameter, 0.33 inch; and it is $\frac{1}{10}$ of an inch thick.

membrane is here flaccid and takes the name of the *membrana flaccida.*

The *arteries* to the membrane are supplied from the tympanic branch of the internal maxillary, the stylo-mastoid branch of the posterior auricular, the Vidian, and the internal carotid.

Ossicula Auditus. The three small bones in the tympanum are named after their fancied resemblance to certain implements, the *malleus*, *incus* and *stapes*. They are articulated to each other by perfect joints, and are so placed that the chain somewhat resembles the letter Z. Their use is to transmit the vibrations of the membrana tympani to the membrane of the fenestra ovalis, and, through it, to the fluid contained within the vestibule. But they have another use, which would be incompatible with a single bone—namely, to permit the tightening and relaxation of the membrane, and thus adapt it either to resist the impulse of a very loud sound, or to favor a more gentle one.

FIG. 202.

THE OSSICLES OF THE RIGHT TYMPANUM.

A. Malleus. B. Incus.
C. Stapes. It lies horizontally and forms a right angle with the long process of the incus.

The *malleus* (Fig. 202, A) or hammer bone, consists of an upper part or *head*, which is suspended from the roof of the tympanum by the suspensory ligament, and articulates posteriorly with the incus. Below the head is a narrow constriction, the *neck*, which is continued on into a long, somewhat curved, tapering process, the *manubrium* or *handle:* it is nearly vertical, and is attached along its whole length to the upper half of the membrana tympani, passing between its inner and middle layers. The *processus gracilis* projects at a right angle below the neck, runs into the Glaserian fissure, and receives the insertion of the laxator tympani. The *processus brevis* is a stunted projection, situated at the junction of the processus gracilis and manubrium, and touches the membrana tympani; it receives the insertion of the tensor tympani.

The *incus*, or anvil bone (Fig. 202, B), is shaped like a tooth, with two unequal, widely separated fangs. Its broad part or *body* presents a concavo-convex articulation in front for the head of the malleus; its *long process* runs nearly parallel with the handle

of the malleus, and articulates with the stapes through the intervention of a small bone, the *os orbiculare*, which, in adult life, forms part of the long process, but in fœtal life is a separate bone; its *short process* is directed horizontally backward, and its point is fixed in a small hollow at the commencement of the mastoid cells.

The *stapes*, or stirrup bone (Fig. 202, c), lies horizontally. Its *head* articulates with the long process of the incus. Below the head is a constriction, the *neck*, which receives at its posterior part the insertion of the stapedius. Two *diverging crura* pass from the head to an oval plate of bone, the *base*, which is attached to the membrane covering the fenestra ovalis.

The tympanic bones are maintained in their positions by various ligaments. The *anterior ligament of the malleus* passes from the head of this bone to the anterior wall of the tympanum; the *suspensory ligament* descends from the roof of the tympanum outward to the head of the malleus, and the *posterior ligament of the incus* passes from the short process to the posterior wall near the mastoid cells. The ossicles are connected by an imperfect *capsular ligament*, which passes from the long process of the incus to the head of the stapes; and by another which passes from the head of the malleus to the incus. The base of the stapes is attached to the margin of the fenestra ovalis by an *annular ligament*. The surfaces of the bones forming these two little joints are covered with cartilage. The joints have also synovial membranes.

Muscles of the Tympanum. The muscles, by moving the tympanic bones, tighten or relax the membrana tympani.

The *tensor tympani* runs in a canal above and parallel to the Eustachian tube, from the cartilaginous part of which it arises, as well as from the apex of the petrous portion of the temporal bone. It passes backward, and terminates in a round tendon, which enters the front wall of the tympanum through a special bony canal, and making a sharp bend outward, is *inserted* into the root of the handle of the malleus. Its nerve comes from the otic ganglion. Its action is to draw inward the head of the malleus, and thus render the membrane tense.

The *laxator tympani arises* from the spinous process of the

sphenoid, and the cartilaginous portion of the Eustachian tube, and is *inserted* into the neck of the malleus close to the root of the processus gracilis. It is supplied by a branch of the facial nerve.[1] Its action is to relax the membrana tympani.

The *stapedius arises* from the hollow of the pyramid, and its tendon, emerging through the aperture in the apex, runs forward to be *inserted* into the neck of the stapes.[2] Its nerve is derived from the facial. By its action it increases the tension upon the fluid in the vestibule.

The *mucous membrane* of the tympanum is continuous with that of the pharynx. It covers the ossicles, muscles and nerves, and is prolonged into the mastoid cells. The membrane is pale and thin, and lined with columnar ciliated epithelium, except on the promontory, the membrana tympani, and the ossicles, where there is only a single layer of flattened cells.

A branch (*chorda tympani*) of the facial nerve enters the tympanum through a foramen, *foramen chordæ posterius*, at the base of the pyramid; it then crosses the tympanum beneath the handle of the malleus and the long process of the incus, leaves the tympanum through a foramen, *foramen chordæ anterius*, and then traverses a canal (*canal of Huguier*), which runs close to the Glaserian fissure. It eventually joins the submaxillary ganglion (p. 160).

The *arteries* supplying the *tympanum* are: (1) the *tympanic* branch of the internal maxillary, which enters through the fissura Glaseri; (2) the *stylo-mastoid* branch of the posterior auricular; (3) small branches from the ascending pharyngeal, which enter with the Eustachian tube; (4) branches from the internal carotid artery; and (5) the petrosal branch of the arteria meningea media.

The *veins* open into the middle meningeal and the pharyngeal veins.

The mucous membrane is supplied with branches from the tympanic plexus, which is formed by filaments from the tympanic

[1] This is usually regarded as a muscle, and is described here as such; no muscular fibres, however, can be traced in it, so that it is probably only ligamentous in structure—a fact borne out in the lower animals.

[2] There is a little sheath, lined with synovial membrane, to facilitate the play of the tendon in the pyramid.

branch of the glosso-pharyngeal nerve, from the carotid sympathetic plexus, and from the large and small superficial petrosal nerves.

Internal Ear. This, in consequence of its complexity, is called the *labyrinth.* It consists of cavities excavated in the most compact part of the temporal bone, and it is divided into three parts: a middle one, called the *vestibule*, being the common cavity in which all communicate; an anterior, named, from its resemblance to a snail's shell, the *cochlea;* and a posterior, consisting of *three semicircular canals;* it communicates externally with the tympanum by means of the fenestra ovalis and rotunda, and internally with the meatus auditorius internus. These cavities are filled with a clear fluid, called the *endolymph*, and contain a membranous expansion, the *membranous labyrinth*, upon which the filaments of the auditory nerve are expanded.

Vestibule. The *vestibule*, or central chamber, is an irregular oblong cavity, about one-fifth of an inch in its widest part, which is at its antero-posterior and at its vertical diameters. On its *outer wall* is the fenestra ovalis, which is closed in the recent state by the base of the stapes; on its *inner wall*, at the front part, is a shallow, round depression, the *fovea hemispherica*, which is perforated at its lower by numerous foramina, *macula cribrosa*, for the transmission of the filaments of the auditory nerve. Posteriorly, this pit is bounded by a ridge, the *crista vestibuli*, and in some subjects there is behind this eminence the opening of a small canal, called the *aquæductus vestibuli.* It leads to the posterior surface of the pars petrosa, and transmits a small vein. In the *roof* is a transverse oval depression, the *fovea hemi-elliptica*, which lodges the utricle; *posteriorly*, the five openings of the semicircular canals open into it; and, *in front*, is a large opening through which it communicates with the scala vestibuli of the cochlea.

Semicircular Canals. The *semicircular canals*, three bony canals, are situated above and rather behind the vestibule. Each canal forms about two-thirds of a circle, is compressed laterally, and is about $\frac{1}{10}$ of an inch in diameter. The canals are not of equal diameter throughout; each presents at one end a dilatation termed the *ampulla*, about $\frac{1}{10}$ of an inch in diameter.

This dilatation corresponds to a similar dilatation of the membranous sac, upon which the auditory nerve expands. The canals open at each extremity into the vestibule by five openings, since one of the apertures is common to the extremities of two canals. Each canal differs in its direction; they are named accordingly, superior, posterior and external.

The *superior semicircular canal* (Fig. 203, 1) is the most anterior of the three; its direction is vertical, and runs across the petrous bone. It rises up higher than any other portion of the labyrinth, and its ampulla is at the outer and anterior extremity, and opens into the upper part of the vestibule; its non-ampulla-

FIG. 203.

1. The superior semicircular canal.
2. The posterior semicircular canal.
3. The external semicircular canal.
4. Common opening of the superior and posterior semicircular canals.

5. Aquæductus vestibuli.
6. Aquæductus cochleæ.
7. Fovea hemi-elliptica.
8. Fovea hemispherica.
9. Scala tympani.
10. Scala vestibuli.

OSSEOUS LABYRINTH OF THE RIGHT SIDE (Sömmering).

(Two and a half times its natural size.)

ted extremity opens by a common orifice with the posterior semicircular canal into the back part of the vestibule.

The *posterior semicircular canal* (Fig. 203, 2) is also vertical, and runs parallel to the posterior surface of the petrous bone, consequently at right angles to the preceding. It is the longest of the three canals, and its ampullated extremity is at the lower end, opening into the lower and back part of the vestibule. Its upper non-dilated end joins with that of the superior semicircular canal.

The *external semicircular canal* (Fig. 203, 3), the shortest of the three, is horizontal in position, with the convexity of the arch diverted backward; it opens by its extremities directly into the

back of the vestibule; the ampulla is at the outer end and opens into the vestibule just above the fenestra ovalis.

Cochlea.

The *cochlea* is the most anterior part of the osseous labyrinth; it very closely resembles a common snail's shell, and is placed nearly horizontally, so that its first coil is directed forward and outward, and corresponds with the promontory; while its base corresponds to the bottom of the meatus auditorius internus, and is perforated by apertures for the transmission of the cochlear branches of the auditory nerve. The diameter of its base, and also of its height, is about the same, namely, a quarter of an inch. It consists of a gradually-tapering spiral tube, which winds round a central pillar, called the *modiolus* or *columella*. The spiral canal is divided into two parallel tubes, *scalæ*, by a delicate lamina, partly bony, partly membranous, which is called the *lamina spiralis*. In the dry condition this partition is only partial; but in the recent state it is completed by a membrane.

The *spiral canal* (Fig. 203, 9, 10) is about an inch and a half long, and about one-tenth of an inch in diameter, lessening as it approaches the summit. After making two turns and a half, it terminates at the apex of the cochlea in a rounded dome—the *cupola*. The coil at the base is the widest, the second being a very small one. The canal has in it three openings; thus it communicates with the vestibule by an oval opening; with the tympanum by the fenestra rotunda, but which in the recent state is closed by the *membrana secundaria;* and, lastly, there is the aperture of the *aquæductus cochleæ*, which transmits a small vein from the cochlea to the internal jugular vein.

The *modiolus* or *columella* (Fig. 204, 4) is the central pillar of the cochlea around which the spiral canal coils, and it passes from the base to the apex. It is of considerable thickness at its base, but gradually tapers toward the apex, where at the last half coil it terminates in a half funnel-shaped curved lamella, called the *infundibulum*. Here the partition disappears, and is called the *helicotrema*, so that the scalæ vestibuli and cochleæ communicate with each other in this situation. The interior of the modiolus is composed of cancellous bone, and is traversed by numerous canals, which transmit small vessels and nerves to the lamina spiralis.

One of these canals, larger than the others, runs up the centre of the modiolus nearly to the apex, and transmits a small artery, the *arteria centralis modioli.*

On making a vertical section through the cochlea, we observe that the spiral canal is divided into three tubes, termed *scalæ;* the lower and largest is the *scala tympani* (Fig. 205, s t); the upper is the *scala vestibuli* (Fig. 205, s v), which is subdivided by an oblique membrane to form an outer or third tube—the *scala media* or *canalis cochleæ* (Fig. 205, d c).

The *lamina spiralis* (Fig. 204, 3) is the projecting partition which divides the spiral canal into two tubes or scalæ; is composed on the inner half, of bone—*lamina spiralis ossea*—and on the outer half of membrane—*membrana basilaris.* The lamina spiralis ossea ends at the cupola in a hook-like process—the *hamulus.*

FIG. 204.

THE OSSEOUS COCHLEA. (MAGNIFIED FIVE TIMES.)

Winding round the modiolus, close to the attachment of the lamina spiralis ossea, is a small canal—the *canalis spiralis modioli* —which is filled by the gangliform swelling of the cochlear nerve, called the *ganglion spirale.* The osseous lamina spiralis is seen, on a vertical section, to be composed of two plates, between which the structure is spongy, and presents a number of small canals for the passage of the small filaments of the cochlear division of the auditory nerve, in their course to the membranous part of the lamina.

On examining the membranous continuation of the lamina spiralis, it is seen, not far from its attachment to the osseous zone, to be thickened into an elongated crest—the *limbus laminæ spiralis* (Fig. 205, *lls*)—which overhangs a groove, called the *sulcus*

spiralis (Fig. 205, *ss*); the upper horn of the groove is called the *labium vestibulare;* the lower, the *labium tympanicum.* From the labium tympanicum, the *basilar membrane* is continued outward to be attached to the outer wall of the canal, and thus completes the septum. At the point of attachment of the basilar membrane with the outer wall of the cochlea may be seen a triangular projection, which, formerly described as a muscle (*cochlearis muscle*), is now recognized to be a collection of connective-tissue cells, and called the *ligamentum spirale* (Fig. 205, *lsp*). The structure of the limbus consists of firm connective tissue, on the under part of which are

FIG. 205.



DIAGRAMMATIC SECTION OF A COIL OF THE COCHLEA. (From Quain.)

- s v. Scala vestibuli.
- d c. Ductus cochlearis.
- s t. Scala tympani.
- m r. Membrane of Reissner.
- *l l s*. Limbus laminæ spiralis.
- *m* t. Membrana tectoria.
- *r* c. Rods of Corti.
- *m b*. Membrana basilaris.
- *l s p*. Ligamentum spirale.
- *s s*. Sulcus spiralis.
- *g s*. Ganglion spirale.

found numerous cells. Close to the junction of the limbus with the basilar membrane are a series of regularly arranged apertures, looking upward to the sulcus spiralis; these are ovoid apertures for the exit of branches of the cochlear nerve.

The *basilar membrane* forms, at the base of the cochlea, but a small breadth of the septum, the broadest part being composed of bone; but it gradually increases in breadth toward the cupola, where it constitutes nearly the entire septum. It consists of a firm, fibrillated tissue, which is probably formed, at any rate on its upper surface, of a structure closely resembling the organ of Corti.

It has been stated that in the bony cochlea there is a partial septum dividing the spiral tube into two incomplete scalæ. In the recent condition the basilar membrane completes the septum, dividing the upper tube into an upper canal, the scala vestibuli, and a lower, the scala tympani. The upper scala is subdivided by an oblique membrane, ***membrane of Reissner***, into two canals —an inner, the *scala vestibuli*, and an outer, the *canalis cochleæ*, the ***scala media*** or the *ductus cochlearis* (Fig. 205, D C). The canalis cochleæ terminates at the helicotrema in a cul-de-sac; inferiorly, it is connected with the saccule by a long, narrow duct, called the *canalis reuniens*.

The *membrane of Reissner* (Fig. 205, M R) is the oblique membrane which separates the scala vestibuli and the canalis cochleæ. It is a delicate, almost structureless, layer, composed of connective tissue, continuous with the periosteum lining the upper surface of the lamina spiralis. It is smooth on its vestibular surface, and is lined with flattened connective-tissue cells; on its cochlear surface it is covered with squamous epithelium.

The *inner wall* of the canalis cochleæ is formed by the membrane of Reissner covered with pavement epithelium. The *outer wall*, the periosteum, is thickened by a quantity of retiform connective tissue lined with columnar epithelial cells. An increase of this tissue is seen a little above the ligamentum spirale as a conical eminence, in which runs a small vessel, the *vas spirale;* midway between the vas and the attachment of Reissner's membrane is another thickening, consisting also of numerous blood vessels, *stria vascularis*, which form anastomosing loops. The lower wall is formed by the limbus spiralis and the basilar membrane; upon the latter is placed the complex structure, called the organ of Corti.

The *organ of Corti*, placed upon the upper surface of the membrana basilaris, presents a slight triangular elevation outside the limbus, and winds spirally throughout the cochlea, from its base to its summit. The central part of the organ (Fig. 206, T C) consists of two sets, an inner and an outer, of slanting rods—*rods of Corti* —which rest against each other at their upper extremities, thus forming a triangular tunnel, called the *tunnel of Corti*, filled in the recent state with endolymph. The inner and the outer rods are

similar in structure, but differ in shape—the inner are shorter, less oblique, and have the shape of the human ulna, the outer resemble a swan's head, the head being received into the concavity of the inner rod, the part resembling the bill looking horizontally outward. Both have a broad nucleated base, and present a fibrillar appearance. The inner rods are more numerous than the outer.[1]

FIG. 206.



VERTICAL SECTION OF THE FIRST TURN OF THE COCHLEA, SHOWING THE MEMBRANOUS COCHLEA AND THE POSITION OF THE ORGAN OF CORTI. (AFTER WALDEYER AND QUIN.)

On the inner side of the inner rods is a single row of broad epithelial cells tipped with stiff ciliated processes, called the *inner hair cells;* and on the outer side of the outer rods, resting on cells which are placed on the basilar membrane, are four to six rows of similar cells, termed the *outer hair cells* (206, o c). The

[1] According to Waldeyer, in the proportion of 6000 of the inner to 4500 of the outer rods.

bases of the outer hair cells present on one side a rounded bulge, while from the other are long processes which pass downward to be attached to the membrana basilaris. The outer rods are placed upon numerous fusiform nucleated cells, *cells of Deiters*, whose bases rest upon the basilar membrane, and whose summits taper off into fine, long, cubicular processes, *phalangeal*, which pass between the outer hair cells to be connected to the phalanges of the reticular lamina.

The *lamina reticularis* is the net-like membrane surmounting the summits of the outer hair cells. It is an open network, of a fiddle-shape pattern, consisting of four rows of fiddle-shaped cells termed *phalanges*, through which the ciliated processes of the hair cells project. To these phalanges, as before stated, are attached the phalangeal processes of the cells of Deiters.

The *tectorial membrane* (Fig. 205, M T) is the only remaining membrane to be described, and lies above and parallel to the basilar membrane, but does not extend more than half-way over it. It is connected on its inner side with the limbus spiralis, and is then continued outward, overlying and resting upon the rods of Corti; at its origin it is thin; subsequently it thickens, and then gradually tapers off to end in a free extremity. It is a strong, elastic membrane, distinctly fibrous, especially upon its inner and thicker part.

Membranous Labyrinth.

If the bony labyrinth just described be properly understood, there will not be much difficulty in comprehending the shape of the membranous labyrinth in its interior—a structure supporting the ultimate ramifications of the auditory nerve. It has the general form and shape of the vestibule and the semicircular canals, although smaller, and it is separated from the osseous labyrinth by a quantity of fluid called *perilymph* or *liquor Cotunnii*, which is secreted by the delicate serous membrane lying in the bony labyrinth.

The membranous labyrinth is a sac, contained partly in the vestibule and partly in the semicircular canals; that situated in the vestibule is termed the *vestibular portion;* that in the bony canals, the *membranous semicircular canals.*

The sac in the vestibule is so constructed as to form two sacs

of unequal size, which indirectly communicate with each other.[1] The *utricle* or *common sinus*, the larger of the two, is oval and compressed laterally, and communicates with the five openings of the membranous semicircular canals. It is lodged in the fovea hemi-elliptica, and its wall is thickest (*macula acustica*) close to the crista vestibuli, where the branches from the auditory nerve enter it. The *saccule*, the smaller, is globular and flattened, and lies in the fovea hemispherica, in front of the utricle. It is connected with the membranous canal of the cochlea by a small, short duct, termed the *canalis reuniens*. From the saccule there passes downward, along the aquæductus vestibuli, a narrow prolongation, which terminates in a pyriform dilatation, *saccus endolymphaticus;* this canal is joined, at an acute angle, by a short, narrow canal from the front of the utricle, so that there is a communication existing throughout the entire length of the membranous labyrinth.

The utricle and the saccule contain on their inner wall a minute mass of calcareous matter in connection with nerve-ends, called by Breschet the *otoliths* or *otoconia*. They are crystals of carbonate of lime, and are present in the labyrinth of all mammalia. From their greater hardness and size in aquatic animals, there is reason to believe that they perform the office of rendering the vibrations of sound sharper and more distinct.[2]

Membranous Semicircular Canals.

The membranous semicircular canals present the same dilatations or ampullæ as the bony ones at one end, and at this part they nearly fill their bony cases; but in the rest of their extent the diameter of the membranous canal is not more than one-third to one-fifth that of the bony. At the ampullated extremity the sac is connected on its outer aspect, by blood vessels and nerves, to the periosteum, forming on section a transverse projection, called

[1] From the utricle there proceeds a small canal, which lies in the aquæductus vestibuli; this is joined close to its commencement by a similar canal from the saccule; thus forming the indirect communication above alluded to.

[2] For a detailed description of the relation of the otoliths with the hair-like processes of the nerve filaments, the student is referred to an article by Dr. Urban Pritchard, in the *Quarterly Journal of Microscopic Science*, October, 1876, entitled "The Termination of the Nerves in the Vestibule, and Semicircular Canals of Mammals."

the *septum transversum* or *crista acustica*, which forms a partial septum.

The membranous semicircular canals consist of three layers, an outer or *fibrous layer*, which is connected with the periosteum by blood vessels and contains irregular pigment cells; a middle or *tunica propria*, clear and structureless; and an inner or *epithelial layer*, which lines the inner space of the tunica propria. At the ampullæ the epithelial layer is composed of the columnar variety, upon which are arranged cells of a spindle shape, having delicate ciliated processes (*auditory hairs*) projecting into the endolymph.

The membranous labyrinth is protected, inside and out, by fluid. The fluid in the interior is termed the *endolymph* or *liquor Scarpæ*, and the thin layer between it and the bone the *perilymph* or *liquor Cotunnii;* thus the delicate nervous membrane is placed between two layers of fluid.

Distribution of the Auditory Nerve. The *auditory nerve*, or the eighth cranial nerve, passes down the meatus auditorius internus, and at the bottom of it divides into an anterior and posterior branch which, after breaking up into numerous fasciculi, are distributed to the cochlea and to the vestibule.

The *vestibular nerve* divides into five branches, which proceed to the utricle, the saccule, and the three ampullæ of the semicircular canals, respectively; those for the utricle and the superior and external semicircular canals enter the vestibule along the crista vestibuli; that for the saccule enters through the fovea hemispherica, and that for the posterior semicircular canal is continued along a bony canal to its termination. The nerves to the semicircular canals enter the ampullæ by a forked swelling which corresponds to each septum transversum.

The *cochlear division* of the auditory nerve is a short, thick branch, which breaks up into numerous filaments at the bottom of the meatus auditorius internus. These enter the canals in the base of the modiolus, and then arch outward between the plates of the lamina ossea. In their course outward between the plates, they pass through the spirally arranged ganglionic cord, *ganglion spirale*, beyond which they form a wide plexus. They are collected together close to the free border of the osseous zone, forming a very

minute nerve plexus, whose filaments interlace freely; they then enter the membranous zone, to be connected with the inner hair cells of the organ of Corti.[1]

The vessels which supply the cochlea are from ten to twelve in number, and are derived from the auditory artery; they, like the nerves, enter the bony canals of the modiolus, and then turn outward to ramify upon the osseous zone, supplying its periosteum. The plexus formed by these branches communicates with a vessel known as the *vas spirale*, which runs longitudinally in the ligamentum spirale to the outer attachment of the membrana basilaris. The veins from the cochlea terminate in the superior petrosal sinus, having previously joined those of the vestibule and semicircular canals.

DISSECTION OF THE MAMMARY GLAND.

The form, size, position, and other external characters of the mammary gland in the female vary more or less in different persons. The longest diameter of the gland is in a direction upward and outward toward the axilla; its thickest part is at the centre, and the fullness and roundness of the gland depend upon the amount of fat about it. Its deep surface is flattened in adaptation to the pectoral muscle, to which it is firmly connected by an abundance of areolar tissue. In its vertical direction the breast corresponds to the space between the third and sixth or seventh ribs; in its lateral direction, to the space between the side of the sternum and the axilla, while the nipple corresponds to the fourth rib, or a little below it.

It is enclosed by a fascia, which not only supports it as a whole, but penetrates into its interior, so as to form a framework for its several lobes; hence it is that, in cases of mammary abscess, the matter is apt to be circumscribed, not diffused.

The *nipple* (*mammilla*) projects a little below the centre; it is surrounded by a colored circle, termed the *areola;* this circle is of a rose-pink color in virgins, but in those who have borne children of a dark brown. It begins to enlarge and grow darker about the second or third month of pregnancy, and these changes

[1] Some anatomists describe filaments as passing between the rods of Corti to end in the outer hair cells.

continue till parturition. The areola is abundantly provided with papillæ, and with subcutaneous sebaceous glands, to lubricate the surface during lactation; the areola as well as the nipple is destitute of fat.

Structure. The gland itself consists of distinct lobes held together by firm connective tissue, and provided with separate lactiferous ducts. Each lobe divides and subdivides into lobules, and the duct branches out accordingly.[1] Traced to their origin, we find that the ducts commence in clusters of minute cells, and that the blood vessels ramify minutely upon these cells; altogether, then, a single lobe might be compared to a bunch of grapes, of which the stalk represents the main duct. The main ducts (*galactophorous ducts*) from the several lobes, from fifteen to twenty in number, converge toward the nipple, and, just before they reach it, become dilated into small sacs or *ampullæ*, two or three lines wide; after this they run up to the apex of the nipple, and, running parallel, terminate in separate orifices.

The vesicles and the galactophorous ducts are lined with columnar epithelium, except at their orifices, where it becomes squamous.

The *arteries* of the gland are derived from the long thoracic, the internal mammary, and the intercostals; the *nerves* come from the anterior and lateral cutaneous branches of the intercostal nerves, and from the descending branches of the cervical plexus. The *veins* diverge from the nipple, and terminate in the axillary and internal mammary veins.

The *lymphatics* run chiefly to the axillary glands, but some pierce the front of the intercostal spaces to join the anterior mediastinal glands.

DISSECTION OF THE SCROTUM AND TESTIS.

Scrotum. The *scrotum* is a pouch of skin for the lodgment of the two testes. They are originally developed in the abdomen, and descend into the scrotum about

[1] It is observed, in some cases, that one or more lobules run off to a considerable distance from the main body of the gland, and lie imbedded in the subcutaneous tissue. This should be remembered when it is necessary to remove the entire gland.

the eighth month of intra-uterine life. In their descent they push before them certain coverings derived from the strata of the abdominal walls, through which they pass, and which constitute, with the layers of the scrotum, the covering of the testes. The scrotum presents in the middle a ridge, the *raphé*, on each side of which it is corrugated into transverse folds or *rugæ*. It is divided by a distinct septum into two lateral halves, of which the left is the longer. The scrotum consists of two layers, the integument and the dartos.

The *integument* is of dark color, thrown into transverse rugæ, and provided with sebaceous glands and hairs.

The *dartos* is a thin layer, consisting of muscular fibres of the involuntary kind, like those of the bladder and intestines. It serves to corrugate the loose and extensible skin of the scrotum, and in a measure to support and brace the testes. It is more abundant in the anterior than the posterior part of the scrotum. Beneath the dartos, and partly intermingled with it, is a large quantity of loose connective tissue, remarkable for the total absence of fat. Together with the dartos it forms a vertical partition between the testes, termed the *septum scroti*, which passes from the raphé to the under aspect to the penis, as far as its root, It is not a complete partition, since air or fluid will pass from one side to the other. The great abundance and looseness of this tissue explains the enormous swelling of the scrotum in cases of anasarca, and in cases where urine is extravasated into it in consequence of rupture or ulceration of the urethra.

The coverings of the testes, in addition to these *two layers of the scrotum*, are the *intercolumnar* or *spermatic fascia*, derived from the pillars of the external abdominal ring; the *cremasteric fascia*, derived from the lower border of the internal oblique of the abdomen; the *infundibuliform fascia*, derived from the fascia transversalis; and, lastly, the *tunica vaginalis*, derived from the parietal layer of the peritoneum.

The spermatic fascia, cremaster muscle and the infundibuliform fascia have been described (pp. 447, 449, 456).

Each of these coverings cannot be demonstrated under ordinary circumstances, because they are so blended together; but they can be shown when hypertrophied in the case of old and large herniæ.

The *arteries* supplying the tissues of the testis are the cremasteric branch of the deep epigastric artery, the superficial and deep external pudic branches of the common femoral artery, and the superficial perineal branch of the internal pudic artery.

The *nerves* are derived from the ilio-inguinal, the genital branch of the genito-crural, the superficial perineal nerves and the inferior pudendal branch of the lesser sciatic nerve.

The *lymphatics* pass to the inguinal glands.

Testis. The *testis* is a gland of an oval shape, with flattened sides, suspended obliquely in the scrotum by the spermatic cord, so that its upper end is directed forward

FIG. 207.



1. Mediastinum testis, containing the rete testis.
2, 2. Trabeculæ.
3. One of the lobules.
4, 4. Vasa recta.
5. Coni vasculosi, forming the "globus major" of the epididymis.
6. Globus minor, or lower end of epididymis.
7. Vas deferens.

DIAGRAM OF A VERTICAL SECTION THROUGH THE TESTICLE.

and outward, its lower end in the reverse direction. The left is generally a little lower of the two. Each testis is from an inch and a half to two inches in length, an inch in breadth and an inch and a quarter from behind forward. Its weight is from six to eight drachms; but few organs present greater variations in size and weight, even in men of the same age; generally speaking, the left is the larger. The front and sides of the testis are convex and smooth, and are covered with the visceral layer of the tunica vaginalis; but the posterior surface is only partially invested, as there is here placed a long, narrow body, termed the epididymis;[1]

[1] From δίδυμος, a testicle.

this is not a part of the testis, but an appendage to it, formed by the convolutions of its long excretory duct.

The *epididymis* consists of a larger upper end called the *globus major* (Fig. 207, 5) and of a lower smaller end, the *globus minor* (Fig. 207, 6), the two being connected together by the *body*. The globus major is connected with the testicle by radiating efferent ducts; the globus minor is only connected with the organ by connective tissue and the tunica vaginalis. The upper and lower ends and the outer surface of the epididymis are covered with serous membrane, as is also the body, except at its anterior border, where the vessels enter and emerge. Situated between the globus major and the body of the testis are one or two small pedunculated bodies, called the *hydatids of Morgagni;* they are formed by pouchings of the tunica vaginalis, and are filled with blood vessels bound together by connective tissue.[1]

A considerable quantity of unstriped muscular tissue exists at the posterior part of the epididymis and testis beneath the infundibuliform fascia, and has been described by Kölliker as the *inner muscular tissue.*

Coverings of the Testicle. The testicle is invested by three coverings, which are: 1. A serous membrane, called the *tunica vaginalis*, to facilitate its movements. 2. A strong fibrous membrane, called the *tunica albuginea*, to support the glandular structure within. 3. A delicate vascular stratum, termed the *tunica vasculosa*, which consists of a layer of minute blood vessels.

The *tunica vaginalis* is a closed serous sac, one part of which, *tunica vaginalis propria*, adheres closely to the testis; the other, *tunica vaginalis reflexa*, is the reflected portion, adherent to the inner surface of the infundibuliform fascia, and loosely surrounds the testicle. On opening the sac, it will be seen that the *visceral layer* completely covers the testicle, except behind, where the vessels and duct are situated (Fig. 207); and that it covers the outer part of the epididymis in front and behind, forming here a pouch called the *digital fossa*. The *parietal layer* extends upward for a variable distance upon the cord and below the testicle. The interior of the sac is smooth and polished, like all other serous

[1] The largest, which lies upon the top of the testis, is stated to be the vestige of the Müllerian duct.

membranes, and lubricated by a little fluid. An excess of this fluid gives rise to the disease termed *hydrocele*.

The portion of the process of peritoneum between the internal abdominal ring and the upper part of the tunica vaginalis testis (the spermatic portion of the tunica vaginalis) becomes, in the process of development, converted into a fibrous cord, which may usually be recognized, but which is sometimes so atrophied as not to be recognized.

The tunica vaginalis testis was originally derived from the peritoneum. In some subjects it still communicates with that cavity by a narrow canal, and is therefore liable to become the sac of a hernia (see diagram, p. 464). Such herniæ are called *congenital* —a misleading term, since they do not necessarily take place at

FIG. 208.

1. Spermatic artery.
2. Vas deferens.
3. Deferential artery.
4. Epididymis.
5. Mediastinum testis.
6, 6. Cavity of tunica vaginalis.

TRANSVERSE SECTION THROUGH THE LEFT TESTICLE.
(The dots show the reflections of the tunica vaginalis.)

birth, but may occur at any period of life, even in very old age.[1] Sometimes the communication continues through a very contracted canal, open to the passage of fluid alone; or the communication may be only partially obliterated, and then one or more isolated serous sacs are left along the cord. Such a one, when distended with fluid, gives rise to *hydrocele of the cord*.

The *tunica albuginea* is a dense, white, inelastic membrane, composed of white fibrous tissue, interlacing in every direction; analogous to the sclerotic coat of the eye. It completely invests

[1] It would be a better term to call this lesion a *hernia in the tunica vaginalis*, denoting thereby its anatomical position; at the same time implying a congenital arrest in development, and without limiting its occurrence to any age of life.

the testis, but not the epididymis. It is covered by the visceral layer of the tunica vaginalis, except behind and at the attachments of the epididymis. At the posterior part of the gland it penetrates into its substance for a short distance, and forms an incomplete vertical septum, termed, after the anatomist who first discovered it, *corpus Highmorianum*, and subsequently by Sir A. Cooper, *mediastinum testis* (Fig. 208, 5).

The mediastinum testis transmits the blood vessels of the organ, and contains also the network of seminal ducts, called the *rete testis*, shown in the diagram (Fig. 207). This septum gives off from its front and sides a number of diverging slender fibrous cords, *trabeculæ testis*,[1] which traverse the interior of the gland, and are attached to the inside of the tunica albuginea. They serve to maintain the general shape of the testicle, to support the numerous lobules of which its glandular substance is composed, and to convey the blood vessels into it. These septa, as well as the mediastinum from which they proceed, are readily seen on making a transverse section through the gland (Fig. 208).

The *tunica vasculosa* (*pia mater testis*) consists of a multitude of minute blood vessels, formed by the ramifications of the spermatic artery, and held together by delicate areolar tissue. It covers the inner surface of the tunica albuginea, and gives off branches, which run with the fibrous septa into the interior of the gland.

Minute structure.—When the testis is cut into, its surfaces become convex, and present a dirty fawn color. The section is soft and pulpy, and is seen to consist of numerous *lobules*, between two hundred and fifty and four hundred in number,[2] of various sizes, the central being the larger, and contained in the compartments formed by the fibrous septa proceeding from the mediastinum testis. A few only of these lobules are shown in the diagram. These lobules are conical in shape, their bases being anterior and their apices at the mediastinum, and are bounded by the septa which pass from the mediastinum. They contain two or more minute convoluted tubes, *tubuli seminiferi*, which may be

[1] Kölliker has demonstrated unstriped muscular fibres upon the septa as well as the mediastinum.

[2] The larger estimate is that by Krause; the smaller, that by Berres.

easily unraveled, in consequence of their tough walls. Their number has been estimated to be between 800 and 900, and each has a length of about two feet and a diameter averaging $\frac{1}{120}$ of an inch. They commence either by communications with other tubes or by cæcal extremities, and they frequently exhibit small bulgings in their course backward. The walls of the tubuli consist of a membrana propria, composed of several layers of flattened cells, and the walls are lined with several irregular layers of cells, between which may, under the microscope, be distinguished seminal filaments in various stages of development. The tubuli seminiferi are connected together by a delicate interstitial tissue, the laminæ of which are surrounded by flattened epithelioid cells, and between them are lymph spaces in direct communication with the lymphatics of the testicle. In this intertubular tissue ramify the minute branches of the spermatic artery which surround the tubules.

After pursuing a convoluted course, the tubules unite in front of the mediastinum into from thirty to fifty straight vessels, *vasa recta*, which penetrate the mediastinum testis, and these form an anastomosing plexus of seminal tubes, called the *rete testis* (Fig. 207). This lies along the back of the gland. From the upper part of the rete, its tubes converge to form twelve to fifteen tubes, termed *vasa efferentia*, which perforate the tunica albuginea, and convey the seminal secretion to the upper part of the epididymis. The vasa efferentia are at first straight, but ultimately form a number of coils termed *coni vasculosi*, which collectively constitute the globus major of the epididymis. The coni vasculosi are about $\frac{1}{30}$ of an inch in diameter, and about six to eight lines long; when unraveled they attain a length of six to eight inches.

At the globus major the smaller tubes terminate in a single duct, the *canal of the epididymis*, which in its descent describes an extremely tortuous coil, constituting the body and globus minor of the epididymis. The length of the canal of the epididymis is, in its natural condition, about three inches, but when unraveled it is nearly twenty feet in length. The diameter of the canal at its commencement is about $\frac{1}{70}$ of an inch; at the globus minor about $\frac{1}{50}$ of an inch, after which it again increases in diameter. It is lined with columnar ciliated epithelium.

Vas Deferens. The *vas deferens* begins at the lower part of the globus minor; at first it is somewhat convoluted, but, as it ascends behind the epididymis, it becomes subsequently straight, and joins the other component parts of the cord. After passing through the inguinal canal, it enters the abdomen through the internal ring. It then winds round the outer side of the deep epigastric artery, and, after crossing over the external iliac artery and vein, it enters the pelvis, curves round the side and lower part of the bladder, and empties itself into the prostatic part of the urethra, after running a course of about two feet. Its course in the abdomen has been previously described (p. 554).

In connection with the anterior aspect of the cord, just above the epididymis, are two or three small masses of convoluted tubes, which are known as the *organ of Giraldès*, or the *parepididymis*. They are lined with squamous epithelium, and are probably the remains of part of the Wolffian body.

The *vas aberrans* is a small convoluted tubule, with a cæcal extremity, found between the epididymis and the cord, and communicating usually with the canal of the epididymis. It is about an inch in length, but, when frayed out, varies from two to twelve inches in length. It, like the organ of Giraldès, is connected with a fœtal structure—the Wolffian body.

The vas deferens consists of an external or connective-tissue coat; a middle or muscular coat, composed of longitudinal and circular fibres intermingled with elastic tissue; and an internal or mucous coat, arranged in longitudinal folds, and lined with columnar epithelium. It can always be recognized from the other constituents of the spermatic cord by its hard, whipcord-like feel.

Spermatic Cord. The *spermatic cord* begins at the internal ring, traverses the inguinal canal, and extends to the testis, where its component parts pass to their respective destinations. It is composed of the spermatic vessels, nerves and lymphatics; of the vas deferens, with the deferential artery, a branch of the superior vesical; of the cremaster muscle, and the cremasteric artery, a branch of the deep epigastric. The coverings of the cord have been described with the anatomy of the parts concerned in inguinal hernia (p. 459).

The *spermatic artery* in its course along the cord becomes remarkably tortuous; it enters the back part of the testicle, and breaks up into a number of fine ramifications, which spread out on the inner surface of the tunica albuginea.

The *spermatic veins* leave the testis at its back part, and, as they ascend along the cord, become extremely tortuous and form a plexus, termed the *pampiniform plexus*. They lie in front of the vas deferens and unite to form a single vein, which on the right side opens into the inferior vena cava, and on the left side into the left renal vein. It is usually stated that these veins are destitute of valves; and this fact is adduced as one of the reasons for the occurrence of varicocele. It is, however, certain that the larger veins do contain valves.

The *lymphatics* of the testis pass through the lumbar glands; hence these glands, and not the inguinal, become affected in malignant disease of the testis.

The *nerves* of the testicle are derived from the sympathetic. They descend from the abdomen with the spermatic arteries, and come from the aortic and renal plexuses, with a few filaments from the hypogastric plexus, which surround the deferential artery (p. 559). This accounts for the ready sympathy of the stomach and intestines with the testicle, and for the constitutional effects of an injury to it.

Descent of the Testis. The testicle is originally developed in the lumbar region, immediately below the kidney, and is loosely attached to the back of the abdomen by a fold of peritoneum, termed the *mesorchium*, along which its vessels and nerves run up to it, as to any other abdominal viscus. From the lower end of the gland a fibrous cord, termed the *gubernaculum testis*,[1] proceeds to the bottom of the scrotum. There is no evidence to warrant the assumption that the gradual contraction of the gubernaculum effects the descent of the testis. The organ begins to descend from the lumbar region about the fifth month of fœtal life, reaches the internal ring about the

[1] Mr. Curling considers the gubernaculum testis to be a muscular cord. See his Observations on the Structure of the Gubernaculum, and on the Descent of the Testis in the Fœtus: *Medical Gazette*, April 10th, 1841. This is denied by Cleland, see *Mechanism of the Gubernaculum Testis*, 1856.

seventh, and about the ninth has entered the scrotum. Its original peritoneal coat is retained throughout; but as it enters the inguinal canal, the peritoneal lining of the abdomen is pouched out before it, and eventually becomes the tunica vaginalis reflexa. Immediately after the descent of the testis, its serous bag communicates with the abdomen, and in the lower animals continues to do so throughout life.[1] But in the human subject the canal of communication soon begins to close. It closes at the upper extremity first,[2] and the closure is generally complete in a child born at its full time.[3] This provides against the occurrence of ruptures, to which man, owing to his erect position, is more exposed than animals. At the end of the first month after birth, the canal is entirely obliterated from the internal ring to the testis. Sometimes, however, this obliteration fails, or is only partial; hence may arise congenital hernia, or hydrocele. The possible existence of a communication between the tunica vaginalis and the peritoneal cavity of the abdomen, is one reason, among many, why caution should be observed in treating hydrocele in children with stimulating injections.

[1] According to Professor Owen, the African orang-outang (*Simia troglodytes*) is the only exception to this rule. In this animal it is interesting to observe that the lower extremities are more fully developed as organs of support, and there is a ligamentum teres in the hip-joint.

[2] The frequency of hernia in the funicular portion of the vaginal process of the peritoneum hardly bears this out.

[3] Camper has shown that the canal on the right side is nearly always open at birth, whereas that on the left is usually closed. This explains the greater frequency of hernia on the right side in children under one year old. Thus, out of 3014 cases of inguinal hernia seen at the City of London Truss Society under one year, 2269 occurred on the right side, and 745 on the left; or in the proportion of 3 to 1.

INDEX.

ABD

ABDOMEN, dissection of, 439
parts exposed on opening, 468
regions of, 440
superficial fascia of, 441
surface marking of, 439
Abdominal muscles, functions of, 453
ring, external, 447
internal, 456
walls, arteries of, 456
nerves of, 454
Abducens, nerve, 35, 78, 763
Accelerator urinæ, 531
Acervulus cerebri, 782
Achillis, tendo, 691
Adami, pomum, 259
Adductor brevis, 647
longus, 645
magnus, 648
Ala cinerea, 785
Alimentary canal, anastomoses on, 624
length of, 475
Ampulla of semicircular canals, 835
of rectum, 550
Amygdala cerebelli, 789
Anal glands, 526
fascia, 548
Anastomoses, alimentary canal, 624
Angular artery, 57
vein, 57
Ani, sphincter, 528
levator, 556
Ankle joint, ligaments of, 723
movements of, 724
Annular ligaments of hand, 378
posterior ligament of forearm, 409
ligament of ankle, 662
Annulus ovalis, 219
Ansa hypoglossi, 102, 128
Antihelix, 826
fossa of, 826
Antitragus, 826
Aorta, abdominal, 502
arch, course of, 193
descending thoracic, 201

ART

Aorta arch, (*continued*)
great sinus of, 195
relations of, 194
ascending part of, 195
descending part of, 196
transverse part of, 195
Aponeurosis, epicranial, 18
gluteal, 632
lumbar, 393
pharyngeal, 245, 249
supra-hyoid, 110
vertebral, 298, 393
Apparatus ligamentosus colli, 319
Aquæductus cochleæ, 837
Fallopii, 288
Sylvii, 783
vestibuli, 835
Arachnoid membrane of brain, 729
cavity of, 25
of spinal cord, 795
Arantius, corpus of, 223
nodules of, 223
Arbor vitæ of cerebellum, 790
uterus, 587
Arch, crural, 446, 636
parts passing under, 637
deep, 639
Arm, upper, cutaneous nerves of, 344
musculo-cutaneous nerves of, 356
surface marking of, 343
Arnold's ganglion, 165
nerve, 22, 167
Arteries:
acromial thoracic, 330
alar thoracic, 337
alveolar, 153
anastomotica magna of brachial, 354
femoral, 657
angular, 57
anterior cerebral, 731
ciliary, 815
communicating, 732
perforating, of foot, 704
peroneal, 697

ART

Arteries anterior (*continued*)
spinal, 733
superior cerebellar, 734
tibial, 665
aorta, 193
abdominal, 502
branches of, 485
arch of, 193
ascending, 194
descending, 196
transverse, 195
descending thoracic, 201
articular, of knee, 683
auditory, 734, 845
auricular anterior, 147
posterior, 22
axillary, 334
azygos, 693
back, 310
basilar, 734
transverse of, 734
bicipital, 197
brachial, 351
brachio-cephalic, 197
of brain, 731
bronchial, 213, 243
buccal, 149
bulb, 538
calcanean internal, 697, 703
capsular, 598
carotid, common, 98, 198
external, 112
internal, 39, 163, 289, 731
carpal of radial, anterior, 367
posterior, 367
ulnar, anterior, 369
posterior, 369, 418
centralis modioli, 838
retinæ, 75, 822
cerebellar, anterior superior, 734
posterior inferior, 733
superior, 734
cerebral anterior, 731
middle, 732
posterior, 734
cervical ascending, 139
superficial, 138, 397
transverse, 137
cervicis princeps, 125
profunda, 138
choroid anterior, 733
posterior, 734
ciliary anterior, 75
long, 75
short, 75
circle of Willis, 734.

ART

Arteries (*continued*)
circulus major, 815
minor, 815
circumflex of arm, anterior, 338
posterior, 338, 401
thigh, external, 655
internal, 655
circumflexa ilii deep, 458, 513
superficial, 443, 628
coccygeal, 678
cœliac axis, 486
colica dextra, 493
media, 493
sinistra, 494
comes nervi ischiatici, 678
mediani, 372
phrenici, 176
communicating anterior, of tibial, 697, 732
posterior, of tibial, 732
coronaria ventriculi, 488
coronary, 226
coronary, of lip, inferior, 55
superior, 55
corpus cavernosum, 539
cremasteric, 458
crico-thyroid, 115
cystic, 488
deep cervical, 141
external pudic, 628
palmar arch, 385
temporal, 154
deferential, 559
dental anterior, 155
inferior, 153
superior, 155
digital, of hand, 377
toes, 704
dorsalis hallucis, 667
indicis, 418
linguæ, 124
pedis, 667
pollicis, 418
scapulæ, 405
dorsal interosseous of hand, 418
of foot, 667
dorsal, of penis, 539
epigastric, deep, 457
superficial, 443
ethmoidal anterior, 75
posterior, 75
facial, 55, 116
femoral, 646
superficial, 647
deep, 653
frontal, 20, 76

ART

Arteries (*continued*)
gastric, 488
gastro-duodenalis, 488
gastro-epiploica dextra, 488
sinistra, 488
gluteal, 559, 673
inferior, 679
helicine, 579
hæmorrhoidal external, 528
middle, 559
superior, 494
hepatic, 488, 598
hyoid inferior, 113
superior, 124
hypogastric, 559
ileo-colic, 493
iliac common, 511
external, 512
internal, 557
ilio-lumbar, 558
inferior labial, 55
infraorbital, 65, 155
infraspinous, 139, 396
innominate, 197
intercostal, 209, 456
anterior, 178
collateral, 210
dorsal of, 210
superior, 140
interlobular, 598
interosseous, anterior, 372
common, of forearm, 369
of hand, 386
posterior, 415
recurrent, 416
intestini tenuis, 493
labial inferior, 55
superior, 56
lachrymal, 74
laryngeal, superior, 114
lateral, of nose, 56
sacral, 558
spinal, 733
lingual, 123
long ciliary, 815
lumbar, 505
malleolar, 666
mammary, internal, 140, 178, 457
masseteric, 154
mastoid, 126
maxillary, external, 55
internal, 152
mediastinal, 178
meningea media, 153
parva, 33, 153
meningeal, 32

ART

Arteries meningeal (*continued*)
anterior, 32
middle, 32
posterior, 32, 125, 733
mental, 66
mesenteric, inferior, 494
superior, 492
metatarsal, 667
middle cerebral, 732
hæmorrhoidal, 559
sacral, 562
temporal, 146
vesical, 559
muscular, to eye, 76
musculo-phrenic, 178, 457
mylo-hyoid, 154
nasal, 76, 155
nutrient, of femur, 657
humerus, 354
obturator, 559, 659
pubic branch of, 559
occipital, 20, 125
œsophageal, 213
of gastric, 488
omental, 489
ophthalmic, 74
orbital, 153
ovarian, 505
palatine, ascending, 116, 162
descending, 155
palmar arch, deep, 385
superficial, 375
interosseous, 386
palmaris profunda, 386
palpebral, inferior, 76
superior, 76
pancreatic, 489
pancreatico-duodenal, inferior, 493
superior, 488
perforating, of hand, 386
foot, 704
thigh, 656
intercostal, 178
pericardiac, 178
perineal, superficial, 530
transverse, 531
plantar, external, 705
internal, 704
peroneal, 697
anterior, 697
petrosal, 32, 153
pharyngeal, ascending, 127, 164
meningeal of, 127
pharyngeal, branches of, 164
phrenic, 503
popliteal, 683, 692

BUR

Bursæ (*continued*)
over olecranon, 408
patella, 652
under coraco-acromial ligament, 401
coraco-brachialis, 350
deltoid, 401
gastrocnemius, 690
gluteus maximus, 672
medius, 672
gracilis, 645
latissimus dorsi, 392
ligamentum patellæ, 716
obturator internus, 675
popliteus, 694
Sartorius, 643
semi-membranosus, 686
subscapularis, 405
teres major, 403
triceps, 406
tuberosity of ischium, 528

CÆCUM, 474
meso-cæcum, 483
Calamus scriptorius, 785
Canal, crural, 639
femoral, 639
of Huguier, 834
Hunter, 653
inguinal, 458
of Nück, 525
Petit, 825
Schlemm, 810
Stilling, 824
Wirsung, 602
Canaliculus, 46, 49
Canalis reuniens, 843
spiralis modioli, 838
Canals, semicircular, 835
Canthi, 46
Capsule, Glisson's, 596
of Tenon, 67, 79, 805
Capsules, supra-renal, 613
Caput gallinaginis, 571
Cardiac nerves, of pneumogastric, 167, 206
sympathetic, 169
plexus, deep, 215
superficial, 215
Carotid artery, common, 98, 198
ligature of, 99
difference of left, 101
external, 112
branches of, 113
internal, 163, 289, 731
curves of, 39
plexus, 40, 171
triangles, 94

CER

Carpal arteries, 417
Carpus, bursal sac of, 384
Cartilages, arytenoid, 261
cornicula laryngis, 261
cricoid, 260
cuneiform, 262
epiglottis, 262
of larynx, 259
of nose, 290
of pinna, 827
of trachea, 236
of Wrisberg, 262
tarsal, 48
thyroid, 259
Cartilago triticea, 259
Caruncula lachrymalis, 46
Carunculæ myrtiformes, 542
Cava, vena, inferior, 507
superior, 192
Cavernous plexus, 40, 172
sinus, 30, 37
structures passing through, 37
Centrum ovale majus, 767
minus, 766
Cephalic vein, 330, 347
Cerebelli, falx, 26
tentorium, 26
Cerebellum, 786
inferior peduncles of, 784
middle peduncles of, 789
superior peduncles of, 783
structure of, 790
inferior vermiform process, 787
superior vermiform process, 787
Cerebral circulation, peculiarities of, 735
Cerebri falx, 25
Cerebro-spinal fluid, 730
of cord, 795
Cerebrum, 746
base of, 756
crura of, 759
development of, 790
hemispheres of, 746
labium of, 755
longitudinal fibres of, 786
peduncular fibres of, 786
structure of, 786
transverse fibres of, 786
ventricles of, 769, 772, 779, 783
Cervical fascia, 82
deep, 83, 87
glands, 106
plexus of nerves, 85
deep, 128
superficial, 85
Cervicalis ascendens artery, 139

CER

Cervicalis profunda, 141
superficialis, 140
Cervicis princeps artery, 125
Cervico-facial nerve, 63
Chambers of the eye, 823
Check ligaments, 319
Chest, surface marking of, 324
Chorda tympani nerve, 160
Chordæ tendineæ of ventricles, 222
vocales, 265
Willisii, 28
Choroid arteries, 733, 734
coat of eye, 810
plexus, 778
Chyli receptaculum, 203
Cilia, 47
Ciliary arteries, anterior, 75
long, 75
short, 75
muscle, 813
nerves, long, 73
processes, 812
Circular sinus, 31
Circulation, fœtal, 232
Circulus major and minor, 815
Circumflex artery, anterior, 338
posterior, 338, 401
nerve, 341, 401
external, 655
iliac artery, deep, 458
superficial, 627
internal, 655
Clitoris, 541
erector of, 541
glans of, 541
prepuce of, 541
Coccygeal gland, 507
Coccygeus, 557
Cochlea, 837
canal of, 838
columella of, 837
modiolus, 837
spiral canal, 837
Collateral circulation after ligature of axillary artery, 339
carotid, 100
iliac, external, 514
common, 512
innominate, 198
femoral, common, 657
superficial, 657
subclavian, 141
Colon, ascending, 474
descending, 474
transverse, 474

COW

Columnæ carneæ, 221
rugarum, 583
Columns of Bertini, 607
Commissures, anterior, 780
great transverse, 767
middle, 780
of cord, 798
of labium, 541
optic, 762
posterior, 780
simplex, 788
Communicantes noni nerves, 128
Complexus, 304
Concha of ear, 827
Coni vasculosi testis, 852
Conjunctiva, 47, 804
Constrictors of pharynx, 246
Conus arteriosus, 221
medullaris, 797
Convolutions, dentate, 755
frontal, 750
occipital, 752
parietal, 751
temporo-sphenoidal, 753
uncinate or hippocampal, 755
Cord, spermatic, 459, 853
Cornea, 807
Cornicula laryngis, 261
Coronary arteries of heart, 226
inferior, 55
superior, 55
plexuses, 216
Coronary sinus, 219, 227
Corpora albicantia, 758
quadrigemina, 782
Corpus, Arantii, 224
callosum, 767
ventricle of, 767
cavernosum penis, 577
dentatum cerebelli, 790
fimbriatum, 775
geniculatum, 779
Highmorianum, 851
luteum of ovary, 592
spongiosum, 579
striatum, 775
Corti, organ of, 840
rods of, 849
tunnel of, 840
Costo-coracoid membrane, 328
Cotunnii liquor, 842
Cotunnius, nerve of, 281
Cotyloid ligament, 712
Cowper's glands, 537
structure of, 573

Cranial nerves, dissection to expose, 33
exit of, 33
origin of, 760
Cremaster muscles, 449
Cribriform fascia, 635
Cricoid cartilage, 260
ligaments of, 260
Crista acustica, 844
vestibuli, 835
Crucial anastomosis, 655
ligaments, 718
Crural arch, 636
deep, 639
canal, 639
nerve, anterior, 518, 658
Crusta of crus cerebri, 762
Crystalline lens, 825
Cuneiform cartilages, 262
Cuneus, 755
Cupola of cochlea, 837
Cuvier, duct of, 189
Cystic duct, 491

DARTOS scroti, 847
Deglutition, mechanism of, 257
Deiters, cells of, 842
Deltoid muscle, 399
parts covered by, 401
Demours, membrane of, 809
Dental artery, inferior, 153
superior, 155
nerve, inferior, 159
Descemet, membrane of, 809
Descent of testis, 854
Diaphragm, 498
arteries of, 500
central tendon, 499
crura of, 498
functions of, 501
nerves of, 500
openings in, 500
Digastric muscle, 110
triangle and contents, 109
Diogenes, cup of, 374
Dissection of the:
abdomen, 439
abdominal viscera, 593
axilla, 331
back, 297
muscles of, connected with arm, 387
brain, 729
to remove, 28
cavernous sinus, 37
cranial cavity, 24
digastric triangle, 109
Dissection (*continued*)
epicranial region, 16
eye, 804
extremity, lower, 626
upper, 324
face, 40
femoral hernia, 635
foot, sole of, 698
forearm, back, 408
front, 359
gluteal region, 669
hand, 373
hearing, organ of, 826
heart, 217
inguinal hernia, 460
larynx, 257
leg, back of, 688
front of, 659
mammary gland, 845
maxillary nerve, superior, 277
neck, 80
nose, 290
orbit, 64
pelvis, side view of, 547
viscera of, 519
female, 581
perineum, male, 525
female, 540
pharynx, 244
pterygoid region, 150
scrotum, 846
shoulder muscles, 398
spinal cord, 792
submaxillary triangle, 109
supra-clavicular region, 91
temporal region, 145
testis, 846
thorax, 174
tongue, 272
Dorsal nerves, 211
Douglas, semilunar fold of, 449
pouch of, 480
Ducts, or duct, of—
Bartholin, 122
biliary, 599
common bile, 491
hepatic, 599
Cowper's gland, 537
Cuvier, 189
cystic, 491, 601
galactophorous, 846
hepatic, 491
nasal, 294
pancreatic, 496, 602
parotid gland, 59
prostate, 570

DUC

Ducts (*continued*)
Rivinus, 122
Steno's, 59
sublingual gland, 122
thoracic, 203
Wharton's, 160
Wirsung, 602
Ductus arteriosus, 214
communis choledochus, 491, 601
ejaculatorius, 554
venosus, fissure of, 594
Duodenum, course of, 470
relations of, 495
Dura mater, 24
nerves of, 25
processes of, 25
sinuses of, 27
spinal cord, 794
Duverney's glands, 542

EAR, 826
muscles, 19
intrinsic, 827
Eighth pair of nerves, 36, 764
Ejaculatory ducts, 554
Elbow joint, 428
synovial membrane of, 430
triangle, 361
veins in front of, 346
Eleventh nerve, 37, 168, 765
Eminence, hypothenar, 373
thenar, 373
Eminentia cinerea, 785
collateralis, 777
Endocardium, 229
Endolymph of labyrinth, 835, 843
Ependyma ventriculorum, 769
Epicardium, 229
Epicranial aponeurosis, 18
Epididymis, 849
Epiglottis, 262
cushion of, 264
ligaments of, 265
mucous folds of, 263
Eustachian tube, 255, 831
valve, 220
Eye, 804
lashes, 47
lids, 46
Exit of the cranial nerves, 33

FACE, dissection of, 40
motor nerves of, 41
sensory nerves of, 41, 64
Facial artery, 55, 116
nerve, 36, 86, 287, 764

FIB

Facial nerve (*continued*)
on face, 61
transverse artery, 58
vein, 57, 112
Fallopian tubes, 589
fimbriæ of, 590
Fallopii aquæductus, 61
Falx cerebelli, 26
cerebri, 25
Fascia, anal, 548
arm, 348
forearm, 361
back of forearm, 409
axillary, 331
buccal, 53
bucco-pharyngeal, 53
cervical, deep, 86
costo-coracoid, 329
cremasteric, 450
cribriform, 635
iliac, 508
infundibuliform, 456
intercolumnar, 447
intermuscular of arm, 348
ischio-rectal, 548
lata of thigh, 632
lumbar, 298
metacarpus, 410
muscular of arm, 348
obturator, 547
orbit, 67
palmar, 374
pelvic, 547
perineal, deep, 534
superficial, 529
pharyngeal, 245
plantar, 703
prevertebral, 87
psoas, 508
recto-vesical, 549
semilunar of biceps, 349
spermatic, 447
temporal, 148
transversalis, 455
Fasciculi graciles, 740
teretes, 785
Faucium isthmus, 251
Femoral artery, 647
in canal, 653
sheath of, 638
ring, 639
Fenestra ovalis, 830
rotunda, 830
Ferrein, pyramids of, 610
Fibro-cartilages, interarticular, lower jaw, 323

FIB

Fibro-cartilages (*continued*)
scapulo-clavicular, 424
semilunar of knee, 718
sterno-clavicular, 422
wrist, 431
Fifth cranial nerve, 34, 156, 277, 762
Filum terminale, 796
Fimbriæ of Fallopian tube, 591
Fissures, Bichat, 775
calcarine of brain, 754
calloso-marginal of brain, 755
horizontal, 788
longitudinal, 768
palpebral, 46
parieto-occipital, 749
præcentral, 748
primary of brain, 748
Rolando, 750
Santorini, 827
sphenoidal, nerves in, 38
Sylvius, 748
triradiate, 752
Flocculus, 789
Fœtal circulation, 232
changes at birth, 234
Fold, ary-epiglottic, 262
glosso-epiglottic, 262, 272
Houston's, 624
hepato-renal, 480
rectal, 624
Follicles, Graafian, 592
Fontana, spaces of, 813
Foot, cutaneous nerve of sole, 704
Forearm, cutaneous nerves of, 360
back of, 408
deep, fascia of, 361
fascia on back of, 409
Foramen chordæ anterius, 834
posterius, 834
cæcum of medulla, 737
Magendie, 730, 784
Monro, 774
ovale of heart, 219
quadratum, 500
Thebesius, 221
Winslow, 482
Fornix, 773
conjunctivæ, 47, 804
crura of, 774
Fossa, ischio-rectal, 527
Mohrenheim's, 81
navicularis of labia, 541
of urethra, 575
ovalis, 219
scaphoidea of ear, 826
Fourchette, 541

GLA

Fourth cranial nerve, 34, 69, 762
ventricle, 783
Fovea centralis of eye, 817
hemi-elliptica, 835
hemispherica, 835
inferior, 785
superior, 785
Frænulum, 783
labiorum, 541
Frontal artery, 20, 76
nerve, 68

GALACTOPHOROUS ducts, 846
Galen, veins of, 30, 778
Gall-bladder, 600
duct of, 601
fissure for, 594
relations of, 476
structure of, 601
Ganglion of Andersch, 163
Arnold, 165, 282
Bochdalek, 279
cardiac, 215
cervical inferior, 173
middle, 173
superior, 170
diaphragmatic, 501
Gasserian, 34
geniculate of facial, 287
impar, 170, 565
jugular, 163, 284
lenticular, 77
Meckel's, 279
ophthalmic, 77
optic, 282
petrous, 163, 284
pneumogastric, root of, 165, 286
trunk, 286
semilunar, 498
spheno-palatine, 279
spirale, 838
submaxillary, 123
Wrisberg, 215
Gasserian ganglion, 34
Gastrocnemius, 689
Genito-crural nerve, 518
Gimbernat's ligament, 447, 636
Giraldès, organ of, 853
Glands, agminate, 621
anal, 526
Bartholin, 542
Brunner's, 620
buccal, 53
coccygeal, 507
Cowper's, 537
Duverney's, 542

GLA

Glands (*continued*)
Havers', 714
lachrymal, 70
Lieberkühn, 620
lingual, 275
Littré, 556
Luschka, 507
Meibomian, 50
molar, 54
parotid, 58
pineal, 781
pituitary, 758
prostate, 555
sublingual, 122
submaxillary, 111
thyroid, 103
tracheal, 237
Tyson, 577
uterine, 589
lymphatic axillary, 333
bronchial, 214
elbow, at, 347
femoral, 630
inguinal, 442, 630
intercostal, 213
lumbar, 502
mediastinal, anterior, 179
neck, deep, 106
superficial, 84
popliteal, 693
submaxillary, 117
Glandula socia parotidis, 59
Glandulæ concatenatæ, 106
Pacchionii, 26
Glans clitoridis, 541
penis, 579
Glisson's capsule, 596
Globus major, 849
minor, 849
Glosso-epiglottic folds, 263
Glosso-pharyngeal nerve, 162, 284, 765
Glottidis rima, 264
Glottis, 264
muscles acting on, 269
Gluteal region, 670
Gluteus maximus, parts seen under, 671
nerve, inferior, 672
superior, 564, 673
Graafian follicles, 592
Groin, dissection of, 627
Gubernaculum testis, 854
Gyri operti, 756
Gyrus fornicatus, 755

HAMSTRING muscles, 684
Hamulus of cochlea, 838

HUM

Hand, dissection of palm, 373
surface marking, 373
Hasner, valve of, 294
Havers, gland of, 714
Heart auricle, left, 224
right, 218
cavities, thickness of, 231
chordæ tendinæ, 222
endocardium, 219
fibrous rings of, 228
lymphatics of, 228
muscular fibres of, 229
musculi papillares, 222
nerves of, 214
openings, size of, 226
position of, 186, 217
valves of, 222
veins of, 227
ventricles, left, 225
right, 221
weight of, 218
Helicotrema, 837
Helix, 826
Henle, looped tubes of, in kidney, 610
Hepatic artery, 488
duct, 491
Hernia, congenital, 465
encysted, 466
femoral, 635
coverings of, 641
seat of stricture, 641
funicular, 465
infantile, 465
inguinal, anatomy of, 460
changes in old, 466
direct, 463
oblique, 462
position of cord in, 466
seat of stricture, 466
tunica vaginalis, 465
lumbar, 467
of ovary, 541
umbilical, 467
ventral, 467
Herophili, torcular, 32, 735
Hesselbach's triangle, 463
Hey's ligament, 634
Highmorianum corpus, 851
Hilton's muscle, 270
Hip-joint, movements at, 712
Hippocampus, major, 777
minor, 777
Houston, valves of, 624
Hugier, canal of, 834
Humor, aqueous, 823
vitreous, 823

HUN

Hunter's canal, 653
Hyaloid membrane, 823
Hydatids of Morgagni, 849
Hymen, 542
  varieties of, 542
Hypogastric plexus, 516
Hypoglossi ansa, 102, 128
Hypoglossal nerve, 37, 121, 169, 766
Hyoides os, 258
  basi-hyal, 258
  cerato-hyals, 258
  ligaments of, 258
  thyro-hyals, 258

ILEO-CÆCAL Valve, 623
  Ileo-colic artery, 493
Ileum, course of, 470
Iliac artery
  common, 511
  external, 513
  internal, 557
  ligature of, 513
  fascia, 508
  veins, 562
Iliacus, 510
Ilio-costalis, 301
  lumbar artery, 557
  tibial band, 632
Impressio colica, 595
  renalis, 595
Incisura cerebelli anterior, 788
  posterior, 788
Incus, 832
Infra clavicular triangle, 328
Infundibuliform fascia, 456
Infundibulum of brain, 786
  cochlea, 837
  heart, 221
Inguinal region, 458
  glands, 628
  hernia, 460
  nerves, 444
Infra-orbital artery, 65, 155
  nerve, 65
Innominate artery, 197
  veins, 191
Interarticular fibro-cartilages
    (*See* Fibro-cartilages)
Intercarotic ganglion, 507
Intercolumnar fascia, 447
Intercostal arteries, 209
    anterior, 178
    collateral, 110
    dorsal, 210
    superior, 140
  glands, 213

LAB

Intercostal (*continued*).
  muscles, 209
  nerves, 212
Intercosto-humeral nerves, 332
Internal ear, 835
Interpleural space, 181
Intervertebral fibro-cartilages, 315
Intestine, large, course of, 471
    relations of, 474
    structure of, 622
  small, course of, 471
    structure of, 619
Intumescentia gangliformis, 287
Iris, 813
  arteries of, 815
  nerves of, 815
  structure of, 814
Ischio-rectal fascia, 548
  fossa, 527
    boundaries of, 540
Iter a tertio ad quartum ventriculum, 783
  ad infundibulum, 779

JACOB'S membrane, 820
  Jacobson, nerve of, 285
  Jaw, lower, ligaments of, 161
Jejunum, course of, 470
Joints (*See* Articulations)
Jugular ganglion, 284
  vein, anterior, 84
    external, 83
    internal, 101
    posterior external, 84

KERKRING, valves of, 619
  Kidneys, 606
  arteries of, 505, 614
  capsule of, 609
  cortical structure of, 607
  lymphatics of, 614
  medullary structure of, 608
  nerves of, 614
  pyramids of, 608
  situation of, 477
  tubuli uriniferi, 609
  veins of, 504, 614
Knee-joints, arteries of, 692
  ligaments of, 716
  movements of, 720
  nerves of, 669

LABIA majora, 540
  minora, 522
Labial arteries, 55
  nerves, 65

LAB

Labium cerebri, 767
tympanicum. 839
vestibulare, 839
Labyrinth of ear, 835
Lachrymal gland, 70
nerve, 70
sac, 49, 294
Lacunæ of urethra, 575
Lacus lachrymalis, 46
Lamina, anterior, of cornea, 809
posterior, of cornea, 809
cinerea, 757
cribrosa, 807
fusca, 807
spiralis, 838
ossea, 838
supra-choroidea, 811
Laminated tubercle of cerebellum, 788
Lancisi, nerves of, 768
Larynx, 258
arteries, 270
cartilages, 259
male and female, 271
mucous membrane of, 263
muscles, 267
nerves, 270
situation of, 257
upper opening of, 264
ventricles of, 266
Laryngotomy, 107
Lateral sinuses, 30
ventricles, 769
Latissimus dorsi, 342, 392
Lawrence on femoral hernia, 641
Leg, dissection of, back, 688
front, 659
Lens, crystalline, 825
capsule of, 825
structure of, 825
suspensory ligament of, 824
Lenticular ganglion, 77
Lieberkühn's glands, 620, 622
Ligamenta alaria, 720
annularia, 383
vaginalia, 383
Ligaments of:
accessory of shoulder, 426
acromio-clavicular, inferior, 424
superior, 424
ankle, anterior, 722
lateral external, 723
internal, 722
annular anterior of ankle, 662
external of ankle, 663
internal of ankle, 663
carpus anterior, 378

LIG

Ligaments of (*continued*)
radius, 430
posterior of wrist, 409
anterior carpal, 433
common, 314
elbow, 429
wrist, 433
arcuate, external, 498
internal, 498
astragalo-scaphoid, 724
atlo-axoid, anterior, 319
posterior, 319
bladder, false, 523, 546
true, 548
Burns', 634
calcaneo-astragaloid, external, 722
interosseous, 722
posterior, 722
calcaneo-cuboid, long, 726
short, 726
internal, 726
superior, 725
carpo-metacarpal, 435
dorsal, 436
palmar, 436
interosseous, 436
carpus, 433
central of cord, 796
check, 319
ciliary of eye, 813
common, anterior vertebral, 314
posterior vertebral, 314
conoid, 424
coraco acromial, 425
clavicular, 424
humeral, 426
coracoid, 425
coronary of knee, 719
liver, 594
costo-clavicular, 422
sternal, 322
transverse, 321
vertebral, 320
cotyloid, 714
crico-arytenoid, 261
crico-thyroid, 260
crucial of knee, 718
cruciata, 383
cruciform, 319
deltoid of ankle, 723
elbow, anterior and posterior, 429
lateral external, 429
internal, 429
falciform of Burns, 634
liver, 475
sacro-sciatic, 710

LIG

Ligaments of (*continued*)
femoral (Hey's), 634
gastro-phrenic, 480
Gimbernat's, 447, 639
glenoid, 427
of fingers, 437
Hey's, 634
hip, 711
hyo-epiglottic, 262
hyoid bone, 258
ilio-femoral, 712
lumbar, 708
incus, 833
interarticular of ribs, 320
intercarpal, 434
interclavicular, 422
intermetatarsal, 728
interosseous of cuneiform bones, 726
interosseous of forearm, 430
leg, 722
tarsus, 724
interspinous, 315
intertransverse, 317
jaw, 322
knee, 715
latum pulmonis, 179
long calcaneo-cuboid, 726
lumbo-sacral, 708
metacarpal, 436
metacarpo-phalangeal, 437
metatarsal, 728
metatarso-phalangeal, 728
mucosum of knee, 720
malleus, 833
nuchæ, 317
oblique of radius, 431
sacro-iliac, 710
occipito-atloid, 317
occipito-axial, 318
odontoid, 319
orbicular of radius, 430
ovary, 590
palpebral, 49
palmar, 383
patella, 651
phalangeal of hand, 438
foot, 748
plantar, long, 726
posterior, common, 314
of Winslow, 686, 717
Poupart's, 446, 636
pterygo-maxillary, 53
pubic, 711
sub-pubic, 711
pubo-femoral, 712
prostatic, 545

LIG

Ligaments of (*continued*)
radio-carpal, 433
ulnar, 726
rhomboid, 422
round, of hip, 713
liver, 476
radius, 431
uterus, 459, 525
sacro-coccygeal, 709
sacro-iliac anterior, 709
posterior, 710
oblique, 710
sacro-sciatic, great, 710
lesser, 710
sacro-vertebral, 708
scapulo-clavicular, 424
shoulder joint, 425
stellate, 320
sterno-clavicular, 421
stylo-hyoid, 162
stylo-maxillary, 88, 118
subpubic, 711
subflava, 315
supraspinous, 315
suspensory of lens, 824
liver, 475
penis, 576
tarso-metatarsal, 727
tarsus, 724
temporo-maxillary, 322
teres of hip, 713
thyro-arytenoid inferior, 265
superior, 265
thyro-epiglottic, 262
thyro-hyoid, 259
tibio-fibular inferior, 722
superior, 720
transverse, of fingers, 375
hip, 705
knee, 719
metacarpal, 419
metatarsal, 727
trapezoid, 424
triangular of abdomen, 447
perineum, 535
uterus, 524
Winslow, 686, 717
wrist, 433
Zinn, 78
Ligamenta subflava, 315
Ligamentum denticulatum, 797
mucosum, 720
nuchæ, 317
patellæ, 716
pectinatum iridis, 810
posticum Winslowii, 717

LIG

Ligamentum (*continued*)
spirale, 837
suspensorium, 319
Ligula, 785
Limbus spiralis, 837
Linea alba, 452
semilunaris, 449
splendens, 796
transverse of fourth ventricle, 784
rectus, 452
Linguæ frænum, 272
dorsalis artery, 124
Lingual artery, 123
glands, 275
nerve, 122
Lithotomy, parts divided in, 536
parts to be avoided in, 536
Littré, glands of, 575
Liver, situation of, 475
arteries of, 598
cells of, 599
coats of, 596
ducts of, 599
fissures of, 594
functions of, 600
ligaments of, 596
lobes of, 595
lobules of, 596
lymphatics of, 599
nerves of, 600
plexus of, 597
structure of, 596
veins of, 597
Lobule of the ear, 826
paracentral, 750
Lobulus caudatus, 596
centralis, 788
quadratus, 596
Spigelii, 595
Lobus cæruleus, 785
niger, 759
perforatus anticus, 757
posticus, 758
Longitudinal sinus, inferior, 30
superior, 27
Lumbar aponeurosis, 393
arteries, 505
fascia, 298
hernia, 467
plexus of nerves, 516
Lumbricales of hand, 385
foot, 702
Lungs, 238
air cells of, 242
arteries of, 242
infundibula of, 241

MED

Lungs (*continued*)
lobes of, 185
lymphatics of, 243
nerves of, 243
position and form of, 185
root of, constituents, 217
shape of, 258
structure of, 235
veins of, 243
Lunulæ of valves, 224
Luschka, gland of, 507
Lymphatics, heart, 228
kidney, 613
liver, 599
lung, 243
pancreas, 602
parotid gland, 58
scalp, 24
spleen, 603
testis, 854
uterus, 585
Lymphatic glands, cervical deep, 106
superficial, 84
Lyra of fornix, 775

MACULA acustica, 843
cribrosa, 835
lutea, 817
Magendie, foramen of, 730, 784
Malleus, 832
ligaments of, 833
Malpighi, pyramids of, 608
Malpighian bodies of kidney, 612
spleen, 605
Mammary gland, 845
arteries of, 846
lymphatics of, 846
internal artery, 140, 178
Mammilla, 845
Marginal convolution, 755
Marshall, vestigial fold of, 221
Maxillary artery, external, 55
internal, 152
nerve, inferior, 35, 156
superior, 34, 277
vein, 156
Meatus auditorius externus, 828
of the nose, 293
urinarius, female, 543
Meckel's ganglion, 279
Median artery, 370
nerve, 355
in palm. 381
vein, 360
Mediastinum, 181
anterior, 182

MED

Mediastinum (*continued*)
middle, 183
posterior, 182, 200
superior, 183
Medulla oblongata, 736
anterior pyramid, 737
fissures of, 737
lateral tracts of, 741
nuclei of, 742, 744
olivary bodies, 738
fasciculus, 741
posterior pyramids, 740
restiform bodies, 739
septum of, 743
Medulli-spinal veins, 799
Meibomian glands, 50
Meissner's plexus of nerves, 622
Membrana basilaris, 838
fusca, 807
limitans of retina, 821
nictitans, 47
pupillaris, 813
sacciformis, 433
tectoria, 842
tympani, 831
arteries of, 832
flaccida, 832
secundaria, 837
Membrane, of aqueous humor, 823
arachnoid of brain, 729
spinal cord, 795
Bruch, 812
choroid, 811
Corti, 839
costa-coracoid, 328
crico-thyroid, 260
Demours, 809
Descemet, 809
hyaloid, 823
interosseous of forearm, 430
leg, 721
Jacob's, 820
Reissner, 840
thyro-hyoid, 258
Schneiderian, 294
Membranes of brain, 24, 729
spinal cord, 795
Membranous labyrinth, 842
semicircular canals, 842
Meningeal arteries, 32
anterior, 32
middle, 32, 153
parva, 33
posterior, 33, 125
small, 153

MUS

Mesenteric artery, inferior, 494
superior, 492
veins, inferior, 494
superior, 493
Mesentery, 483
Meso colon, ascending, 483
descending, 483
transverse, 482
sigmoid, 484
rectum, 484, 550
Mesorchium, 854
Metencephalon, 790
Mitral valves, 225
Modiolus of cochlea, 837
Mohrenheim's fossa, 81
Molar glands, 54
Monro, foramen of, 774
Mons Veneris, 540
Monticulus cerebelli, 788
Morgagni, sinus of, 249
hydatids of, 849
Morsus diaboli, 590
Motor oculi nerve, 77, 762
Mouth, muscles of, 41
Movements of spine, 317
Müller, fibres of, 817
Multifidus spinæ, 305
Muscles:—
abdominal functions of, 453
nerves of, 454
abductor hallucis, 700
indicis, 419
minimi digiti manus, 380
pedis, 700
pollicis, 379
accelerator urinæ, 531
accessorius ad sacro-lumbalem, 301
flexor, 701
adductor brevis femoris, 647
hallucis, 705
longus, 645
magnus, 648
pollicis, 379
anconeous, 413
antitragicus, 826
aryteno-epiglottideus, 268
inferior, 270
superior, 270
arytenoideus, 268
attollens aurem, 19
attrahens aurem, 19
azygos uvulæ, 254
back of, 299
biceps of arm, 349
thigh, 685

MUS

Muscles (*continued*)
biventer cervicis, 304
brachialis anticus, 351
buccinator, 52
cervicalis ascendens, 302
ciliary, 813
coccygeus, 557
cochlearis, 839
complexus, 304
compressor naris, 51
sacculi laryngis, 270
urethræ, 538
in the female, 543
coraco-brachialis, 350
corrugator supercilii, 46
cremaster, 449
crico-arytenoideus lateralis, 268
posticus, 267
crico-thyroideus, 267
crureus, 650
dartos, 847
deltoid, 399
depressor alæ nasi, 51
anguli oris, 43
labii inferioris, 44
diaphragm, 498
digastricus, 110
dilatator naris anterior, 52
posterior, 51
ejaculator urinæ, 531
erector clitoridis, 541
penis, 532
spinæ, 300
external sphincter ani, 528
extensor brevis digitorum, 666
carpi radialis brevior, 411
longior, 410
carpi ulnaris, 413
communis digitorum, 411
indicis, 415
longus digitorum pedis, 664
minimi digiti, 412
ossis met. pollicis, 414
primi internodii pollicis, 414
secundi internodii pollicis, 414
proprius hallucis, 665
flexor accessorius, 702
brevis digitorum, 701
hallucis, 705
minimi digiti manus, 380
pedis, 706
pollicis, 379
carpi radialis, 363
ulnaris, 364
longus digitorum pedis, 684
pollicis, 371

MUS

Muscle flexor (*continued*)
profundus digitorum, 371
proprius hallucis, 695
sublimis digitorum, 364
gastrocnemius, 689
gemellus inferior, 675
superior, 675
genio-hyoglossus, 120
hyoideus, 119
gluteus maximus, 671
medius, 672
minimus, 673
gracilis, 645
helicis major, 827
minor, 828
hyoglossus, 119
iliacus, 510
ilio-costalis, 301
indicator, 415
inferior constrictor of pharynx, 247
infra-spinatus, 402
intercostal, external, 209
internal, 209
interosseous of foot, 707
of hand, 420
interspinales, 306
intertransversales, 306
ischio-cavernosus, 532
kerato-cricoid, 268
latissimus dorsi, 342, 392
laxator tympani, 833
levator anguli oris, 52
scapulæ, 394
ani, 556
female, 581
labii inferioris, 44
superioris alæque nasi, 52
proprius, 52
menti, 44
palati, 253
palpebræ, 48, 71
prostatæ, 557
levatores costarum, 306
lingualis, inferior, 276
superficial, 276
longissimus dorsi, 302
longus colli, 312
lumbricales manus, 385
pedis, 703
masseter, 147
of mastication, 148
middle constrictor of pharynx, 247
mouth, 41
multifidus spinæ, 305
mylo-hyoideus, 17
nose, of the, 51

MUS

Muscles (*continued*)
obliquus auris, 828
externus abdominis, 445
inferior oculi, 80
capitis, 308
internus abdominis, 448
superior oculi, 71
capitis, 308
obturator externus, 676
internus, 674
occipito-frontalis, 18
opponens digiti manus, 380
pollicis, 378
orbicularis oris, 43
palpebrarum, 45
omo-hyoid, 95, 395
palato-glossus, 254
pharyngeus, 254
palmaris brevis, 374
longus, 363
pectineus, 643
pectoralis major, 327
minor, 334
peroneus brevis, 668
longus, 668, 706
tertius, 664
plantaris, 690
platysma myoides, 82
popliteus, 694
prevertebral, 312
pronator quadratus, 372
radii teres, 363
psoas magnus, 509
parvus, 510
pterygoideus externus, 150
internus, 150
pyramidalis, 452
nasi, 51
pyriformis, 674
quadratus femoris, 675
lumborum, 511
menti, 44
quadriceps femoris, 649
recti of the eye, 78
rectus abdominis, 451
capitis anticus major, 313
minor, 314
lateralis, 308
posticus major, 307
minor, 308
externus oculi, 78
femoris, 650
internus oculi, 78
inferior oculi, 78
sternalis, 327
superior oculi, 71

MUS

Muscles (*continued*)
retrahens aurem, 19
rhomboideus major, 395
minor, 395
risorius, 42
rotatores spinæ, 305
sacro-lumbalis, 301
sartorius, 643
scalenus anticus, 129
medius, 129
posticus, 129
semi-membranosus, 685
semi-spinalis colli, 305
dorsi, 304
semi-tendinosus, 685
serratus magnus, 343, 397
posticus inferior, 297
superior, 297
soleus, 690
sphincter ani internus, 550
iridis, 814
vaginæ, 544
vesicæ, 567
spinalis dorsi, 303
splenius capitis, 300
colli, 300
stapedius, 834
sterno-cleido mastoideus, 89
hyoid, 95
thyroid, 95
stylo glossus, 121, 161
hyoideus, 111
pharyngeus, 161
subanconeus, 407
subclavius, 329
subcostal, 209
subcrureus, 651
sublimis digitorum, 364
subscapularis, 342, 403
superior constrictor of pharynx, 247
supinator radii brevis, 415
longus, 365
supra-spinales, 306
supra-spinatus, 402
temporal, 149
tensor fasciæ femoris, 649
palati, 253
tarsi, 50
tympani, 833
teres major, 342, 403
minor, 402
thyro-arytenoideus, 269
epiglottideus, 270
hyoid, 97
tibialis anticus, 663
posticus, 695, 708

MUS

Muscles (*continued*)
trachelo-mastoid, 302
tragicus of ear, 828
transversalis abdominis, 450
colli, 302
pedis, 707
transverso-spinalis, 304
transversus auriculæ, 828
perinei. 532
deep, 533
trapezius, 390
triangularis sterni, 177
triceps extensor cubiti, 358, 406
femoris, 650
of ureters, 570
uvulæ azygos. 254
vastus externus, 650
internus, 650
zygomaticus major, 44
minor, 44
Musculi papillares, 222, 226
pectinati, 221
Musculo-spiral nerve, 358
Mylo-hyoid artery, 154
nerve, 117, 159

NABOTHI ovula, 587
Nasal fossæ, posterior openings, 251
duct, 294
muscles, 51
Naso-lobular nerve, 51, 73
Nates of brain, 782
Neck, central line of, 107
cutaneous nerves of, 85
dissection of, 80
lymphatics of, 84
surface marking of, 80
triangles of, 90
anterior, 93
carotid inferior, 94
superior, 94
digastric, 109
posterior, 91
submaxillary, 109
supra-clavicular, 92
Nerves:
abducens oculi, 78, 763
accessory obturator, 659
acromial, 86, 326
anterior crural, 518, 658
cutaneous of abdomen, 444
tibial, 669
Arnold's, 165
auditory, 764, 844
auriculo-parotidean, 85
temporal, 22, 147, 158

NER

Nerves (*continued*)
auricular branch of pneumogastric, 167
posterior, 22, 126
axillary, plexus of, 340
back, cutaneous of, 309
Bell, nerve of, 145, 343
brachial plexus, 142, 340
buccal branch of facial, 63
inferior maxillary, 156
calcaneo-plantar, 698
cardiac branch of pneumogastric, 167, 206
inferior, 173
middle, 173
superior, 172
carotid of glosso-pharyngeal, 282
spheno-palatine ganglion, 282
cervical, acromial branch of, 86, 326
posterior branches of, 387
superficial, 86
plexus, deep, 128
cervico-facial, 63
chorda-tympani, 160, 288, 834
ciliary, long, 73
short, 75
circumflex, 341, 401
clavicular, 86, 326
cochlear, 844
coccygeal, 563
posterior branch of, 390
communicans noni, 102, 128
communicans peronei, 669, 682
cornea, nerves of, 810
coronary anterior, 216
posterior, 216
cranial, exit of, 33
at base of skull, 284
origin of, 760
crural branch of genito-crural, 631
cutaneous of chest, 325
forearm, 360
neck, 85
thigh, 630, 684
external, 630
internal, 631
middle, 631
dental, anterior, 279
inferior, 159
posterior. 278
descendens noni, 102
dorsal, 211, 388
penis, 540
twelfth cranial, 444, 454
dura mater, 24
eighth cranial, 36

Nerves (*continued*)
eighth cranial, origin of, 764
eleventh cranial, 37, 168, 765
external cutaneous, of musculo-spiral, 406
of peroneal, 661
of thigh, 518, 630
laryngeal, 114
plantar, 704
popliteal, 669
respiratory, of Bell, 145, 343
saphenous, 681, 688
superficial petrosal, 40, 288
facial, 22, 61
cervical branch of, 86
in skull, 287
origin of, 763
fifth cranial, 34
origin of, 762
sacral, 562
first cranial, 296, 760
origin of, 760
lumbar, 517
fourth cranial, 34, 70
pair, origin of, 762
sacral, 563
frontal, 21, 68
genital branch of genito-crural, 518, 631
genito-crural, 518
crural branch of, 518, 631
glosso-pharyngeal, 162, 284
carotid branches of, 163
lingual branches of, 163
origin of, 765
pharyngeal branches of, 163
tonsillar, 163
gluteal, inferior, 565, 673, 678
superficial, 673
superior, 673
great sciatic, 677, 681, 687
occipital, 22, 309
gustatory, 122, 159
hypoglossal, 37, 121, 169
origin of, 766
ilio-hypogastric, 454, 517
ilio-inguinal, 444, 454, 518
inguinal branch of, 632
incisor, 159
inferior dental, 159
hæmorrhoidal, 539
laryngeal or recurrent, 206
maxillary, 35, 156
pudendal, 529, 530
infra-maxillary of facial, 64
orbital of facial, 62
Nerves (*continued*)
infra-orbital, of superior maxillary, 65
trochlear, 73
intercostal, 212
abdominal, 212
anterior cutaneous branches, 326
intercosto humeral, 333, 344
lateral cutaneous of, 326
pectoral, 212
internal cutaneous, of arm, 344
thigh, 631
plantar, 704
popliteal, 683
branches of, 683
interosseous anterior, 373
posterior, 417
iris, 813
Jacobson's or tympanic, 285
labial, 65
lachrymal, 70
Lancisi, 768
kidney, 613
laryngeal, external, 116, 167
inferior, 168, 206, 271
inferior or recurrent, 168, 206
internal, 115, 167
recurrent, 168, 206
superior, 115, 167, 270
lateral cutaneous of abdomen, 443
lesser cutaneous of arm, 344
lingual, 122, 159
of glosso-pharyngeal, 163
long or inferior pudendal, 529
long saphenous, 658
liver, 600
lumbar, plexus of, 516
posterior branches of, 388
sympathetic, 514
lumbo-sacral, 562
malar branch of superior maxillary, 66, 80
malar of facial, 62
maxillary inferior, 156
superior, 277
median, 355, 370
cutaneous branch of, 371
digital branches of, 381
in the palm, 381
mental, 66, 159
masseteric, 157
motor oculi, 77
origin of, 762
musculo-cutaneous, of arm, 356
foot, 669
spiral, 358, 360
mylo-hyoid, 159

NER

Nerves (*continued*)
mylo-hyoidean, 117
nasal, 65, 72
septal branch of, 73
superior, 281
upper, 281
naso-lobular, 51, 65, 73
palatine, 281
neck, cutaneous of, 85
nervi molles, 57
ninth cranial, 36, 162
origin of, 765
noni communicantes, 102, 128
obturator, 519, 658
accessory, 519
internus, 563
occipital, great, 22, 309
small, 23, 85
olfactory, 33, 296
origin of, 761
ophthalmic, 34
optic, 33, 73
origin of, 762
orbital branch of superior maxillary, 65, 80
malar branch of, 66, 80
palatine, anterior, 281
external, 281
nasal branches of, 281
naso-, 281
posterior, 281
palpebral, 65
palmar branch of median, 374
of ulnar, 374
pancreas, 603
pars intermedia, 764
perineal, 539
peroneal, 668, 682
external cutaneous branch of, 661
pes anserinus, 61
petrosal, external, 288
great, 282
small superficial, 285
pharyngeal of glosso-pharyngeal, 163
of pneumogastric, 167
phrenic, 128, 130
in chest, 199
plantar, external, 703
internal, 703
pneumogastric, 37, 165, 205, 286
auricular branch of, 22
in the chest, 205
course of, 205
origin of, 765
popliteal, external, 669

NER

Nerves (*continued*)
popliteal, internal, 682
portio dura, 61
posterior auricular, 22, 62
branches of spinal, 309
interosseous, 417
scapular, 396
tibial, 697
thoracic, 145, 343
pterygoid, 157
pudendal, inferior or long, 529
pudic, 539, 564
pulmonary branches of pneumogastric, 206
radial, 367
recurrent or inferior laryngeal, 168, 206
rhomboid, 144
sacral, 562
fourth, 529
plexus, branches of, 563
posterior branches of, 388
saphenous, long, 658, 661
scalp, of, 20
second cranial, 33, 73
origin of, 761
cervical, 290
seventh cranial, 22, 61, 287
origin of, 763
shoulder, cutaneous of, 399
sixth cranial, 35, 78
origin of, 763
small occipital, 23, 85
sciatic, 565, 677
superficial petrosal, 40
in sphenoidal fissure, 38
spheno-ethmoidal, 75
spinal accessory, 91, 168, 394
origin of, 765
spinal, posterior branches of, 387
splanchnic, great, 208
lesser, 208
smallest, 209
spleen, 606
sternal, 86, 326
subclavius, 143
suboccipital, 290
cutaneous branch of, 23
subscapular, 341
superior gluteal, 564, 673
maxillary, 34, 277
supra-clavicular, 86, 326
maxillary branch of facial, 63
orbital, 22, 62, 69
scapular, 144, 396
trochlear, 21, 65, 69

NER

Nerves (*continued*)
sympathetic, abdomen, 514
cervical, 170
in the chest, 207
in the pelvis, 565
temporal branch of superior maxillary, 22, 80
deep anterior, 157
posterior, 157
branches of facial, 22, 62
temporo-facial, 22, 62
malar, 22, 80
tenth cranial nerve, 37, 165, 205, 286
origin of, 765
testis, 854
third cranial nerve, 34, 77
origin of, 762
thoracic anterior, 330
posterior, 145, 343
tonsillar of glosso-pharyngeal, 163
trifacial, 34
trigeminal, origin of, 762
trochlear, of orbit, 69, 73
tympanic, 285, 288
twelfth cranial, 37, 121, 169
origin of, 766
ulnar, 357, 369
deep palmar branch of, 386
dorsal, cutaneous of, 370
in the palm, 377
deep, 386
uterus, 585
vestibular, 844
Vidian, 282
Wrisberg, 333, 344
Nervi molles, 57
Nodule of cerebellum, 788
Nose, arteries of, 56, 292, 295
cartilages of, 290
dissection of, 290
interior of, 292
meatus of, 293
mucous membrane of, 294
muscles of, 51
nerves of, 296
septum of, 292
veins of, 296
Nück, canal of, 525
Nymphæ, 541

OBEX, 785
Obturator artery, 559, 659
abnormal, 560
externus, 676
fascia, 547
internus, 674

OST

Obturator (*continued*)
nerve, 519, 659
accessory, 519, 659
Obliquus externus, 445
inferior, 80, 308
internus, 448
superior, 71, 308
Occipital artery, 20, 125
sinus, 32
nerve, great, 22, 309
small, 23, 85
vein, 20, 126
Occipito-atloid ligaments, 317
axial ligaments, 318
frontalis, 18
Odontoid ligaments, 319
Oculi tendo, 44
tutamina, 46
Œsophageal arteries, 213
plexus, 207
Œsophagus, 204
structure of, 204
Olfactory bulb, 33
origin of, 760
nerves, 296
Omentum, gastro-colic, 484
hepatic, 484
splenic, 484
great, 484
cavity of, 481
lesser, 484
Openings for aorta in diaphragm, 500
cava, vena, in diaphragm, 500
œsophageal in diaphragm, 500
of heart, 226
upper, of thorax, 176
Operculum, 748
Ophthalmic artery, 74
ganglion, 77
nerve, 34
veins, 76
Optic nerve, 33, 73, 761
Ora serrata, 816
structure of, 822
Orbit, dissection of, 66
contents of, 68
fascia of, 67
nerves at back of, 39
periosteum of, 66
Organ of Corti, 840
Giraldès, 853
Rosenmüller, 593
Ossicula auditus, 832
ligaments of, 833
Ostium abdominale, 590
externum of uterus, 587

OST

Ostium (*continued*)
internum of uterus, 586
internum of Fallopian tube, 589
Otic ganglion, 282
branches of, 282
Otoliths, 843
Ovaries, 590
arteries of, 505
position of, 591
structure of, 591
Oviducts, 589
Ovula of Naboth, 587
Ovum, 592

PACCHIONIAN bodies, 26
Palate, glands of, 256
hard, 256
pillars of, 252
soft, 252
muscles of, 253
Palati circumflexus, 253
levator, 253
tensor, 253
velum pendulum, 251
Palatine artery, ascending, 16
descending, 155
Palmar arch, deep, 385
superficial, 375
profunda artery, 386
Palpebræ, 47
cartilages of, 49
conjunctiva of, 47
Pampiniform plexus, 854
Pancreas, 602
duct of, 602
functions of, 603
lesser, 602
lymphatics of, 603
position of, 477
relations of, 477
structure of, 603
vessels of, 603
Papilla lachrymalis, 46
Papillæ circumvallatæ, 273
filiformes, 274
fungiformes, 274
Parepididymis, 853
Parotid gland, 58
duct, 59
relations of, 58
structure of, 60
structures in, 59
Parotidis glandula socia, 59
Parovarium, 593
Pars ciliaris retinæ, 816
Pecquet, cistern of, 203

PES

Pectiniforme, septum, 578
Pedis dorsal artery 667
Pelvic fascia, 547
female viscera, 581
male viscera, 548
Pelvis, contents of female, 523
male, 520
side view of female, 581
male, 545
Penis, 576
bulb of, 579
artery of, 538
cervix of, 576
corona glandis, 577
corpus cavernosum, 577
artery of, 579
spongiosum, 579
crus, 577
dorsal artery of, 539
erector, 532
glans, 579
glandulæ Tysonii, 577
helicine arteries of, 579
lymphatics of, 580
nerves of, 539, 580
vessels of, 580
Perforating arteries of thigh, 656
Pericardium, 86
objects seen on opening, 189
structure of, 188
vestigial fold of, 189
Perilymph, 842
Perineum, female, 540
male, 525
boundaries of, 525
cutaneous nerves of, 528
raphé of, 526
tendinous centre of, 528
triangular ligament of, 534
parts between, 536
surgery of, 535
Perineal arteries, superficial, 530
fascia, deep, 529
superficial, 529
transverse muscle, 532
deep muscle, 533
Peritoneum, 478
course of, 479
lesser cavity of, 481
parts covered by, 481
partially by, 481
uncovered by, 481
Peroneal artery, 692
anterior, 693
nerve, 669, 682
Pes anserinus, 62



sacral, 562
spermatic, 516
splenic, 515
superficial cardiac, 2
supra-renal, 516
sympathetic of abdo
uterine, 566
vesical, 566
of veins, interlobular,
pampiniform, 854
pterygoid, 156
vaginal, 581
vesico-prostatic, 548
Pleura, 179
outlines of, 184
cavity of, 179
Plica semilunaris, 47
Pneumogastric nerve, 37,
in chest, 205
ganglia of, 286
auricular branch of, 2
Pomum Adami, 259
Pons hepatis, 594
Tarini, 758
Varolii, 744
Popliteal artery, 683, 692
nerve, external, 669
internal, 682
space, 680
vein, 683, 693
Popliteus, 694
Porus opticus, 807, 816
Portal fissure, 594
Portio dura, 61, 764
mollis, 765
Pouch of Douglas, 525
recto-vaginal, 525
vesical, 521
Pouches, laryngeal, 266
Poupart's ligament, 446, 6

PRO

Prosencephalon, 790
Prostate, 555
arteries of, 572
lobes of, 570
lymphatics of, 573
nerves of, 572
position of, 555
relations of, 555
sinus of, 570
structure of, 571
veins of, 572
Psoas fascia, 508
magnus, 509
parvus, 510
Pterygo-palatine artery, 155
maxillary ligament, 53
region, 149
Pudendal, inferior, nerve, 529, 530
Pudenda, 540
Pudic artery, internal, 537, 561, 679
nerve, 539, 564
Pulmonary artery, 213, 242
nerves of pneumogastric, 205
valves, 223
Pulmonis hilum, 185
Puncta lachrymalia, 46, 49
Pupil, 814
Pylorus, 616
Pyramid, anterior, of tympanum, 831
posterior, of tympanum, 830
cerebellum, 788

QUADRATUS lumborum, 611

RADIAL artery, 365
at back of wrist, 417
in palm of hand, 385
nerve, 367
vein, 360
Ranine artery, 124
vein, 124
Ranula, 161
Receptaculum chyli, 208, 502
Recess, lateral of fourth ventricle, 784
Recto-vaginal pouch, 525
Recto-vesical fascia, 548
pouch, 521
Rectum, 622
ampulla of, 550
arteries of, 551
course of, 521
digital examination of, 551
folds in, 624
relations of, 551
Rectus femoris, 650

SCH

Region, precordial, 186
Regions of abdomen, 440
Reil, fillet of, 741
island of, 749
Reissner, membrane of, 840
Restiform bodies, 739
Rete testis, 852
Retina, 816
arteria centralis of, 75
cones and rods of, 820
structure of, 817
Ribs, movements of, 322
Rima glottidis, 264
Ring, abdominal external, 447
internal, 456
femoral, 640
fibrous of heart, 228
Rivini, notch of, 831
Rivinus, ducts of, 122
Rolando, arciform fibres of, 739
fissure of, 749
funiculus of, 739
tubercle of, 739
Rosenmüller, organ of, 593
Rugæ of vagina, 583
Ruyschiana tunica, 812

SAC, lachrymal, 49, 294
of omentum, 484
Saccule of vestibule, 843
Sacculus laryngis, 266
Sacro-lumbalis, 301
iliac articulation, 709
Santorini, cartilages of, 261
fissures of, 827
Saphena, external vein, 688
internal vein, 630
Saphenous, external nerve, 689
opening, 633
posterior vein, 688
Scala media, 838
tympani, 838
vestibuli, 838
Scalp, dissection of, 17
arteries, of, 20
lymphatics of, 24
nerves of, 20
veins of, 20
Scaleni muscles, 129
Scapulæ artery, dorsalis, 337
Scapular artery, posterior, 140, 396
supra, 139, 405
vein, posterior, 140
Scarpa, liquor of, 844
triangle of, 643
Schlemm, canal of, 810

SCH

Schneiderian membrane, 294
Sciatic artery, 561, 678
nerve, great, 677, 681, 687
small, 565, 678
Sclerotic coat of eye, 806
structure of, 807
Scrotum, 846
dartos of, 847
lymphatics of, 848
septum of, 847
vessels of, 848
Semicircular canals, 837
Semilunar cartilages of knee, 718
ganglia, 498
valves, 223
Semimembranosus, 685
Semispinalis colli, 837
dorsi, 836
Semitendinosus, 685
Septum, artery of nasal, 56
auricularum, 219
lucidum, 772
pectiniforme, 578
scroti, 847
tongue, 277
transversum, 844
ventriculorum, 221
Serratus magnus, 343, 397
posticus inferior, 297
superior, 298
Seventh cranial nerve, 36, 287, 764
Sheath, axillary vessels, 329
femoral vessels, 638
rectus muscle, 452
Sheath for extensor tendons of hand, 409
Sheaths for flexor tendons of hand, 382
Shoulder, cutaneous nerves of, 398
joint, 425
movements of, 428
muscles in relation with, 428
synovial membrane of, 427
Sigmoid flexure of colon, 474
Sinus circularis iridis, 810
coronary, 227
Morgagni, 249
pocularis, 571
prostaticus, 570
venous, cavernous, 30, 37
circular, 31
coronary of heart, 220
dura mater, 27
lateral, 30
longitudinal, inferior, 30
superior, 27
occipital, 32
petrosal, inferior, 30

SPL

Sinus (*continued*)
petrosal, superior, 30
straight, 30
transverse, 32
venosus, 219
Sinuses, great, of aorta, 195
Valsalva, 196
Sixth cranial nerve, 35, 764
Small intestines, 618
Socia parotidis, glandula, 59
Sœmmering, foramen of, 816
Solar plexus, 515
Soleus, 690
Space, interpleural, 181
popliteal, 679
subdural, 25
Spaces of Fontana, 810
subarachnoid, 730
Spermatic cord, 459, 853
arteries of, 459, 505
lymphatics of, 459
nerves of, 459
veins of, 459
Sphenoidal fissure, structures passing through, 38
Spheno-palatine artery, 155
ganglion, 281
branches of, 282
Spigelii lobulus, 595
Spinæ rotatores, 305
Spinal-accessory nerve, 37, 91, 168, 394, 765
Spinal cord, 797
arteries of, 803
central canal of, 800
fissures of, 798
functions of, 803
membranes of, 795
structure of, 799
Spinal lateral artery from vertebral, 138
Spinal nerves, origin of, 800
Spine, ligaments of, 314
movements of, 317
Splanchnic nerves, 208
great, 208
lesser, 208
smallest, 209
Spleen, 603
artery of, 489, 605
functions of, 606
lymphatics of, 606
Malpighian corpuscles of, 605
nerves of, 606
omenta of, 604
pulp of, 604
relations of, 476

SPL

Spleen (*continued*)
tunics of, 606
veins of, 489, 606
Splenium of corpus collosum, 769
Splenius capitis, 300
colli, 300
Stapedius, 834
Stapes, 833
Steno's duct, 59
Sterno-mastoid muscle, 89
artery, middle, 115
superior, 125
parts beneath, 98
Stilling, canal of, 824
Stomach, 614
relations of, 476
structure of, 617
Straight sinus, 30
Striæ acusticæ, 785
laterales, 768
longitudinales, 785
medullares, 785
Stylo-glossus, 121, 161
Stylo-hyoid ligament, 162
muscle, 111
Stylo-maxillary ligament, 70, 118
Subarachnoid fluid of brain, 730
of spinal cord, 795
spaces of, 730
Subclavian artery, left, 133, 198
right, 131
ligature of, 134
vein, 142
Subdural space, 25
Sublingual artery, 124
gland, 122
Suboccipital nerve, 23, 290
triangle, 308
Submaxillary ganglion, 123
gland, 111
triangle, 109
Subperitoneal fat, 458
Subscapular artery, 336
nerves, 341
Subscapularis, 342, 403
Sulci of brain, 747
Sulcus spiralis, 838
Supercillii corrugator, 349
Supra-clavicular nerves, 85
triangle, 92
Supra-renal capsules, 613
arteries of, 614
nerves of, 614
relations of, 478
structure of, 614
Supra-orbital artery, 20, 75

THA

Supra-orbital nerve, 21, 64, 69
Supra-scapular artery, 139, 396
nerve, 144, 396
vein, 140
Supra-trochlear nerve, 21, 64, 67
Supra-spinales, 306
Supra-spinous ligament, 315
Sylvius, aqueduct of, 783
fissure of, 749
Sympathetic nerves:
cervical, 169, 173, 174
cranial, 171
lumbar, 514
nervi molles, 57
pelvic, 565
thoracic, 207
Symphysis pubis, 711
sacro-iliac, 709

TÆNIA, fourth ventricle, 785
hippocampi, 775
Tænia semicircularis, 777
Tarsal cartilages and ligaments, 48
Tarsi, tensor, 50
Tarsus, synovial membranes of, 727
Tectorial membrane, 842
Tegmentum, 759
Temporal artery, 146
superficial, 20
deep, 147, 154
fascia, 148
muscle, 149
nerves, deep, 157
veins, 146
superficial, 20
Temporo-fascial nerve, 62
Temporo-maxillary ligaments, 324
Tendo-Achillis, 691
palpebrarum, 45
Tenon, capsule of, 67, 79, 805
Tenth cranial nerve, 37, 165, 205, 765
Tentorium cerebelli, 26
Teres major, 342, 403
minor, 402
Testes cerebri, 783
muliebres, 590
Testis, 848
arteries of, 848
coverings of, 848
descent of, 854
gubernaculum, 854
lymphatics of, 854
mediastinum, 851
nerves of, 854
structure of, 851
Thalamencephalon, 790

THA

Thalami optici. 778
Thebesii foramina, 221
valve, 211, 227
veins of, 227
Theca vertebralis, 795
Third cranial nerve, 34, 77, 762
Thoracic aorta, 193
Thoracic artery, alar, 336
long, 337
superior, 330
nerves, anterior, 330
posterior, 145, 343
Thoracic duct, 203, 502
Thoracico-acromialis artery, 330
Thorax, base of, 176
dissection of, 174
osseous measurements, 174
upper opening of, 176
Thymus gland, 176
Thyro-arytenoideus, 268
Thyro-epiglottideus, 268
Thyro hyoid, 97
Thyroid artery, superior, 113
inferior, 139
vein, 115
cartilage, 259
gland, 103
arteries of, 104
lymphatics of, 105
nerves of, 105
structure of, 105
veins of, 104
Tibial artery, anterior, 665
posterior, 696
nerve, anterior, 669
posterior, 696
Tibialis anticus, 663
posticus, 695
Tomentum cerebri, 731
Tongue, 272
arteries of, 277
foramen, cæcum of, 272
glands beneath, 275
mucous membrane of, 272
muscular fibres of, 275
nerves of 277
papillæ of, 273
raphé of, 272
septum of, 277
Tonsillar artery, 116
Tonsils, 254
Torcular Herophili, 32
Trachea, 235
cartilages, 238
glands of, 237
mucous membrane of, 237

ULN

Trachea (*continued*)
muscular fibres of, 237
relations of, 235
Tragus, 826
Transversalis fascia, 455
abdominis, 450
Trapezium of pons Varolii, 744
Trapezius, 390
Triangle, carotid inferior, 94
superior, 94
digastric, 109
elbow, 361
Hesselbach's, 463
infraclavicular, 328
neck, anterior, 93
posterior, 97
occipital, 91
Scarpa's, 643
subclavian, 92
submaxillary, 109
suboccipital, 308
supraclavicular, 92
Triangular ligament of abdomen, 447
perineum, 535
Triangularis sterni, 177
Triceps extensor cubiti, 358, 406
femoris, 650
Tricuspid valves, 221
Trifacial nerve, 34, 762
Trigonum vesicæ, 569
Trochlea of orbit, 71
Trochlearis nerve, 34, 762
Tube, Eustachian, 255
Fallopian, 589
Tuber annulare, 744
cinereum, 757
Tubercula quadrigemina, 782
Tuberculum acusticum, 785
Tubes, bronchial, 236
Tunica albuginea, 850
Ruyschiana, 812
vaginalis, 849
vasculosa, 851
Tutamina oculi, 46
Twelfth cranial nerve, 37, 121, 766
Tympani laxator, 833
membrana, 831
tensor, 833
Tympanic nerve of facial, 288
glosso pharyngeal, 284
Tympanum, 829
arteries of, 834
Tyson's glands, 577

ULNAR artery, 368
nerve, 357, 368

ULN

Ulnar nerve (*continued*)
in palm, 377
deep, in hand, 387
veins, 350
Umbilical hernia, 467
Umbilicus, 452
Urachus, 468
Ureter, 553
course of, 553
muscles of, 569
orifices of, 569
Urethra, female, 543
male, 573
bulbous portion of, 575
fossa navicularis of, 575
lacunæ of, 575
membranous part of, 556
spongy part of, 575
structure of, 575
Uterus, 583
arbor vitæ of, 587
arteries of, 584
cervix of, 587
glands of, 589
lymphatics of, 585
masculinus, 571
nerves of, 585
os of, 586
position of 523
round ligament of, 525
structure of, 585
veins of, 585
Utricle of vestibule, 843
Utriculus, 571
Uvea, 814
Uvula of bladder, 567
cerebellum, 788
palate, 252
Uvulæ azygos, 254

VAGINA, 544, 582
arteries of, 585
bulb of, 544
structure of, 582
venous plexus of, 581
Vagus nerve, 37, 165, 205, 752
Vallecula of cerebellum, 788
Vallum of tongue, 273
Valsalva, sinuses of, 195
Valve, coronary, 220
Eustachian, 220
Hasner, 294
ileo-cæcal, 624
Kerkring, 619
Thebesius, 221, 227
Vieussens, 778

VEI

Valves, cardiac, position of, 180
aortic, 226
mitral, 225
pulmonary, 223
semilunar, 223
tricuspid, 222
Valvulæ conniventes, 619
Varolii pons, 744
Vas aberrans, 883
deferens, 459, 554, 853
spirale, 844
Vastus externus, 650
internus, 650
Veins:
angular, 57
auricular, 127
axillary, 339
azygos, left upper, 202
major, 202
minor, 202
basilic, 347
median, 346
basi-vertebral, 793
brachial, 354
brachio-cephalic, 191
bronchial, 243
capsular, 598
cardiac anterior, 226
great, 226
posterior, 226
cava, inferior, 507
superior, 192
cephalic, 330, 347
median, 346
circumflex iliac, 438
coronary, of heart, 226
of stomach, 490
dorsal of penis, 539
dorsi-spinal, 793
elbow in front of, 346
epigastric, 457
facial, 57, 112
femoral, 644
frontal, 20
of Galen, 30, 778
gastric, 488
hæmorrhoidal, 494, 549
hepatic, 488, 598
iliac, common, 512
external, 513
internal, 549
inferior cava, 503
innominate, 191
intercostal, superior, 202
interlobular, 598
intralobular, 598

VEI

Veins (*continued*)
jugular, anterior, 84
external, 83
internal, 102
posterior external, 84
kidney of, 505
lumbar, 506
mammary internal, 178
maxillary internal, 152
median, 360
deep, 360
medulli-spinal, 794
meningeal, 32
meningo-rachidian, 794
mesenteric inferior, 494
superior, 493
oblique, of Marshall, 221
occipital, 20, 125
ophthalmic, 77
pharyngeal, 127
phrenic, 503
popliteal, 683, 692
portal, 598
profunda femoris, 654
pudic, external, 629
internal, 561
pulmonary, 243
radial, 360
ranine, 124
rectum, 551
renal, 503, 612
sacra media, 506
salvatella, 360
saphenous external, 689
internal, 629
scalp of, 20
scapular posterior, 140
supra, 139
spermatic, 507, 854
spinal longitudinal anterior, 793
posterior, 793
posterior external, 793
splenic, 489, 606
subclavian, 142
sublobular, 597
supra-orbital, 20
renal, 503
scapular, 139
temporal, 146
superficial, 20
Thebesii, 227
thyroid superior, 115
tibial posterior, 697
tympanum, of, 834
ulnar anterior, 360
posterior, 360

ZYG

Veins (*continued*)
umbilical, 232, 234
uterine, 584
vaginal, 584
of liver, 598
vertebral, 138
Velum interpositum, 777
medullary anterior, 783
posterior, 789
pendulum palati, 251
Venæ vorticosæ, 811
Ventral hernia, 467
Ventricle of brain, fifth, 772
fourth, 783
lateral, 769
third, 779
Ventricle of heart, left, 225
right, 221
muscular fibres of, 230
Ventricle of larynx, 266
Vermiform process, inferior, 787
superior, 788
Vertebral aponeurosis, 298, 393
artery, 137, 733
Vincula tendinum, 384
Verumontanum, 571
Vesico-prostatic plexus, 548
Vesicula seminalis, 554
structure of, 573
vessels of, 573
Vestibule of ear, 835
vagina, 544
Vestigial fold of pericardium, 189
Vibrissæ, 295
Vidian artery, 155
nerve, 282
Vieussens, valve of, 783
Vitreous body, 823
Vocal cords, inferior or true, 265
superior or false, 265

WHARTON'S duct, 160
Willis, circle of, 734
cords of, 28
Winslow, foramen of, 482
posterior ligament of, 686
Wirsung, canal of, 496
Wrisberg, cartilages of, 262
ganglion of, 215
nerve of, 344
Wrist joint, synovial membranes of, 437
triangular fibro-cartilage of, 431

ZINN, ligament of, 78
zone of, 824
Zygomaticus major, 44
minor, 44

ALOGUE NO. 1. NOVEMBER, 1884.

CATALOGUE

OF

MEDICAL, DENTAL, PHARMACEUTICAL

AND

SCIENTIFIC PUBLICATIONS,

PUBLISHED BY

P. BLAKISTON, SON & CO.,

(SUCCESSORS TO LINDSAY & BLAKISTON)

1012 WALNUT STREET,

PHILADELPHIA.

These publications may be had through Booksellers in all the principal cities of the United tes and Canada, or any book will be sent, postpaid, by the publisher, upon receipt of price, or ll be forwarded by express C. O. D. upon receiving a remittance of 25 per cent. of the amount dered to cover express charges.

NOW READY, THE SEVENTH REVISED EDITION.

MEIGS AND PEPPER, ON CHILDREN.

THE MOST THOROUGH, COMPLETE AND PRACTICAL WORK ON THE SUBJECT NOW BEFORE THE PROFESSION.

A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. By J. FORSYTH MEIGS, M.D., one of the Physicians to the Pennsylvania Hospital, Consulting Physician to the Children's Hospital, etc., and WILLIAM PEPPER, M.D., Professor of Clinical Medicine, University of Pennsylvania, Provost and ex-officio President of the Faculty, Physician to the Philadelphia Hospital, Fellow of the College of Physicians, etc., etc. The Seventh Revised and Improved Edition. In one volume of over 1100 royal octavo pages.

Price, handsomely bound in Cloth, $6.00; Leather $7.00.

The rapid sale of six large editions of Drs. Meigs and Pepper's work on Children, and the demand for the new edition now ready, is sufficient evidence of its great popularity. The large practice, of many years' standing, of the authors, imparts to it a value unequaled, probably, by any other on the subject now before the profession.

The entire work has been now again subjected to an entire and thorough revision, some articles have been rewritten, many additions made, and great care observed by the authors, that it should be most effectually brought up to the light, pathological and therapeutical, of the present day.

The publishers have very many favorable notices of the previous editions, received from numerous sources, foreign and domestic. They append a few from leading journals, which will give a general idea of the value placed upon it, both as a *Text-Book* for the Student and a work of reference for the *General Practitioner*.

"It is the most complete work upon the subject in our language; it contains at once the results of personal and the experience of others; its quotations from the most recent authorities, both at home and abroad, are ample, and we think the authors deserve congratulations for having produced a book unequaled for the use of the student, and indispensable as a work of reference for the practitioner."—*American Medical Journal.*

"But as a scientific guide in the diagnosis and treatment of the diseases of children, we do not hesitate to say that we have seldom met with a text-book so complete, so just, and so readable, as the one before us, which in its new form cannot fail to make friends wherever it shall go, and wherever great erudition, practical tact, and facile and agreeable diction are appreciated."—*American Journal of Obstetrics.*

"It is only three years since we had the pleasure of recommending the Fifth Edition of this excellent work. With the recent additions it may safely be pronounced one of the best and most comprehensive works on diseases of children of which the American Practitioner can avail himself, for study or reference."—*N. Y. Med. Journal.*

"It is not necessary to say much, in the way of criticism, of a work so well known. But it is clinical. Like so many other good American medical books, it marvelously combines a *résumé* of all the best European literature and practice, with evidence throughout of good personal judgment, knowledge, and experience. The book also abounds in exposition of American experience and observation in all that relates to the diseases of children. We are glad to add it to our library. There are few diseases of children which it does not treat of fully and wisely, in the light of the latest physiological, pathological, and therapeutical science."—*London Lancet.*

P. BLAKISTON, SON & CO., PUBLISHERS,

Successors to LINDSAY & BLAKISTON,

1012 WALNUT STREET, PHILADELPHIA.

Mr. Presley Blakiston having on January 1st, 1882, purchased all the interest of the late firm of Lindsay & Blakiston, continues the publication and sale of MEDICAL AND SCIENTIFIC BOOKS at No. 1012 Walnut Street, Philadelphia, having associated with him his son, Kenneth M. Blakiston, and Frank W. Robinson, under the firm-name of

P. BLAKISTON, SON & CO.

MEDICAL, DENTAL, SCIENTIFIC

AND

PHARMACEUTICAL BOOKS

PUBLISHED BY

P. BLAKISTON, SON & CO. PHILADELPHIA.

☞ Any book in this catalogue can be had from or through booksellers in the principal cities in the United States, or will be forwarded free, by mail or express, upon receipt of the price by the publisher.

AMERICAN HEALTH PRIMERS.

Edited by W. W. KEEN, M.D. Complete in 12 volumes, handsomely bound. Price, in cloth binding, 50 cents; paper covers, 30 cents.

I. **Hearing and How to Keep It.** With illustrations. By CHAS H. BURNETT, M.D.

II. **Long Life, and How to Reach It.** By J. G. RICHARDSON, M.D.

III. **The Summer and Its Diseases.** By JAS. C. WILSON, M.D.

IV. **Eyesight, and How to Care for It.** With illustrations. By GEORGE C. HARLAN, M.D.

V. **The Throat and the Voice.** With illustrations. By J. SOLIS COHEN, M.D.

VI. **The Winter and Its Dangers.** By HAMILTON OSGOOD, M.D.

VII. **The Mouth and the Teeth.** With illustrations. By J. W. WHITE, M.D., D.D.S.

VIII. **Brain Work and Overwork.** By H. C. WOOD, JR., M.D.

IX. **Our Homes.** With illustrations. By HENRY HARTSHORNE, M.D.

X. **The Skin in Health and Disease.** By L. D. BULKLEY, M.D.

XI. **Sea Air and Sea Bathing.** By JOHN H. PACKARD, M.D.

XII. **School and Industrial Hygiene.** By D. L. LINCOLN, M.D.

LIBRARY EDITION, IN 4 VOLS., CLOTH, EACH $1.25.

"In their practical teachings, learning, and sound sense, these volumes are worthy of all the compliments they have received. They teach what every man and woman should know, and yet what nine-tenths of the intelligent class are ignorant of, or at best, have but a smattering knowledge of."—*Chicago Inter-Ocean.*

"These handbooks of practical suggestion deserve hearty commendation. They are prepared by men whose professional competence is beyond question, and for the most part, by those who have made the subject treated the specific study of their lives."—*New York Sun.*

AMERICAN PSYCHOLOGICAL JOURNAL.

Issued by the National Association for the Protection of the Insane and Prevention of Insanity. Edited by JOSEPH PARRISH, M.D.

Single numbers 50 cents; per annum, $2.00

ACTON, THE REPRODUCTIVE ORGANS.

The Functions and Disorders of the Reproductive Organs in Childhood, Youth, Adult Age and Advanced Life, considered in their Physiological, Social and Moral Relations. By WILLIAM ACTON, M.D., M.R.C.S. Sixth Edition. 8vo. Cloth, $2.00

"In the work now before us, all essential detail upon its subject matter is clearly and scientifically given. We recommend it accordingly, as meeting a necessary requisition of the day, refusing to join in that opinion which regards the consideration of the topics in question as beyond the duties of the medical practitioner."—*The London Lancet.*

"On the subjects of Impotence and Spermatorrhœa, those bugbears of so many weak and foolish persons, and sources of inexhaustible wealth to the quack fraternity, Mr. Acton discourses with good sense, and indignantly exposes the nefarious tricks of the scoundrels who, on the pretence of curing a disease which often exists only in imagination, extract enormous sums from their unwary victims. He seems to regard the spermatorrhœa phobia, as we may term it, to be a species of monomania; but he judiciously advises that to a patient laboring under this form of mental malady, the tone adopted should be one of sympathy and attention; and that by the employment of appropriate moral and therapeutical means, a healthy and hopeful tone of mind be restored."—*The Medical Times.*

AGNEW, ON THE PERINEUM AND FISTULA.

Lacerations of the Female Perineum and Vesico-vaginal Fistula. Their History and Treatment. With many Illustrations. By D. HAYES AGNEW, M.D., Professor of Surgery, University of Pennsylvania. 8vo. Cloth, Price $1.25

So many applications having been made for these papers, as originally issued, the author has thought best, after a thorough revision, to place them before the profession in book form.

ALLBUTT. VISCERAL NEUROSES.

The Gulstonian Lectures for March, 1884, on Neuralgia of the Stomach and Allied Disorders. By T. CLIFFORD ALLBUTT, M.D., F.R.S., Consulting Physician to the Leeds General Infirmary and the Leeds Hospital for Women and Children. Octavo. Cloth, $1.50

ARMATAGE. VETERINARY REMEMBRANCER.

The Veterinarian's Pocket Remembrancer. Containing concise directions for the Treatment of Urgent or Rare Cases, embracing Semeiology, Diagnosis, Prognosis, Surgery, Therapeutics, Detection of Poisons, Hygiene, etc. New Revised Edition. 18mo. Cloth, $1.25

ALLAN, FEVER NURSING.

Notes on Fever Nursing. Addressed to nurses in hospital and private life. By JAMES W. ALLAN, M.D. 12mo. Illustrated. Price .75

ALLINGHAM, DISEASES OF THE RECTUM. Illustrated.

Fistula, Hæmorrhoids, Painful Ulcer, Stricture, Prolapsus, and other Diseases of the Rectum, their Diagnosis and Treatment. By WILLIAM ALLINGHAM, F.R.C.S. Fourth Edition, enlarged. Price, Paper covers, .75; Cloth, $1.25

London Edition, thick paper and larger type, $2.00.

"He is in charge of the only hospital in the world (St. Marks) devoted exclusively to diseases of the rectum, and he is recognized, both in this country and in Europe, as the highest authority upon diseases of this class."—*Louisville Medical Herald.*

"This book has always been a great favorite, and deservedly so. It is practical in tone and character, magisterial in its teaching, and valuable in showing operative results. It is by an author who, as an authority, has no superior."—*Gaillard's Medical Journal.*

"No book on this special subject can at all approach Mr. Allingham's in precision, clearness and practical good sense."—*London Medical Times and Gazette.*

"It is, as indeed the verdict of the profession has already pronounced it, one of the very best works on Diseases of the Rectum."—*American Journal of Medical Science.*

ALTHAUS, MEDICAL ELECTRICITY.

A Treatise on Medical Electricity, Theoretical and Practical, and its Use in the Treatment of Paralysis, Neuralgia, and other Diseases. By JULIUS ALTHAUS, M.D. Third Edition, Enlarged. 246 Illustrations. 8vo. Price $6.00

In revising this new edition the author has carefully brought each section up with the latest knowledge of the subject.

ANSTIE, STIMULANTS AND NARCOTICS.

With special researches on the Action of Alcohol, Ether and Chloroform on the Vital Organism. By FRANCIS E. ANSTIE, M.D. 8vo. Price $3.00

"He is an original worker and independent thinker. His opinions and conclusions are valuable, and cannot be neglected."—*American Medical Journal.*

ATTHILL, DISEASES OF WOMEN.

Clinical Lectures on Diseases Peculiar to Women. By LOMBE ATTHILL, M.D. 5th edition, revised and enlarged, with numerous illustrations. 12mo. Cloth. Price $1.25

"It is the concentrated essence of the knowledge of one who has become wise by reason of long and well-digested experience in the subjects treated."—*American Journal of Medical Science.*

"The work is one of great value to the general practitioner."—*American Journal of Obstetrics.*

AITKEN'S PRACTICE OF MEDICINE. New Edition.

The Science and Practice of Medicine. By WILLIAM AITKEN, M.D., F.R.S. London, Professor of Pathology in the Army Medical School, etc. Seventh Edition. To a large extent rewritten; enlarged, remodeled and carefully revised throughout. In Two Volumes. 196 Engravings on Wood, and a Map showing the Geographical Distribution of Diseases, and Copious Index. Octavo. Cloth, $12.00; Leather, $14.00

BALFOUR, ON THE HEART AND AORTA.

Clinical Lectures on Diseases of the Heart and Aorta. By G. W. BALFOUR, M.D. Illustrated. 2d Edition. Price $5.00

"The whole work reflects much credit on its author, and firmly establishes his reputation as an authority on the important diseases of which he treats."—*London Practitioner.*

BARTH AND ROGER, AUSCULTATION AND PERCUSSION.

A Manual for the Student. By M. BARTH and M. HENRI ROGER. Translated from the 6th French Edition. 12mo. Price $1.00

BIBLE HYGIENE;

Or, Health Hints. By a Physician. This book has been written, first, to impart in a popular and condensed form the elements of Hygiene; second, to show how varied and important are the Health Hints contained in the Bible, and third, to prove that the secondary tendency of modern Philosophy runs in a parallel direction with the primary light of the Bible. 12mo. Paper, .50; Cloth, $1.00

"The scientific treatment of the subject is quite abreast of the present day, and is so clear and free from unnecessary technicalities that readers of all classes may peruse it with satisfaction and advantage."—*Edinburgh Medical Journal.*

BIDDLE, MATERIA MEDICA. Ninth Edition.

(*Contains all the changes in the Sixth Revision of the New Pharmacopœia.*)

Materia Medica. For the Use of Students and Physicians. By the late PROF. JOHN B. BIDDLE, M.D., Professor of Materia Medica in Jefferson Medical College, Philadelphia. The Ninth Edition, thoroughly revised, and in many parts rewritten, by his son, CLEMENT BIDDLE, M.D., Assistant Surgeon, U. S. Navy, assisted by HENRY MORRIS, M.D. Containing all the additions and changes made in the last revision of the United States Pharmacopœia. The Botanical portions have been curtailed or left out, and the other sections, on the Physiological action of Drugs, greatly enlarged. Octavo. Cloth, $4.00; Leather, $4.75

"The additions are valuable, and we must congratulate the author upon having improved what was already so useful a work, both to the student and physician."—*Phila. Medical and Surgical Reporter.*

"It has been the design of the author to present in his work a text-book for the student. It is brief, and yet sufficiently comprehensive. His style is clear and yet succinct. He covers the ground—covers it well, and cumbers his work with nothing superfluous."—*Atlanta Medical and Surgical Journal.*

"One thing that particularly recommends this work to the student is, that the book is not so large as to discourage and cause him to feel that it is impossible for him to get over it and so much else in the short time before him."—*St. Louis Medical and Surgical Journal.*

"It contains, in a condensed form, all that is valuable in materia medica, and furnishes the medical student with a complete manual on this subject."—*Canada Lancet.*

BREWING, DISTILLING, ETC.

The Brewer, Distiller and Wine Manufacturer; a Handbook for all interested in the Manufacture and Trade of Alcohol and its Compounds. Edited by John Gardner, Fellow of the Chemical Society of London. Illustrated. Cloth, $1.75

SYNOPSIS OF CONTENTS.—Alcohol, its Preparations, etc.; Alcoholometry; Brewing and Beers; Varieties of Malt Liquors; Malt; Raw Grain; Sugar; Hops; Arrangement of a Brewery; Different Processes; Chemical Changes during Washing, Boiling, Cooling, Fermentation, etc., etc.; Storing and Clarifying, Porters, Ales; Analysis of Beers, Ciders, Perry, Mum; Liquors and Cordials, giving over 80 preparations. Other sources of Spirituous Liquors; Distillation of Alcoholic Liquors, including Rums, Brandies, Whiskies, Gins, etc.; Wine and Wine Making; Tests for Adulterations; Remarks on the Cultivation of Grapes, etc.; Imitation of Wines.

BLOXAM. CHEMISTRY, Inorganic and Organic. Fifth Edition.

With Experiments. By CHARLES L. BLOXAM, Professor of Chemistry in King's College, London, and in the Department for Artillery Studies, Woolwich. Fifth edition. With nearly 300 Engravings. Cloth, $3.75; Leather, $4.75

A most complete Text-Book for Schools and Colleges.

"Professor Bloxam has given us a most excellent and useful practical treatise. His 666 pages (now 700) are crowded with facts and experiments, nearly all well chosen, and many quite new, even to scientific men. It is astonishing how much information he often conveys in a few paragraphs. We might quote fifty instances of this."—*Chemical News.*

BLOXAM. LABORATORY TEACHING. Fourth Edition.

Progressive Exercises in Practical Chemistry. By CHARLES L. BLOXAM, Professor of Chemistry in King's College, London, etc. Fourth edition. With 89 engravings. 12mo. Price $1.75

This work is intended for use in the Chemical Laboratory, by those who are commencing the study of Practical Chemistry. It contains:—

1. A series of simple Tables for the analysis of unknown substances of all kinds. 2. A brief description of all the practically important single substances likely to be met with in ordinary analysis. 3. Simple directions and illustrations relating to Chemical Manipulation. 4. A system of Tables for the detection of unknown substances with the aid of the Blowpipe. 5. Short instructions upon the purchase and preparation of the tests intended for those who have not access to a Laboratory.

"A great amount of valuable practical information is here condensed into a book of 260 pages, such as only a practical teacher could prepare."—*New England Journal of Education.*

BRUEN. PHYSICAL DIAGNOSIS. Second Edition.

A Pocket-Book of Physical Diagnosis, for Physicians and Students. By EDWARD T. BRUEN, M.D., Demonstrator of Clinical Medicine, University of Penn'a. Illustrated by Original Wood Engravings. 12mo. 2d Ed. Cloth, $1.50

"We consider the description of the manner and rules governing the art of percussion well given. The subject is always a difficult one for beginners, and requires to be well handled in order to be properly understood."—*American Journal of Medical Sciences.*

"The volume before us is intended as a guide to the student and practitioner in making a diagnosis of Diseases of the Lungs and Heart, and as such is an excellent book, full of practical hints and valuable points."—*Philadelphia Medical Times.*

BENNETT. NUTRITION IN HEALTH AND DISEASE.

A Contribution to Hygiene and Clinical Medicine. By J. HENRY BENNETT, M.D. Third Edition, Revised and Enlarged. Cloth. Price $2.50

BEALE ON SLIGHT AILMENTS. New Edition. Just Ready.

Slight Ailments, Their Nature and Treatment. By LIONEL S. BEALE, M.D., F.R.S., Professor of Practice, King's Medical College, London. Second Edition. Enlarged and Illustrated. Price, Cloth, $1.25; Paper covers, .75 cents. Fine Edition, Heavy Paper. Extra Cloth, Price $1.75

OUTLINE OF CONTENTS.

Introductory. The Tongue in Health and Slight Ailments. Appetite. Nausea. Thirst. Hunger. Indigestion, its Nature and Treatment. Constipation, its Treatment. Diarrhœa. Vertigo. Giddiness. Biliousness. Sick Headache. Neuralgia. Rheumatism. The Feverish and Inflammatory State. Of the Actual Changes in Fever and Inflammation. Common Forms of Slight Inflammation, etc., etc.

"We venture to say that among the numerous medical publications issued during 1880, there has been none which will prove more useful to the young general practitioner, for whom it is really intended, than this volume, while the time of the older physician might be much more unprofitably spent."—*American Journal of Medical Science*

BY SAME AUTHOR.

ON LIFE AND VITAL ACTION IN HEALTH AND DISEASE.

12mo. Price $2.00

THE USE OF THE MICROSCOPE IN PRACTICAL MEDICINE.

For Students and Practitioners, with full directions for examining the various secretions, etc., in the Microscope. Fourth Edition. 500 Illustrations. Much enlarged. 8vo. Price $7.50

"We have before us Prof. BEALE'S work, *The Microscope in Medicine*, a book which it gives us pleasure to recommend to every student of microscopy, whether he be a physician or naturalist."—*Journal of the Franklin Institute, Philadelphia.*

"As a microscopical observer, and a histological manipulator, his (Dr. BEALE) skill and eminence are generally conceded."—*Popular Science Monthly.*

HOW TO WORK WITH THE MICROSCOPE.

A Complete Manual of Microscopical Manipulation, containing a full description of many new processes of investigation, with directions for examining objects under the highest powers, and for taking photographs of microscopic objects. Fifth Edition. Containing over 400 Illustrations, many of them colored. Octavo. Price $7.50

"The Encyclopædic character of this last edition of Dr. BEALE'S well known work on the *Microscope* renders it impossible to present an abstract of its contents; suffice it to say, that anything in his department upon which the physican can desire such information will be found here, and much more in addition. It is, moreover, a storehouse of facts, most valuable to the physician, and is indispensable to every one who uses the microscope."—*American Journal of Medical Science.*

BIOPLASM.

A Contribution to the Physiology of Life, or an Introduction to the Study of Physiology and Medicine, for Students. With numerous Illustrations. Price $2.25

PROTOPLASM; or MATTER AND LIFE.

Third Edition, very much enlarged. Nearly 350 pages. Sixteen Colored Plates. Part I. DISSENTIENT. Part II. DEMONSTRATIVE. Part III. SUGGESTIVE. One volume.

LIFE THEORIES; Their Influence upon Religious Thought.

Six Colored Plates. Price $2.00

ONE HUNDRED URINARY DEPOSITS,

On eight sheets, for the Hospital, Laboratory, or Surgery. New Edition. 4to, paper. Price $2.00

BERNAY, CHEMISTRY.

Notes for Students in Chemistry. Compiled from Fowne's and other manuals. By ALBERT J. BERNAY, PH.D. Sixth Edition. 12mo. Price $1.25

BENTLEY'S STUDENTS' BOTANY.

The Students' Guide to Structural and Physiological Botany. By PROFESSOR ROBERT BENTLEY. Illustrated by nearly 500 Wood Engravings.

In Preparation.

BEASLEY. THE BOOK OF PRESCRIPTIONS.

Containing over 3100 Prescriptions, collected from the Practice of the most Eminent Physicians and Surgeons—English, French and American; a Compendious History of the Materia Medica, Lists of the Doses of all Officinal and Established Preparations, and an Index of Diseases and their Remedies. By HENRY BEASLEY. Sixth Edition, Revised and Enlarged. Price $2.25

BY SAME AUTHOR.

THE DRUGGIST'S GENERAL RECEIPT-BOOK.

Comprising a copious Veterinary Formulary; numerous Recipes in Patent and Proprietary Medicines, Druggists' Nostrums, etc.; Perfumery and Cosmetics; Beverages, Dietetic Articles and Condiments; Trade Chemicals, Scientific Processes, and an Appendix of Useful Tables. Eighth Edition. Price $2.25

THE POCKET FORMULARY and Synopsis of the British and Foreign Pharmacopœias.

Comprising Standard and Approved Formulæ for the Preparations and Compounds Employed in Medical Practice. Tenth Edition. 511 pp. 18mo. Price $2.25

BENTLEY AND TRIMEN'S MEDICINAL PLANTS.

A New Illustrated Work, containing full botanical descriptions, with an account of the properties and uses of the principal plants employed in medicine, especial attention being paid to those which are officinal in the British and United States Pharmacopœias. The plants which supply food and substances required by the sick and convalescent are also included. By R. BENTLEY, F.R.S., Professor of Botany, King's College, London, and H. TRIMEN, M.B., F.H.S., Department of Botany, British Museum. Each species illustrated by a colored plate drawn from nature. In Forty-two parts. Eight colored plates in each part.

Price $2 each, or handsomely bound in 4 volumes, Half Morocco, $90.00

"It would be impossible to enumerate all the new plants that are here delineated. The result is a work which, from all points of view, is a credit to the scientific literature of the day."—*London Lancet.*

"It is an indispensable work of reference to every one interested in pharmaceutical Botany."—*London Pharmaceutical Journal.*

"This work may be recommended as a most useful one to druggists, and all who desire to be familiar with the Botany of Medicinal Plants."—*Druggists' Circular.*

"The work when complete (it is now complete) will be the most valuable compend of Medical Botany ever published."—*Boston Journal of Chemistry.*

BRUBAKER, PHYSIOLOGY.

A Compend of Physiology specially adapted for the use of Students and Physicians. "No. 4, ? Quiz-Compend Series?" 12mo, Cloth. Price $1.00

"Dr. Brubaker deserves the hearty thanks of medical students for his Compend of Physiology. He has arranged the fundamental and practical principles of the science in a particularly inviting and accessible manner. I have already introduced the work to my class."—*Maurice N. Miller, M.D., Demonstrator of Physiology, Medical Department University of the City of New York.*

BYFORD. DISEASES OF WOMEN. New Revised Edition.

The Practice of Medicine and Surgery, as applied to the Diseases of Women. By W. H. BYFORD, A.M., M.D., Professor of Obstetrics and The Diseases of Women and Children, in the Chicago Medical College. Third Edition. Revised and Enlarged, much of it rewritten, with numerous additional illustrations.

Price, in Cloth $5.00; Leather, $6.00

"The treatise is as complete a one as the present state of our science will admit of being written. We commend it to the diligent study of every practitioner and student, as a work calculated to inculcate sound principles and lead to enlightened practice.—*New York Medical Record.*

"The author is an experienced writer, an able teacher in his department, and has embodied in the present work the results of a wide field of practical observation. We have not had time to read its pages critically, but freely commend it to all our readers, as one of the most valuable practical works issued from the American press."—*Chicago Medical Examiner.*

BY SAME AUTHOR.

ON THE UTERUS. The Chronic Inflammation and Displacement of the Unimpregnated Uterus.

An Enlarged Edition, with Illustrations. 8vo. Price $2.50

"A good book from a good man."—*American Journal Medical Science.*

"It is a sensible, practical work, and cannot fail to be read with interest and profit."—*Boston Medical and Surgical Journal.*

BRAUNE, TOPOGRAPHICAL ANATOMY.

An Atlas of Topographical Anatomy. Thirty-four Full-page Plates, Photographed on Stone, from Plane Sections of Frozen Bodies, with many other illustrations. By WILHELM BRAUNE, Professor of Anatomy at Leipzig. Translated and Edited by EDWARD BELLAMY, F.R.C.S., Lecturer on Anatomy, Charing Cross Hospital, London. Quarto. Price, Cloth, $8.00; Half Morocco, $10.00

"As a whole the work cannot fail to meet with a hearty reception by every progressive student of the human body. To the surgeon it is a contribution to the study of topographical anatomy which needs to be known to be properly appreciated To such practitioners who reside in large cities, where anatomy can be studied upon the cadaver, it will afford a valuable aid, while to those who are without such means of study it is an almost indispensable addition to a working library."—*New York Medical Record.*

"We commend the book most heartily to the Profession."—*American Journal of Medical Science.*

BUCKNILL AND TUKE ON INSANITY.

A Manual of Pyschological Medicine: containing the Lunacy Laws, the Nosology, Œtiology, Statistics, Description, Diagnosis, Pathology (including morbid Histology), and Treatment of Insanity. By JOHN CHARLES BUCKNILL, M.D., F.R.S., and DANIEL HACK TUKE, M.D., F.R.C.P. Fourth Edition, much enlarged, with twelve lithographic plates, and numerous illustrations. Octavo. Price $8.00

"We have read no book in any language, and certainly none in English, which ought to be preferred to this for a text book, by those who wish to make a thorough study of the subject.—*Edinburgh Medical Journal.*

"We can heartily commend the work.—*American Journal of Insanity.*

BURDETT, HOSPITALS.

Pay Hospitals and Paying Wards throughout the World. Facts in support of a rearrangement of the system of Medical Relief. By HENRY C. BURDETT. 8vo. Price $2.25

"Mr. Burdett displays and discusses the whole scheme of Hospital accommodation with a comprehensive understanding of its nature and extent.—*American Practitioner.*

BY SAME AUTHOR.

COTTAGE HOSPITALS.

General, Fever, and Convalescent: their Progress, Management, and Work. Second Edition, rewritten and much Enlarged, with many Plans and Illustrations. Crown 8vo. Price $4.50

CONTENTS.—CHAP.—1. Origin and Growth of the Cottage Hospital System. 2. Comparative Success of Treatment in large and small Hospitals. 3. Finance. 4. Cottage Hospital Construction and Sanitary Arrangements. 5. The Medical and Nursing Departments. 6. Domestic Supervision and General Management. 7. Cottage Hospital Appliances and Fittings. 8. Cottage Fever Hospitals. 9. Midwifery in Cottage Hospitals. 10. Remunerative Paying Patients. 11. Convalescent Cottages. 12. Cottage Hospitals in America. 13. Mortuaries. 14. A more Detailed Account of certain Cottage Hospitals, with Plans and Elevations. 15. Selected and Model Plans criticised and compared, with a detailed description of various Hospitals. 16. Peculiarities and Special Features in the Working of Cottage Hospitals. With an Appendix containing much statistical and useful information.

"Mr. Burdett's book contains a mass of information, statistical, financial, architectural, and hygienic, which has already proved of great practical utility to those interested in cottage hospitals, and we can confidently recommend this second edition to all who are in search of the kind of information which it contains."—*Lancet.*

BUZZARD, NERVOUS DISEASES.

Clinical Lectures on Diseases of the Nervous System. By THOS. BUZZARD, M.D. Illustrated. Octavo. Price $5.00

CARPENTER, THE MICROSCOPE. Sixth Edition.

The Microscope and its Revelations. By W. B. CARPENTER, M.D., F.R.S. Sixth Edition. Revised and Enlarged, with over 500 Illustrations. Price $5.50

"Not only the student of medicine, but amateurs, and others interested in the study of natural history, will find this volume one of great practical value."—*New York Medical Journal.*

"It is by far the most complete and useful treatise now accessible to the student."—*The Technologist.*

"As a text book of Microscopy in its special relation to natural history and general science, the work before us stands confessedly first, and is alone sufficient to supply the wants of the ordinary student."—*American Journal of Microscopy.*

CARTER, EYESIGHT. New Edition now ready.

Eyesight, Good and Bad. A Treatise on the Exercise and Preservation of Vision. By ROBERT BRUDENELL CARTER, F.R.C.S. Second Edition, with 52 Illustrations, Test Types, etc. 12mo. Price, Cloth, $1.25

"It is written in a lucid and agreeable style, conveying an easily comprehensible account of the structure of the eye and the function of vision, and gives a description of the principal anomalies of the latter, at the same time inculcating such salutary advice as may be beneficial for the preservation of sight."—*London Medical Times and Gazette.*

"There is much wholesome advice given on the 'Care of the Eyes in Infancy and Childhood,' and on this account, if no other, the book should be in the hands of every parent and teacher."—*St. Louis Courier of Medicine.*

CARTER, PRACTICE OF MEDICINE.

Elements of Practical Medicine. By ALFRED H. CARTER, M.D., London, Member of the Royal College of Physicians; Physician to the Queen's Hospital, Birmingham, etc. Crown 8vo. Price $3.00

CULLINGWORTH, ON NURSING. Illustrated.

A Manual of Nursing, Medical and Surgical. By CHARLES J. CULLINGWORTH, M.D., Physician to St Mary's Hospital, Manchester, England. With eighteen Illustrations. 12mo. Cloth, $1.00

BY THE SAME AUTHOR.

ON MONTHLY NURSING.

A Manual for Monthly Nurses. 32mo. Price, Cloth, .75

CAZEAUX AND TARNIER'S MIDWIFERY. New Revised Edition. Twelve Full-page Plates.

The Theory and Practice of Obstetrics; including the Diseases of Pregnancy and Parturition, Obstetrical Operations, etc. By P. CAZEAUX, Member of the Imperial Academy of Medicine, Adjunct Professor in the Faculty of Medicine in Paris. Remodeled and rearranged, with revisions and additions, by S. TARNIER, M.D., Professor of Obstetrics and Diseases of Women and Children in the Faculty of Medicine of Paris. A New American, from the Eighth French and First Italian Edition. Edited and Enlarged by ROBERT J. HESS, M.D., Physician to the Northern Dispensary, Phila., etc. About 1100 pages quarto, with 12 Full-page Plates (five of which are beautifully colored) and over 175 Wood Engravings.

Sold by subscription only. Circulars and information will be sent, upon application to the Publishers.

For many years "Cazeaux's Obstetrics" has been one of the leading text-books in America and England, as well as in France, and the recent decision of four of the foremost Professors of Italy, Drs. Chiara, Morisani, Porro, and Tibone, to translate it into their own language, is a further proof of its usefulness, and that it still stands highest as a practical guide to the general practitioner. Another and perhaps a stronger proof of the value of this great work, is the fact that by comparing many of the text-books issued since the first edition of Cazeaux was published, it will be found that the later authors have, to more or less extent, used it as a foundation for their work: drawing cases, conclusions and illustrations from its mass of facts and the experience of its author. With these important proofs in their mind, the publishers determined to issue a new edition; which would represent the advances made in the science of midwifery. Written expressly for the use of practitioners and students of medicine, and those of midwifery especially, its teachings are plain and explicit, presenting a condensed summary of the leading principles established by the masters of the obstetric art, and such clear, practical directions for the management of the pregnant, parturient and puerperal states, as have been sanctioned by the most authoritative practitioners, and confirmed by the author's own experience. The publishers selected for editor a gentleman who has had an extended experience in private practice and through his connection with one of the largest public dispensaries in the country. The revision has to some extent been based on the last French edition, and on the revisions and additions made by the Italian translators. The writings of other well-known men have also been drawn from, as well as the experience of the editor, an experience gained in every-day work. New chapters or sections have been written, new illustrations inserted, old ones recut, and twelve full-page plates added, five of which are colored.

CHARTERIS, PRACTICE OF MEDICINE.

Hand-Book of the Practice of Medicine. By M. CHARTERIS, M.D., Member of Hospital Staff and Professor in University of Glasgow. With Microscopic and other illustrations. Price $1.25

"We have not often met with a book which can be so confidently recommended to physicians *or men* in general practice."—*Lancet.*

"The style in which it is written is clear and attractive. The illustrations are a marked feature in it. It can be recommended as a very reliable, handy book, well adapted for ready reference."—*New Remedies.*

CHAVASSE ON CHILDREN.

The Mental Culture and Training of Children. By PYE HENRY CHAVASSE. 12mo. Price, Paper covers, .50; Cloth, $1.00

The mental culture and training of children is of immense importance. Many children are so wretchedly trained, or rather not trained at all, and so mismanaged, that a few thoughts on this subject cannot be thrown away, even upon the most careful.

CLAY ON OBSTETRIC SURGERY. Third Edition.

A complete Hand-Book of Obstetric Surgery, with Rules for every Emergency and Descriptions of the more difficult as well as the every day operations. By CHARLES CLAY, M.D., with numerous Illustrations. From the Third London Edition. 12mo. Paper Covers, .75; Cloth, $1.25

"It is a useful and convenient book of reference; the illustrations are good, and the book will be found of value to the student and young practitioner, as well as to the skilled Obstetrician."—*American Journal of Obstetrics.*

CLEVELAND, POCKET DICTIONARY.

A Pronouncing Medical Lexicon, containing correct Pronunciation and Definition of terms used in medicine and the collateral sciences. By C. H. CLEVELAND, M.D. Thirty-first Edition. 16mo.
Price, Cloth, 75 cents; Tucks with Pocket, $1.00

This is a most convenient size for the pocket, and contains all the principal words in use, together with rules for pronunciation, abbreviations used in prescriptions, list of poisons, their antidotes, etc.

COHEN, INHALATION. Enlarged Edition.

Inhalation, its Therapeutics and Practice, including a Description of the Apparatus Employed, etc. By J. SOLIS COHEN, M.D. With cases and Illustrations. A New Enlarged Edition. 8vo. Price $2.50

"The book has the merit of containing much information that cannot be found elsewhere."—*N. Y. Medical Journal.*

"One of the best treatises we have seen on this subject."—*Medical Times and Gazette.*

COOPER ON SYPHILIS.

Syphilis and Pseudo-Syphilis. By ALFRED COOPER, F.R.C.S., Surgeon to the Lock Hospital, to St. Marks and to the West London Hospitals. Octavo.
Cloth, $3.50

COBBOLD, PARASITES.

A Treatise on the Entozoa of Man and Animals, including some account of the Ectozoa. By T. SPENCER COBBOLD, M.D., F.R.S. With 85 illustrations. 8vo. Price $5.00

COLES, THE MOUTH. Third Edition, just ready.

Deformities of the Mouth, Congenital and Acquired, with Their Mechanical Treatment. By OAKLEY COLES, D.D.S. Third Edition. 83 Wood Engravings and 96 Drawings on Stone. 8vo. Price $4.50

"Altogether we must heartily congratulate Mr. Coles on this creditable completion of a work which cannot but redound to his credit wherever it is known."—*British Journal of Dental Science.*

"We recommend this book to the study of both surgeons and dentists."—*London Lancet.*

BY SAME AUTHOR.

CHAPMAN. THE CIRCULATION OF THE BLOOD.

A History of the Discovery of the Circulation of the Blood. By HENRY C. CHAPMAN, M.D., Professor of Institutes of Medicine and Medical Jurisprudence in Jefferson Medical College, Philadelphia. Octavo. Cloth, $1.00

THE DENTAL STUDENT'S NOTE-BOOK.

A new Edition. 16mo. Price $1.00

CORMACK, CLINICAL STUDIES.

Illustrated by Cases Observed in Hospital and Private Practice. By Sir JOHN ROSE CORMACK, M.D., K.B., etc. Illustrated. 2 vols. 1,127 pp. Price $5.00

COURTY, THE UTERUS, OVARIES, ETC. *Subscription only.*

A Practical Treatise on Diseases of the Uterus, Ovaries, and Fallopian Tubes. By Prof. A. COURTY, of Montpellier, France. Translated from the Third Edition by his pupil and assistant, AGNES MCLAREN, M.D., M.R.Q.C.P.I. With a Preface by J. MATTHEWS DUNCAN, M.D., LL.D., F.R.S., Obstetric Physician to Saint Bartholomew's Hospital, London. With 431 Illustrations. One Vol., 8vo. Price, in Handsome Cloth, $6.00; Full Sheep, Raised Bands, $7.00

OUTLINE OF CONTENTS.

INTRODUCTION.—On the Anatomy, Physiology, and Teratology of the Organs of Generation. PART I.—GENERAL SURVEY OF UTERINE DISEASES. Diagnosis of Uterine Diseases in General; Treatment of Uterine Diseases in General; General Characteristics of Uterine Diseases. PART II.—UTERINE DISEASES IN DETAIL. Functional Disorders; Changes of Position; Morbid States without Neoplasm; Organic Alterations; Diseases of the Uterine Appendages; Pelvic Hemorrhages and Peri-uterine Hæmatocele; Cyst of the Ovary and Genito-pelvic Tumor; Sterility, etc., etc. INDEX.

"Courty's work has, since its first publication, been recognized everywhere. In France, its position is attested by the sale of two editions, numbering, I am told, ten thousand copies, and by the appearance of another, the third edition. I recommend to the careful study of my professional brethren a book which has already been crowned by the Institute of France."—*J. Matthews Duncan.*

CURLING, ON THE TESTIS.

A Practical Treatise on the Diseases of the Testis, Spermatic Cord, and Scrotum. By T. B. CURLING, M.D., F.R.S. Fourth Edition, Enlarged and Illustrated. 8vo. Price $5.50

"We believe this work to be the most trustworthy that can be consulted in this Department of Surgery, ... his pages abound with valuable suggestions and cautions that mark his intimate knowledge of the subject."—*London Practitioner.*

COOPER'S SURGICAL DICTIONARY.

A Dictionary of Practical Surgery and Encyclopædia of Surgical Science. By SAMUEL COOPER. New Edition, brought down to the present time. By SAMUEL A. LANE, F.R.C.S., assisted by various eminent Surgeons. In two vols. Price $12.00

COTTLE, ON THE HAIR.

The Hair in Health and Disease. By E. W. COTTLE, M.D. Partly from the notes of the late GEORGE NAYLER. 18mo.

CORFIELD, DWELLING HOUSES.

The Sanitary Construction and Arrangement of Dwelling Houses. By W. H. CORFIELD, M.A., M.D. Enlarged Edition, with Plans and Illustrations. 12mo. Price $1.25

COULSON, THE BLADDER. Sixth Edition.

Diseases of the Bladder and Prostate Gland. By WALTER J. COULSON, F.R.C.S. Sixth Edition. Revised and Enlarged, with 22 Engravings. 8vo. Price $6.40

CRIPPS, THE RECTUM.

Cancer of the Rectum. Its Pathology, Diagnosis and Treatment. By W. HARRISON CRIPPS, F.R.C.S. Illustrated by Plates. 8vo. Price $2.40

DAY ON CHILDREN. Second Edition. Just Ready.

The Diseases of Children. A Practical and Systematic Treatise for Practitioners and Students. By WM. H. DAY, M.D. Second Edition. Rewritten and very much Enlarged. 8vo. 752 pp. Price, Cloth, $5.00; Sheep, $6.00

"Believing the work well adapted to meet the wants of the Student as well as the Practitioner, I will recommend it to the classes of Rush Medical College."—*DeLaskie Miller*, M.D., *Chicago*.

"On the whole, we must confess we are pleased with this book and can heartily recommend it—a recommendation which it does not appear to need, as it has already reached its second edition."—*American Journal of Medical Science*.

"Dr. Day brings to his task a large experience, and evidences a very thorough knowledge of the literature, native and foreign, pertaining to this special branch of medicine. The book has been written with great care, and the author is a good writer. The publisher's part of the task has also been excellently performed."—*Boston Medical and Surgical Journal*.

DAY ON HEADACHES. Fourth Edition.

The Nature, Causes, and Treatment of Headaches. Fourth Edition. Illustrated. By WM. HENRY DAY, M.D. Octavo.
Paper Covers, 75 cents; Cloth, $1.25

SUMMARY OF CONTENTS.—Headache from Cerebral Anaemia, Cerebral Hyperæmia, Sympathetic, Congestive, Dyspeptic or Bilious Headaches, Headache from Plethora, from Exhaustion, from Change in Cerebral Tissue, from Affections of the Periosteum, Nervous and Nervo-Hyperæmic Headache, Toxæmic, Rheumatic, Arthritic or Gouty Headache, Neuralgic Headache, and Headaches of Childhood, Early and Advanced Life.

"Well worth reading. The remarks on treatment are very sensible."—*Boston Medical and Surg. Journal*.

DALBY, ON THE EAR.

The Diseases and Injuries of the Ear. By W. B. DALBY, M.D., Surgeon and Lecturer on Aural Surgery, St. George's Hospital. With Illustrations. 12mo. Price $1.50

"A safe and readable introduction to aural surgery."—*Medical Press and Circular*.

"Dr. Dalby has presented us with a very readable little book, which is destined to render much service in the saving of ears."—*N. Y. Medical Journal*.

"The lectures occupy 226 pages, are clearly and concisely written, contain a number of good illustrations, and are well worth the careful study of both student and practitioner. To aurists the work will be most welcome and valuable."—*Specialist*.

DILLINGBERGER, WOMEN AND CHILDREN'S DISEASES.

A Hand-Book of the Treatment of the Diseases Peculiar to Women and Children. By Dr. EMIL DILLINGBERGER. 12mo. Price $1.50

"It is a *magnum in parvo*. The style is simple, clear, lucid, and free from theoretical discussion. No one will regret the small outlay for this volume.—*Richmond and Louisville Medical Journal*.

DUNCAN ON STERILITY.

Sterility in Women; being the Gulstonian Lectures delivered in the Royal College of Physicians, February, 1883. By J. MATHEWS DUNCAN, M.D., LL.D., Obstetric Physician to St. Bartholomew's Hospital, etc. Octavo. Cloth, $2.00

DURKEE, VENEREAL DISEASES. Sixth Edition.

Gonorrhœa and Syphilis. By SILAS DURKEE, M.D. Sixth Edition. Revised and Enlarged, with Portrait and Eight Colored Illustrations. 8vo. Price $3.50

"We may, finally, recommend Dr. Durkee's book as eminently practical, well written, full of excellent counsel, and worthy of being consulted by every member of the profession. A late number of the *London Medical Times and Gazette* also speaks of the book in terms of the highest approval."—*Boston Medical and Surgical Journal*.

DAGUENET, OPHTHALMOSCOPY.

A Manual of Ophthalmoscopy, for the Use of Students. By Dr. DAGUENET. Translated from the French, by Dr. C. S. JEAFFRESON, F.R.C.S.E. Illustrated. 12mo. Price $1.50

"Its portable size, the condensed nature of its text, and the admirably systematic arrangement of its contents, render it extremely useful as a pocket manual for Students.—*Translator's Preface.*

DOBELL, WINTER COUGH AND CATARRH.

On Winter Cough, Catarrh, Bronchitis, Emphysema, Asthma, etc. By HORACE DOBELL, M.D., Lecturer at the Royal Hospital for Diseases of the Chest. Third Edition. With Colored Plates. 8vo. Price $3.50

BY SAME AUTHOR.

ON LOSS OF WEIGHT. Revised Edition.

Blood Spitting and Lung Disease. Colored Frontispiece of Lung. Tabular Map, etc. Second Edition Enlarged. 8vo. Price $4.00

DOMVILLE, ON NURSING.

A Manual for Hospital Nurses and others engaged in attending to the sick. 4th Edition. With Recipes for Sick Room Cookery, etc. Price .75

DRUITT'S MODERN SURGERY. Eleventh Edition.

The Surgeon's Vade Mecum; a Manual of Modern Surgery. By ROBERT DRUITT, F.R.C.S. Eleventh Enlarged Edition, with 369 Illustrations. 864 pp. 1878. Price $5.00

This is a most complete, accurate, and trustworthy Hand, or Text-Book of Surgery. Unrivaled as a book for the Student. Fully illustrated, and brought up to the present state of the science. In use in many Medical Colleges.

DULLES, ACCIDENTS.

What to Do First, In Accidents and Poisoning. By C. W. DULLES, M.D. Second Edition, Enlarged, with new Illustrations. Cloth, .75

"Its usefulness entitles it to a wide and permanent circulation."—*Boston Gazette.*

"A complete guide for sudden emergencies.—*Philadelphia Ledger.*

"So plain and sensible that it ought to be introduced into every female seminary.—*Evening Chronicle, Pittsburgh.*

EDWARDS, BRIGHT'S DISEASE. New Edition.

How a Person Affected with Bright's Disease Ought to Live. By JOS. F. EDWARDS, M.D. Second Edition. 12mo. Price .75

"Physicians, as well as laymen, will find the work interesting, and will obtain many valuable hints as to the proper hygiene to be observed in this disease."—*Cincinnati Medical News.*

BY SAME AUTHOR.

CONSTIPATION. New Edition.

Plainly Treated and Relieved Without the Use of Drugs. Second Edition. 12mo. Price .75

MALARIA.

Malaria: What It Means; How to Escape It; Its Symptoms; When and Where to Look for It. 12mo. Price .75

VACCINATION AND SMALL-POX.

Showing the Reasons in favor of Vaccination, and the Fallacy of the Arguments Advanced against it, with Hints on the Management and Care of Small-Pox patients. 16mo. Price .50

These are invaluable little treatises upon subjects that enter painfully into the life experiences of a large majority of the human family. Dr. Edwards shows not only how they may be avoided, but in plain and simple language he tells those already afflicted with them how they may find relief.

ELLIS, DISEASES OF CHILDREN.

A Practical Manual of the Diseases of Children, with a Formulary. By EDWARD ELLIS, M.D. Late Physician to the Victoria Hospital for Children, London. Fourth Edition Enlarged. Now Ready. Price $3.00

BY SAME AUTHOR.

WHAT EVERY MOTHER SHOULD KNOW.

12mo. Price .75

"It is only too true that our children have to dodge through the early part of life as through a labyrinth. We must be thankful to meet with such a sensible guide for them as Dr. Ellis."—*Pall Mall Gazette.*

FLÜCKIGER, THE CINCHONA BARKS.

The Cinchona Barks Pharmacognostically Considered. By Professor FRIEDRICH FLÜCKIGER, of Strasburg. Translated by FREDERICK B. POWER, PH.D., formerly Professor of Chemistry, Philadelphia College of Pharmacy, now Professor of Materia Medica and Pharmacy, University of Wisconsin. With 8 Lithographic Plates. Royal Octavo. Cloth, $1.50

FENNER, ON VISION. Second Edition, Enlarged.

Vision; Its Optical Defects, the Adaptation of Spectacles, Defects of Accommodation, etc. By C. S. FENNER, M.D. With Test Types and 74 Illustrations. Second Edition, Revised and Enlarged. 8vo. Price $3.50

FENWICK, THE PRACTICE OF MEDICINE.

Outlines of the Practice of Medicine. With Appropriate Formulæ and Illustrations. By SAMUEL FENWICK, M.D., Physician to the London Hospital. 12mo.

"This little work displays a sound judgment in the arrangement of its subject matter, and an intimate acquaintance with the practice of medicine possessed by but few writers, and should have been elaborated into a more comprehensive work. Of all the hand-books we have seen, this is certainly one of the best."—*Medical Herald.*

"It is an eminently practical little treatise, pervaded with much common sense, and will doubtless be found useful, particularly by advanced students."—*Boston Medical and Surgical Journal.*

BY SAME AUTHOR.

ON THE STOMACH.

Atrophy of the Stomach and Its Effect on the Nervous Affections of the Digestive Organs. 8vo. Price $3.20

FOTHERGILL, ON THE HEART. Second Edition.

The Heart and Its Diseases. With Their Treatment. Including the Gouty Heart. By J. MILNER FOTHERGILL, M.D., Associate Fellow of the College of Physicians of Philadelphia. Second Edition, Entirely Re-written. Octavo. Price $3.50

"It is the best, as well as the most recent work on the subject in the English language."—*Medical Press and Circular.*

"The most interesting chapter is undoubtedly that on the gouty heart, a subject which Dr. Fothergill has specially studied, and on which he entertains views such as are likely, we think, to be generally accepted by clinical physicians, although they have not before been stated, so far as we are aware, with the same breadth of view and extended illustration."—*British Medical Journal.*

"To many an earnest student it will prove a light in darkness; to many a practitioner cast down with a sense of his powerlessness to cope with the rout and demoralization of Nature's forces, a present help in time of trouble."—*Philadelphia Medical Times.*

"The work throughout is a masterpiece of graphic, lucid writing, full of good, sound teaching, which will be appreciated alike by the practitioner and the student."—*Students' Journal.*

FULTON, ON PHYSIOLOGY.

A Text-Book of Physiology. By J. FULTON, M.D., Professor at Trinity Medical College, Toronto. Second Edition, Illustrated and Revised. 8vo. Price $4.00

FLOWER, DIAGRAMS OF THE NERVES.

Diagrams of the Nerves of the Human Body. Exhibiting their Origin, Divisions, and Connections, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. By WILLIAM H. FLOWER, F.R.C.S., F.R.S., Hunterian Professor of Comparative Anatomy, and Conservator of the Museum of the Royal College of Surgeons. Third Edition, thoroughly revised. With six Large Folio Maps, or Diagrams. Royal Quarto. Price $3.50

"Admirably arranged, and will be of incalculable aid to the student of anatomy. Each of the large and beautiful plates is accompanied with explanatory text."—*N. Y. Medical Record.*

"The nerves and ganglia are clearly represented. The impressions are well made, and no doubt the diagrams will prove useful."—*Medical and Surgical Reporter.*

FLAGG, PLASTIC FILLING.

Plastics and Plastic Filling; As Pertaining to the Filling of all Cavities of Decay in Teeth below Medium in Structure, and to Difficult and Inaccessible Cavities in Teeth of all Grades of Structure. With some beautifully executed Illustrations. By J. FOSTER FLAGG, D.D.S., Professor of Dental Pathology and Therapeutics in Philadelphia Dental College. Octavo, 2d Edition. Price $4.00

FOX, WATER, AIR AND FOOD.

Sanitary Examinations of Water, Air and Food. By CORNELIUS B. FOX, M.D. 94 Engravings. 8vo. Price $4.00

FOSTER, CLINICAL MEDICINE.

Lectures and Essays on Clinical Medicine. By BALTHAZAR FOSTER, M.D. Illustrated. 8vo. Price $3.00

"No one can peruse the thoughtful comments of our author upon every subject he considers, without feeling himself a wiser man for his pains."—*N. Y. Medical Journal.*

"It is the record of honest work, such as Dr. Foster may be proud of; we can recommend it to the profession, it may be read with profit and advantage by both practitioner and student.—*Edinburgh Medical Journal.*

FOX, ATLAS OF SKIN DISEASES.

Complete in Eighteen Parts, each containing Four Chromo-Lithographic Plates, with Descriptive Text and Notes upon Treatment. In all 72 large colored Plates. By TILBURY FOX, M.D., F.R.C.P., Physician to the Department for Skin Diseases in University College Hospital. Folio Size.

Price $1.00 each, or complete, bound in cloth, $20.00

No Atlas of Skin Diseases has been issued in this country for many years, and no complete work of the kind is now procurable by the Profession. This one, brought out under the editorial supervision and care of Dr. Tilbury Fox (the most distinguished writer on Cutaneous Medicine now in the English language), is partly based upon the classical work of Willan and Bateman (now entirely out of print), but completely remodeled, so as to represent fully the Dermatology of the present day.

"Preference will be given to this work over Hebra; not simply, however, because it is a home production, but by reason of the manner of its execution, the excellent delineation of disease, and the natural coloring of the plates. . . . The letter-press is entirely new. In the accuracy of the latter the subscriber may have the fullest confidence, since it is from the pen of Dr. Tilbury Fox."—*British and Foreign Medico-Chirurgical Review.*

FRANKLAND, WATER ANALYSIS.

Water Analysis, For Sanitary Purposes, with Hints for the Interpretation of Results. By E. FRANKLAND, M.D., F.R.S. Illustrated. 12mo. Price $1.00

"The author's world-wide reputation will commend this manual to all sanitarians, and they will not be disappointed in finding all the essentials of the important subject of which it treats."—*The Sanitarian.*

"The work is one which physicians practicing in the country and in villages and towns remote from medical centres cannot afford to be without."—*Medical and Surgical Reporter.*

GRANVILLE. NERVE VIBRATION AND EXCITATION.

Nerve Vibration and Excitation as Agents in the Treatment of Functional Disorder and Organic Disease. By J. MORTIMER GRANVILLE, M.D. Illustrated. 8vo. Price $2.00

GALLABIN. DISEASES OF WOMEN.

The Student's Guide to the Diseases of Women. By A. LEWIS GALLABIN, M.A., M.D., F.R.C.P. Illustrated with 63 Engravings. 12mo. Price $1.25

BY SAME AUTHOR.

A MANUAL OF MIDWIFERY.

For Students and Practitioners. Illustrated. *In Press.*

***Prof. Gallabin is Obstetric Physician to Guy's Hospital, London, and occupies the chair of Midwifery in that Institution. His work in this department has been noted for its perfection and practical character.

GAMGEE. WOUNDS AND FRACTURES.

The Treatment of Wounds and Fractures. Clinical Lectures by SAMPSON GAMGEE, F.R.S.E., Consulting Surgeon to Queen's Hospital, Birmingham. 34 Engravings. Second Edition. Octavo. Price $3.50

GARDNER. BREWING, DISTILLING, ETC.

The Brewer, Distiller and Wine Manufacturer; a handbook for all interested in the Manufacture and Trade of Alcohol and Its Compounds. Edited by JOHN GARDNER, Fellow of the Chemical Society of London, Editor of "Cooley's Cyclopædia," etc. Illustrated. Price $1.75

BY THE SAME AUTHOR.

THE DYER AND BLEACHER,

The Dyer and Bleacher; being the second volume of the Technological Handbooks. Octavo. Price $1.75

GIBBES. STUDENT'S PATHOLOGY.

Practical Histology and Pathology. By HENEAGE GIBBES, M.B. 12mo. Cloth. Second Edition. Price $1.50

GILL. ON INDIGESTION. Third Edition.

Indigestion: What It Is; What It Leads To; and a New Method of Treating It. By JOHN BEADNELL GILL, M.D. Third Edition. 12mo. Price $1.25

GILLIAM'S PATHOLOGY. Illustrated.

The Essentials of Pathology; a Handbook for Students. By D. TOD GILLIAM, M.D., Professor of Physiology, formerly Professor of Pathology, Starling Medical College, Columbus, O. With 47 Illustrations. 12mo. Cloth, $2.00

GLISAN. TEXT-BOOK OF MODERN MIDWIFERY.

A Text-Book of Modern Midwifery. By RODNEY GLISAN, M.D., Emeritus Professor of Midwifery and Diseases of Women and Children, in the Medical Department of Willamette University, Portland, Oregon, and Late President of the Oregon State Medical Society. With 129 Illustrations. One Volume. Octavo. 624 pp. Price, in Cloth, $4.00; in Leather, $5.00

GOODHART. THE DISEASES OF CHILDREN.

The Student's Guide to the Diseases of Children. By J. F. GOODHART, M.D., F.R.C.P., Physician to Evelina Hospital for Children, Demonstrator of Morbid Anatomy at Guy's Hospital. *In Press.*

GORGAS. DENTAL MEDICINE.

A Manual of Dental Medicine, Materia Medica and Therapeutics. By FERDINAND J. S. GORGAS, M.D., D.D.S., Professor of the Principles of Dental Science, Dental Surgery and Dental Mechanism, in the Dental Department of the University of Maryland. Octavo. Price $3.00

GROSS. BIOGRAPHY OF JOHN HUNTER.

John Hunter and His Pupils. By S. D. GROSS, M.D., Professor of Surgery in Jefferson Medical College, Philadelphia. With a beautifully executed full length Portrait of the Author in his Study. A Handsome Octavo volume. Bound in Beveled Cloth. Paper, .75; Cloth, $1.25

GODLEE'S ATLAS OF HUMAN ANATOMY.

Illustrating most of the Ordinary Dissections and many not usually practiced by the Student. Accompanied by References and an Explanatory Text. Complete. Folio Size. 48 Colored Plates. By RICKMAN JOHN GODLEE, M.D., F.R.C.S. Forming a large Folio Volume, with References, and an Octavo Volume of Letter-press.

Price of the two Volumes, Atlas and Letter-press, Cloth, $20.00

"It is likely to prove as useful to the physician and surgeon as to the anatomist."—*Medical Times and Gazette.*

"The explanatory text is concise, well written, and contains many valuable suggestions for the surgeon."—*London Lancet.*

GOWERS, SPINAL CORD.

Diagnosis of Diseases of the Spinal Cord. With Colored Plates and Engravings. A Second Edition, Revised and Enlarged. By WILLIAM R. GOWERS, M.D., Assistant Professor Clinical Medicine, University College, London. 8vo. Third Edition. Price $1.50

BY SAME AUTHOR.

OPHTHALMOSCOPY.

A Manual and Atlas of Medical Ophthalmoscopy. With 16 Colored Autotype and Lithographic Plates and 26 Wood Cuts, comprising 112 Original Illustrations of the Changes in the Eye in Diseases of the Brain, Kidneys, etc. 8vo. Price $6.00

EPILEPSY AND ITS TREATMENT.

Epilepsy and other Chronic Convulsive Diseases: Their Causes, Symptoms, and Treatment. Octavo. Price, Cloth, $4.00

NERVOUS DISEASES.

A Manual of Diseases of the Nervous System, for Practitioners and Students. *In Press.*

"Dr. Gowers, while profoundly conversant with the literature of his subject, has not allowed himself to be influenced to an undue extent by the writings of others, but while fairly stating their views, where this is necessary, he at the same time brings to bear upon them the experience derived from his own extensive observations, and when, consequently, they receive confirmation or not at his hands, they are all the more valuable as being the outcome of the most searching and unbiased criticism. It would be impossible, within the limits of a short review, to convey an adequate idea of the extent of Dr. Gowers' work."—*Edinburgh Medical Journal.*

GREENHOW, BRONCHITIS.

On Chronic Bronchitis, especially as connected with Gout, Emphysema, and Diseases of the Heart. By E. HEADLAM GREENHOW, M.D. 12mo. Price $1.50

BY SAME AUTHOR.

ADDISON'S DISEASE.

Being the Croonian Lectures, delivered before the Royal College of Physicians, London. Revised and Illustrated by Plates and Reports of Cases. 8vo. Price $3.00

"The book forms a most interesting and valuable monograph, comprehensive and exhaustive."—*British Medical Journal.*

HUGHES, COMPEND OF THE PRACTICE OF MEDICINE.

A Compend of Practice. By DANIEL E. HUGHES, M.D., Demonstrator of Clinical Medicine at Jefferson Medical College, Philadelphia. In two parts—

PART I.—Continued, Eruptive, and Periodical Fevers, Diseases of the Stomach Intestines, Peritoneum, Biliary Passages, Liver, Kidneys, etc., and General Diseases, etc.

PART II.—Diseases of the Respiratory System, Circulatory System, and Nervous System; Diseases of the Blood, etc.

Price of each Part, in Cloth, $1.00; interleaved for the addition of Notes, $1.25

*** These little books can be regarded as a full set of notes upon the Practice of Medicine, containing the Synonyms, Definitions, Causes, Symptoms, Prognosis, Diagnosis, Treatment, etc., of each disease, and including a number of new prescriptions. They have been compiled from the lectures of prominent Professors, and reference has been made to the latest writings of Professors FLINT, DA COSTA, REYNOLDS, BARTHOLOW, ROBERTS and others.

HABERSHON, ON THE STOMACH.

On Diseases of the Stomach—The Varieties of Dyspepsia—Their Diagnosis and Treatment. By S. O. HABERSHON, M.D., F.R.C.P., Senior Physician to, and Late Lecturer on, the Principles and Practice of Medicine at Guy's Hospital. Third Edition, Revised. Crown 8vo.

"As an expression of the results of long personal experience in both hospital and private practice, conveyed in agreeable though not always perspicuous diction, this contribution of Dr. Habershon's has special value of its own, and is so far entitled to the favorable consideration of the practitioner, as is already testified by a demand for a third edition."—*American Journal of Medical Sciences.*

HALE, ON CHILDREN.

The Management of Children in Health and Disease. A Book for Mothers. By MRS. AMIE M. HALE, M.D. Abounding in valuable information and common sense advice. New Enlarged Edition. 12mo. Price .75

"We shall use our influence in the introduction of this work to families under our care, and we urge the profession generally to follow our example."—*Buffalo Medical and Surgical Journal.*

HORWITZ, COMPEND OF SURGERY.

A Compend of Surgery, including Minor Surgery, Amputations, Fractures, Ligatures, Dislocations, Surgical Diseases, etc., with Differential Diagnosis and Treatment. By ORVILLE HORWITZ, B.S., M.D., with Illustrations. 12mo.
Cloth, $1.00

HARDWICKE, MEDICAL EDUCATION.

Medical Education and Practice in All Parts of the World. Containing Regulations for Graduation at the Various Universities throughout the World. By HERBERT JUNIUS HARDWICKE, M.D., M.R.C.P. 8vo. Price $3.00

"Dr. Hardwicke's book will prove a valuable source of information to those who may desire to know the conditions upon which medical practice is or may be pursued in any or every country of the world, even to the remotest corners of the earth. The work has been compiled with great care, and must have required a vast amount of labor and perseverance on the part of its author."—*Dublin Medical Journal.*

HARLEY, ON THE LIVER. Illustrated.

On Diseases of the Liver, with or without Jaundice. Diagnosis and Treatment. By GEORGE HARLEY, M.D. Author of the Urine and Its Derangements. With Colored Plates and Numerous Illustrations. Royal Octavo.
Price, Cloth, $5.00; Leather, $6.00.

"It is one of the *freshest, most readable,* and *most instructive* medical books that have been laid upon our table during the present decade. . . . In conclusion, we commend again most heartily Dr. Harley's extremely valuable book."—*Philadelphia Medical Times.*

"The work is far in advance, in original and practical information, of any treatise on the subject with which we are acquainted, and is worth many times its cost to any physician treating hepatic troubles."—*Chicago Medical Times.*

"The whole subject-matter is treated in a masterly manner, and the work is destined to find a place among the classics."—*Medical Herald, Louisville, Ky.*

"It is the outcome of a mind that went to its task amply equipped therefor. It is the product of long thinking and ripe judgment. . . . We must content ourselves with this bare statement, hoping that those who read the book will derive as much benefit as ourselves."—*New Orleans Medical and Surgical Journal.*

HOLDEN, HUMAN OSTEOLOGY. Sixth Edition.

Comprising a Description of the Bones, with Colored Delineations of the Attachments of the Muscles. The General and Microscopical Structure of Bone and its Development. By the Author and A. DORAN, F.R.C.S., with Lithographic Plates, etc. By LUTHER HOLDEN, F.R.C.S. Numerous Illustrations. Sixth Edition, carefully Revised. Price $6.00

BY SAME AUTHOR.

ANATOMY.

Manual of Dissections of the Human Body. Fifth London Edition. With 200 Illustrations. *In Press.*

LANDMARKS.

Landmarks, Medical and Surgical. Third London Edition. Revised and Enlarged. Price $1.00

"Mr. Holden is the happy possessor of the faculty of writing interesting works on Anatomy. A part of the charm consists in the frequent references to practical points, and in the explanation of the advantages and objects of details of structures."—*Boston Medical and Surgical Journal.*

HEATH'S OPERATIVE SURGERY.

A Course of Operative Surgery, consisting of a Series of Plates, each plate containing Numerous Figures, Drawn from Nature by the Celebrated Anatomical Artist, M. Léveillé, of Paris, Engraved on Steel and Colored by Hand, under his immediate superintendence, with Descriptive Text of Each Operation. By CHRISTOPHER HEATH, F.R.C.S., Surgeon to University College Hospital, and Holme Professor of Clinical Surgery in University College, London. One Large Quarto Volume. Second Edition, Revised and Enlarged. *Subscription.*

The author has embodied in this work the experience gained by him during twenty years of surgical teaching. It comprises all the operations that are required in ordinary surgical practice. He has selected for illustration and description those methods which appear to give the best results in practice, referring to the errors likely to occur and the best methods of avoiding them.

BY SAME AUTHOR.

THE STUDENT'S GUIDE TO SURGICAL DIAGNOSIS.

12mo. Price $1.25

"Mr. Heath is so well known, both as a practical surgeon, teacher and writer, that anything from his pen requires no introduction from the hands of reviewers, and scarcely any notice but the announcement of the fact that he has written a book."—*Medical Record.*

A MANUAL OF MINOR SURGERY AND BANDAGING.

Sixth Edition, Revised and Enlarged. With 115 Illustrations. 12mo. Price $2.00

"This excellent work should not be termed a 'Minor' Surgery, but it really consists of the sum and substance of Practical surgery. We would not exchange it for any book in our possession."—*Southern Clinic.*

HEATH'S PRACTICAL ANATOMY. Fifth London Edition.

Practical Anatomy. A Manual of Dissections. Fifth London Edition. 24 Colored Plates, and nearly 300 other Illustrations. Price $5.00

INJURIES AND DISEASES OF THE JAWS.

The Jacksonian Prize Essay of the Royal College of Surgeons of England, 1867. Third Edition. Revised, with over 150 Illustrations. Octavo.

HOOD, ON GOUT AND RHEUMATISM.

A Treatise on Gout, Rheumatism, and the Allied Affections. Their Treatment, Complications, and Prevention. By PETER HOOD, M.D. Second Edition. Revised and Enlarged. With some Considerations on Longevity. Octavo. Price $3.50

"The Observations on Treatment are specially to be commended."—*London Lancet.*

HOLDEN, THE SPHYGMOGRAPH.

The Sphygmograph. Its Physiological and Pathological Indications. By EDGAR HOLDEN, M.D. Illustrated by Three Hundred Engravings on Wood. 8vo. Price $2.00

HOLMES, THE LARYNGOSCOPE.

A Guide to the Use of the Laryngoscope in General Practice. By GORDON HOLMES, M.D., Physician to the Throat and Ear Infirmary. 12mo. Price $1.00

BY SAME AUTHOR.

VOCAL PHYSIOLOGY.

Vocal Physiology and Hygiene. With reference to the Cultivation and Preservation of the Voice. Illustrated. 12mo.

HOFF, ON HÆMATURIA.

Hæmaturia as a Symptom of the Diseases of the Genito-Urinary Organs. By O. HOFF, M.D. Illustrated. 12mo. Price .75

HUNTER, MECHANICAL DENTISTRY.

A Practical Treatise on the Construction of the Various kinds of Artificial Dentures, with Formulæ, Receipts, etc. By CHARLES HUNTER, D.D.S. 100 Illustrations. 12mo. Price $1.50

"It is the outcome of his own experience of some twenty years as a Mechanical Dentist, and contains, moreover, much derived from practical knowledge of other dentists. The value of the book is also much added to by illustrations. It will be very useful to the Dental Student, and to all Mechanical Dentists."—*London Medical Times and Gazette.*

HUTCHINSON'S ILLUSTRATIONS OF CLINICAL SURGERY. First Volume Complete.

Consisting of Plates, Photographs, Woodcuts, Diagrams, etc. Illustrating Surgical Diseases, Symptoms, and Accidents; also Operations and other Methods of Treatment. With Descriptive Letter-press. By JONATHAN HUTCHINSON, F.R.C.S., Senior Surgeon to the London Hospital, Surgeon to the Moorfields Ophthalmic Hospital, and to the Hospital for Diseases of the Skin, Blackfriars. In Quarterly Fasciculi. Imperial 4to. Volume 1. (Ten Fasciculi) bound complete in itself. Price $25.00. Parts Eleven to Sixteen of Volume 2, Now Ready. Each, $2.50

HEWITT, DISEASES OF WOMEN. Fourth Edition.

The Diagnosis, Pathology, and Treatment of Diseases of Women, Including the Diagnosis of Pregnancy. Founded on a Course of Lectures Delivered at St. Mary's Hospital Medical School. By GRAILY HEWITT, M.D., Lond., M.R.C.P., Physician to the British Lying-in Hospital; Lecturer on Midwifery and Diseases of Women and Children at St. Mary's Hospital Medical School; Honorary Secretary to the Obstetrical Society of London, etc. The Fourth American Edition. Revised and Enlarged, with New Illustrations. Octavo.

Price, Paper, $1.50; Cloth, $2.50

"Readers of the former editions will not require to be told that the additions now made are of the highest possible excellence."—*Times and Gazette.*

"It is one of the most useful, practical, and comprehensive works upon the subject in the English language, a true guide to the student, and an invaluable means of reference for the teacher."—*N. Y. Medical Record.*

"The excellent work of Dr. Hewitt presents—in a form well adapted to conduct the student to a knowledge of the Diseases of Women, and to assist the young practitioner in his study of these diseases at the bedside of the patient—a very full and clear exposition of the views entertained by the most authoritative teachers as to their pathological treatment and their correct Diagnosis."—*Amer. Med. Journal.*

HAY, SARCOMATOUS TUMOR.

History of a Case of Recurring Sarcomatous Tumor of the Orbit in a Child. By THOMAS HAY, M.D. Illustrated. Paper. Price .50

HEWSON, EARTH IN SURGERY.

Earth as a Topical Application in Surgery, Being a Full Exposition of its Use in Cases Requiring Topical Applications. By ADDINELL HEWSON, M.D. Illustrated. 8vo. Price $2.50

HODGE, ON ABORTION.

On Fœticide or Criminal Abortion. By HUGH L. HODGE, M.D.

Price, Paper, .30; Cloth, .50

HODGE, CASE-BOOK.

Note-Book for Cases of Ovarian Tumors. By H. LENNOX HODGE, M.D. With Diagrams. Price, Paper, .50

HIGGINS, DISEASES OF THE EYE. Now Ready.

A Hand-Book of Ophthalmic Practice. By CHARLES HIGGINS, F.R.C.S. Ophthalmic Assistant Surgeon at Guy's Hospital. Second Edition. 16mo.

Price .50

CONTENTS.—SECTION I. Discharge from the Eyes. II. Intolerance of Light. III. Iritis and Glaucoma. IV. Diseases of the Eyelids. V. Watering of the Eye. VI. Acuteness of Vision, Field of Vision, Anomalies of Refraction, Astigmatism, Accommodation, Presbyopia. VII. Disturbance of Vision, Use of the Ophthalmoscope, Normal and Morbid Appearances. VIII. Injuries.

"We have rarely seen so much important information condensed in so short a space."—*American Medical Journal.*

HARRIS, THE PRACTICE OF DENTISTRY. Tenth Edition.

The Principles and Practice of Dentistry. Tenth Revised Edition. In great part Rewritten, Rearranged, and with many new and important Illustrations. By CHAPIN A. HARRIS, M.D., D.D.S. Edited by P. H. AUSTEN, M.D., Professor of Dental Science and Mechanism in the Baltimore College of Dental Surgery. With nearly 400 Illustrations. Royal Octavo. Price, Cloth, $6.50; Leather, $7.50

This new edition of Dr. Harris' work has been thoroughly revised in all its parts, more so than any previous edition. So great have been the advances in many branches of dentistry that it was found necessary to rewrite the articles on subjects, and this has been done in the most efficient manner by Professor Austen, for many years an associate and friend of Dr. Harris, assisted by Professor Gorgas and Thomas S. Latimer, M.D. The publishers feel assured that it will now be found the most complete text-book for the student, and guide for the practitioner in the English language.

BY SAME AUTHOR.

MEDICAL AND DENTAL DICTIONARY. Fourth Edition.

A Dictionary of Medical Terminology, Dental Surgery, and the Collateral Sciences. Fourth Edition, Carefully Revised and Enlarged. By FERDINAND J. S. GORGAS, M.D., D.D.S., Professor of Dental Surgery in the Baltimore College, etc. Royal Octavo. Price, Cloth, $6.50; Leather, $7.50

This Dictionary, having passed through *three* editions, and been for some time out of print, has been again carefully revised by F. J. S. Gorgas, M.D., Dr. Harris' successor as Professor of Dental Surgery in the Baltimore College of Dental Surgery. In his preface to this new edition, the editor says:—

"The object of the reviser has been to bring the book thoroughly up to the present requirements of the profession, the *Medical* portion having been as carefully revised and added to as that devoted more especially to *Dental Science*, while a number of obsolete terms and methods have been omitted. In nearly every one of the seven hundred and forty-three pages of the former edition corrections and additions have been made, and many new processes, terms and appliances described, some of which are not found in any other work published."

HANDY, ANATOMY.

Text-Book of Anatomy and Guide to Dissections. For the Use of Students. By W. R. HANDY, M.D. 312 Illustrations. Price $3.00

HILLIER, DISEASES OF CHILDREN.

A Clinical Treatise on the Diseases of Children. By THOMAS HILLIER, M.D. 8vo. Price $2.00

HUFELAND, LONG LIFE.

The Art of Prolonging Life. By C. W. HUFELAND. Edited by ERASMUS WILSON, M.D. 12mo. Price $1.00

"We wish all doctors and all their intelligent clients would read it, for surely its perusal would be attended with pleasure and benefit."—*American Practitioner.*

"It certainly should be in the library of every physician."—*Medical Brief.*

HUNTER, PORTRAIT OF.

Portrait of John Hunter. From Sharp's well-known Engraving; a copy of Sir Joshua Reynold's Portrait. For Framing. Large size, 9 x 11; sheet 16 x 20. Price, in the Sheet, sent free by mail, 50 cents; or, Handsomely Framed, Price $2.00

HEADLAND, THE ACTION OF MEDICINES. Ninth Edition.

On the Action of Medicines in the System. By F. W. HEADLAND, M.D. Ninth American Edition, Revised and Enlarged. 8vo. Price $3.00

"It displays in every page the evidence of extensive knowledge and of sound reasoning; it will be useful alike to those who are just commencing their studies, and to those who are engaged in the active pursuits of professional life."—*Medical Times.*

"The very favorable opinion which we were amongst the first to pronounce upon this essay has been fully confirmed by the general voice of the profession, and Dr. Headland may now be congratulated on having produced a treatise which has been weighed in the balance, and found worthy of being ranked with our standard medical works."—*London Lancet.*

JAMES, SORE THROAT.

On Sore Throat, Its Nature, Varieties and Treatment, Including its Connection with other Diseases. By PROSSER JAMES, M.R.C.P. Fourth Edition, Revised and Enlarged. With Colored Plates and Numerous Wood-cuts. 12mo. Price $1.25

"We can confidently recommend his therapeutic teachings as well worthy of the careful consideration of the Profession, for they set forth the practice of an enthusiastic worker, whose special experience has been large and lengthened."—*British Medical Journal.*

"The practitioner who buys Dr. James' unpretending little book will provide himself with a wise and practical clinical commentary, and with a well arranged digest of long and varied experience."—*Westminster Review.*

JONES, AURAL ATLAS.

An Atlas of Diseases of the Membrana Tympani. Being a Series of Colored Plates, containing 62 Figures. With appropriate Letter-press and Explanatory Text. By H. MACNAUGHTON JONES, M.D., Surgeon to the Cork Ophthalmic and Aural Hospital. 4to. Price $4.00.

"The cases are well selected, the drawings executed from life, highly artistic and very conscientious, and the commentaries indicate familiarity with the subject and good judgment in dealing with it."—*British Medical Journal.*

BY SAME AUTHOR.

AURAL SURGERY.

A Practical Hand-book on Aural Surgery. Illustrated. Second Edition, Revised and Enlarged, with new Wood Engravings. 12mo. Cloth. Price $2.75

JONES, SIEVEKING AND PAYNE, PATHOLOGICAL ANATOMY.

A Manual of Pathological Anatomy. By C. HANDFIELD JONES, M.D., and EDWARD H. SIEVEKING, M.D., Physician to St. Mary's Hospital. A New Enlarged Edition. Edited by J. F. PAYNE, M.D., Lecturer on Morbid Anatomy at St. Thomas' Hospital. With Numerous Illustrations. Demi 8vo. Price $5.50.

JONES, ON SIGHT AND HEARING.

The Defects of Sight and Hearing, their Nature, Causes, and Prevention. By T. WHARTON JONES, M.D. Second Edition. 16mo. Price .50.

KIRBY, ON PHOSPHORUS. Fifth Edition.

Phosphorus as a Remedy for Functional Diseases of the Nervous System. By E. A. KIRBY, M.D. Fifth Edition. 8vo. Price $1.00

BY THE SAME AUTHOR.

SELECTED REMEDIES.

A Pharmacopœia of Selected Remedies, with Therapeutic Annotations, Notes on Alimentation in Disease, Air, Massage, Electricity, and other Supplementary Remedial Agents, and a Clinical Index; arranged as a Handbook for Prescribers. By EDMUND A. KIRBY, M.D., M.R.C.S., late Physician to the London City Dispensary. Sixth Edition, Enlarged and Revised. 4to. Price $2.25

KOLLMEYER, KEY TO CHEMISTRY.

Chemia Coartata, or Key to Modern Chemistry. By A. H. KOLLMEYER, M.D. With Numerous Tables, Tests, etc. Price $2.25

KIRKE, PHYSIOLOGY. Revised and Enlarged.

A Hand book of Physiology. By KIRKE. Eleventh London Edition. By W. MORRANT BAKER, M.D. 420 Illustrations. *Now Ready.* Price $5.00

"This is undoubtedly the best work for students, on Physiology, extant."—*Cincinnati Med. News.*

KANE, THE OPIUM, MORPHINE AND SIMILAR HABITS.

Drugs that Enslave. The Opium, Morphine, Chloral, Hashisch and Similar Habits. By H. H. KANE, M.D., of New York. With Illustrations. Price $1.25

"It contains a large amount of information collected with much labor and presented in a systematic manner. The subject of the chloral habit has not been investigated by any one, we believe, so thoroughly as by Dr. Kane."—*Medical Record.*

"It deserves to be read by those who feel an interest in discouraging the use of these dangerous drugs. The book is embellished by an excellent phototype frontispiece of Laocoön."—*American Journal of Pharmacy.*

"A work of more than ordinary ability and careful research . . . For the first time, reliable statistics on the use of chloral are classified and published, . . . and it is shown that the use of chloral causes a more *complete and rapid ruin of mind and body* than either opium or morphine."—*Druggists' Circular and Chem.*

KIDD, THERAPEUTICS.

The Laws of Therapeutics; or, the Science and Art of Medicine. By JOSEPH KIDD, M.D. 12mo. Cloth. Price $1.25.

"Dr. Kidd acknowledges two laws—that of *contraria contrariis* and *similia similibus*; but the cases he gives in his chapter on *ars medica* show that, like a sensible practitioner, he does not allow himself blindly to follow either the one or the other, but seeks out the cause of disease, and tries by rational measures to remove it. The cases are the most valuable part of the book."—*London Practitioner.*

LANDIS, A COMPEND OF OBSTETRICS. Illustrated.

A Compend of Obstetrics; especially adapted to the Use of Students and Physicians. By HENRY G. LANDIS, M.D., Professor of Obstetrics and Diseases of Women in Starling Medical College, Columbus, Ohio. Illustrated. 12mo. Cloth. Price $1.00: interleaved for the addition of Notes, $1.25

"The questions are well chosen, the answers clear, concise, and well up to the present state of obstetrical science. It will be a handy book for reference for practitioner as well as student."—*Prof. E. O. F. Roler, Chicago Medical College.*

"I have observed no statement to the correctness of which I could take exception. There are very few practitioners who cannot be instructed by its perusal."—*David Wark, M.D., U. S. Medical College, New York.*

"It is complete, accurate and scientific; the very best book of its kind."—*Prof. J. S. Knox, Rush Medical College, Chicago.*

"I have been teaching in this department for many years, and am free to say that this will be the best assistant I ever had. It is accurate and comprehensive, but brief and pointed."—*Prof. P. D. Vest, St. Louis.*

LEGG, ON THE URINE.

Practical Guide to the Examination of the Urine, for Practitioner and Student. By J. WICKHAM LEGG, M.D. Fifth Edition, Enlarged. Illustrated. 12mo. Price .75

This little work is intended to supply the Physician or Student with a concise guide to the recognition of the different characteristics of the urine, and though small and well adapted to the pocket, contains, probably, everything that could be gleaned from a larger work.

LEARED, IMPERFECT DIGESTION.

The Causes and Treatment of Imperfect Digestion. By ARTHUR LEARED, M.D. The 7th Edition. Revised and Enlarged. 12mo. Price $2.00

LIEBREICH, ATLAS OF OPHTHALMOSCOPY.

An Atlas of Ophthalmoscopy, containing 12 Full-page Chromo-Lithographic Plates, with 59 Figures. By R. LIEBREICH, M.D. Second Edition, Enlarged. Large Quarto.

LIVEING, ON SICK HEADACHE.

Megrim, or Sick Headache and Some Allied Disorders. By EDWARD LIVEING, M.D. With Plates, Tables, etc. 8vo. Price $5.50

LEBER AND ROTTENSTEIN, DENTAL CARIES.

Dental Caries and Its Causes. An Investigation into the Influence of Fungi in the Destruction of the Teeth. By Drs. LEBER and ROTTENSTEIN. Illustrated. 8vo. Paper Cover 75 cents; Cloth, $1.25

"The work gives the result of patient observation, presents the deductions of its authors with a perspicuity and modesty calculated to secure for its positions a thoughtful consideration. We heartily commend it as an educational work."—*Dental Cosmos.*

LEWIN, ON SYPHILIS.

The Treatment of Syphilis. By Dr. GEORGE LEWIN, of Berlin. Translated by CARL PROEGLER, M.D., and E. H. GALE, M.D., Surgeons U. S. Army. Illustrated. 12mo. Price $1.25

"When such authorities as Dr. Drysdale (as we quoted a few weeks ago) condemn the use of mercury in syphilis as "too dangerous," while, on the other hand, eminent surgeons, such as Professor Gross, will not treat a case without that drug, general practitioners will gladly welcome any *media via* which gives us all the good effects of mercurials without any danger of their ill results appearing. This is what is accomplished by Dr. Lewin."—*Philadelphia Medical and Surgical Reporter.*

LIZARS, ON TOBACCO.

The Use and Abuse of Tobacco. By JOHN LIZARS, M.D. 12mo. Price .50

LONGLEY, POCKET MEDICAL LEXICON.

Students' Pocket Medical Dictionary, Giving the Correct Definition and Pronunciation of all Words and Terms in General Use in Medicine and the Collateral Sciences, with an Appendix, containing Poisons and their Antidotes, Abbreviations Used in Prescriptions, and a Metric Scale of Doses. By ELIAS LONGLEY. 24mo. Price, Cloth, $1.00; Tucks and Pocket $1.25

This is an entirely new Medical Dictionary, containing some 300 compactly printed 24mo pages, very carefully prepared by the author, who has had much experience in the preparation of similar works, assisted by the Professors of Chemistry and of Botany in one of our leading medical colleges.

"This little book will be welcomed by students in medicine and pharmacy as a convenient pocket companion, giving the pronunciation, acceptation, and definition of medical, pharmaceutical, chemical and botanical terms."—*American Journal of Pharmacy.*

"It would seem to be just the book for dental and medical students."—*Dental Advertiser.*

"It is, we believe, also the only lexicon in existence in which the pronunciation of words is fully and distinctly marked."—*Canada Medical Review.*

"This is a very compact and complete little dictionary. We commend it as particularly useful to students."—*New York Medical Journal.*

LEFFMANN. ORGANIC AND MEDICAL CHEMISTRY.

A Compend of Organic Chemistry, including Medical Chemistry, Urine Analysis and the Analysis of Water and Food. By HENRY LEFFMANN, M.D., Professor of Clinical Chemistry and Hygiene at the Philadelphia Polyclinic and College for Graduates in Medicine. 12mo. Cloth, Price, $1.00; Interleaved for the addition of Notes, $1.25

THE POLYCLINIC.

A Monthly Journal of Medicine and Surgery, conducted by the Faculty of the Philadelphia Polyclinic and School for Graduates in Medicine. *Sample copies free.* Terms, per Annum, $1.00

MACDONALD, MICROSCOPICAL EXAMINATION OF WATER AND AIR.

A Guide to the Microscopical Examination of Drinking Water, with an Appendix on the Microscopical Examination of Air. By J. D. MACDONALD, M.D. With Twenty-five Full-page Lithographic Plates, Reference Tables, etc. Second Edition, Revised. 8vo. Price $2.75

"The volume is an excellent hand-book and will greatly facilitate the study of the subject."—*Popular Science Monthly.*

MAYS, THE THERAPEUTIC FORCES;

Or, The Action of Medicine in the Light of the Doctrine of Conservation of Force. By THOMAS J. MAYS, M.D. 12mo. Price $1.25

MACKENZIE, ON THE THROAT AND NOSE. Ready.

Including the Pharynx, Larynx, Trachea, Œsophagus, Nasal Cavities, and Neck. By MORELL MACKENZIE, M.D., London, Senior Physician to the Hospital for Diseases of the Chest and Throat, Lecturer on Diseases of the Throat at London Hospital Medical College, etc., etc.

VOL. I. Including the Pharynx, Larynx, Trachea, etc. 112 Illustrations. Price, Cloth, $4.00; Leather, $5.00

VOL. II. Including the Œsophagus, Nasal Cavities, Neck, etc. Illustrated. Price, Cloth, $3.00; Leather, $4.00

THE TWO VOLUMES TAKEN TOGETHER, CLOTH, $6.00; LEATHER, $7.50.

Author's Edition, issued under his supervision, containing all the original Wood Engravings, and the essay on "Diphtheria, Its Causes, Nature, and Treatment," formerly published separately. Each volume sold separately.

"We have long felt the want of a thoroughly practical and systematic treatise on diseases of the throat and nasal passages. Admirable essays have from time to time appeared; no standard work has been written. Any one familiar with laryngoscopic work must appreciate the valuable addition now made to this special department in the work before us. The entire work will include the consideration of affections of the pharynx, larynx, trachea, œsophagus, nasal cavities, and neck. The matter now presented complete for the first time is the result of the author's large and unrivaled experience, both in hospital and private practice, extending over a period of twenty years. There can be but one verdict of the profession on this manual—it stands without any competitor in medical literature, as a standard work on the organs it professes to treat of."—*Dublin Journal.*

"It is both practical and learned; abundantly and well illustrated; its descriptions of disease are graphic, and the diagnoses the best we have anywhere seen. To give examples of the thoroughness of Dr. Mackenzie's book, we may cite the chapter on diphtheria, which embraces 47 pages. The chapter on non-malignant tumors of the larynx would appear to be absolutely exhaustive. Nowhere else have we seen so elaborate a statement of the subject. We can predict for this work a high position, and congratulate its distinguished author upon its appearance."—*Philadelphia Medical Times.*

BY SAME AUTHOR.

THE PHARMACOPŒIA of the Hospital for Diseases of the Throat and Nose.

The Fourth Edition, much enlarged, containing 250 Formulæ, with Directions for their Preparation and Use. 16mo. Price $1.25

GROWTHS IN THE LARYNX.

Their History, Causes, Symptoms, etc. With Reports and Analysis of one Hundred Cases. With Colored and Other Illustrations. 8vo. Price $2.00

HAY FEVER: ITS ETIOLOGY AND TREATMENT.

A Lecture delivered at the London Hospital Medical College Octavo. Price, Paper covers, .50

MACNAMARA, DISEASES OF THE EYE.

A Manual of the Diseases of the Eye. By C. MACNAMARA, M.D. Fourth Edition, Carefully Revised; with Additions and Numerous Colored Plates, Diagrams of Eye, Wood-cuts, and Test Types. Demi 8vo. Price $4.00

"As a book of ready reference on diseases of the eye it has no superior, and we may safely say, no equal in our language."—*Cincinnati Lancet and Observer.*

MADDEN, HEALTH RESORTS.

Health Resorts for the Treatment of Chronic Diseases. A Hand-Book, the result of the author's own observations during several years of health travel in many lands, containing also remarks on climatology and the use of mineral waters. By T. M. MADDEN, M.D. 8vo. Price $2.50

"Rarely have we encountered a book containing so much information for both invalids and pleasure seekers."—*The Sanitarian.*

MEDICAL DIRECTORY OF PHILADELPHIA.

A Directory of the Physicians, Pharmacists, Dentists, Nurses, Veterinary Surgeons, etc., of Philadelphia. Compiled from the Registrar's Records in the Court of Common Pleas. Containing information concerning Medical Colleges, Hospitals, Asylums, Charities, etc., etc. By S. B. HOPPIN, M.D. 12mo.

Price, Cloth, $1.50

MARSHALL & SMITH, ON THE URINE.

The Chemical Analysis of the Urine. By JOHN MARSHALL, M.D., and EDGAR F. SMITH, M.D., of the Chemical Laboratory, Medical Department, University of Pennsylvania. Illustrated by Phototype Plates. 12mo. Price $1.00

MARSHALL, ANATOMICAL PLATES;

Or Physiological Diagrams. Life Size (7 by 4 feet) and Beautifully Colored. By JOHN MARSHALL, F.R.S. An Entirely New Edition, Revised and Improved, Illustrating the Whole Human Body.

The Set, Eleven Maps, in Sheets, Price $50.00
" " handsomely Mounted on Canvas, with Rollers, and Varnished, Price $80.00
An Explanatory Key to the Diagrams, Price .50

No. 1. The Skeleton and Ligaments. No. 2. The Muscles, Joints, and Animal Mechanics. No. 3. The Viscera in Position—The Structure of the Lungs. No. 4. The Organs of Circulation. No. 5. The Lymphatics or Absorbents. No. 6. The Digestive Organs. No. 7. The Brain and Nerves. No. 8. The Organs of the Senses and Organs of the Voice, Plate 1. No. 9. The Organs of the Senses, Plate 2. No. 10. The Microscopic Structure of the Textures, Plate 1. No. 11. The Microscopic Structure of the Textures, Plate 2.

MARSDEN, ON CANCER.

A New and Successful Mode of Treating Certain Forms of Cancer. By ALEXANDER MARSDEN, M.D. Second Edition. Colored Plates. 8vo. Price $3.00

MARTIN, MICROSCOPIC MOUNTING.

A Manual of Microscopic Mounting. With Notes on the Collection and Examination of Objects, and upwards of 150 Illustrations. By JOHN H. MARTIN. Second Edition, Enlarged. 8vo. Price $2.75

MORRIS, ON THE JOINTS.

The Anatomy of the Joints of Man. Comprising a Description of the Ligaments, Cartilages, and Synovial Membranes; of the Articular Parts of Bones, etc. By HENRY MORRIS, F.R.C.S. Illustrated by 44 Large Plates and Numerous Figures, many of which are Colored. 8vo. Price $5.50

MUTER, MEDICAL AND PHARMACEUTICAL CHEMISTRY.

An Introduction to Pharmaceutical and Medical Chemistry. PART ONE.—Theoretical and Descriptive. PART TWO.—Practical and Analytical. Arranged on the principle of the Course of Lectures on Chemistry as delivered at, and the Instruction given in the Laboratories of, the South London School of Pharmacy. By JOHN MUTER, M.D., President of the Society of Public Analysts. A Second Edition, Enlarged and Rearranged. The Two Parts bound in one large octavo volume. Price $6.00

PART TWO.—Practical and Analytical. Bound Separately, for the Special Convenience of Students. Large 8vo. Cloth. Price $2.50

MAC MUNN, THE SPECTROSCOPE.

The Spectroscope in Medicine. By CHAS. A. MAC MUNN, M.D. With 3 Chromo-lithographic Plates of Physiological and Pathological Spectra, and 13 Wood Cuts. 8vo. Price $3.00

"This book is, without question, the best that has yet been published on the subject; to those not familiar with Physiological Spectroscopy it will prove interesting, while to those who are working in this field it is a necessity."—*New York Medical Journal.*

MERRELL'S DIGEST OF MATERIA MEDICA.

A Digest of Materia Medica and Pharmacy; forming a complete Pharmacopœia for the use of Physicians, Pharmacists and Students. By ALBERT MERRELL, M.D. Octavo. Price, Half dark Calf, $4.00

MANN, PSYCHOLOGICAL MEDICINE.

A Manual of Psychological Medicine and Allied Nervous Diseases. Their Diagnosis, Pathology, Prognosis and Treatment, including their Medico-Legal Aspects; with chapter on Expert Testimony, and an abstract of the laws relating to the Insane in all the States of the Union. By EDWARD C. MANN, M.D., of New York. With Illustrations of Typical Faces of the Insane, Handwriting of the Insane, and Micro-Photographic Sections of the Brain and Spinal Cord. Octavo. Cloth, $5.00; Full Leather, $6.00

FROM THE LONDON LANCET, APRIL 19TH, 1884.

"A perusal of this volume reveals the fact that an enormous amount of research has been carried on by the author in order to present to the profession a digest of most of what is known of insanity in its medical, legal and social aspects. The lengthy preface, which is written in admirable style, shows the thorough grasp and liberal spirit with which the writer has prosecuted his work. At the outset an energetic protest is entered against the pernicious system of forcing the intellect at the expense of the natural development of the nervous system. This comes as a timely warning when the tendency is strong, both in America and this country, to educate children unsuggestive of their physical powers and mental receptivity. The opening chapter is devoted to a consideration of the history and classification of insanity. Others follow on ætiology, diagnosis and treatment, on irresponsibility, general paralysis, idiocy and dementia; criminal responsibility; morbid anatomy and pathology; treatment; expert testimony; codification of the criminal common law; evolution of insanity and neurosis; and, lastly, the psychology of crime. Of the four appendices, the first, which occupies forty pages of closely printed matter, consists of a resumé of the legal enactments and procedure as adopted in the several States of the Union; whilst the last is a copious index of the literature of diseases of the mind. As a matter of course, the author condemns the practice of restraint in the treatment of lunatics. He argues strongly in favor of the appointment of expert commissioners, who shall exercise a supervision of asylums, and determine the medical questions in cases of alleged criminal irresponsibility. The letter-press is good, and the phototypes of the physiognomy of some of the more pronounced of mental disorders well executed."

FROM THE ALIENIST AND NEUROLOGIST.

"It will in no way detract from the merits of other contemporary works to say that Dr. Mann's book will supply a want which no other has yet filled, viz.: a manual of plain rules for guidance in the practical consideration of insanity and the treatment of the various allied nervous affections."

MEADOWS, OBSTETRICS, Revised Edition.

A Text-Book of Midwifery. Including the Signs and Symptoms of Pregnancy, Obstetric Operations, Diseases of the Puerperal State, etc. By ALFRED MEADOWS, M.D. Third American, from Fourth London Edition. Revised and Enlarged. With 145 Illustrations. 8vo. Price $2.00

"It is with great gratification that we are enabled to class Dr. Meadows' Manual as a rare exception, and in performance it is an accurate, practical, and credible work, and we unhesitatingly recommend it to [illegible] and practitioner."—*American Journal of Obstetrics.*

"We cannot but feel that every teacher of Obstetrics has great cause to congratulate himself on being able to put in the hands of the student a book which contains so much valuable and reliable information."—*Boston Medical Times.*

"On all questions of treatment, whether by medicines, by hygienic regimen, or by mechanical or operative appliances, this treatise is as satisfactory as a work of manual size could be; students and practitioners can hardly do better than adopt it as their vade mecum."—*The Practitioner.*

"The systematic arrangement of subjects, and the concise, practical style in which it is written, make the work especially valuable as a student's manual."—*Chicago Medical Examiner.*

MEARS, PRACTICAL SURGERY.

Practical Surgery. Including: Part I.—Surgical Dressings; Part II.—Bandaging; Part III.—Ligations; Part IV.—Amputations. With 227 Illustrations. By J. EWING MEARS, M.D., Demonstrator of Surgery in Jefferson Medical College, and Professor of Anatomy and Clinical Surgery in the Pennsylvania College of Dental Surgery. 12mo. Price $2.00

"[illegible] a convenient and useful [illegible] We can most cordially endorse [illegible] made in its modest [illegible] Times and Clinic.

"It contains a great deal of information upon the subjects of which it treats, in a convenient and condensed form. Each division is well illustrated, thereby rendering the text doubly clear."—*New York Medical Record.*

[illegible] OF ALCOHOL.

[illegible] Its Place and Power. By JAMES MILLER, F.R.C.S. 12mo. Price .50

[illegible] ALCOHOL AND TOBACCO.

[illegible] Its Place and Power. By JAMES MILLER, F.R.C.S.; and, Tobacco, [illegible] By JOHN LIZARS, M.A. The two essays in one volume. Price $1.00

MENDENHALL, VADE MECUM.

The Medical Student's Vade Mecum. A Compend of Anatomy, Physiology, Chemistry, The Practice of Medicine, Surgery, Obstetrics, etc. By GEORGE MENDENHALL, M.D. Eleventh Edition. 224 Illustrations. 8vo. Price $2.00

MEIGS AND PEPPER, DISEASES OF CHILDREN.

A Practical Treatise on the Diseases of Children. By J. FORSYTH MEIGS, M.D., Fellow of the College of Physicians of Philadelphia, etc., etc., and WILLIAM PEPPER, M.D., Physician to the Philadelphia Hospital, Provost University of Pennsylvania. Seventh Edition, thoroughly Revised and Enlarged. A Royal Octavo Volume of over 1000 pages. Price, Cloth, $6.00; Leather, $7.00

"With the recent additions it may safely be pronounced one of the best and most comprehensive works on Diseases of Children."—*New York Medical Journal.*

"Must be regarded as the most complete work on Diseases of Children in our language."—*Edinburgh Medical Journal.*

"We have seldom met with a text-book so complete, so just and so readable as the one before us."—*American Journal of Obstetrics.*

MATHIAS, LEGISLATIVE MANUAL.

A Rule for Conducting Business in Meetings of Societies, Legislative Bodies, Town and Ward Meetings, etc. By BENJ. MATHIAS, A.M. Sixteenth Edition. 16mo. Price .50

MORTON, REFRACTION OF EYE.

The Refraction of the Eye. Its Diagnosis and the Correction of its Errors. With Chapter on Keratoscopy. By A. STANFORD MORTON, M.B., F.R.C.S. 12mo. Second Edition. Price $1.00

"The author has not only given very thorough rules for the objective and subjective examinations of the eye in the various conditions of refraction which present themselves, but has entered into an explanation of the phenomena observed, which is at once scientific and elementary."—*Edinburgh Medical Journal.*

MEDICAL TIMES AND GAZETTE (LONDON).

A Weekly Journal of 32 pages, containing Clinical Lectures, Hospital Reports, Leading Articles, News, Notes, Book Reviews, Correspondence, etc., etc. Subscription, per Annum, $5.00

OPHTHALMIC REVIEW.

A Monthly Record of Ophthalmic Science. Edited by KARL GROSSMANN, M.D., of Liverpool, PRIESTLEY SMITH, M.D., of Birmingham, and JOHN B. STORY, M.D., of Dublin. The only periodical representing the advancement of this science in England. *Now in its third year.* Subscription, per Annum, $3.00

OVERMAN, MINERALOGY.

Practical Mineralogy, Assaying, and Mining, with a Description of the Useful Minerals, etc. By FREDERICK OVERMAN, Mining Engineer. 11th Edition. 12mo. Cloth. Price $1.00

OGSTON, MEDICAL JURISPRUDENCE.

Lectures on Medical Jurisprudence. By Drs. FRANCIS and FRANCIS OGSTON, JR. With Copper-plate Illustrations. 8vo. Price $6.00

"We have a high appreciation of Dr. Ogston's lectures, and can cordially recommend the work as accomplishing all that the distinguished author promised for it."—*American Journal of Medical Science.*

OLDBERG, PRESCRIPTION BOOK. 300 New Prescriptions.

Three Hundred Prescriptions, Selected Chiefly from the Best Collections of Formulæ used in Hospital and Out-patient-practice, with a Dose Table, and a Complete Account of the Metric System. By OSCAR OLDBERG, PHAR. D., Late Medical Purveyor, United States Marine Hospital Service; Professor of Materia Medica, National College of Pharmacy, Washington, D. C.; Member of the American Pharmaceutical Association, and of the Sixth Decennial Committee of Revision and Publication of the Pharmacopœia of the United States. 12mo. Price, Paper Covers, .75; Cloth, $1.25

With over 350 Illustrations and Appropriate Letter-press Exp
erences.

Price, Two Volumes, Text and Plates, separate,
" One " " " bound toge
" " " " " " "

Adopted as a Text-book at Yale College. and used at other
America and England.

"Recognizing the fact that Physiology is emphatically an experimental science, it furnishes minute instructions for performing a great variety of experiments. A student could scarcely desire a better guide."
—*Boston Medical and Surgical Journal.*

"We confidently recommen
who are interested in the wid
siological research."—*New*
"This is a most superb boo
every physiological student
Medical Journal.

SANDERSON, PHYSIOLOGY. Second Edition.

A Syllabus of a Course of Lectures on Physiology. By J.
SON, M.D. For the Use of Students. Second Edition. 8vo.

SANDERSON, PRACTICAL EXERCISES IN P

8vo. Illustrated.

SANSOM, PHYSICAL DIAGNOSIS. Third Editi

The Physical Diagnosis of Diseases of the Heart. Includ
Sphygmograph and Cardiograph. By ARTHUR ERNEST SA
Edition. Revised and Enlarged. With Illustrations. 12mo.

BY SAME AUTHOR.

DISEASES OF THE HEART.

The Lettsomian Lectures on the Treatment of some of the
Disease of the Heart. Illustrated. 12mo.

BY SAME AUTHOR.

ON CHLOROFORM.

Chloroform. Its Action and Administration. 12mo.

SMITH. DYSMENORRHŒA.

Its Pathology and Treatment. By HEYWOOD SMITH, M.D
Hospital for Women and to the British Lying-in Hospital. 1

SMITH, RINGWORM.

SMITH, ON CHILDREN.

Clinical Studies of Diseases in Children. By EUSTACE SMITH, M.D. Second Revised Edition. Price $2.50

MEDICAL HERESIES, HISTORICALLY CONSIDERED.

A Series of Critical Essays on the Origin and Evolution of Sectarian Medicine, embracing a Special Sketch and Review of Homœopathy, Past and Present. By GONZALVO C. SMYTHE, A.M., M.D. Professor of the Principles and Practice of Medicine, College of Physicians and Surgeons, Indianapolis, Indiana. 12mo. Cloth. Price $1.25

"This book gives, in a small compass, an excellent history of medicine, from its earliest day to the present time."—*Buffalo Medical and Surgical Journal.*

"Cannot fail to be of interest, not only to the medical profession, but to the general reader."—*Baltimore Gazette.*

"The work is pleasantly written, in an easy, familiar style, and has cost the writer much literary research."—*New York Medical Journal.*

"Students and others interested in the subject of medicine will find a digest of the entire controversy (between the various schools of medicine) presented in this volume."—*Journal of Education.*

"Professor Smythe has succeeded in writing a brief, clear, and interesting sketch of the evolution of medical eccentricities, and of modern homœopathy, its facts and fallacies."—*Philadelphia Medical Times.*

SAVAGE, FEMALE PELVIC ORGANS. Author's Edition.

The Surgery, Surgical Pathology and Surgical Anatomy of the Female Pelvic Organs. In a Series of Colored Plates taken from Nature, with Commentaries, Notes and Cases. By HENRY SAVAGE, M.D., F.R.C.S. New Edition. Issued by arrangement with the Author, from the original Plates. Quarto. Price $12.00

SMITH. WASTING DISEASES OF CHILDREN.

The Wasting Diseases of Infants and Children. By EUSTACE SMITH, M.D., F.R.C.P., Physician to the East London Children's Hospital. Fourth London Edition, Enlarged. Octavo. Price $3.00

SCHULTZE, OBSTETRICAL PLATES.

Obstetrical Diagrams. Life Size. By Prof. B. S. SCHULTZE, M.D., of Berlin. Twenty in the Set. Colored.

Price, in Sheets, $15.00; Mounted on Rollers $25.00

SCANZONI, DISEASES OF WOMEN.

A Practical Treatise on the Diseases of the Sexual Organs of Women. By Dr. F. W. VON SCANZONI. Translated by A. K. GARDINER, M.D. 8vo. Price $5.00

SIEVEKING, LIFE ASSURANCE.

The Medical Adviser in Life Assurance. By E. H. SIEVEKING, M.D. 12mo. Second Edition, Revised. Price $2.00

SHEPPARD, ON MADNESS.

Madness, in its Medical, Social and Legal Aspects. A series of Lectures delivered at King's Medical College, London. By EDGAR SHEPPARD, M.D. 8vo. Price $2.25

STOCKEN, DENTAL MATERIA MEDICA. Third Edition.

The Elements of Dental Materia Medica and Therapeutics with Pharmacopœia. By JAMES STOCKEN, D.D.S. Third Edition. 12mo. $2.50

The first edition of this book was disposed of in a little less than four months. In making this revision the author has endeavored to make it still more useful by the addition of considerable new matter.

SUTTON, VOLUMETRIC ANALYSIS. Fourth Edition.

A Systematic Handbook of Volumetric Analysis, or the Quantitative Estimation of Chemical Substances by Measure, Applied to Liquids, Solids, and Gases. By FRANCIS SUTTON, F.C.S. Fourth Edition. Revised and Enlarged, with Illustrations. 8vo. Price $5.00

SEWELL, DENTAL ANATOMY AND SURGERY.

A Manual of Dental Anatomy and Surgery, Including the Extraction of the [illegible]. By H. E. SEWELL, D.D.S., M.D. With 77 Illustrations. 12mo. [illegible]

"A valuable book for the general Practitioner who [illegible] a work of a practical manual nature especially to [illegible] of the teeth."—*Medical Brief.*

"It will be found useful to the general [illegible] the management of many incidental affections with the teeth and mouth, which [illegible] handed over to the specialist."—*[illegible]*

STILLE, ON MENINGITIS.

Epidemic Meningitis, or Cerebro-spinal Meningitis. By ALFRED STILLE, Professor of Practice at the University of Pennsylvania. 8vo. [illegible]

"The name of the author is a sufficient guarantee that this monograph is elegant in style, exhaustive [illegible] and rich with practical suggestions."—*Philadelphia Medical and Surgical Reporter.*

STOKES, DISEASES OF THE HEART.

The Diseases of the Heart and Aorta. By WILLIAM STOKES, M.D. [illegible] 8vo. [illegible]

SWERINGEN, REFERENCE BOOK.

A Pharmaceutical Lexicon or Dictionary of Pharmaceutical Science. Containing explanations of the various subjects and terms of Pharmacy, with appropriate definitions from the Collateral Sciences. Formulæ for Officinal, Empirical and General Preparations, etc., etc. By HIRAM. V. SWERINGEN, M.D. 8vo.

Price, Cloth, $3.00; Leather

"[illegible] of a welcome, and sure of a ready recognition of its merits."—*London Pharmaceutical [illegible]*

"[illegible] will be found of great service to the pharmaceutical student, apprentice, druggist and physician [illegible] reference and as an aid to the study of scientific works."—*American Journal of Pharmacy.*

SOLLY. COLORADO SPRINGS FOR HEALTH.

Colorado Springs and Manitou as Health Resorts. By S. EDWIN SOLLY, M.D., Edin., including an article descriptive of the scenery and resources of the State. 12mo.

Paper covers, 25

TEMPERATURE CHARTS.

Charts for Recording Temperature, Respiration, Pulse, Day of Disease, [illegible] Symptoms, Name, etc.

Put up in pads, each 50

THOMPSON. MANUAL OF PHYSICS.

[illegible] Manual of Physics. By SILVANUS P. THOMPSON, B.A., [illegible] of Experimental Physics in University College, Bristol. Ex-

Pre[illegible]

THOMPSON LITHOTOMY AND LITHOTRITY.

[illegible] and Lithotrity; or, an Inquiry into the best Mode [illegible] from the Bladder. By SIR HENRY THOMPSON, F.R.C.S., [illegible] Clinical Surgery in University College. Third Edition. [illegible]

Price

[illegible] those in which Bigelow's operation is discussed, and the final [illegible] cases of male adults under the author's care. Such a [illegible]"—*Lancet.*

BY SAME AUTHOR.

[illegible] ORGANS. Seventh Edition.

[illegible] Organs. Clinical Lectures. Seventh London Edition. [illegible]

Price, Cloth, $1.25; Paper

[illegible] ESTATE.

[illegible] Their Pathology and Treatment. Fifth Edition. [illegible] Plates. Price, Cloth, $1.25; Paper, .75.

[illegible] DISEASES.

[illegible] of Calculous Disease, and the Use of Sounds [illegible] 12mo.

Price;

[illegible] labor of others, cautious in all his deductions, rejecting a [illegible] useful results, as clever with his pen as he is with the [illegible] deemed the master that he is."—*American Journal of M[illegible]*

THOMPSON, COUGHS AND COLDS.

The Causes, Nature, and Treatment of Coughs and Colds. By E. S. THOMPSON, M.D. 16mo. Price .60

THOROWGOOD, MATERIA MEDICA.

The Student's Guide to Materia Medica. By JOHN C. THOROWGOOD, M.D. Illustrated. 318 pages. 12mo. Price $2.00

BY SAME AUTHOR.

ON ASTHMA.

The Forms, Nature, and Treatment of Asthma. 2d Edition. Cloth, $1.75

TUSON, VETERINARY PHARMACOPŒIA.

A Pharmacopœia, Including the Outlines of Materia Medica and Therapeutics. For the Use of Students and Practitioners of Veterinary Medicine. By RICHARD V. TUSON, F.C.S. Third Edition. 12mo. Price $2.50

"Not only practitioners and students of veterinary medicine, but chemists and druggists will find that this book supplies a want in veterinary literature."—*Druggist and Chemist.*

THUDICHUM ON THE URINE. Second Edition.

The Pathology of the Urine and Complete Guide to Analysis. By JOHN L. W. THUDICHUM, M.D. Second Edition. Enlarged and Illustrated. 8vo. Price $5.00

"The treatise of Dr. Thudichum is well known as one of the medical classics of the language, and in completeness, thoroughness, and originality, the volume before us has few rivals in any branch of our science. For the specialist, for the physiological chemist, for the physiologist, the volume of Dr. Thudichum is a *sine qua non*, and to such the new edition must be a most welcome guest."—*Philadelphia Medical Times.*

TROUSSEAU, CLINICAL MEDICINE.

Lectures on Clinical Medicine, Delivered at the Hôtel Dieu, Paris, by A. TROUSSEAU, Professor of Clinical Medicine to the Faculty of Medicine, Paris, etc., etc. Translated from the Third Revised and Enlarged Edition by P. VICTOR BAZIRE, M.D., London and Paris; and JOHN ROSE CORMACK, M.D., Edinburgh, F.R.S., etc. With a full Index, Table of Contents, etc. 2 vols. 8vo. Sold by Subscription only.

Trousseau's Lectures have attained a reputation, both in England and in this country, far greater than any work of a similar character heretofore written. In order to bring the work within the reach of all the profession, the publishers now issue an American edition, containing all the lectures as contained in the five-volume Sydenham edition, at a much lower price. Below are a few only of the many favorable opinions expressed of the work:—

"A clever translation of Prof. Trousseau's admirable and exhaustive work; the best book of reference upon the Practice of Medicine."—*Indiana Medical Gazette.*

"We scarcely know of any book better fitted for presentation to a young man when entering upon the practical work of his life."—*London Medical Times and Gazette.*

TEST TYPES.

Selections from Snellen's Test Types mounted upon heavy card board; suitable for hanging in the office. Price 50 cents

TIDY, MODERN CHEMISTRY.

A Hand-Book of Modern Chemistry. Organic and Inorganic. By C. MEYMOTT TIDY, M.D. 8vo. Price $5.00

"We doubt if any other chemical work containing so large an amount of information could be procured."—*Dublin Medical Journal.*

MEMORANDA OF POISONS.

A Memoranda of Poisons and their Antidotes and Tests. from the Last London Edition. Revised and Enlarged.

This most complete Toxicological Manual should be within rea ...ons and pharmacists, and as an addition to every family libra ...means of saving life and allaying pain when the delay of sending would prove fatal.

SACTIONS OF THE AMERICAN SURGICAL ASSO-TION.

olume I. Illustrated. Edited by J. EWING MEARS, M.D., Recorder of the ociation. Royal 8vo. Cloth, $3.50

NSACTIONS OF THE COLLEGE OF PHYSICIANS.

The Transactions of the College of Physicians of Philadelphia. New Series. ls. I, II, III, IV, V. 8vo, Price, per volume, $2.50

Vol. VI. Containing Articles and Discourses by Drs. Atlee, Da Costa, Mills, V. Meigs, H. C. Wood, Cohen; Profs. Tyson, Gross, Bartholow, Allen, Leeds d others. Cloth, Gilt Top, $3.50

ON, BRIGHT'S DISEASE AND DIABETES.

A Treatise on Diabetes and Bright's Disease. With Especial Reference to athology and Therapeutics. By JAMES TYSON, M.D., Professor of Pathology nd Morbid Anatomy in the University of Pennsylvania. With Colored Plates and many Wood Engravings. 8vo. Price $3.50

his volume is the outcome of some fifteen years' l study and observation, and will be found to be y well prepared monograph. . . . . His direc- re clear and minute.—*Med. and Surg. Reporter.*

"The symptoms are clearly defined, and the treatment is exceedingly well described, so that every one reading the book must be profited."—*Cincinnati Lancet and Clinic.*

BY SAME AUTHOR.

UIDE TO THE EXAMINATION OF URINE.

A Practical Guide to the Examination of Urine. For the use of Physicians and Students. With Colored Plates and Numerous Illustrations Engraved on Wood. Fourth Edition. 12mo. Price $1.50

Advantage has been taken, in bringing out a new edition of this work, not only to orrect the previous one, but to make such additions of new Facts and Processes as ould add to its value without materially increasing its size.

"Dr. Tyson commences with a short account of the theory of renal secretion, the physical and chemical characrs of the urine, and the reagents and apparatus used in its analysis. Excellent rules are then given for detecting he presence of albumen, sugar, coloring-matters, bile, urea, uric acid, chlorides, phosphates and sulphates; and minute instructions for approximative and quantitative determination of most of those ingredients by volumetric analysis are supplied."—*Philadelphia Medical Times.*

"We have experienced both pleasure and profit from the perusal of this book. It is agreeably written, contains much practical information, and is, we believe, a reliable and satisfactory guide to the clinical examination of urine. We can recommend Dr. Tyson's book as one that amply supplies the clinical needs of the physician."—*Dublin Journal of Medical Science.*

THE CELL DOCTRINE. Second Edition.

The Cell Doctrine. Its History and Present State. With a Copious Bibliography of the subject. Illustrated by a Colored Plate and Wood Cuts. Second Edition. 8vo. Price $2.00

TURNBULL, ARTIFICIAL ANÆSTHESIA.

The Advantages and Accidents of Artificial Anæsthesia; Its Employment in the Treatment of Disease; Modes of Administration; Considering their Relative Risks; Tests of Purity; Treatment of Asphyxia; Spasms of the Glottis; Syncope, etc. By LAURENCE TURNBULL, M.D., PH.G., Aural Surgeon to Jefferson College Hospital, etc. Second Edition. Revised and Enlarged. With 27 Illustrations of Various Forms of Inhalers, etc. 12mo. Price $1.50

"Anæsthesia is a subject of great interest and importance to physicians and dentists, and everything that will aid them in better understanding the subject is sought with great avidity. This work we regard as the best aid in the study of the subject, and it presents the subject up to the present hour."—*Dental Register.*

TUKE. SLEEP-WALKING.

Sleep-Walking and Hypnotism. By D. HACK TUKE, M.D., LL.D., F.R.C.P., Co-Editor of the Journal of Mental Diseases. 8vo. Cloth, $1.75

VAN HARLINGEN, ON SKIN DISEASES.

A Practical Manual on Diseases of the Skin, with Diagnosis and Treatment. For Students and Practitioners. By ARTHUR VAN HARLINGEN, M.D., Vice-President of the American Dermatological Association. Including Formulæ. Illustrated by two Colored Plates. 12mo. Cloth. *In Press.*

VALENTIN. QUALITATIVE ANALYSIS.

A Course of Qualitative Chemical Analysis. By WM. G. VALENTIN, F.C.S. Sixth Edition, Enlarged. With over 260 Illustrations. Octavo. Cloth. *In Press.*

VACHER, CHEMISTRY.

A Primer of Chemistry, Including Analysis. By ARTHUR VACHER. 18mo. Price .50

VIRCHOW, POST-MORTEM EXAMINATIONS. Second Edition.

Post-mortem Examinations. A Description and Explanation of the Method of Performing them in the Dead House of the Berlin Charité Hospital, with especial reference to Medico-legal Practice. By Prof. VIRCHOW. Translated by Dr. T. P. SMITH. Second Edition. 12mo. With 4 Plates. Price $1.25

"A most useful manual from the pen of a master. . . . For thorough and systematic method in the performance of post-mortem examinations, there is no guide like it."—*Lancet.*

"Its low price and portability make it accessible and convenient to every surgical registrar and practitioner."—*British Medical Journal.*

WAGSTAFFE, HUMAN OSTEOLOGY.

The Student's Guide to Human Osteology. By WILLIAM WARWICK WAGSTAFFE, F.R.C.S. With 23 Lithographic Plates of the Bones, Showing Muscle Attachments, and 60 Wood Engravings. 12mo. Price $3.00

WICKES. SEPULTURE.

Sepulture: Its History, Methods and Sanitary Requisites. By STEPHEN WICKES, A.M., M.D., Author of a History of Medicine and Medical Men of New Jersey, etc. Octavo. Price $1.50

WEST. ON THE CHEST.

How To Examine the Chest. A Practical Guide for the use of Students. By SAMUEL WEST, M.D. Oxon., M.R.C.P., Physician to the City of London Hospital for Diseases of the Chest. Illustrated. 32mo. Cloth, $1.75

WOOD. BRAIN WORK.

Brain Work and Overwork. By Prof. H. C. WOOD, JR. 32mo. Price, Paper cover, .30; Cloth, .50

WATTS. CHEMISTRY.

A Manual of Chemistry, Physical and Inorganic. By HENRY WATTS, B.A., F.R.S., Editor of the Journal of the Chemical Society; Author of "A Dictionary of Chemistry," etc. With Colored Plate of Spectra and other Illustrations. 12mo. 595 pages. Price, Cloth, $2.25

This volume commences with a short sketch of the more important Elementary Bodies, the principal Laws of Chemical Combination, and the representation of the constitution and reactions of bodies by Symbolic Notation, followed by a section on CHEMICAL PHYSICS, including the determination of Densities, the mechanical properties of Gases, and the chief phenomena of Heat, Light, Electricity and Magnetism. The next section contains a description of the Non-metallic Elements, and the more important compounds which they form with one another; followed by a discussion of the general principles of Chemical Philosophy. In this part of the work the Laws of Chemical Combination and Decomposition and the principles of the Atomic Theory, briefly noticed in the Introduction, are more fully developed. The last section is devoted to the Chemistry of the Metals. A comparison of the Centigrade and Fahrenheit Scales of temperature is given at the end of the volume.

WEST, THE DISEASES OF WOMEN. Fourth Edition.

Lectures on the Diseases of Women. By CHARLES WEST, M.D. Fourth London Edition. Revised and in part re-written by the Author. With Numerous Additions by J. MATHEWS DUNCAN, M.D., Obstetric Physician to St. Bartholomew's Hospital 8vo. Price $5.00

Drs. West and Duncan are, perhaps, the most celebrated London physicians giving attention to the Diseases of Women, and together have made a most complete work, either for the physician or student.

WILKS, PATHOLOGICAL ANATOMY.

Lectures on Pathological Anatomy. By SAMUEL WILKES, F.R.S. Second Edition. Revised and Enlarged by WALTER MOXON, M.D., F.R.S., Physician to and Lecturer at Guy's Hospital, London. 8vo. Price $6.00

BY SAME AUTHOR.

DISEASES OF THE NERVOUS SYSTEM.

Lectures on Diseases of the Nervous System, Delivered at Guy's Hospital, London. New Edition, with Additions, Numerous Illustrative Cases, etc. 8vo. Cloth, $6.00

"A book of great value, embodying as it does the results of the experience and observation of one of the most accomplished of the London Hospital Physicians."—*American Journal of Medical Science.*

WRIGHT, ON HEADACHES. Ninth Thousand.

Headaches, their Causes, Nature and Treatment. By HENRY G. WRIGHT, M.D 12mo. Price .50

WILSON, ON DRAINAGE.

Drainage for Health; or, Easy Lessons in Sanitary Science, with Numerous Illustrations. By JOSEPH WILSON, M.D., Medical Director United States Navy. One Vol. Octavo. Price $1.00

"Dr. Wilson is favorably known as one of the leading American writers on hygiene and public health. The book deserves popularity."—*Medical and Surgical Reporter.*

"Well written and well illustrated. Attention to its teachings may save much disease and perhaps many lives."—*Cincinnati Gazette.*

"Interesting as well as useful."—*Philadelphia Ledger.*

"Easily understood, and briefly and concisely presented."—*Providence Journal.*

"Will be found of value."—*Boston Transcript.*

"Worthy of praise as a popular statement of the subject."—*Boston Journal of Chemistry.*

"Will be sure to be a harbinger of good in every family whose good fortune it may be to possess a copy."—*Builder and Wood Worker.*

BY SAME AUTHOR.

NAVAL HYGIENE.

Naval Hygiene, or, Human Health and Means for Preventing Disease. With Illustrative Incidents derived from Naval Experience. Illustrated. Second Edition. 8vo. Price $3.00

WILSON, HOW TO LIVE.

Health and Healthy Homes. A Guide to Personal and Domestic Hygiene. By GEORGE WILSON, M.D., Medical Officer of Health. Edited by JOS. G. RICHARDSON, M.D., Professor of Hygiene at the University of Pennsylvania. 314 pages. 12mo. Price $1.00

CHAPTER I.—Introductory, page 17. II. The Human Body, 33. III. Causes of Disease, 66. IV. Food and Diet, 119. V. Cleanliness and Clothing, 169. VI. Exercise, Recreation and Training, 187. VII. Home and Its Surroundings, Drainage, Warming, etc., 221. VIII. Infectious Diseases and their Prevention, 269.

"A most useful, and in every way, acceptable book."—*New York Herald.*

"Marked throughout by a sound, scientific spirit, and an absence of all hasty generalizations, sweeping assertions, and abuse of statistics in support of the writer's particular views. . . . We cannot speak too highly of a work which we have read with entire satisfaction."—*Medical Times and Gazette.*

BY SAME AUTHOR.

A HAND-BOOK OF HYGIENE

And Sanitary Science. With Illustrations. Fifth Edition. Revised and Enlarged. 8vo. Price $2.75

WILSON, HUMAN ANATOMY. Tenth Edition.

The Anatomist's Vade-Mecum. General and Special. By Prof. ERASMUS WILSON. Edited by GEORGE BUCHANAN, Professor of Clinical Surgery in the University of Glasgow; and HENRY E. CLARK, Lecturer on Anatomy at the Royal Infirmary School of Medicine, Glasgow. Tenth Edition. With 450 Engravings (including 26 Colored Plates). Crown 8vo. Price $6.00

Recommended as a Text-book at Rush Medical College, Chicago; Bellevue Hospital, New York; St. Louis Medical College; Yale and Dartmouth Schools, and many other Colleges.

BY SAME AUTHOR.

HEALTHY SKIN. Eighth Edition.

A Practical Treatise on the Skin and Hair; their Preservation and Management. Eighth Edition. 12mo. Paper. Price $1.00

WILSON, SEA VOYAGES FOR HEALTH.

The Ocean as a Health Resort. A Hand-book of Practical Information as to Sea Voyages, for the Use of Tourists and Invalids. By WM. S. WILSON, L.R.C.P. Lond., M.R.C.S.E. With a Chart showing the Ocean Routes, and Illustrating the Physical Geography of the Sea. Crown 8vo. Price $2.50

WELCH. ENTERIC FEVER.

Enteric Fever: Its Prevalence and Modifications; Ætiology, Pathology and Treatment, as illustrated by army data at home and abroad. By FRANCIS H. WELCH, F.R.C.S., Surgeon and Major A.M.D. Being the Alexander Prize Essay, Modified and Revised. 8vo. Price, Cloth, $2.00

WELLS, OVARIAN AND UTERINE TUMORS.

The Diagnosis and Surgical Treatment of Ovarian and Uterine Tumors. By T. SPENCER WELLS, M.D. Illustrated. 8vo. Price, Cloth, $7.00

So long a time having elapsed since Dr. Wells has collected the results of his large experience in book form, the present volume will be eagerly looked for by all interested in this very important subject.

WOLFE, ON DISEASES OF THE EYE.

A Practical Treatise on Diseases and Injuries of the Eye. Being a Course of Systematic and Clinical Lectures to Students and Medical Practitioners. By M. WOLFE, F.R.C.P.E., Senior Surgeon to the Glasgow Ophthalmic Institution, etc. With 10 Colored Plates, and numerous other Illustrations. Octavo. Price $7.00

WALKER, INTERMARRIAGE.

Intermarriage, or, The Mode in which, and the Causes why, Beauty, Health and Intellect result from certain Unions; and Deformity, Disease and Insanity from others. Illustrated. 12mo. Price $1.00

WARD'S COMPEND OF CHEMISTRY. Revised Edition.

A Compend of Chemistry for Chemical and Medical Students. By G. MASON WARD, M.D., Demonstrator of Chemistry in Jefferson Medical College, Philadelphia. Containing a Table of Elements and Tables for the Detection of Metals in Solutions of Mixed Substances, etc. 12mo. Cloth.

Interleaved for the addition of Notes, $1.25; plain, $1.00

WOODMAN and TIDY, MEDICAL JURISPRUDENCE.

Forensic Medicine and Toxicology. By W. BATHURST WOODMAN, M.D., Physician to the London Hospital, and CHARLES MEYMOTT TIDY, F.C.S., Professor of Chemistry and Medical Jurisprudence at the London Hospital. With Chromo-Lithographic Plates, representing the Appearance of the Stomach in Poisoning by Arsenic, Corrosive Sublimate, Nitric Acid, Oxalic Acid; the Spectra of Blood and the Microscopic Appearance of Human and other Hairs; and 116 other Illustrations. Large octavo. Sold only by Subscription.

Price, Cloth, $7.50; Medical Sheep, $8.50; Law Leather, $8.50

WOAKES, ON DEAFNESS AND GIDDINESS.

On Deafness, Giddiness and Noises in the Head; or, The Naso-Pharyngeal Aspect of Ear Disease. By EDWARD WOAKES, M.D., Senior Aural Surgeon to the Hospital for Diseases of the Throat and Chest. Third Edition. Revised and Enlarged, with Additional Illustrations. 12mo.

"No brief summary of his views could do full justice to the cogency and subtlety of his reasons. We prefer to commend the whole work to the thoughtful perusal of all intelligent medical practitioners who desire to rise above the level of mere routine empiricism."—*Lancet.*

BY THE SAME AUTHOR.

WOAKES, ON NASAL CATARRH.

Catarrh and Diseases of the Nose, Causing Deafness. 12mo. Illustrated.

Cloth, $1.50

WYTHE, ON THE MICROSCOPE.

The Microscopist. A Manual of Microscopy and Compendium of the Microscopic Sciences, Micro-Mineralogy, Micro-Chemistry, Biology, Histology, and Practical Medicine. By JOSEPH H. WYTHE, A.M., M.D. Fourth Edition. 252 Illustrations. 8vo. Price, Cloth, $3.00; Leather, $4.00

An Index and Glossary have been combined in this edition, so as to be a source of valuable information. Notices of recent additions to the microscope, together with the genera of microscopic plants, have been given in an Appendix.

"From what we knew of the author of this work, as a skilled practical Microscopist, a successful teacher of the science, and a practitioner of medicine and surgery of long and varied experience, we had a right to expect a good book from his hands. Our expectations are fully realized in the volume before us. The style is clear and distinct, and one reads the book with the utmost facility of comprehension. It is the more valuable to the physician and medical student on account of its closer application of the microscope to medical subjects than we find elsewhere. The numerous plates, many of which are beautifully colored, are not to be excelled. We feel proud of it as an American production."—*Pacific Medical and Surgical Journal.*

"This is one of the most valuable text-books on microscopy ever offered to students or practitioners of medicine. This edition has been greatly enhanced in value by the addition of chapters on the use of the microscope in pathology, diagnosis, and etiology, and numerous new illustrations, some of which are from Rindfleisch.

"The author very carefully brings out every necessary fact and principle relating to the use of the microscope, and now that this instrument has become an essential part of every practitioner's armamentarium, a practical guide and reference book is also a necessity, and we are fully warranted in reiterating the statement that this is one of the most valuable text-books ever offered to students and practitioners of medicine."—*The Cincinnati Lancet and Clinic.*

BY SAME AUTHOR.

DOSE AND SYMPTOM BOOK. Eleventh Edition.

The Physician's Pocket Dose and Symptom Book. Containing the Doses and Uses of all the Principal Articles of the Materia Medica, and Original Preparations. Eleventh Revised Edition.

Price, Cloth, $1.00; Leather, with Tucks and Pocket, $1.25

"The chapter on Dietetic Preparations will be found useful to all practicing physicians, most of whom have but little acquaintance with the mode of preparing the various articles of diet for the sick."—*Boston Medical and Surgical Journal.*

"Many a hard-worked practitioner will find it a useful little work to have on his study table."—*Canada Medical and Surgical Journal.*

YEO. A MANUAL OF PHYSIOLOGY.

A Manual of Physiology; being a Text-book for Students of Medicine. By GERALD F. YEO, M.D., F.R.C.S., Professor of Physiology in King's College, London. With over 300 carefully printed Illustrations. A Glossary and Complete Index. Crown Octavo. Price, Cloth, $4.00; Leather, $5.00

"This work * * * is the legitimate successor of the similar treatise by Dr. Carpenter, which, excellent as that was at the time it was written, has, we suppose, been found so defective as to require more time and trouble to renovate than would be demanded to write a new work. We think the publishers have done well in selecting Dr. Gerald Yeo for the author of their new manual. This gentleman occupies the physiological chair at King's College, has a good reputation as a teacher, and is known as having ably assisted Dr. Ferrier in the interesting researches made by that physician in the localization of function in the brain.

"The plan followed in the work is, to give, in the first place, a general view of the animal structures, then to describe the chemical basis of the body and the vital characteristics of animal organisms, and next to consider, in succession, and in the following order, the functions of digestion, absorption, circulation, respiration, nutrition, animal heat, muscle, voice and speech, the nervous system, special senses, reproduction and development. In dealing with his subject, Dr. Yeo has introduced in many places a brief account of the histological features of the parts of which he is about to show the uses. * * * The mode in which Dr. Yeo deals with disputed points has, in every instance we have met, been remarkably sensible and straightforward. * * * Upon the whole, although we have taken exception to one or two points, we consider Dr. Yeo's book to be a very good one, it is most worthy, and clearly and intelligently written. We cannot conclude our notice of this work without referring to the excellence of the wood-cuts. These are not only very numerous and generally well executed, but to a large extent novel."—*London Lancet, May 24th, 1884.*

ILLUSTRATED BOOKS.

MEDICINAL PLANTS.

Being Descriptions, with original Figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By ROBERT BENTLEY, F.L.S., Professor of Botany in the King's College, and to the Pharmaceutical Society, and HENRY TRIMENS, M.B., F.L.S., late Lecturer on Botany at St. Mary's Hospital Medical School. In 42 Parts, each, $2.00, or in 4 vols., large 8vo, with 306 Colored Plates, bound in half morocco, gilt edged. $90.00

AN ATLAS OF TOPOGRAPHICAL ANATOMY.

After Plane Sections of Frozen Bodies. By WILLIAM BRAUNE, Professor of Anatomy in the University of Leipzig. Translated by EDWARD BELLAMY, F.R.C.S., Surgeon to and Lecturer on Anatomy at Charing Cross Hospital. With 34 Photo-lithographic Plates and 46 Wood cuts. Large imp. 8vo. $8.00

ATLAS OF SKIN DISEASES.

Consisting of a Series of Illustrations, with Descriptive Text and Notes upon Treatment. By TILBURY FOX, M.D., F.R.C.P., late Physician to the Department for Skin Diseases in University College Hospital. With 72 Colored Plates. In 18 Parts, each, $1.00 or, 1 Vol., Royal 4to, Cloth. $20.00

AN ATLAS OF HUMAN ANATOMY.

Illustrating most of the ordinary Dissections, and many not usually practiced by the Student. By RICKMAN J. GODLEE, M.S., F.R.C.S., Assistant Surgeon to University College Hospital, and Senior Demonstrator of Anatomy in University College. With 48 imp. 4to Colored Plates (112 Figures), and a volume of Explanatory Text. $20.00

A COURSE OF OPERATIVE SURGERY.

By CHRISTOPHER HEATH, F.R.C.S., Home Professor of Clinical Surgery in University College, and Surgeon to the Hospital. With 20 Plates drawn from Nature by M. LÉVEILLÉ, and colored by hand under his direction. Second Edition, Enlarged. 4to. *Sold only by Subscription.* $14.00

ILLUSTRATIONS OF CLINICAL SURGERY.

Consisting of Plates, Photographs, Wood cuts, Diagrams, etc., etc., illustrating Surgical Diseases, Symptoms, and Accidents; also Operative and other Methods of Treatment, with Descriptive Letterpress. By JONATHAN HUTCHINSON, F.R.C.S., Senior Surgeon to the London Hospital. Vol. I, containing fasciculi I to X, bound, with Appendix and Index. $25.00

Fasciculi XI to XVI. Ready. Each, $2.50

NOW READY.

Diseases of the Liver.

BY GEORGE HARLEY, M.D., F.R.S., Etc.,

Author of "The Urine and Its Derangements," and "Diabetes, Its Various Forms and Treatment."

On Fine Paper, from Good Type, with Colored Plates and Thirty-six Wood-cuts. Bound in Heavy Cloth, Beveled Edges, $5.00; Leather, with Raised Bands, $6.00.

THE Publishers call special attention to this work, the only thorough book now before the profession. The reputation of its distinguished author is a guarantee of its merits.

THE AUTHOR, IN HIS PREFACE, SAYS:

THIS NEW TREATISE, which I have thought fit to entitle Diseases of the Liver, with and without Jaundice, with special application to Diagnosis and Treatment, embodies within it the whole substance of my original monograph on Jaundice and Diseases of the Liver; though greater than it, both as regards its scope and materials, and the large amount of clinical and scientific data that has never before been collected together into one volume; while in a great many instances it gives a new rendering to old clinical facts, by presenting them to the reader in the light of modern pathological science.

As I think time is quite of as much value to the professional as it is to the mercantile man, I have endeavored to condense my materials to the utmost, without running the risk of endangering their perspicuity. Added to which, as this treatise has not been penned either for the use of the tyro or the dilettante in medicine, but for that of my qualified brethren, I shall neither waste time by entering into detailed accounts of the literature, nor give tedious, and probably at the same time profitless, discussions of the theories of the mechanism of jaundice in hepatic derangements. Taking care, however, in order that it may carry more weight with it in the eyes of the reader, to illustrate it freely with cases reported by independent observers, both at home and abroad. While, in order, again, that the reader may be able to see for himself, at a glance, how many of the old-fashioned theories of the pathology of jaundice have been abandoned, as well as how many new ones have been espoused, I have put my views, in accordance with the facts and arguments expressed throughout the body of the volume, into a concise and diagrammatic tabular form.

I WOULD DIRECT the special attention of my readers to the chapter devoted to treatment, as well as that at the end of the book, entitled Hints on Diagnosis.

SYNOPSIS OF THE CONTENTS.

Introduction, giving a general view of the scope of the volume, and the application of Physiological Chemistry to the diagnosis and treatment of Hepatic affections.

Chemistry, Physics and Physiology of the Liver and its secretions.

Etiology of Jaundice—different kinds—causes producing them—treatment.

Signs and Symptoms of Liver Diseases.

General remarks on all kinds of Hepatic Remedies.

Special Hepatic Medicines; their modes of action and uses.

Mineral Waters, Wines and Foods; treatment of Pyrexia, Cerebral complications, etc.

Congenital and Hereditary Liver Diseases, Biliousness; Its Varieties and Treatment.

Jaundice from Enervation, all its forms explained and their different Treatments

Different forms of Inflammation of the Liver and their Treatments.

Jaundice caused by Disease Germs, Yellow Fevers, Contagious and Epidemic Jaundice, different kinds and their Treatments.

Jaundice of Pregnancy.

Different forms of Hepatic Atrophy and Ascites.

Biliary Concretions, Inspissated Bile, Gall-Stones of every kind and form, direct and indirect effects of, their Symptoms and Treatment, very fully gone into.

Different kinds of Colics, etc.

Catarrhal Jaundice.

Jaundice from Poisons.

Different kinds of Jaundice from Permanent Obstructions.

Physiological Chemistry of the Excretions, Urine and Stools, as a Guide to Diagnosis and Treatment.

All kinds of Abscess, Tropical, Pyæmic, Metastatic, etc.

Different kinds of Cancers of the Liver and its Appendages.

Hydatid and Cystic Diseases of the Liver; Syphilitic and Fibroid Diseases of the Liver.

Embolisms, Fatty, Amyloid and other Degenerations of the Liver.

Traumatic Diseases of the Liver.

Diseases of the Gall Bladder.

A concluding chapter, entitled Hints on Differential Diagnosis.

Index.

This work is now ready, and will be sent by mail, postpaid, upon receipt of price. (Cloth, $5.00. Leather, $6.00.

P. BLAKISTON, SON & CO., 1012 Walnut St., Philadelphia.

☞ *This has been entirely rewritten and remodeled, and brought up to the present day.*

BIDDLE'S
Materia Medica.

NINTH REVISED EDITION.

(Contains all the Changes in the New Pharmacopœia.)

Recommended as a Text-book at Yale College, University of Michigan, College of Physicians and Surgeons, Baltimore, Baltimore Medical College, Louisville Medical College, and a number of other Colleges throughout the United States.

BIDDLE'S MATERIA MEDICA. For the Use of Students and Physicians. By the late PROF. JOHN B. BIDDLE, M.D., Professor of Materia Medica in Jefferson Medical College, Philadelphia. The Ninth Edition, thoroughly revised, and in many parts rewritten, by his son, CLEMENT BIDDLE, M.D., Assistant Surgeon, U. S. Navy, assisted by HENRY MORRIS, M.D. Containing all the additions and changes made in the last revision of the United States Pharmacopœia. Octavo.

Ready.

Bound in Cloth. Price $4.00; Leather, $4.75.

RECOMMENDATIONS.

"It will be found a useful handbook by students, especially, who may be under the instruction of its able and accomplished author."—*American Medical Journal.*

"In short, it is just the work for a student, embracing as it does what will be discussed in a course of lectures on materia medica."—*Cincinnati Medical News.*

"In truth, the work is well adapted to the wants of students."—*The Clinic.*

"Nothing has escaped the writer's scan. All the new remedies against disease are duly and judiciously noted. Students will certainly appreciate its shapely form, grace of manner, and general *multum in parvo* style."—*American Practitioner.*

"Biddle's 'Materia Medica' is well known to the profession, being a standard text book in several leading colleges."—*New York Medical Journal.*

"It contains, in a condensed form, all that is valuable in materia medica, and furnishes the medical student with a complete manual on this subject."—*Canada Lancet.*

"The necessity for a new edition of this work in so short a time is the best proof of the value in which it is held by the profession."—*Medical and Surgical Reporter.*

"The standard 'Materia Medica' with a large number of medical students is Biddle's."—*Buffalo Medical and Surgical Journal.*

"The larger works usually recommended as text-books in our medical schools are too voluminous for convenient use. This work will be found to contain in a condensed form all that is most valuable, and will supply students with a reliable guide."—*Chicago Medical Journal.*

*** This Ninth Edition contains all the additions and changes in the U. S. Pharmacopœia, Sixth Revision.

P. BLAKISTON, SON & CO., Publishers and Booksellers.
1012 WALNUT STREET, PHILADELPHIA.

BYFORD'S
Diseases of Women.

THIRD ENLARGED EDITION.

Recommended as a Text-book at Rush Medical College, Chicago; Woman's Medical College, Chicago; National Medical College, Washington, D. C., and other Medical Schools.

THE PRACTICE OF MEDICINE AND SURGERY, APPLIED TO THE DISEASES AND ACCIDENTS INCIDENT TO WOMEN. By WM. H. BYFORD, A.M., M.D., author of "A Treatise on the Chronic Inflammation and Displacement of the Unimpregnated Uterus," and Professor of Obstetrics and Diseases of Women and Children in the Chicago Medical College, and in the Woman's Medical College, Chicago. Third Edition, Revised. In One Volume. Octavo. 164 Illustrations.

Price, Cloth, $5.00; Leather, $6.00.

RECOMMENDATIONS.

"I have read with pleasure your book on Diseases of Women; it gratifies me to indorse its teachings."—*Prof. H. P. C. Wilson, Jr., Baltimore.*

"I am well acquainted with the author, and regard him an excellent practitioner and teacher."—*Dr. J. A. Ireland, Professor of Obstetrics and Gynæcology, Louisville Medical College.*

"It is much improved and is still more what it always was, a valuable practical assistant to the practitioner."—*Prof. Henry G. Landis, Starling Medical College, Columbus.*

"I find, from its being so well brought up to the present state of the science it treats of, that it is particularly adapted to the requirements of the medical student."—*J. Fred Prioleau, M. D., Professor of Gynæcology, Medical College of South Carolina, Charleston.*

"The author has presented to the profession a concise and comprehensive treatise on the subject of diseases of females. We congratulate Dr. Byford upon his success, and feel satisfied that the effort will be duly appreciated by those whom the work is intended to instruct. It is one of the best treatises on the subject ever submitted to the American profession, and will early be acknowledged as such. It is well written, and where necessary, is appropriately illustrated."—*Medical and Surgical Reporter, Philadelphia.*

"The present work being particularly of use where questions of etiology and general treatment are concerned. * * * This volume, hence, will commend itself both to those who possess former editions and to those who do not. It cannot fail to be useful to many, and will certainly prove interesting to all."—*The American Journal of Obstetrics.*

"These important topics are all handled vigorously and practically. The work cannot fail to be in great demand, and to exert a material influence on clinical practice."—*New York Medical Journal.*

P. BLAKISTON, SON & CO., Publishers and Booksellers.
1012 WALNUT STREET, PHILADELPHIA.

MEDICAL JOURNALS

Published by P. BLAKISTON, SON & CO.

THE LONDON MEDICAL TIMES
AND GAZETTE.

832 Pages Weekly for $5.00 per Annum. Post-F

Commencing with the number for January 5th, 1884, the London "Medical Times and Gazette" underwent an alteration in the size of page that makes it much more convenient for handling and binding. A clearer type, better arrangement of the matter employed, and an increase in the number of pages made, thus increasing the amount of matter. The contents will be printed on the first page of reading matter, enabling them to be bound up with the volume, and a reduction in the price of subscription makes it one of the cheapest and best Weekly Medical Papers now published.

CONTENTS.—The contents of each number consist of several original Lectures or Communications from the most London or British Government Hospitals, Editorial Notes on current topics, Reviews and Notices of Treatises just Issued, New Discoveries, Remedies, &c., leading articles, Reviews and Extracts from Current Literature, Reports of the London and Foreign Medical Societies, Medical News and Notes.

As a circulating medium the London Journal the "Medical Times" is among the first in the world, compared with other foreign journals of the same size, stands it within the reach of every physician who wishes to keep acquainted with the progress of Medical Science at the least cost to a home.

THE POLYCLINIC.

A monthly Journal of Medicine and Surgery, conducted by the Faculty of the Philadelphia Polyclinic and College for Graduates in Medicine. HENRY LEFFMANN, Editor in Chief. Published on the 15th day of each month. Now in its second volume.

SUBSCRIPTION, $2.00 PER ANNUM.

CONTENTS OF DECEMBER NUMBER.—I. On Some Skin Diseases, by Dr. [illegible], of the London Hospital. II. Ophthalmology, Prof. Roberts Bartholow. Nervous Prostration, Dr. Mitchell and [illegible]. Nervous Diseases, Dr. [illegible]. Diseases of the Skin, Dr. [illegible], of the American Dermatological Society, in New York. [illegible], Dr. [illegible]. Diseases of the Skin, Dr. [illegible]. Prof. Pancoast, Jefferson Hospital, Philadelphia. Dr. [illegible]. Prof. Roberts and Dr. [illegible], Surgery and Hospital Reports, Dr. [illegible], Syphilis, Cancer, Pediatrics, Dr. [illegible], Obstetrics, Dr. [illegible], Pathology, Dr. Formad, Dr. James T. Ferris, Dr. [illegible], Dr. [illegible].

[illegible]

LANE MEDICAL LIBRARY

To avoid fine, this book should be returned on or before the date last stamped below.

APR


E34
H726
1885
Holden, L. Holden's anatomy.
6199
NAME
DATE DUE
Univ. Oregon
APR 23 '53
MAY 13 1931